<http://www.e-journals.net>

ISSN: 0973-4945; CODEN ECJHAO

E-Journal of Chemistry

Vol. 5, No.2, pp.347-354, April 2008

Syntheses of 1, 2, 4 Triazole Derivatives and their Biological Activity

FREDDY H. HAVALDAR* and ABHAY R. PATIL

Nadkarny-Sacasa Research Laboratory,
Department of Chemistry, St. Xavier's College, Mumbai - 400 001
drfreddy_11@yahoo.co.in

Received 14 October 2007; Accepted 10 December 2007

Abstract: Cyclisation of [4-(4-oxo-2-phenyl-4*H*-quinazolin-3-yl)-phenoxy]-acetic acid-*N'*-(substituted phenyl)-thiosemicarbazides with sodium metal in 99.0 % ethanol gave 3-[4-(4-substituted phenyl-5-thioxo-4,5-dihydro-1*H*-1,2,4 triazol-3-yl-methoxy)-phenyl]-2-phenyl-3*H*-quinazolin-4-one. The structures of the newly synthesized compounds have confirmed by IR, ¹H NMR and Mass spectra. The compounds have also been screened for their biological activity.

Keywords: 2-Phenyl-4-(3*H*)-quinazolin-4-one, 1,2,4- Triazole derivatives, IR, Spectral studies, Biological activity.

Introduction

The chemistry of heterocycles is an interesting branch in organic chemistry. Quinazolinone derivatives derived from 2-phenyl-4*H*-3,1-benzoxazin-4-one are important compounds in chemistry and pharmacology. They have drawn much attention due to their broad range of pharmacological properties¹, which include anticancer², anti-inflammatory³, anticonvulsant⁴, and antidiuretic⁵ activities. Also quinazolinone such as tryptanthrin have been found to have remarkable antimalarial activity⁶.

1,2,4-Triazoles are associated with diverse pharmacological activities such as analgesic, antiasthmatic, diuretic, antihypersensitive, anticholinergic, antibacterial, antifungal and anti-inflammatory activity⁷⁻¹⁰. These biological data prompted us to synthesize some new quinazolin-4-one derivatives containing 1,2,4- triazole ring.

Experimental

General procedures

Melting points were taken in open capillaries and are uncorrected. IR spectra (KBr in cm⁻¹) were recorded on Jasco 410 plus FTIR spectrophotometer. ¹H NMR spectra were recorded on a JEOL 300 MHz FT-NMR spectrophotometer using DMSO-d₆ as solvent and TMS as

internal standard (chemical shifts in δ ppm). The mass spectra of compounds were determined with Shimadzu model No.QP 2010. The elemental analysis was carried out on a Perkin Elmer C,H,N analyzer and sulphur analysis was obtained by oxygen-flask method. The purity of the compounds was monitored by thin layer chromatography. TLC was carried out on precoated 0.2 mm silica gel $60F^{254}$ plates.

(4-Acetyl-amino-phenoxy)-acetic acid ethyl ester (2)

A solution of 4-(hydroxy phenyl)-acetamide **1** (15.1 g, 0.1 mole) in 500 mL *N,N*-dimethyl formamide was heated at 80-85 °C with ethyl chloroacetate (13.47 mL, 0.11 mole) in an oil bath in presence of anhydrous potassium carbonate (13.8 g, 0.11 mole). The reaction mixture was cooled and 1500 mL of cold water was added. A white-coloured crystalline solid separated was filtered, washed thoroughly first with cold water and then with cold ethanol, and finally recrystallised from 99 % ethanol, yield 70 %, m.p. 104 °C; IR (KBr) 3347 (NH), 1750 (C=O ester), 1589, 1508, 1477 (C=C, aromatic).

(4-Amino-phenoxy)-acetic acid (3)

Compound **2** (23.7 g, 0.1 mole) was heated with 35 mL concentrated hydrochloric acid and 100 mL water for about 2 hours. A clear solution obtained was filtered and neutralized with 40% of sodium acetate solution. The white crystalline solid precipitated out was filtered, washed thoroughly with water and recrystallised from DMF: ethanol (1:1), yield 72 %, m.p. 248 °C ; IR (KBr) 3464 (NH₂), 1650 (C=O), 1558, 1508, 1445 (C=C, aromatic).

(4-Amino-phenoxy)-acetic acid ethyl ester (4)

Compound **3** (11.85 g, 0.05 mole) was suspended in 250 mL ethanol and cooled to 10 °C. To this solution, sulphuric acid (3.0 mL, 0.055 mole) was added dropwise with stirring, maintaining the temperature at 10 -15 °C. The reaction mixture was then refluxed for about 2 hours. The excess ethanol was distilled under reduced pressure and then to one-third of the total volume of the solution, 100 mL water was added followed by 20 % of potassium carbonate solution for neutralization. The product was extracted with 3 x 25 mL of dichloromethane. The dichloromethane layer was washed with 3 x 20 mL of water. The compound **4** obtained on distillation of dichloromethane at atmospheric pressure was dried in an oven at 40 °C and recrystallised from benzene, yield 80 %, m.p. 58 °C; IR (KBr) 3382 (NH₂), 1740 (C=O ester), 1558, 1509, 1476 (C=C, aromatic).

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid ethyl ester (5)

A mixture of 2-phenyl-4H-3,1-benzoxazin-4-one (4.46 g, 0.02 mole) and (4-amino-phenoxy) acetic acid ethyl ester **4** (4.29 g, 0.022 mole) in 100 mL pyridine was refluxed for about 12 hours. The reaction mixture was then allowed to cool to room temperature and then added slowly 100 mL of methanol. The product **5** separated was filtered, washed with ethanol and crystallized from ethanol, yield 82 %, m.p. 180 °C; IR (KBr) 1735 (C=O ester), 1650 (C=O), 1606 (C=N), 1577, 1544, 1508, 1448 (C=C, aromatic); ¹H NMR (DMSO-*d*₆) δ 1.24 (t, 3H, CH₃), 4.1 (q, 2H, CH₂), 4.7 (s, 2H, OCH₂C), 6.9-8.6 (m, 13H, ArH); Anal.Calc'd. for C₂₄H₂₀N₂O₄ (400): C, 72.00; H, 5.00; N, 7.00. Found: C, 71.96; H, 5.02; N, 6.94.

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid hydrazide (6)

Compound **5** (4.0 g, 0.01 mole) and 2 mL of 99 % hydrazine hydrate in ethanol was refluxed for about 8 hours. The reaction mixture was then allowed to cool to room temperature. The separated white coloured solid was filtered, washed with ethanol and crystallized from

ethanol, yield 80 %, m.p. 220 °C; IR (KBr) 3315, 3272 (NH·NH₂), 1674 (C=O amide), 1652 (CO·N), 1603 (C=N), 1585, 1508, 1450 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 4.3 (s, 2H, NH₂), 4.4 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 13H, ArH), 9.3 (s, 1H, NH); Anal.Calc'd. for C₂₂H₁₈N₄O₃ (386): C, 68.39; H, 4.66; N, 14.50. Found: C, 68.41; H, 4.65; N, 14.52.

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid-N'-(substituted phenyl)-thiosemicarbazides (7a-d)

Compound 6 (0.386g. 0.001 mole) was refluxed with various aromatic phenyl isothiocyanates (0.0011 mole) in isopropyl alcohol (40 mL) with continuous stirring over a period of 16 hours. The crystalline solid obtained on cooling the reaction mixture was filtered, washed with isopropyl alcohol and finally crystallized from ethanol to afford compounds (7a-d).

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)- phenoxy]-acetic acid- N'-(phenyl thiosemi-carbazide (7a)

Colorless solid, yield 84 %, m.p. 195°C; IR (KBr) 3230 (NH), 1678 (CO·N), 1662 (CH₂·CO), 1600 (C=N), 1558, 1508, 1496, 1446 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 4.6 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 18H, ArH), 9.6 (s, 1H, NH), 9.7 (s, 1H, NH), 10.3 (s, 1H, NH); Anal.Calc'd. for C₂₉H₂₃N₅O₃S (521): C, 66.79; H, 4.41; N, 13.43; S, 6.14. Found: C, 66.74; H, 4.38; N, 13.52; S, 6.10.

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid-N'-(4-fluoro-phenyl)-thiosemicarbazide (7b)

Colorless solid, yield 86%, m.p. 200°C; IR (KBr) 3213 (NH), 1697 (CH₂·CO), 1683 (CO·N), 1602 (C=N), 1585, 1508, 1489 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 4.6 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 17H, ArH), 9.6 (s, 1H, NH), 9.7 (s, 1H, NH), 10.3 (s, 1H, NH); Anal.Calc'd. for C₂₉H₂₂FN₅O₃S (539): C, 64.56; H, 4.08; N, 12.77; S, 5.93. Found: C, 64.59; H, 4.11; N, 12.82; S, 6.00.

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid-N'-(4-nitro-phenyl)-thiosemicarbazide (7c)

Pale yellow coloured solid, yield 80%, m.p. 180°C; IR (KBr) 3286 (NH), 1701(CH₂·CO), 1647 (CO·N), 1597 (C=N), 1587, 1556, 1508, 1496 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 4.6 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 17H, ArH), 10.0 (s, 1H, NH), 10.1 (s, 1H, NH), 10.4 (s, 1H, NH); Anal.Calc'd. for C₂₉H₂₂N₆O₅S (566): C, 61.48; H, 3.88; N, 14.84; S, 5.65. Found: C, 61.45; H, 4.39; N, 14.82; S, 5.70.

[4-(4-Oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid-N'-(2,4-difluoro-phenyl)-thiosemicarbazide (7d)

Colorless solid, yield 78 %, m.p. 202°C; IR (KBr) 3250 (NH), 1662 (CH₂·CO), 1635(CO·N), 1600 (C=N), 1583, 1558, 1508, 1485 (C=C, aromatic); ¹H NMR (DMSO-d₆)δ 4.6 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 16H, ArH), 9.7 (s, 1H, NH), 9.8 (s, 1H, NH), 10.3 (s, 1H, NH); Anal.Calc'd. for C₂₉H₂₁F₂N₅O₃S (557): C, 62.47; H, 3.77; N, 12.56; S, 5.74. Found : C, 62.51; H, 3.76; N, 12.62; S, 5.78.

3-[4-(4- Substituted phenyl-5-thioxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl-methoxy)-phenyl]- 2-phenyl- 3H-quinazolin-4-ones (8a-d)

To the hot reaction mixture of compounds (7a-d, 0.0005 mole) in 50 mL of ethanol was added sodium metal (0.001 mole) pieces and the reaction mixture was refluxed with continuous stirring over a period of 4 hours. The excess ethanol was distilled under

atmospheric pressure till 3-4 mL was left behind. The viscous liquid obtained on cooling was poured into 20 mL of cold water, followed by neutralized with 1:1 hydrochloric acid under cold condition. The product obtained was filtered, washed thoroughly with water and crystallised from ethanol to give compounds (**8a-d**).

3-[4-(4-Phenyl-5-thioxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl-methoxy)-phenyl]-2-phenyl-3H-quinazolin-4-one (8a)

Colorless solid, yield 56 %, m.p. 265 °C; IR (KBr) 3429 (NH), 1667 (CO·N), 1602 (C=N), 1585, 1508, 1450 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 5.0 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 18H, ArH), 14.0 (s, 1H, NH); MS (m/z): 503 [M⁺], 371, 327, 313, 223, 205, 179, 166, 146, 118, 109, 105, 90, 77; Anal.Calcd. for C₂₉H₂₁N₅O₂S (503): C, 69.18; H, 4.17; N, 13.91; S, 6.36. Found: C, 69.12; H, 4.19; N, 13.98; S, 6.40.

3-[4-[4-(4-Fluoro-phenyl)-5-thioxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl-methoxy]-phenyl]-2-phenyl-3H-quinazolin-4-one (8b)

Colorless solid, yield 65 %, m.p. 298 °C; IR (KBr) 3326 (NH), 1637 (CO·N), 1603 (C=N), 1587, 1558, 1508, 1450 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 5.0 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 17H, ArH), 14.0 (s, 1H, NH); MS (m/z): 521 [M⁺], 371, 327, 313, 223, 205, 179, 166, 146, 118, 109, 105, 90, 77; Anal.Calcd. for C₂₉H₂₀FN₅O₂S (521): C, 66.79; H, 3.83; N, 13.43; S, 6.14. Found: C, 66.75; H, 3.80; N, 13.49; S, 6.21.

3-[4-[4-(4-Nitro-phenyl)-5-thioxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl-methoxy]-phenyl]-2-phenyl-3H-quinazolin-4-one (8c)

Pale yellow coloured solid, yield 59 %, m.p. 185 °C; IR (KBr) 3314 (NH), 1648 (CO·N), 1600 (C=N), 1587, 1508, 1449 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 4.7 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 17H, ArH), 14.0 (s, 1H, NH); MS (m/z): 548 [M⁺], 371, 327, 313, 223, 205, 179, 166, 146, 118, 109, 105, 90, 77; Anal.Calcd. for C₂₉H₂₀N₆O₄S (548): C, 63.50; H, 3.64; N, 15.32; S, 5.83. Found: C, 63.52; H, 3.66; N, 15.39; S, 5.80.

3-[4-[4-(2,4-Difluoro-phenyl)-5-thioxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl-ethoxy]-phenyl]-2-phenyl-3H-quinazolin-4-one (8d)

Colorless solid, yield 61 %, m.p. 240 °C; IR (KBr) 3332 (NH), 1647(CO·N), 1602 (C=N), 1587, 1509, 1449 (C=C, aromatic); ¹H NMR (DMSO-d₆) δ 5.0 (s, 2H, O·CH₂·C), 6.9-8.6 (m, 16H, ArH), 14.0 (s, 1H, NH); MS (m/z): 539 [M⁺], 371, 327, 313, 223, 205, 179, 166, 146, 118, 109, 105, 90, 77; Anal.Calcd. for C₂₉H₁₉F₂N₅O₂S (539): C, 64.56; H, 3.52; N, 12.98; S, 5.93. Found: C, 64.51; H, 3.49; N, 13.05; S, 5.90.

Results and Discussion

4-Hydroxy-phenyl acetamide **1** on electrophilic substitution with ethyl chloroacetate gave (4-acetylamino-phenoxy)-acetic acid ethyl ester **2** which on hydrolysis with aqueous hydrochloric acid gave 4-(amino-phenoxy)-acetic acid **3**. This on further esterification with sulphuric acid gave 4-(amino-phenoxy)-acetic acid ethyl ester¹¹ **4**.

The 2-phenyl-4H-3,1-benzoxazin-4-one^{12,13} which was prepared from 2-amino anthranilic acid and benzoyl chloride at 20 -25 °C in pyridine was allowed to react with 4-(amino-phenoxy)-acetic acid ethyl ester **4** to give [4-(4-oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid ethyl ester **5**. The structure of compound **5** was established by analytical and spectral data. The compound **5** was then condensed with hydrazine hydrate in ethanol to afford [4-(4-oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid hydrazide **6**.

in good yields. The compound **6** on reaction with various phenyl isothiocyanates in isopropyl alcohol gave [4-(4-oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetic acid-N'-(substituted phenyl)-thiosemicarbazides (**7a-d**). The presence of NH group is confirmed by IR spectra which showed characteristic band at 3300 cm^{-1} and ^1H NMR signal at δ 9.6, 9.7 and 10.3 corresponding to $\text{O}=\text{C}\text{NHNH}\text{C}=\text{S}\text{NH}$ protons. The compounds (**7a-d**) on Cyclisation with sodium metal in ethanol gave 3-[4-(4-substituted phenyl-5-thioxo-4,5-dihydro-1H-[1,2,4]-triazol-3-yl)-methoxy)-phenyl]-2-phenyl-3H-quinazolin-4-ones (**8a-d**) [Scheme 1]. IR spectra of compounds (**8a-d**) showed the presence of NH group between at 3300 cm^{-1} and 3100 cm^{-1} and also the ^1H NMR signal at δ 14.0 indicates the presence of NH group for structures of compounds (**8a-d**).

Scheme 1

Biological Activity

Antibacterial activity

All the newly synthesized compounds (**8a-d**) were screened *in vitro* for their antibacterial activity against *Staphylococcus aureus*, *Escherichia coli* and *Bacillus subtilis* by the ditch-plate technique²⁰ using concentrations of 50 $\mu\text{g/mL}$. Nutrient agar was employed as culture media and DMF was used as solvent control for antibacterial activity.

Antifungal activity

The compounds (**8a-d**) synthesized were screened for their antifungal activity against *Aspergillus niger*, *Candida albicans* and *Cryptococcus neoformans* by paper-disc diffusion method²¹ at concentrations of 50 µg/mL. Nutrient agar was employed as culture media and DMF was used as solvent control for antifungal activity.

The known compounds, such as ampicillin, amoxicillin, norfloxacin, penicillin and Griseofulvin were used for comparison purpose. The diameter of zone of inhibition was measured in mm. The antibacterial and antifungal screening data are recorded in Table 1.

Table 1. Biological activity data

Compounds	Zone of inhibition inmm					
	Antibacterial activity			Antifungal activity		
	<i>S. aureus</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>A. niger</i>	<i>C. albicans</i>	<i>C. neoformans</i>
8a	11	10	11	12	14	16
8b	13	14	16	14	11	15
8c	17	12	18	16	14	18
8d	16	15	13	11	15	12

From the Table 1, it can be seen that the compounds **8c** and **8d** showed remarkable activity against *Staphylococcus aureus*, compounds **8b** and **8c** showed remarkable activity against *Bacillus subtilis*. The compounds **8c** showed good activity against *A. niger* and the compounds **8a** and **8c** showed maximum activity against *Cryptococcus neoformans*.

Antimalarial activity

The effort to find new antimalarial activity is still a high priority given to the increasing malarial emergency due to chloroquine resistant *Plasmodium falciparum* strains. The chloroquine-resistant *Plasmodium falciparum* malarial parasite was cultured *in vitro* and the sensitivity of parasite to the newly synthesized compounds was evaluated using the tritiated Hypoxanthine incorporation assay¹⁶. The compounds (**8a-d**) were tested for antimalarial activity and the only compound 3-[4-[4-(4-fluoro-phenyl)-4H-[1,2,4]triazol-3-yl-methoxy]-phenyl]-2-phenyl-3H-quinazolin-4-one **8b** was found to be most active against *Plasmodium falciparum* strains and its 50 % inhibitor concentration (IC₅₀) values were 1.2µM.

Acknowledgement

The authors are thankful to TIFR and Institute of Science, Mumbai for ¹H NMR and mass spectra and Dr. (Mrs.) Vivien Amonkar, Head, Department of Microbiology, St. Xavier's College, Mumbai for providing biological activity.

References

- (a) Wolfe J F, Rathman T L, Sleevi MC, Campbell J A and Greenwood T D, *J. Med. Chem.* 1990, **33**, 161. (b) Padia J K, Field M, Hilton J, Meecham K, Pablo J, Pinnock R, Roth B D, Singh L, Suman-Chauhan N, Trivedi B K and Webdale L, *J. Med. Chem.* 1998, **41**, 1042.
- Xia Y, Yang Z Y, Hour M J, Kuo S C, Xia P, Bastow K F, Nakanishi Y, Nampoothiri P, Hackl T, Hamel E and Lee K H, *Bioorg. Med. Chem. Lett.* 2001, **11**, 1193.
- Kenichi O, Yoshihisa Y, Toyonari O, Toru I and Yoshio I, *J. Med. Chem.* 1985, **28**, 568.
- Buchanan J G, Sable H Z and Thygarajan B S Ed.; Wiley-Interscience: New York, 1972, **2**, pp.1-96.

5. Lyle F R, US Pat. 5,973,257. *Chem. Abstr.* 1985, **65**, 2.
6. Ridley R G, *Sciences* 1999, **285**, 1502.
7. Birendra N G, Jiban C S K and Jogendra N B, *J. Heterocycl. Chem.* 1984, **21**, 1225.
8. (a) Kothari P J, Mehlhoff M A, Singh S P, Parmar S S and Stenberg V I *J. Heterocycl. Chem.* 1980, **17**, 1369. (b) Kothari P J, Singh S P, Parmar S S and Stenberg V I, *J. Heterocycl. Chem.* 1980, **17**, 1393.
9. Sengupta A K and Misra H K, *J. Indian Chem. Soc.* 1981, **8**, 508.
10. Sarmah S C and Bahel S C, *J. Indian Chem. Soc.* 1982, **59**, 877.
11. Walter A, Heidelberger, Jacobs and Michael, *J. American Chem. Soc.* 1917, **39**, 2196.
12. Tiwari S S and Satsangi R K, *J. Indian Chem. Soc.* 1975, **55**, 736.
13. Zentmyer D T and Wagner E C, *J. Org. Chem.* 1949, **14**, 967.
14. Collins C H and Lyne P M, *Microbiological Methods*; 3rd Ed; Butterworths, London, 1970, p. 424.
15. Seeley H W and Van Denmark P J, *Microbes in Action*, W.H. Freeman and Co. 1972, USA.
16. Desjardins R E, Canfield C J, Hayens J D and Chulay J D, *Antimicrob Agents Chemother* 1979, **16**, 710.