Synthesis and Antimicrobial Studies of a New Series of *Bis*-Heterocycles

B. JAYASHANKARA and K.M. LOKANATHA RAI*

Department of Studies in Chemistry, University of Mysore,
Manasagangothri, Mysore-570006, India.

jai_shan321@yahoo.co.in

Received 7 September 2007; Revised 28 October 2007; Accepted 14 November 2007

Abstract: Series of novel *bis*-heterocycles bearing isoxazoline and imidazole moieties have been synthesized *via* 1,3-dipolarcycloaddition reactions. ¹HNMR, ¹³CNMR, IR and elemental analyses characterized the newly synthesized compounds. All the synthesized compounds were evaluated for their antimicrobial activity and were compared with the standard drugs. The compounds demonstrated potent to weak antimicrobial activity.

Keywords: *Bis*-Heterocycles; Antimicrobial activity; Chloramine-T.

Introduction

Among five membered heterocycles, isoxazoline and imidazole are represents a class of compounds of great importance in biological chemistry. For instance, isoxazoline posses biological activities like^{1,2} (insecticidal, antibacterial, antibiotic, antitumour, antifungal). Isoxazoline also serves as anti-influenza virus activity³, inhibition of human leukocyte elastase and cathepsin G⁴. In fact, Valdecocib an isoxazoline derivative is now widely used in the market as anti-inflammatory drug⁵. Imidazole derivatives are gaining synthetic interest in recent years due to their broad spectrum of biological activities like anti-inflammatory⁶, analgesic⁷, antibacterial⁸ and antifungal⁹. 2-*n*-Butyl-4-chloro-5-farmyl-imidazole is a key intermediate for the synthesis of Losartan a nonpeptide angiotensin antagonist, which is an orally active antihypertensive drug¹⁰. Literature survey reveals that *bis*-heterocycles bearing isoxazoline^{11,12} was synthesized *via* 1,3-dipolar cycloaddition of aldoxime to divinyl ketone / sulfone using chloramine-T as dehydrating agent.

1,3-Dipolar cycloaddition reactions are useful tools for constructing biologically potent five membered heterocycles² and nitrile oxides serves as excellent 1,3-dipoles. Cycloaddition of nitrile oxide to olefinic compounds are of synthetic interest, since the product isoxazoline obtained are the versatile intermediate for the synthesis of bifunctional

compounds.¹³ Nitrile oxides can be generated by dehydrogenation of aryl aldoximes with mercuric acetate¹⁴, manganese dioxide¹⁵, *tert*-butyl hypo chlorite¹⁶, chloramine-T *etc.* In our laboratory Rai *et.al*¹⁷ extensively used chloramine-T for the generation of nitrile oxide and nitrile imine from aldoxime and aldehyde hydrazone respectively. With this background, it is considered worthwhile to prepare *bis*-heterocycles starting from 2-n-butyl-4-chloro-(*N*-substituted)-imidazole-5-carbaldehyde and screen them for antimicrobial activity.

Scheme 1.

Experimental

¹H NMR spectra were recorded on a Bruker AM 300 MHz spectrometer using CDCl₃ as solvent and tetramethylsilane as internal standard. ¹³C NMR spectra were measured on Jeol 400 (100MHz) instrument. The chemical shifts are expressed in δ and following abbreviations were used: s = singlet, d = doublet, t = triplet and m = multiplet. Infrared (IR) spectra were measured on Shimadzu 8300 spectrometer. Elemental analyses were obtained on a Vaio-EL instrument. Thinlayer chromatography (TLC) was done with pre-coated silica gel G plates using chloroform-acetone as eluent.

Antimicrobial activity

All the synthesized compounds were evaluated for antimicrobial activity by the disc diffusion method¹⁸ and microdilution method.¹⁹ Five bacteria and five fungal species were used as the antimicrobial test strains namely: *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas fluorescens*, *Xanthomonas campestris pvs*, *Xanthomonas oryzae*, *Aspergillus niger*, *Aspergillus flavus*, *Fusarium oxysporium*, *Trichoderma species*, *Fusarium monaliforme*. Streptomycin and tetracycline were used as standard drugs against bacteria and nystatin was used against fungi. In all the determinations tests were performed in triplicate and the results were taken as a mean of at least three determinations.

Preparation of 4'-(2-Butyl-4-chloro-5-formyl-imidazol-1-yl)methyl-4,5-dimethoxy-benzene (2)

A mixture of 2-Butyl-5-chloro-3H-imidazole-4-carbaldehyde **1** (1.0 g, 5.37 mmol) and anhy. K_2CO_3 (0.90 g, 6.52 mmol) in dimethylformamide (10 mL) was stirred for 15 min at rt. Bromomethyl-4,5-dimethoxy-benzene (1.66 g, 5.35 mmol) was then added and the mixture was stirred at rt for 6 hr. After completion of the reaction (TLC toluene-ethylacetate; 7:3), the reaction mass was diluted with 25 mL water and the product was extracted with dichloromethane (25 mL). The extract was washed with water (10 mL) and then dried (Na_2SO_4) the solvent was evaporated and the remaining pale yellow oil was crystallized from ethanol to gave **2** as a white crystalline solid to yield 1.83 g (82%), m.p 86-87 °C.

Synthesis of 4,5-dihydro-3-(substituted-imidazole)-5-phenylisoxazole (5a)

A mixture of substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4a** (0.265 g, 2.55 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) in ethanol (20 mL) was warmed on a water bath for 2-3 hr. the progress of the reaction was monitored by TLC. After completion of the reaction the solvent was evaporated in vacuum. The residual mass was extracted into ether (25 mL), washed successively with water (2 x 20 mL), 5% NaOH (1x10 mL), brine solution (2 x 15 mL) and dried over anhydrous sodium sulphate. Evaporation of the solvent afforded crude oily substance, which was purified by column chromatography using chloroform-acetone (9:1) as eluent to gave the product as yellow solid (0.94 g, 69% yield), m.p.124-126 °C.

4,5-Dihydro-3-(substituted-imidazole)-5-methyl-5-phenylisoxazole (5b)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4b** (0.30 g, 2.55 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow solid (0.99 g, 71% yield), and m.p.130-132 °C.

4,5-Dihydro-3-(substituted-imidazole)isoxazole-5-carbonitrile (5c)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4c** (0.135 g, 2.55 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow oil (0.78 g, 62% yield).

5-(Chloromethyl)-4,5-dihydro-3-(substituted-imidazole)isoxazole (5d)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4d** (0.194 g, 2.57 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow solid (0.84 g, 65% yield), m.p.138-140 °C.

5-(Bromomethyl)-4,5-dihydro-3-(substituted-imidazole)isoxazole (5e)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4e** (0.31 g, 2.56 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow solid (1.02 g, 73% yield), and m.p.144-146 °C.

(4,5-Dihydro-3-(substituted-imidazole)isoxazol-5-yl)methanol (5f)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4f** (0.148 g, 2.55 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow oil (0.78 g, 63% yield).

4,5-Dihydro-3-(substituted-imidazole)isoxazole-5-yl acetate (5g)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4g** (0.22 g, 2.55 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow solid (1.0 g, 76% yield), m.p.148-150 °C.

4,5-Dihydro-3-(substituted-imidazole)isoxazole-5-yl propionate (5h)

Obtained from substituted-imidazole aldoxime **3** (1.0 g, 2.55 mmol), **4h** (0.255 g, 2.55 mmol) and chloramine-T trihydrate (0.72 g, 2.56 mmol) as yellow solid (1.05 g, 78% yield), m.p.158-160 °C.

Results and Discussion

The general synthetic pathway discussed hereafter is depicted in **Scheme**. Compound **1** was alkylated using bromomethyl-4,5-dimethoxy benzene and potassium carbonate in DMF. Then formyl function of **2** was converted into the aldoxime **3**. When oxidative dehydrogenation of **3** by chloramine-T (CAT) afforded nitrile oxide, which was *in situ* trapped by the different olefins **4 (a-h)** under refluxing condition in ethanol. Thus produced compound was identified by NMR spectroscopy as 4,5-dihydro-3-(substituted-imidazole)-5-substituted-isoxazoline **5 (a-h)** in good quality and yield. The starting substrate 2-*n*-butyl-4-chloro-(*N*-substituted)-imidazole-5-carbaldehyde **1** was prepared according to literature procedure.²⁰ Compound **3** was prepared by standard procedure²¹.

Antimicrobial activity

Antimicrobial activity of all the compounds was shown in **Table 1** and **2**. Among the series of synthesized compounds, **5d** and **5e** shown better inhibition. Remaining compounds shown moderate inhibition. The better inhibition shown by **5d** and **5e** may be due to the presence of chloro and bromo group in the compound.

Table 1. Minimal inhibitory concentration in $\mu\text{g mL}^{-1}$ [X] and Inhibitory zone in (diameter) mm [Y] of the synthesized compounds against tested bacterial strains by micro dilution method and disc diffusion method respectively

Compound	<i>Bacillus subtilis</i>		<i>Escherichia coli</i>		<i>Pseudomona fluorescens</i>		<i>Xanthomonas campestris pvs</i>		<i>Xanthomonas oryzae</i>	
	X	Y mm	X	Y mm	X	Y mm	X	Y mm	X	Y mm
5a	21	07	26	13	23	16	25	12	23	12
5b	22	11	26	08	23	10	25	07	21	10
5c	19	10	17	11	20	13	19	10	23	10
5d	17	14	14	12	13	15	11	09	15	10
5e	18	10	15	12	15	13	12	10	15	09
5f	22	11	26	08	23	10	25	07	21	10
5g	21	07	26	13	23	16	25	12	23	12
5h	22	10	21	11	25	14	22	11	24	11
Streptomycin	19	13	13	14	12	17	-	-	-	-
Tetracycline	-	-	-	-	-	-	09	12	13	12

Streptomycin sulfate (25 μg per disc); Tetracycline (25 μg per disc) were used as positive reference standard antibacterial discs, Synthesized compounds (25 μg per disc).

Table 2. Minimal inhibitory concentration in $\mu\text{g mL}^{-1}[\text{X}]$ and Inhibitory zone in (diameter) mm [Y] of the synthesized compounds against tested fungal strains by micro dilution method and disk diffusion method respectively

Compound	<i>Aspergillus niger</i>		<i>Aspergillus flavus</i>		<i>Fusarium oxysporium</i>		<i>Trichoderma species</i>		<i>Fusarium monalifome</i>	
	X	Y mm	X	Y mm	X	Y mm	X	Y mm	X	Y mm
5a	18	08	18	09	14	12	15	14	14	11
5b	19	07	18	07	18	11	17	12	15	09
5c	16	08	16	10	13	13	14	15	11	11
5d	14	09	13	12	10	16	11	17	10	14
5e	15	09	15	11	11	14	13	16	10	12
5f	18	07	18	07	18	11	17	12	15	09
5g	17	07	16	08	15	11	16	13	13	10
5h	20	06	17	07	16	10	18	11	14	08
Nystatin	15	08	15	10	11	14	12	16	09	12

Nystatin (25 μg per disc) was used as positive reference standard antifungal discs, Synthesized compounds (25 μg per disc).

Spectral data of the compounds

Compound 2: ^1H NMR CDCl_3 : δ 0.94 (t, 3H, CH_3), 1.32 (m, 2H, CH_2), 1.64 (m, 2H, CH_2), 2.61 (t, 2H, CH_2), 3.64 (s, 6H, OCH_3) 4.86 (s, 2H, CH_2), 6.45 (s, 1H, ArH), 6.82 (s, 1H, ArH), 9.50 (s, 1H, CH), ^{13}C NMR CDCl_3 : δ 13.9 (C), 23.2 (C), 24.5 (C), 33.1 (C), 33.9 (C), 57.5 (2C), 116.1 (C), 117.6 (C), 118.2 (C), 134.2 (C), 136.6 (C), 140.9 (C), 147.2 (C), 148.4 (C), 156.2 (C), 189.2 (C). IR (KBr pellets cm^{-1}) ν 3069, 2961, 1764, 1667, 1485. Anal.Calcld. For $\text{C}_{17}\text{H}_{20}\text{BrClN}_2\text{O}_3$: C, 49.12, H, 4.85, N, 6.74%. Found: C, 49.19, H, 4.87, N 6.70 %.

Compound 5a: ^1H NMR CDCl_3 : δ 0.97 (t, 3H, CH_3), 1.35 (m, 2H, CH_2), 1.64 (m, 2H, CH_2), 2.57 (t, 2H, CH_2), 3.21 (dd, 1H, $J=8.2$, 4-H), 3.28 (dd, 2H, $J=8.2$, 4-H), 3.73 (s, 6H, OCH_3), 5.02 (s, 2H, CH_2), 5.20 (dd, 1H, $J=4.0$, 5-H), 6.42 (s, 1H, ArH), 6.78 (s, 1H, ArH), 7.21 (m, 5H, ArH), ^{13}C NMR CDCl_3 : δ 14.2 (C), 23.0 (C), 25.7 (C), 33.4 (C), 34.8 (C), 41.6 (C), 56.3 (2C), 81.1 (C), 116.3 (C), 117.7 (C), 118.5 (C), 122.2 (C), 126.2 (C), 127.2 (2C), 127.8 (C), 128.9 (2C), 134.5 (C), 140.7 (C), 148.0 (C), 148.7 (C), 149.0 (C), 164.7 (C). Anal.Calcld. For $\text{C}_{25}\text{H}_{27}\text{BrClN}_3\text{O}_3$: C, 56.35; H, 5.11; N, 7.89; Found: C, 56.37; H, 5.13; N, 7.87 %.

Compound (5b): ^1H NMR CDCl_3 : δ 0.95 (t, 3H, CH_3), 1.33 (m, 2H, CH_2), 1.59 (s, 3H, CH_3), 1.62 (m, 2H, CH_2), 2.55 (t, 2H, CH_2), 3.18 (s, 2H, 4- CH_2), 3.74 (s, 6H, OCH_3), 5.02 (s, 2H, CH_2), 6.40 (s, 1H, ArH), 6.76 (s, 1H, ArH), 7.19 (m, 5H, ArH), ^{13}C NMR CDCl_3 : δ 14.3 (C), 23.1 (C), 25.6 (C), 29.7 (C), 33.5 (C), 37.7 (C), 56.3 (2C), 87.1 (C), 116.3 (C), 117.5 (C), 118.6 (C), 122.2 (C), 126.1 (2C), 126.2 (2C), 126.4 (C), 128.5 (2C), 134.3 (C), 148.2 (C), 148.7 (C), 149.0 (C), 150.1 (C), 164.6 (C). Anal. Calcld. For $\text{C}_{26}\text{H}_{29}\text{BrClN}_3\text{O}_3$ C, 57.10; H, 5.34; N, 7.68; Found: C, 57.12, H, 5.32, N, 7.63 %.

Compound (5c): ^1H NMR CDCl_3 : δ 0.94 (t, 3H, CH_3), 1.32 (m, 2H, CH_2), 1.60 (m, 2H, CH_2), 2.53 (t, 2H, CH_2), 3.42 (dd, 1H, $J=8.0$, 4-H), 3.49 (dd, 2H, $J=8.0$, 4-H), 3.79 (s, 6H, OCH_3), 4.96 (s, 2H, CH_2), 5.24 (dd, 1H, $J=4.0$, 5-H), 6.34 (s, 1H, ArH), 6.70 (s, 1H, ArH). ^{13}C NMR CDCl_3 : δ 14.2 (C), 23.0 (C), 25.7 (C), 33.4 (C), 33.7 (C), 40.8 (C), 56.3 (2C), 68.8

(C), 116.4 (C), 117.4 (C), 118.2 (C), 118.6 (C), 122.1 (C), 126.7 (C), 134.5 (C), 148.2 (C), 148.6 (C), 149.1 (C), 164.7 (C). Anal.Calcld. For $C_{20}H_{22}BrClN_4O_3$; C, 49.86; H, 4.60; N, 11.63; Found: C, 49.89; H, 4.67; N, 11.53 %.

Compound (5d): 1H NMR $CDCl_3$: δ 0.97 (t, 3H, CH_3), 1.34 (m, 2H, CH_2), 1.64 (m, 2H, CH_2), 2.56 (t, 2H, CH_2), 3.38 (dd, 1H, $J=8.4$, 4-H), 3.42 (dd, 2H, $J=8.4$, 4-H), 3.46 (dd, 1H, $J=7.6$, CH_2Cl), 3.69 (dd, 1H, $J=7.6$, CH_2Cl), 3.75 (s, 6H, OCH_3), 5.02 (s, 2H, CH_2), 5.12 (m, 1H, 5-H), 6.40 (s, 1H, ArH), 6.78 (s, 1H, ArH). ^{13}C NMR $CDCl_3$: δ 14.1 (C), 22.8 (C), 25.5 (C), 33.4 (C), 33.7 (C), 37.8 (C), 51.8 (C), 56.2 (2C), 69.8 (C), 116.4 (C), 117.5 (C), 118.6 (C), 121.9 (C), 126.2 (C), 134.5 (C), 148.1 (C), 148.6 (C), 149.1 (C), 164.7 (C). Anal.Calcld. For $C_{20}H_{24}BrCl_2N_3O_3$; C, 47.55; H, 4.79; N, 8.32; Found: C, 47.53; H, 4.78; N, 8.35 %.

Compound (5e): 1H NMR $CDCl_3$: δ 0.95 (t, 3H, CH_3), 1.33 (m, 2H, CH_2), 1.62 (m, 2H, CH_2), 2.55 (t, 2H, CH_2), 3.35 (dd, 1H, $J=8.4$, 4-H), 3.40 (dd, 2H, $J=8.4$, 4-H), 3.44 (dd, 1H, $J=7.2$, CH_2Br), 3.64 (dd, 1H, $J=7.2$, CH_2Br), 3.73 (s, 6H, OCH_3), 5.00 (s, 2H, CH_2), 5.04 (m, 1H, 5-H), 6.38 (s, 1H, ArH), 6.75 (s, 1H, ArH). ^{13}C NMR $CDCl_3$: δ 14.3 (C), 22.8 (C), 25.6 (C), 33.6 (C), 33.8 (C), 38.8 (C), 41.0 (C), 56.4 (2C), 70.9 (C), 116.4 (C), 117.4 (C), 118.7 (C), 122.2 (C), 126.3 (C), 134.4 (C), 148.2 (C), 148.8 (C), 149.1 (C), 164.7 (C). Anal.Calcld. For $C_{20}H_{24}Br_2ClN_3O_3$; C, 43.70; H, 4.40; N, 7.64; Found: C, 43.71; H, 4.42; N, 7.62 %.

Compound (5f): 1H NMR $CDCl_3$: δ 0.97 (t, 3H, CH_3), 1.34 (m, 2H, CH_2), 1.63 (m, 2H, CH_2), 2.32 (dd, 1H, OH), 2.57 (t, 2H, CH_2), 3.28 (dd, 1H, $J=8.0$, 4-H), 3.34 (dd, 1H, $J=8.0$, 4-H), 3.52-3.79 (m, 2H, CH_2), 3.74 (s, 6H, OCH_3), 4.98 (s, 2H, CH_2), 5.04 (m, 1H, 5-H), 6.41(s, 1H, ArH), 6.77 (s, 1H, ArH). ^{13}C NMR $CDCl_3$: δ 14.2 (C), 23.0 (C), 25.7 (C), 33.5 (C), 33.8 (C), 36.8 (C), 56.3 (2C), 70.6 (C), 77.5 (C), 116.3 (C), 117.4 (C), 118.4 (C), 121.9 (C), 126.2 (C), 134.5 (C), 148.1 (C), 148.7 (C), 149.0 (C), 164.7 (C). Anal.Calcld. For $C_{20}H_{25}BrClN_3O_4$; C, 49.35; H, 5.18; N, 8.63; Found: C, 49.33; H, 5.19; N, 8.64 %.

Compound (5g): 1H NMR $CDCl_3$: δ 0.94 (t, 3H, CH_3), 1.35 (m, 2H, CH_2), 1.64 (m, 2H, CH_2), 2.04 (s, 3H, CH_3), 2.57 (t, 2H, CH_2), 3.30 (dd, 1H, $J=8.2$, 4-H), 3.37 (dd, 1H, $J=8.2$, 4-H), 3.78 (s, 6H, OCH_3), 5.02 (s, 2H, CH_2), 5.68 (dd, 1H, $J=3.8$, 5-H), 6.39 (s, 1H, ArH), 6.77 (s, 1H, ArH). ^{13}C NMR $CDCl_3$: δ 14.3 (C), 21.1 (C), 23.0 (C), 25.6 (C), 33.4 (C), 33.8 (C), 56.3 (2C), 69.5 (C), 96.3 (C), 116.3 (C), 117.5 (C), 118.6 (C), 121.9 (C), 126.2 (C), 134.4 (C), 148.0 (C), 148.6 (C), 149.1 (C), 164.6 (C), 170.4 (C). Anal.Calcld. For $C_{21}H_{25}BrClN_3O_5$; C, 48.99; H, 4.89; N, 8.16; Found: C, 48.99; H, 4.88; N, 8.17 %.

Compound (5h): 1H NMR $CDCl_3$: δ 0.95 (t, 3H, CH_3), 1.13 (t, 3H, CH_3), 1.33 (m, 2H, CH_2), 1.64 (m, 2H, CH_2), 2.28 (q, 2H, CH_2), 2.54 (t, 2H, CH_2), 3.31 (dd, 1H, $J=8.0$, 4-H), 3.38 (dd, 1H, $J=8.0$, 4-H), 3.75 (s, 6H, OCH_3), 4.99 (s, 2H, CH_2), 5.66 (dd, 1H, $J=4.0$, 5-H), 6.35 (s, 1H, ArH), 6.74 (s, 1H, ArH). ^{13}C NMR $CDCl_3$: δ 9.5 (C), 14.1 (C), 22.8 (C), 25.6 (C), 30.1 (C), 33.4 (C), 33.7 (C), 56.3 (2C), 69.4 (C), 96.5 (C), 116.4 (C), 117.4 (C), 118.4 (C), 122.0 (C), 126.3 (C), 134.3 (C), 148.0 (C), 148.7 (C), 149.1 (C), 164.5 (C), 173.4 (C). Anal.Calcld. For $C_{22}H_{27}BrClN_3O_5$; C, 49.97; H, 5.15; N, 7.95; Found: C, 49.99; H, 5.13; N, 7.93 %.

References

1. Huisgen R, *Angew.Chem. Int.Ed.Engl.* 1963, **2**, 565.
2. Caramella P and Gruinanger P in 1,3-Dipolar Cycloaddition Chemistry, vol **1**, Edited by Padwa A (wiley interscience, New york) 1984, 337.
3. Hiroyuki Kai, Hiroshi Matsumoto, Naohiko Hattori, Akira Takase, Tamio Fujiwara and Hirohiko Sugimoto. *Bioorg. Med. Chem.Lett.* 2001, **11**, 1997.

4. William C. Groutas, Lee S.Chang, Jeffery B. Epp, Radhika Venkataraman, Michael J.Brubaker, Michael A. Stanga, Eun-Hong Kim and Charles E. Keller. *Bioorg. Med. Chem. Lett.* 1992, **2**, 1571.
5. Dannahardt G, Kiefer W, Kramer G, Maehrlein S, Nowe U and Fiebich B. *Eur. J. Med. Chem.*, 2000, **35**, 499.
6. Slee D H, Romano S J, Yu J, Nguyen T N, John J K, Raheja N K. Axe F U, Jones T K and Ripka W C, *J. Med. Chem.*, 2001, **44**, 2094.
7. Umit U, Nalan G, Karaburun and Ihan L. *II Farmaco*, 2001, **56**, 285.
8. Khabnadideh S, Rezaei Z, Khalafi-Nezhad A, Bahrinajafi R, Mahamadi R and Farrokhriz A A. *Bioorg. Med. Chem. Lett.* 2003, **13**, 2863.
9. Gunnay N S, Ulusoy G N, Ergenc N, Otuk G and Kaya D, *II Farmaco*, 1999, **54**, 826.
10. Carini D J, Duncia J V, Aldrich P E, Chiu A T, Johnson A L, Pierce M E, Price W A and Santella III J B, Wells G J, Wexler P C, Yoo S W, Timmermans P B M W M. *J. Med. Chem.*, 1991, **34**, 2525.
11. Padmavathi V, Jagan Mohan Reddy B, Chandra Obula Reddy B and Padmaja A. *Tetrahedron*, 2005, **61**, 2407.
12. Padmavathi V, Venugopal Reddy K, Padmaja A and Bhaskar Reddy D. *Phosphorus, Sulfur and Silicon*, 2003, **178**, 171.
13. (a) Lokanath Rai K M and Hassner A. *Indian J. Chem.* 1997, **36B**, 242. (b) Grundman C and Grunanger P, In *The Nitrile Oxide*. Springer Verlag; New York, 1971.
14. Lokanatha Rai K M, Linganna N, Hassner A, Murthy C A. *Org. Prep. Proc. Int.* 1992, **24**, 91.
15. Keigiel J, Paplawaska M, Jozwik J, Kosior M and Jurezak J. *Tetrahedron*, 1999, **40**: 5605.
16. Osamu Moriya, Hideo Takenaka, Masaichi Iyoda, Yoshikiyo Urata and Takeshi Endo. *J. Chem. Soc., Perkin Trans.I* 1994, 413.
17. (a) Lokanath Rai K M and Hassner A, *Indian J. Chem.* 1997, **36B**, 242. (b) Lokanath Rai K M, Hassner A, *Synth. Commun.* 1997, **27**, 467. (c) Hassner A and Lokanath Rai K M, *Synthesis* 1989, 57. (d) Lokanath Rai K M and Hassner A, *SynthCommun.* 1989, **19**, 2799-2807.
18. Lemriss S, Marquet B, Ginestet H, Lefeuvre L, Fassouane A and Boiron P, *J. Mycol. Med.* 2003, **13**, 189.
19. Zgoda J R and Porter J R, *Pharmaceutical Biology*. 2001, **39**, 221
20. Griffiths G J, Hauck M B, Imwinkelried R, Kohr J, Roten C A and Stucky G C, *J. Org. Chem.*, 1999, **64**, 8084.
21. Vogel. A. I. *A Textbook of practical Organic chemistry*, 5th Ed. Longman's Green and Co. Ltd: London 1989; 1258.