



Simultaneous Estimation of Gemcitabine Hydrochloride and Capecitabine Hydrochloride in Combined Tablet Dosage Form by RP-HPLC Method

V.RAJESH, B.ANUPAMA^{§*}, V.JAGATHI[§] and P.SAI PRAVEEN[§]

M.I.C College of Technology, Vijayawada-521180, A.P, India

[§]K.V.S.R.Siddhartha College of Pharmaceutical Sciences
Siddhartha Nagar, Vijayawada - 520 010, A.P., India

valluru70787@gmail.com

Received 26 September 2010; Accepted 27 November 2010

Abstract: A new reverse phase high performance liquid chromatography (RP-HPLC) method has been developed for the simultaneous estimation of gemcitabine hydrochloride and capecitabine hydrochloride in combined tablet dosage form. An inertsil ODS-3 C-18 column having dimensions of 250 × 4.6 mm and particle size of 5 μm, with mobile phase containing a mixture of acetonitrile : water : triethylamine in the ratio of (70 : 28 : 2v/v) was used. The pH of mobile phase was adjusted to 4.0 with *ortho*-phosphoric acid. The flow rate was 1 mL/min and the column effluents were monitored at 260 nm. The retention time for gemcitabine hydrochloride and capecitabine hydrochloride was found to be 2.76 and 2.3 min respectively. The proposed method was validated in terms of linearity, accuracy, precision, limit of detection, limit of quantitation and robustness. The method was found to be linear in the range of 10-50 μg/mL and 4-24 μg/mL for gemcitabine hydrochloride and capecitabine hydrochloride, with regression coefficient $r = 0.999$ and $r = 0.999$, respectively.

Keywords: Gemcitabine hydrochloride, Capecitabine hydrochloride, RP-HPLC

Introduction

Gemcitabine hydrochloride (GTHC) is chemically a 2'-deoxy-2',2'-difluorocytidine monohydrochloride is a nucleoside metabolic inhibitor that exhibits antitumor activity. Spectroscopic^{1,3}, HPTLC⁵⁻⁶ and RP-HPLC method have been reported for the estimation of gemcitabine individually and in combination with other drugs.

Capecitabine hydrochloride (CTHC); pentyl [1-(3,4-dihydroxy-5-methyl-oxolan-2-yl)-5-fluoro-2-oxo-pyrimidin-4-yl]aminoformate is given for cancer treatment for metastatic⁷ breast

cancer and colorectal⁸ cancer, as well as adjuvant therapy for stage III. Various methods such as LC-MS⁹, RP-HPLC^{10,11} and spectrophotometric method have been reported for the estimation of capecitabine hydrochloride.

Literature survey reveals that no method has been reported so far for the estimation of these two drugs simultaneously in combined dosage forms. Hence, in the present study, a new reverse phase high performance liquid chromatography method was developed and validated for the simultaneous estimation of GTHC and CTHC in combined dosage forms.




Figure 1. Gemcitabine hydrochloride




Figure 2. Capecitabine hydrochloride

Experimental

The GTHC and CTHC were obtained as gift samples from Swarup Exim (India), Nagpur, Maharashtra. Acetonitrile (HPLC grade), water (HPLC grade), triethylamine and orthophosphoric acid were of AR grade. The market formulation of this combination (Label claim: 200 mg), gemget & capget (Getwell Life Sciences India Pvt. Ltd.) was purchased from the local market.

Instrumentation

A Water HPLC 2695 separation module with Water 2996-Photodiode array detector and Inertsil ODS-3 C-18 column having dimensions of 250×4.6 mm and particle size of 5 μm was used.

Chromatographic condition

The mobile phase containing acetonitrile: water: triethylamine (70: 28: 2v/v) with pH 4.0 adjusted by using orthophosphoric acid was selected as the optimum composition of mobile phase, because it was found that this solvent system resolved both the components ideally. The flow rate was set to 1 mL/min and UV detection was carried out at 260 nm. The mobile phase and samples were degassed by ultrasonication for 20 min and filtered through 0.45 μm nylon 66 (N66) membrane filter paper. All determinations were performed at constant column temperature (25 °C).

Preparation of stock solutions

10 mg of standard gemcitabine hydrochloride and 10 mg capecitabine hydrochloride were weighed accurately and transferred to two separate 100 mL volumetric flasks. Both the drugs were dissolved in 50 mL of mobile phase with shaking and then volume was made up to the mark with mobile phase to get standard stock solution of each drug. These stock solutions were filtered through 0.2 μm nylon 66 (N66) membrane filter paper and having concentration of gemcitabine hydrochloride as 100 μg/mL and as 100 μg/mL capecitabine hydrochloride.

Calibration curve

For each drug, appropriate aliquots were pipetted out from each standard stock solution into a series of 10 mL volumetric flasks. The volume was made up to the mark with mobile phase to obtain concentrations of 10, 20, 30, 40 and 50 $\mu\text{g/mL}$ of GTHC and 4, 8, 12, 16, 20 and 24 $\mu\text{g/mL}$ of CTHC. The solutions were injected in triplicates for each concentration using a 20 μL loop system and chromatographed under the conditions as described earlier. Peak areas were recorded for all the peaks at 260 nm and a standard calibration curve of peak area against concentration was plotted.




Figure 3. RP-HPLC chromatogram of a mixture of gemcitabine hydrochloride- (40 $\mu\text{g/mL}$) hydrochloride and capecitabine hydrochloride (20 $\mu\text{g/mL}$) analysis of tablet formulation

Twenty tablets were weighed and their average weight was determined and these were finely powdered. The powder equivalent to 5 mg of GTHC and 2 mg of CTHC was accurately weighed and transferred to 50 mL volumetric flask and dissolved in 25 mL mobile phase as diluent and the flask was kept in ultrasonicator for 10 min. The flask was shaken and volume was made up to the mark with mobile phase. The solution was filtered through Whatman filter paper No. 41 and it contains final concentration of 100 $\mu\text{g/mL}$ of GTHC and 40 $\mu\text{g/mL}$ of CTHC. A 20 μL volume of sample mixture was injected into the sample injector of HPLC system for six times and their chromatograms were recorded under the same chromatographic conditions as described above.

Validation method

Linearity

The standard curve was obtained in the concentration range of 10-50 $\mu\text{g/mL}$ for gemcitabine hydrochloride and 4-24 $\mu\text{g/mL}$ for capecitabine hydrochloride. The linearity of these methods were evaluated by linear regression analysis, using least squares method.

Precision

Procedure for determination of intra-day precision

In intra-day precision, the sample mixture containing 30 $\mu\text{g/mL}$ of gemcitabine hydrochloride and 16 $\mu\text{g/mL}$ of capecitabine hydrochloride was analyzed six times at different time intervals on the same day.

Procedure for determination of inter-day precision

In inter-day precision, a set of six sample mixtures containing 30 µg/mL of gemcitabine hydrochloride and 16 µg/mL of capecitabine hydrochloride were prepared and analyzed at same time on different days. The variation of the results on different days was analyzed and statistically validated.

Accuracy

Recovery studies were carried out by applying the method to drug sample present in tablet dosage form to which known amount of gemcitabine hydrochloride and capecitabine hydrochloride corresponding to 80%, 100% and 120% of label claim was added (standard addition method). After the addition of the standards, the contents were transferred to 100 mL volumetric flask and dissolved in 50 mL mobile phase and the content was kept in ultrasonicator for 25 min. Finally the volume was made up to the mark with mobile phase. The solution was filtered through Whatman filter paper No. 41. The mixed sample solutions were analyzed.

Results and Discussion

The proposed chromatographic conditions were found to be satisfactory for the determination of GTHC and CTHC in combined dosage form. The results of the assay of the marketed formulation are presented in Table 1. The method was validated statistically and validation parameters are summarized.

Table 1. Assay results of gemcitabine hydrochloride and capecitabine hydrochloride in combined dosage form

Drug	Label claim	% Drug found \pm SD*	RSD, %*
Gemcitabine hydrochloride	200 mg	98.69 \pm 0.41	0.46
Capecitabine hydrochloride	200 mg	98.17 \pm 0.45	0.45

* $n = 6$, SD; Standard deviation, RSD; Relative standard deviation

Method validation

The developed analytical method was subjected to validation as per the ICH guidelines.

Specificity

The specificity of the RP-HPLC method was determined by comparison of the chromatogram of standard solutions and sample solutions. The retention time of standard GTHC and CTHC were compared with that of sample solution. Good correlation was obtained between the retention time of standard and sample of GTHC and CTHC.

Linearity

Linearity was established by least square regression analysis of the calibration curve. The linearity range for the GTHC and CTHC were found to be 10-50 µg/mL and 4-24 µg/mL, respectively. Peak areas of GTHC and CTHC were plotted against their respective concentrations and linear regression analysis was performed on the resultant curves. The regression equations were found to be: $y = 26960x + 13795$ ($r = 0.999$) for GTHC and $y = 13786x - 1072$ ($r = 0.999$) for CTHC, respectively.

Limit of detection (LOD) and limit of quantitation (LOQ)

LOD and LOQ were determined based on the standard deviation of response and slope of calibration curve. LOD and LOQ were found to be 0.0018 and 0.0056 for GTHC and 0.0046 and 0.014 for CTHC, respectively.

Precision

For intra-day studies, five concentrations were injected into the HPLC system three times on the same day and for inter-day studies, five concentrations were injected into the HPLC system for three days. The data showed that RSD was found to be less than 2% for both; intra-day and inter-day studies, which shows that method is precise (Table 2).

Table 2. Intra-day and inter-day precision of the method

Drug	Concentration added, $\mu\text{g mL}^{-1}$	Intra-day precision		Inter-day precision	
		Mean amount found, $\mu\text{g mL}^{-1}$ (n = 6)	% RSD (n = 6)	Mean amount found, $\mu\text{g mL}^{-1}$ (n = 3)	%RSD (n = 3)
Gemcitabine hydrochloride	30	29.56 \pm 0.63	0.46	29.81 \pm 0.27	0.82
Capecitabine hydrochloride	16	16.05 \pm 0.45	0.73	15.47 \pm 0.53	0.76

Accuracy

Recovery studies were performed to determine the accuracy of the method. Recovery experiments were performed at three levels, in which the preanalyzed sample solutions were spiked with gemcitabine hydrochloride and capecitabine hydrochloride at 80%, 100% and 120% of the label claim. Three replicate samples of each concentration levels were prepared and the percentage recovery at each level was determined (Table 3).

Table 3. Accuracy studies

% Level of recovery	Amount present, $\mu\text{g/mL}$		Amount of standard drug added, $\mu\text{g/mL}$		Mean \pm S.D Amount recovered (mg) (N=3)		Mean \pm S.D % of Recovery	
	GTHC	CTHC	GTHC	CTHC	GTHC	CTHC	GTHC	CTHC
	80	10	4	10	4	19.45 \pm 0.21	7.76 \pm 0.37	100 \pm 0.23
100	10	4	20	8	30.07 \pm 0.54	11.86 \pm 0.21	100.17 \pm 05	98.82 \pm 0.3
120	10	4	30	12	39.84 \pm 0.84	15.76 \pm 0.58	99.05 \pm 0.4	98.74 \pm 0.6

Robustness

The robustness study was done by making small changes in the optimized method parameters like ± 0.1 change in mobile phase composition, ± 0.1 change in flow rate and ± 0.1 change in column temperature. There was no significant impact on the retention time. The system suitability parameters were given in Table 4.

Table 4. Summary of system suitability parameters of gemcitabine hydrochloride and capecitabine hydrochloride

Parameters	Gemcitabine hydrochloride	Capecitabine hydrochloride
Retention time	2.304	2.769
Resolution factor	-	2.7
Tailing factor	1.3	1.1
Theoretical plate	4000	3009

Conclusion

The developed RP-HPLC method can be used for routine analysis as a method for the simultaneous estimation of gemcitabine hydrochloride and capecitabine hydrochloride in pharmaceutical dosage form. The method was validated and found to be simple, accurate and precise. Statistical analysis of the developed method has been carried out, which shows good accuracy and precision.

Acknowledgment

The authors are thankful to K.V.S.R.Siddhartha College of Pharmaceutical Sciences for providing the necessary facilities and Swarup Exim (India), Nagpur, Maharashtra for providing gift samples of Gemcitabine hydrochloride and Capecitabine hydrochloride.

References

1. Budawari S, (Ed)., in; The Merck Index, 13th Edn., Merck & Co., Inc., Whitehouse Station, NJ, 2001, 221.
2. Joel G H, Lee E L, Perry B M, Raymond W R and Alfred G G, (Ed.), Goodman Gilman's The Pharmacological Basis of Therapeutics. 9th Eds., New Jersey, McGraw Hill, 1996,108.
3. Sweetman S C, Martindale: The Complete Drug Reference, 33rd Edn., Pharmaceutical Press, London, 2002 ,205.
4. <http://www.drugbank.ca/drugs/DB00376>
5. Saudagar R B, Swarnlata S and Saraf S, *Indian J Pharm Sci.*, 2007, **69**, 149-152.
6. Sejal P K and Patel N J, *Indian J Pharm Sci.*, 2009, **71(5)**, 545-547.
7. <http://www.drugbank.ca/drugs/DB00831>
8. Capka V, Xu Y and Chen Y H, *J Pharma Biomed Anal.*, 1999, **21(3)**, 507-517.
9. Miller J C and Miller J N, Statistics for Analytical Chemistry, 2nd Edition, Chichester, UK: Ellis Horwood, 1992, 127- 129.
10. Demuth J E, Basic Statistics and Pharmaceutical Statistical Applications, Newyork, Marcel Dekker, 1999, 501- 512.
11. James Swarbrick and James C Boylan, Encyclopedia of Pharmaceutical Technology, New York: Marcel Dekker; 1988, 217-224.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

