Research Article

Synthesis, Characterization, and Antimicrobial Studies of Novel Series of 2,4-Bis(hydrazino)-6-substituted-1,3,5-triazine and Their Schiff Base Derivatives

Hessa H. Al-Rasheed,¹ Essam N. Sholkamy,²
Monirah Al Alshaikh,¹ Mohammed R. H. Siddiqui ,¹
Ahmed S. Al-Obaidi,² and Ayman El-Faham ^{1,3}

¹Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

²Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

³Chemistry Department, Faculty of Science, Alexandria University, P.O. Box 426, Ibrahimia, Alexandria 12321, Egypt

Correspondence should be addressed to Ayman El-Faham; aymanel.faham@hotmail.com

Received 15 September 2017; Accepted 8 February 2018; Published 20 March 2018

Academic Editor: José M. G. Martinho

Copyright © 2018 Hessa H. Al-Rasheed et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The present work represents the synthesis, characterization, and antimicrobial studies of novel series of 2,4-bis(hydrazino)-6-substituted-1,3,5-triazine and their Schiff base derivatives. IR, NMR (¹H and ¹³C), elemental analysis, and LC-MS characterized the prepared compounds. The biological activity of the target products was evaluated as well. Twenty-two of the prepared compounds were selected according to their solubility in aqueous DMSO. Only eight compounds showed good activity against the selected pathogenic bacteria and did not show antagonistic effect against fungus *Candida albicans*. Two compounds **4k** and **5g** have wide-range effect presently in Gram-positive and Gram-negative bacteria while other compounds (**4f**, **4i**, **4m**, **5d**, **6i**, and **6h**) showed specific effect against the Gram-negative or Gram-positive bacteria. The minimum inhibitory concentration (MIC, µg/mL) of **4f**, **4i**, **4k**, and **6h** compounds against *Streptococcus mutans* was 62.5 µg/mL, 100 µg/mL, 31.25 µg/mL, and 31.25 µg/mL, respectively. The MIC of **4m**, **4k**, **5d**, **5g**, and **6h** compounds against *Staphylococcus aureus* was 62.5 µg/mL, 31.25 µg/mL, 31.25 µg/mL, 100 µg/mL, and 62.5 µg/mL, respectively. The MIC of **4k**, **5g**, and **6i** compounds against *Salmonella typhimurium* was 31.25 µg/mL, 100 µg/mL, and 62.5 µg/mL, respectively. The MIC of **6i** compound against *Escherichia coli* was 62.5 µg/mL.

1. Introduction

Hydrazones are considered as extraordinary class of compounds in the Schiff bases family, characterized by the presence of -C=N-N-. The two connected nitrogen atoms are unlike nature and the C-N double bond that is conjugated with a lone pair of the terminal nitrogen atom is mainly responsible for their physical and chemical properties [1–7]. This class of compounds is usually used as polydentate chelating agents that can form a variety of complexes with a range of transition and inner transition metals which have attracted the attention of many researchers [8, 9]. Huge derivatives of hydrazones and their metal complexes have been studied and used in different applications, such as metal ions extraction and microdetermination of metal ion [10, 11].

In addition, they have different biological activity, for example, antimicrobial activity [12–14], antifungal activity [15, 16], antitumor activity [17, 18], and insecticides [19].

On the other hand, the growing interest in s-triazine derivatives specially its chlorinated derivative (cyanuric chloride) and its efficient reaction with hydrazine to render mono-, di-, or tri- substituted hydrazino derivatives has increased [20–22]. Several prepared and well explored hydrazino-s-triazine derivatives were reported and considered with special interest in coordination chemistry and supramolecular chemistry [23, 24] as well as their enormous potential in the field of material chemistry [25, 26], complexation with large metal ions [27–36], and pharmaceutical industry [37–43].

In view of this, the present work represents the synthesis and characterization of novel series of Schiff base derived from 2,4-bis(hydrazino)-6-substituted-s-triazine and their antibacterial and antifungal activity.

2. Experimental Section

2.1. Chemistry

2.1.1. Materials. The solvents used were of analytical reagent grade. NMR (^1H and ^{13}C) spectra were recorded on a JEOL 400, 600 MHz spectrometer at room temperature. The chemical shifts were measured using internal standard $\delta = 0$ ppm. Elemental analyses were performed on Perkin-Elmer 2400 elemental analyzer, and the values found were within $\pm 0.3\%$ of the theoretical values. Melting points were recorded on a Mel-Temp apparatus in an open capillary and are uncorrected. Fourier transform infrared spectroscopy (FTIR) spectra were recorded on Nicolet 560 spectrometer from KBr discs. The reaction was followed up and the purity was checked using TLC (silica gel-protected aluminum sheets type 60 GF254, Merck). 2-(4-Iodophenyl)-3-(4-nitrophenyl)-5-phenyltetrazolium chloride (INT) (2 mg/mL) was purchased from Sigma and used for the bacterial growth test.

2.1.2. General Method for the Synthesis of 2,4-Dihydrazino-6-substituted-1,3,5-triazine Derivatives 3a–e. Compounds 3a–e were prepared according to the reported procedure with some modifications [21, 44, 45]. Hydrazine hydrate (80%) in 20 mL acetonitrile was added to a solution of 2a–e (20 mmol) in acetonitrile (50 mL) at room temperature. The reaction mixture refluxed for 3 h, and then the excess of hydrazine and solvent was removed under vacuum and the crude white precipitate was filtered off, washed with acetonitrile and ether, and then dried to afford the desired products with high yields and purities. The products were used directly in the next step.

2.1.3. General Procedure for Synthesis of 2,4-Bis(2-benzylidenehydrazinyl)-6-substituted-1,3,5-triazine Derivatives 4–8. 2,4-Dihydrazino-6-substituted-1,3,5-triazine derivatives 3a–e (2 mmol) were added portion-wise to a hot solution of different aldehyde or ketone (4 mmol) in 30 mL ethanol in the presence of 2–3 drops of acetic acid. The reaction mixture was refluxed for 3–4 hours and the progress of the reaction was followed by TLC using ethyl acetate-hexane 2:1. The reaction was left to cool down to room temperature and then the product was filtered and dried.

(1) *N*-Benzyl-4,6-bis(2-benzylidenehydrazinyl)-1,3,5-triazine-2-amine (**4a**, Supporting Information, Figure S1 in Supplementary Data). Beige solid in yield 95%; mp 150–152°C; IR (KBr, cm^{-1}): 3233 (NH), 1679 (C=N), 1587, 1517 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.52$ (d, 2H, $J = 6.0$ Hz, CH_2NH), 7.18 (t, 1H, $J = 7.2$ Hz, Ar), 7.27 (t, 2H, $J = 8.0$ Hz, Ar), 7.30–7.42 (m, 8H, Ar), 7.63 (d, 4H, $J = 8.0$ Hz, Ar), 8.14 (s, 1H, CH), 11.12 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.8$, 126.9, 127.6, 128.6, 130.1, 136.8, 140.7, 143.0, 157.9, 164.6 ppm; Anal. Calc. for $\text{C}_{24}\text{H}_{22}\text{N}_8$ (422.50): C, 68.23; H, 5.25; N, 26.52. Found: C, 68.51; H, 5.12; N, 26.37.

(2) 3,3'-((2,2'-(6-(Benzylamino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol (**4b**, Supporting Information, Figure S2 in Supplementary Data). White solid in yield 80%; mp 273–275°C; IR (KBr, cm^{-1}): 3371 (OH), 3304 (NH), 1579 (C=N), 1520, 1388 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.54$ (d, 2H, $J = 6.4$ Hz, CH_2NH), 6.77 (dd, 2H, $J^4 = 2.4$, $J^3 = 8.2$ Hz, Ar), 7.04 (d, 2H, $J = 8.0$ Hz, Ar), 7.08 (s, 2H, Ar), 7.22 (t, 3H, $J = 8.0$ Hz, Ar), 7.32 (t, 2H, $J = 7.6$ Hz, Ar), 7.38 (s, 2H, Ar), 8.06 (s, 1H, CH), 9.54 (s, 1H, OH), 10.82 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.8$, 112.9, 116.8, 118.3, 126.9, 127.6, 128.6, 130.1, 136.8, 140.7, 143.0, 157.9, 164.6, 166.6 ppm; Anal. Calc. for $\text{C}_{24}\text{H}_{22}\text{N}_8\text{O}_2$ (454.48): C, 63.43; H, 4.88; N, 24.66. Found: C, 63.21; H, 5.01; N, 24.81.

(3) *N*-Benzyl-4,6-bis(2-(3-methoxybenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4c**, Supporting Information, Figure S3 in Supplementary Data). White solid in yield 97%; mp 173–175°C; IR (KBr, cm^{-1}): 3253 (NH), 1584 (C=N), 1515, 1382 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.80$ (d, 6H, $J = 3.8$ Hz, 2OCH_3), 4.53 (d, 2H, $J = 6.0$ Hz, CH_2NH), 6.90 (dd, 2H, $J^4 = 2.4$, $J^3 = 8.4$ Hz, Ar), 7.17–7.24 (m, 3H, Ar), 7.28–7.36 (m, 6H, Ar), 7.38 (s, 2H, Ar), 8.10 (s, 1H, CH), 10.90 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.8$, 55.5, 111.4, 115.4, 119.7, 127.0, 127.6, 128.6, 130.2, 136.9, 140.7, 142.7, 159.9, 164.7, 166.5 (2-HN-C=N-) ppm; Anal. Calc. for $\text{C}_{26}\text{H}_{26}\text{N}_8\text{O}_2$ (482.54): C, 64.72; H, 5.43; N, 23.22. Found: C, 64.99; H, 5.68; N, 23.00.

(4) *N*-Benzyl-4,6-bis(2-(4-methylbenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4d**, Supporting Information, Figure S4 in Supplementary Data). Beige solid in yield 97%; mp 153–155°C; IR (KBr, cm^{-1}): 3345 (NH), 1681 (C=N), 1604, 1534 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.33$ (s, 6H, 2CH_3), 4.55 (d, 2H, $J = 5.6$ Hz, CH_2NH), 7.22 (d, 4H, $J = 9.2$ Hz, Ar), 7.31 (t, 3H, $J = 8.0$ Hz, Ar), 7.38 (s, 2H, Ar), 7.56 (d, 4H, $J = 8.0$ Hz, Ar), 8.13 (s, 1H, CH), 11.05 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 21.4$, 43.8, 126.9, 127.1, 127.7, 128.6, 129.8, 132.4, 139.0, 139.5, 142.6, 159.9, 164.7, 166.5 ppm; Anal. Calc. for $\text{C}_{26}\text{H}_{26}\text{N}_8$ (450.54): C, 69.31; H, 5.82; N, 24.87. Found: C, 69.01; H, 5.89; N, 25.00.

(5) *N*-Benzyl-4,6-bis(2-(4-methoxybenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4e**, Supporting Information, Figure S5 in Supplementary Data). White solid in yield 99%; mp 152–154°C; IR (KBr, cm^{-1}): 3346 (NH), 1677 (C=N), 1607, 1509 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.78$ (s, 6H, 2OCH_3), 4.51 (d, 2H, $J = 6.8$ Hz, CH_2NH), 6.97 (d, 4H, $J = 8.0$ Hz, Ar), 7.31 (t, 1H, $J = 8.0$ Hz, Ar), 7.37 (s, 2H, Ar), 7.57 (d, 4H, $J = 8.0$ Hz, Ar), 8.08 (s, 1H, CH), 10.70 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.8$, 55.6, 112.9, 114.6, 126.9, 128.0, 128.3, 128.5, 140.8, 142.7, 159.9, 164.7, 166.5, 165.0, 172.6 ppm; Anal. Calc. for $\text{C}_{26}\text{H}_{26}\text{N}_8\text{O}_2$ (482.54): C, 64.72; H, 5.43; N, 23.22. Found: C, 64.98; H, 5.65; N, 23.47.

(6) 3,3'-((2,2'-(6-(Benzylamino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol (**4f**, Supporting Information, Figure S6 in Supplementary Data). Beige solid in yield 85%; mp 240–242°C; IR (KBr, cm^{-1}): 3433 (OH), 3251 (NH), 1679 (C=N), 1605, 1503 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.55$ (d, 2H, $J = 5.6$ Hz, CH_2NH), 6.82 (d, 4H,

$J = 8.0$ Hz, Ar), 7.23 (t, 1H, $J = 8.0$ Hz, Ar), 7.32 (t, 2H, $J = 7.6$ Hz, Ar), 7.36 (s, 2H, Ar), 7.52 (d, 4H, $J = 8.0$ Hz, Ar), 8.08 (s, 1H, CH), 9.91 (s, 1H, OH), 11.0 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 47.5, 112.3, 114.8, 116.1, 126.9, 128.7, 129.1, 136.9, 142.8, 160.5, 166.5$ ppm; Anal. Calc. for $\text{C}_{24}\text{H}_{22}\text{N}_8\text{O}_2$ (454.48): C, 63.43; H, 4.88; N, 24.66. Found: C, 63.68; H, 4.77; N, 24.91.

(7) *N*-Benzyl-4,6-bis(2-(4-bromobenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4g**, Supporting Information, Figure S7 in Supplementary Data). Beige solid in yield 97%; mp 167–169°C; IR (KBr, cm^{-1}): 3348 (NH), 1680 (C=N), 1608, 1540 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.55$ (d, 2H, $J = 6.0$ Hz, CH_2NH), 6.76 (d, 4H, $J = 7.6$ Hz, Ar), 7.12 (s, 4H, Ar), 7.23 (t, 1H, $J = 7.2$ Hz, Ar), 7.32 (t, 2H, $J = 8.0$ Hz, Ar), 7.36 (s, 2H, Ar), 8.0 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.9, 122.8, 127.1, 127.6, 128.6, 128.7, 132.1, 133.4, 134.5, 140.3, 142.3, 161.2, 164.2$ ppm; Anal. Calc. for $\text{C}_{24}\text{H}_{20}\text{Br}_2\text{N}_8$ (454.48): C, 49.68; H, 3.47; N, 19.31. Found: C, 49.83; H, 3.61; N, 19.52.

(8) *N*-Benzyl-4,6-bis(2-(4-chlorobenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4h**, Supporting Information, Figure S8 in Supplementary Data). White solid in yield 90%; mp 155–157°C; IR (KBr, cm^{-1}): 3350 (NH), 1676 (C=N), 1610, 1542 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.53$ (d, 2H, $J = 6.0$ Hz, CH_2NH), 7.21 (t, 1H, $J = 7.6$ Hz, Ar), 7.31 (t, 2H, $J = 7.6$ Hz, Ar), 7.36 (s, 2H, Ar), 7.49 (d, 4H, $J = 8.0$ Hz, Ar), 7.65 (d, 4H, $J = 8.0$ Hz, Ar), 8.12 (s, 1H, CH), 11.02 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 45.2, 122.6, 127.0, 128.4, 129.3, 132.3, 134.5, 141.0, 142.7, 161.0, 164.3$. Anal. Calc. for $\text{C}_{24}\text{H}_{20}\text{Cl}_2\text{N}_8$ (490.12): C, 58.66; H, 4.10; N, 22.80. Found: C, 58.43; H, 4.32; N, 23.03.

(9) *N*-Benzyl-4,6-bis(2-(4-fluorobenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4i**, Supporting Information, Figure S9 in Supplementary Data). Beige solid in yield 94%; mp 172–174°C; IR (KBr, cm^{-1}): 3356 (NH), 1690 (C=N), 1606, 1504 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.55$ (d, 2H, $J = 6.0$ Hz, CH_2NH), 7.21–7.35 (m, 7H, Ar), 7.37 (s, 2H, Ar), 7.71 (s, 4H, Ar), 8.15 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.9, 116.2, 127.1, 127.6, 128.6, 128.9, 131.1, 140.2, 142.5, 160.9, 163.5, 163.9$ ppm; Anal. Calc. for $\text{C}_{24}\text{H}_{20}\text{F}_2\text{N}_8$ (458.47): C, 62.87; H, 4.40; N, 24.44. Found: C, 62.64; H, 4.55; N, 24.69.

(10) 4,4'-((2,2'-(6-(Benzylamino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))bis(*N,N*-dimethylaniline) (**4j**, Supporting Information, Figure S10 in Supplementary Data). Yellow solid in yield 95%; mp 190–191°C; IR (KBr, cm^{-1}): 3334 (NH), 1681 (C=N), 1605, 1527 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.96$ (d, 12H, $J = 9.6$ Hz, $2\text{N}(\text{CH}_3)_2$), 4.55 (d, 2H, $J = 6.4$ Hz, CH_2NH), 6.73 (d, 4H, $J = 7.2$ Hz, Ar), 7.22 (t, 1H, $J = 7.2$ Hz, Ar), 7.31 (t, 2H, $J = 7.2$ Hz, Ar), 7.37 (s, 2H, Ar), 7.49 (d, 4H, $J = 8.0$ Hz, Ar), 8.02 (s, 1H, CH), 10.85 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.8, 112.2, 122.5, 127.2, 127.5, 128.7, 129.9, 138.2, 151.6, 152.4, 160.5$ ppm; Anal. Calc. for $\text{C}_{28}\text{H}_{32}\text{N}_{10}$ (508.62): C, 66.12; H, 6.34; N, 27.54. Found: C, 66.37; H, 6.55; N, 27.67.

(11) 4,4'-((2,2'-(6-(Benzylamino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))bis(benzene-1,2-diol) (**4k**, Supporting Information, Figure S11 in Supplementary Data). Beige solid in yield 99%; mp 200–202°C; IR (KBr, cm^{-1}): 3385 (OH), 3341 (NH), 1681 (C=N), 1615, 1513 (C=C); ^1H NMR (DMSO- d_6): $\delta = 4.53$ (d, 2H, $J = 5.6$ Hz, CH_2NH), 7.21 (t, 1H, $J = 8.0$ Hz, Ar), 7.31 (t, 2H, $J = 7.6$ Hz, Ar), 7.37 (s, 2H, Ar), 7.55–7.66 (m, 6H, Ar), 8.11 (s, 1H, CH), 9.50 (s, 1H, OH), 10.85 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.9, 114.3, 116.1, 120.1, 122.3, 127.2, 127.7, 128.7, 140.0, 146.0, 147.8, 149.3, 160.9, 172.5$ ppm; Anal. Calc. for $\text{C}_{24}\text{H}_{22}\text{N}_8\text{O}_4$ (486.48): C, 59.25; H, 4.56; N, 23.03. Found: C, 59.02; H, 4.66; N, 23.25.

(12) *N*-Benzyl-4,6-bis(2-(1-(*p*-tolyl)ethylidene)hydrazinyl)-1,3,5-triazine-2-amine (**4l**, Supporting Information, Figure S12 in Supplementary Data). White solid in yield 87%; mp 103–105°C; IR (KBr, cm^{-1}): 3335 (NH), 1691 (C=N), 1587, 1499 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.26$ (s, 6H, 2CH_3), 2.32 (s, 6H, 2CH_3), 4.56 (d, 2H, $J = 6.0$ Hz, CH_2NH), 7.19–7.23 (m, 5H, Ar), 7.30 (t, 2H, $J = 7.2$ Hz, Ar), 7.40 (d, 2H, $J = 7.6$ Hz, Ar), 7.71 (s, 4H, Ar), 9.58 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 14.0, 21.2, 43.8, 126.4, 126.9, 127.8, 128.5, 129.2, 136.4, 138.4, 140.9, 148.3, 165.5, 166.7$ ppm; Anal. Calc. for $\text{C}_{28}\text{H}_{30}\text{N}_8$ (478.59): C, 70.27; H, 6.32; N, 23.41. Found: C, 70.50; H, 6.44; N, 23.67.

(13) 4,4'-((2,2'-(6-(Benzylamino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(ethan-1-yl-1-ylidene))diphenol (**4m**, Supporting Information, Figure S13 in Supplementary Data). Beige solid in yield 91%; mp. 260–262°C; IR (KBr, cm^{-1}): 3422 (OH), 3317 (NH), 1673 (C=N), 1608, 1502 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.23$ (s, 6H, 2CH_3), 4.56 (d, 2H, $J = 5.6$ Hz, CH_2NH), 6.78 (d, 4H, $J = 8.8$ Hz, Ar), 7.21 (t, 1H, $J = 7.2$, Ar), 7.30 (t, 2H, $J = 7.2$, Ar), 7.38 (d, 2H, $J = 7.2$, Ar), 7.66 (s, 4H, Ar), 9.70 (s, 1H, OH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 14.1$ (2CH_3), 43.9 (CH_2NH), 115.4, 127.1, 127.8, 128.1, 128.6, 129.9, 140.2, 149.3, 158.7, 163.9 ppm; Anal. Calc. for: $\text{C}_{26}\text{H}_{26}\text{N}_8\text{O}_2$ (482.54): C, 64.72; H, 5.43; N, 23.22. Found: C, 64.55; H, 5.31; N, 23.00.

(14) *N*-Benzyl-4,6-bis(2-benzylidenehydrazinyl)-1,3,5-triazine-2-amine (**5a**, Supporting Information, Figure S14 in Supplementary Data). White solid in yield 69%; mp 138–140°C; IR (KBr, cm^{-1}): 3225 (NH), 1675 (C=N), 1588, 1516 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.06$ (s, 3H, CH_3N), 4.83 (s, 2H, CH_2N), 7.22–7.46 (m, 11H, Ar), 7.62 (brs, 4H, Ar), 8.01 (s, 1H, CH), 11.20 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 34.2, 51.3, 126.9, 127.5, 127.9, 128.9, 129.2, 129.7, 135.4, 138.6, 142.8, 164.6, 165.9$ ppm; Anal. Calc. for $\text{C}_{25}\text{H}_{24}\text{N}_8$ (436.51): C, 68.79; H, 5.54; N, 25.67. Found: C, 69.10; H, 5.33; N, 25.98.

(15) *N*-Benzyl-*N*-methyl-4-(2-(4-methylbenzylidene)hydrazinyl)-6-(2-(4-methylbenzylidene)hydrazinyl)-1,3,5-triazine-2-amine (**5b**, Supporting Information, Figure S15 in Supplementary Data). White solid in yield 86%; mp 253–255°C; IR (KBr, cm^{-1}): 3229 (NH), 1608 (C=N), 1553, 1527 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.33$ (s, 6H, 2CH_3), 3.10 (s, 3H, CH_3N), 4.87 (s, 2H, CH_2N), 7.21–7.26 (m, 4H, Ar), 7.31–7.38

(m, 5H, Ar), 7.53 (d, 4H, $J = 8.0$ Hz, Ar) 8.11 (s, 1H, CH), 11.90 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 21.4$, 34.1, 51.1, 126.8, 127.4, 128.1, 128.8, 129.8, 132.8, 138.7, 139.1, 142.8, 164.7, 165.9 ppm; Anal. Calc. for $\text{C}_{27}\text{H}_{28}\text{N}_8$ (464.56): C, 69.80; H, 6.08; N, 24.12. Found: C, 70.03; H, 6.33; N, 24.42.

(16) *N*-Benzyl-4-(2-(4-methoxybenzylidene)hydrazinyl)-6-(2-(4-methoxybenzylidene)hydrazinyl)-*N*-methyl-1,3,5-triazine-2-amine (**5c**, Supporting Information, Figure S16 in Supplementary Data). Beige solid in yield 58%; mp 245–247°C; IR (KBr, cm^{-1}): 3354 (NH), 1678 (C=N), 1605, 1511 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.13$ (s, 3H, CH_3N), 3.80 (s, 3H, OCH_3), 3.81 (s, 3H, OCH_3), 4.90 (s, 2H, CH_2N), 6.98 (d, 4H, $J = 8.0$ Hz, Ar), 7.21–7.27 (m, 1H, Ar), 7.30–7.39 (m, 4H, $J = 8.0$ Hz, Ar), 7.57 (d, 4H, $J = 8.0$ Hz, Ar), 8.12 (s, 1H, CH), 10.80 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 34.2$, 55.8, 114.7, 127.4, 128.9, 129.6, 132.8, 139.5, 142.8, 164.7, 172.6. Anal. Calc. for $\text{C}_{27}\text{H}_{28}\text{N}_8\text{O}_2$ (496.58): C, 65.31; H, 5.68; N, 22.57. Found: C, 65.55; H, 5.81; N, 22.79.

(17) 4,4'-(2,2'-(6-(Benzyl(methyl)amino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene)diphenol (**5d**, Supporting Information, Figure S17 in Supplementary Data). Beige solid in yield 63%; mp 216–218°C; IR (KBr, cm^{-1}): 3407 (OH), 3345 (NH), 1680 (C=N), 1599, 1512 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.05$ (s, 3H, CH_3N), 4.82 (s, 2H, CH_2N), 6.77 (brs, 4H, Ar), 7.23–7.36 (m, 5H, Ar), 7.44 (s, 4H, Ar), 8.01 (brs, 1H, CH), 9.78 (s, 1H, NH), 10.70 (s, 1H, OH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 34.2$, 51.3, 116.0, 126.5, 127.5, 127.8, 128.5, 128.9, 138.8, 143.2, 150.3, 159.0, 164.5, 172.5 ppm; Anal. Calc. for $\text{C}_{25}\text{H}_{24}\text{N}_8\text{O}_2$ (468.51): C, 64.09; H, 5.16; N, 23.92. Found: C, 64.36; H, 5.00; N, 24.10.

(18) *N*-Benzyl-4-(2-(4-bromobenzylidene)hydrazinyl)-6-(2-(4-bromobenzylidene)hydrazinyl)-*N*-methyl-1,3,5-triazine-2-amine (**5e**, Supporting Information, Figure S18 in Supplementary Data). White solid in yield 99%; mp 227–229°C; IR (KBr, cm^{-1}): 3227 (NH), 1548 (C=N), 1524, 1404 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.10$ (s, 3H, CH_3N), 4.86 (s, 2H, CH_2N), 7.21–7.26 (m, 1H, Ar), 7.31–7.39 (m, 4H, Ar), 7.59–7.66 (m, 8H, Ar), 8.11 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 34.2$, 51.2, 122.6, 127.5, 128.0, 128.7, 128.8, 132.2, 134.8, 138.6, 141.5, 164.7, 165.9 ppm; Anal. Calc. for $\text{C}_{25}\text{H}_{22}\text{Br}_2\text{N}_8$ (594.30): C, 50.52; H, 3.73; N, 18.85. Found: C, 50.19; H, 3.99; N, 18.61.

(19) *N*-Benzyl-4-(2-(4-chlorobenzylidene)hydrazinyl)-6-(2-(4-chlorobenzylidene)hydrazinyl)-*N*-methyl-1,3,5-triazine-2-amine (**5f**, Supporting Information, Figure S19 in Supplementary Data). White solid in yield 82%; mp 225–227°C; IR (KBr, cm^{-1}): 3332 (NH), 1584 (C=N), 1514, 1370 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.10$ (s, 3H, CH_3N), 4.87 (s, 2H, CH_2N), 7.21–7.27 (m, 1H, Ar), 7.30–7.38 (m, 4H, Ar), 7.47 (d, 4H, $J = 8.0$ Hz, Ar), 7.65 (d, 4H, $J = 8.0$ Hz, Ar), 8.13 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 34.2$, 51.2, 127.5, 128.0, 128.4, 128.9, 129.3, 133.9, 134.4, 138.6, 141.4, 164.7, 165.9 ppm. Anal. Calc. for $\text{C}_{25}\text{H}_{22}\text{Cl}_2\text{N}_8$ (505.41): C, 59.41; H, 4.39; N, 22.17. Found: C, 59.66; H, 4.49; N, 22.00.

(20) *N*-Benzyl-4-(2-(4-fluorobenzylidene)hydrazinyl)-6-(2-(4-fluorobenzylidene)hydrazinyl)-*N*-methyl-1,3,5-triazine-2-amine (**5g**, Supporting Information, Figure S20 in Supplementary Data). White solid in yield 69%; mp 210–212°C; IR (KBr, cm^{-1}): 3332 (NH), 1602 (C=N), 1500, 1372 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.10$ (s, 3H, CH_3N), 4.87 (s, 2H, CH_2N), 7.21–7.29 (m, 5H, Ar), 7.31–7.38 (m, 4H, Ar), 7.69 (s, 4H, Ar), 8.14 (s, 1H, CH), 11.0 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 34.2$, 51.3, 116.2, 127.4, 128.1, 128.8, 128.9, 130.6, 138.7, 141.6, 162.1, 163.8, 164.7, 165.0 ppm. Anal. Calc. for $\text{C}_{25}\text{H}_{22}\text{F}_2\text{N}_8$ (472.50): C, 63.55; H, 4.69; N, 23.72. Found: C, 63.87; H, 4.89; N, 23.99.

(21) *N*-Benzyl-*N*-methyl-4-(2-(1-(*p*-tolyl)ethylidene)hydrazinyl)-6-(2-(1-(*p*-tolyl)ethylidene)hydrazinyl)-1,3,5-triazine-2-amine (**5h**, Supporting Information, Figure S21 in Supplementary Data). White solid in yield 46%; mp 150–152°C; IR (KBr, cm^{-1}): 3353 (NH), 1711 (C=N), 1586, 1494 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.25$ (s, 6H, 2CH_3), 2.32 (s, 6H, 2CH_3), 3.11 (s, 3H, CH_3N), 4.90 (s, 2H, CH_2N), 7.19–7.27 (m, 5H, Ar), 7.31–7.36 (m, 2H, Ar), 7.67–7.75 (m, 8H, Ar), 11.0 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 13.9$, 21.2, 34.1, 51.1, 126.3, 127.4, 128.1, 128.8, 129.2, 136.4, 138.4, 138.8, 147.9, 165.5, 166.0 ppm; Anal. Calc. for: $\text{C}_{29}\text{H}_{32}\text{N}_8$ (492.62): C, 70.71; H, 6.55; N, 22.75. Found: C, 70.99; H, 6.76; N, 22.98.

(22) 4,4'-(2,2'-(6-(Benzyl(methyl)amino)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(ethan-1-yl-1-ylidene)diphenol (**5i**, Supporting Information, Figure S22 in Supplementary Data). Beige solid in yield 89%; mp 205–207°C; IR (KBr, cm^{-1}): 3445 (OH), 3347 (NH), 1674 (C=N), 1597, 1503 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.25$ (s, 6H, 2CH_3), 3.11 (s, 3H, CH_3N), 4.90 (s, 2H, CH_2N), 6.78 (d, 4H, $J = 8.0$ Hz, Ar), 7.25–7.27 (m, 1H, Ar), 7.30–7.38 (m, 4H, Ar), 7.68 (d, 4H, $J = 8.0$ Hz, Ar), 9.70 (s, 1H, OH), 11.0 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 21.5$, 34.3, 51.3, 115.4, 127.5, 127.8, 128.0, 128.9, 131.1, 140.5, 145.4, 158.7, 166.2, 172.5 ppm; Anal. Calc. for $\text{C}_{27}\text{H}_{28}\text{N}_8\text{O}_2$ (496.56): C, 65.31; H, 5.68; N, 22.57. Found: C, 65.63; H, 5.33; N, 22.89.

(23) 4-(4,6-bis(2-Benzylidenehydrazinyl)-1,3,5-triazine-2-yl)morpholine (**6a**, Supporting Information, Figure S23 in Supplementary Data). White solid in yield 88%; mp 273–275°C; IR (KBr, cm^{-1}): 3228 (NH), 1596 (C=N), 1559, 1510 (C=C); ^1H NMR (DMSO- d_6): $\delta = 3.66$ (d, 4H, $J = 4.4$ Hz, 2OCH_2 -), 3.78 (s, 4H, 2NCH_2 -), 7.36 (t, 2H, $J = 7.2$ Hz, Ar), 7.42 (t, 4H, $J = 7.2$ Hz, Ar), 7.65 (d, 4H, $J = 7.6$ Hz, Ar), 8.14 (s, 1H, CH), 11.0 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): $\delta = 43.8$, 66.5, 126.9, 129.2, 129.6, 135.5, 142.8, 164.8, 165.3 ppm; Anal. Calc. for $\text{C}_{21}\text{H}_{22}\text{N}_8\text{O}$ (402.45): C, 62.67; H, 5.51; N, 27.84. Found: C, 62.88; H, 5.71; N, 28.00. (m/z) Calcd: 402.45; LC-MS [$M + H$]; Found: 403 (Supporting Information, Figure S24 in Supplementary Data).

(24) 4-(4,6-Bis(2-(4-methylbenzylidene)hydrazinyl)-1,3,5-triazine-2-yl)morpholine (**6b**, Supporting Information, Figure S25 in Supplementary Data). White solid in yield 92%; mp 298–300°C; IR (KBr, cm^{-1}): 3225 (NH), 1601 (C=N), 1559, 1503 (C=C); ^1H NMR (DMSO- d_6): $\delta = 2.32$ (s, 6H, 2CH_3),

3.64 (brs, 4H, 2 OCH₂-), 3.77 (brs, 4H, 2 NCH₂-), 7.23 (d, 4H, *J* = 8.4 Hz, Ar), 7.55 (d, 4H, *J* = 7.2 Hz, Ar), 8.10 (s, 1H, CH), 10.90 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 20.4, 43.4, 66.1, 126.6, 129.4, 132.2, 138.9, 143.3, 164.0 ppm; Anal. Calc. for: C₂₃H₂₆N₈O (430.51): C, 64.17; H, 6.09; N, 26.03. Found: C, 64.00; H, 6.21; N, 26.29.

(25) 4-(4,6-Bis(2-(4-methoxybenzylidene)hydrazinyl)-1,3,5-triazine-2-yl)morpholine (**6c**, Supporting Information, Figure S26 in Supplementary Data). White solid in yield 89%; mp. 253–255°C; IR (KBr, cm⁻¹): 3226 (NH), 1605 (C=N), 1565, 1514 (C=C); ¹H NMR (DMSO-d₆): δ = 3.65 (brs, 4H, 2 OCH₂-), 3.78 (brs, 4H, 2 NCH₂-), 3.81 (s, 6H, 2OCH₃), 6.98 (d, 4H, *J* = 8.0 Hz, Ar), 7.58 (d, 4H, *J* = 8.8 Hz, Ar), 8.10 (s, 1H, CH), 10.80 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 43.3, 55.2, 66.1, 114.2, 127.6, 127.9, 142.3, 160.1, 164.2, 165.0 ppm; Anal. Calc. for C₂₃H₂₆N₈O₃ (462.51): C, 59.73; H, 5.67; N, 24.23. Found: C, 59.92; H, 5.80; N, 24.47.

(26) 4,4'-(2,2'-(6-Morpholino-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene)bis(methanylylidene))diphenol (**6d**, Supporting Information, Figure S27 in Supplementary Data). White solid in yield 94%; mp 275–277°C; IR (KBr, cm⁻¹): 3435 (OH), 3349 (NH), 1677 (C=N), 1582, 1509 (C=C); ¹H NMR (DMSO-d₆): δ = 3.64 (brs, 4H, 2 OCH₂-), 3.76 (s, 4H, 2 NCH₂-), 6.82 (d, 4H, *J* = 5.2 Hz, Ar), 7.49 (s, 4H, Ar), 8.05 (s, 1H, CH), 9.85 (brs, 1H, OH), 11.16 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 43.0, 66.0, 115.7, 125.8, 128.3, 145.0, 149.9, 159.0, 163.5 ppm; Anal. Calc. for: C₂₁H₂₂N₈O₃ (434.45): C, 58.06; H, 5.10; N, 25.79. Found: C, 58.33; H, 5.31; N, 25.50. (*m/z*) Calcd: 434.45; LC-MS (*M* + *H*); Found: 435.2.

(27) 4-(4,6-Bis(2-(4-Bromobenzylidene)hydrazinyl)-1,3,5-triazine-2-yl)morpholine (**6e**, Supporting Information, Figure S28 in Supplementary Data). White solid in yield 83%; mp. 308–310°C; IR (KBr, cm⁻¹): 3225 (NH), 1593 (C=N), 1561, 1515 (C=C); ¹H NMR (DMSO-d₆): δ = 3.65 (brs, 4H, 2 OCH₂-), 3.77 (brs, 4H, 2 NCH₂-), 7.58 (m, 8H, Ar), 8.10 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 43.5, 66.0, 122.2, 128.3, 131.8, 134.2, 141.3, 164.2, 164.8 ppm; Anal. Calc. for: C₂₁H₂₀Br₂N₈O (560.24): C, 45.02; H, 3.60; N, 20.00. Found: C, 45.31; H, 3.43; N, 20.28. (*m/z*) Calcd: 560.24; LC-MS [*M* + *H*]; Found: 561.1.

(28) 4-(4,6-Bis(2-(4-Chlorobenzylidene)hydrazinyl)-1,3,5-triazine-2-yl) (**6f**, Supporting Information, Figure S29 in Supplementary Data). White solid in yield 96%; mp. 305–307°C; IR (KBr, cm⁻¹): 3226 (NH), 1596 (C=N), 1563, 1515 (C=C); ¹H NMR (DMSO-d₆): δ = 3.65 (brs, 4H, 2 OCH₂-), 3.78 (brs, 4H, 2 NCH₂-), 7.48 (d, 4H, *J* = 8.0 Hz, Ar), 7.66 (d, 4H, *J* = 8.8 Hz, Ar), 8.12 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 43.3, 66.0, 128.0, 128.8, 133.4, 133.9, 141.1, 164.3, 164.8 ppm; Anal. Calc. for: C₂₁H₂₀Cl₂N₈O (471.34): C, 53.51; H, 4.28; N, 23.77. Found: C, 53.77; H, 4.01; N, 23.56. (*m/z*) Calcd: 471.34; LC-MS [*M* + *H*]; Found: 471.2.

(29) 4-(4,6-Bis(2-(4-Fluorobenzylidene)hydrazinyl)-1,3,5-triazine-2-yl)morpholine (**6g**, Supporting Information, Figure S30 in Supplementary Data). White solid in yield 91%; mp. 301–303°C; IR (KBr, cm⁻¹): 3229 (NH), 1601 (C=N), 1568, 1515

(C=C); ¹H NMR (DMSO-d₆): δ = 3.65 (brs, 4H, 2 OCH₂-), 3.78 (brs, 4H, 2 NCH₂-), 7.25–7.34 (m, 4H, Ar), 7.69–7.74 (m, 4H, Ar), 8.14 (s, 1H, CH), 11.10 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 43.3, 66.1, 115.8, 128.8, 131.4, 141.4, 161.5, 163.9 ppm; Anal. Calc. for C₂₁H₂₀F₂N₈O (438.43): C, 57.53; H, 4.60; N, 25.56. Found: C, 57.87; H, 4.32; N, 25.23.

(30) 4-(4,6-Bis(2-(1-(*p*-Tolyl)ethylidene)hydrazinyl)-1,3,5-triazine-2-yl)morpholine (**6h**, Supporting Information, Figure S31 in Supplementary Data). The product was obtained as a yellow solid in yield 92%; mp. 235–236°C; IR (KBr, cm⁻¹): 3358 (NH), 1643 (C=N), 1563, 1442 (C=C); ¹H NMR (DMSO-d₆): δ = 2.27 (s, 6H, 2CH₃CN), 2.32 (s, 6H, 2CH₃), δ 3.65 (t, 4H, *J* = 4.0 Hz, 2 OCH₂-), δ 3.80 (t, 4H, *J* = 4.4 Hz, 2 NCH₂-), 7.20 (d, 4H, *J* = 8.0 Hz, Ar), δ 7.72 (d, 4H, *J* = 7.2 Hz, Ar), δ 11.18 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 13.6, 20.8, 43.3, 66.1, 125.9, 128.8, 135.9, 138.0, 149.9, 165.0 ppm; Anal. Calc. for C₂₅H₃₀N₈O (458.56): C, 65.48; H, 6.59; N, 24.44. Found: C, 65.71; H, 6.33; N, 24.29.

(31) 4,4'-(2,2'-(6-Morpholino-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(ethan-1-yl-1-ylidene))diphenol (**6i**, Supporting Information, Figure S32 in Supplementary Data). White solid in yield 93%; mp. 265–267°C; IR (KBr, cm⁻¹): 3430 (OH), 3360 (NH), 1673 (C=N), 1593, 1503 (C=C); ¹H NMR (DMSO-d₆): δ = 2.28 (s, 6H, 2CH₃CN), 3.66 (brs, 4H, 2 OCH₂-), 3.82 (brs, 4H, 2 NCH₂-), 6.80 (d, 4H, *J* = 8.0 Hz, Ar), 7.69 (d, 4H, *J* = 8.4 Hz, Ar), 9.80 (brs, 1H, OH), 11.80 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 14.3, 44.1, 66.4, 115.6, 128.2, 129.0, 152.0, 159.1, 165.5, 172.5 ppm; Anal. Calc. for C₂₃H₂₆N₈O₃ (462.50): C, 59.73; H, 5.67; N, 24.23. Found: C, 59.99; H, 5.83; N, 24.51.

(32) 2,4-Bis(2-benzylidenehydrazinyl)-6-(piperidin-1-yl)-1,3,5-triazine (**7a**, Supporting Information, Figure S33 in Supplementary Data). White solid in yield 94%; mp. 247–249°C; IR (KBr, cm⁻¹): 3228 (NH), 1677 (C=N), 1550, 1512 (C=C); ¹H NMR (DMSO-d₆): δ = 1.54 (s, 4H, 2CH₂), 1.63 (s, 2H, CH₂), 3.80 (s, 4H, 2CH₂N), 7.43 (brs, 6H, Ar), 7.70 (brs, 4H, Ar), 8.18 (s, 1H, CH), 11.0 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 24.7, 25.9, 44.3, 127.1, 129.2, 129.7, 134.9, 142.5, 165.1 ppm; Anal. Calc. for C₂₂H₂₄N₈ (400.49): C, 65.98; H, 6.04; N, 27.98. Found: C, 65.79; H, 6.19; N, 28.14. (*m/z*) Calcd: 400.49; LC-MS [*M* + *H*]; Found: 401 (Supporting Information, Figure S50 in Supplementary Data).

(33) 2,4-Bis(2-(4-methylbenzylidene)hydrazinyl)-6-(piperidin-1-yl)-1,3,5-triazine (**7b**, Supporting Information, Figure S34 in Supplementary Data). White solid in yield 89%; mp. 282–284°C; IR (KBr, cm⁻¹): 3235 (NH), 1673 (C=N), 1567, 1509 (C=C); ¹H NMR (DMSO-d₆): δ = 1.58 (s, 4H, 2CH₂), 1.62 (d, 2H, *J* = 4.4 Hz, CH₂), 2.32 (s, 6H, 2CH₃), 3.78 (s, 4H, 2CH₂N), 7.24 (d, 4H, *J* = 7.2 Hz, Ar), 7.54 (d, 4H, *J* = 7.6 Hz, Ar), 8.10 (s, 1H, CH), 10.90 (brs, 1H, NH) ppm; ¹³C NMR (DMSO-d₆): δ = 20.9, 24.3, 25.5, 43.6, 126.4, 129.3, 132.2, 138.7, 142.4, 164.0 ppm; Anal. Calcd for C₂₄H₂₈N₈ (428.53): C, 67.27; H, 6.59; N, 26.15. Found: C, 67.43; H, 6.71; N, 26.38.

(34) 2,4-Bis(2-(4-methoxybenzylidene)hydrazinyl)-6-(piperidin-1-yl)-1,3,5-triazine (**7c**, Supporting Information, Figure S35 in Supplementary Data). White solid in yield 94%; mp. 257–259°C; IR (KBr, cm^{-1}): 3341 (NH), 1665 (C=N), 1570, 1512 (C=C); ^1H NMR (DMSO- d_6): δ = 1.52 (s, 4H, 2CH₂), 1.62 (s, 2H, CH₂), 3.79 (s, 6H, 2OCH₃), 3.81 (s, 4H, 2CH₂N), 7.0 (d, 4H, J = 8.8 Hz, Ar), 7.60 (d, 4H, J = 8.0 Hz, Ar), 8.09 (s, 1H, CH), 10.90 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 24.2, 25.4, 43.8, 55.3, 114.2, 126.5, 128.2, 140.8, 160.4, 161.6 ppm.

(35) 4,4'-((2,2'-(6-(Piperidin-1-yl)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol (**7d**, Supporting Information, Figure S36 in Supplementary Data). Beige solid in yield 99%; mp. 331–333°C; IR (KBr, cm^{-1}): 3435 (OH), 3349 (NH), 1677 (C=N), 1590, 1511 (C=C); ^1H NMR (DMSO- d_6): δ = 1.54 (s, 4H, 2CH₂), 1.63 (s, 2H, CH₂), 3.78 (s, 4H, 2CH₂N), 6.86 (s, 4H, Ar), 7.56 (s, 4H, Ar), 8.09 (s, 1H, CH), 9.80 (s, 1H, OH), 10.30 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 24.6, 25.9, 44.5, 116.2, 125.8, 129.3, 142.7, 159.5, 164.0 ppm; Anal. Calcd for: C₂₂H₂₄N₈O₂ (432.48): C, 61.10; H, 5.59; N, 25.91. Found: C, 61.33; H, 5.78; N, 25.72. (m/z) Calcd: 432.48; LC-MS [M + H]; Found: 433 (Supporting Information, Figure S51 in Supplementary Data).

(36) 2,4-Bis(2-(4-Bromobenzylidene)hydrazinyl)-6-(piperidin-1-yl)-1,3,5-triazine (**7e**, Supporting Information, Figure S37 in Supplementary Data). White solid in yield 96%; mp. 288–290°C; IR (KBr, cm^{-1}): 3240 (NH), 1673 (C=N), 1569, 1512 (C=C); ^1H NMR (DMSO- d_6): δ = 1.52 (s, 4H, 2CH₂), 1.63 (d, 2H, J = 4.4 Hz, CH₂), 3.78 (s, 4H, 2CH₂N), 7.60 (d, 4H, J = 8.0 Hz, Ar), 7.64 (d, 4H, J = 8.0 Hz, Ar), 8.11 (s, 1H, CH), 11.15 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 24.3, 25.5, 43.5, 122.1, 128.2, 131.7, 134.3, 140.9, 164.3 ppm; Anal. Calcd for C₂₂H₂₂Br₂N₈ (558.27): C, 47.33; H, 3.97; N, 20.07. Found: C, 47.61; H, 3.77; N, 20.25.

(37) 2,4-Bis(2-(4-chlorobenzylidene)hydrazinyl)-6-(piperidin-1-yl)-1,3,5-triazine (**7f**, Supporting Information, Figure S38 in Supplementary Data). White solid in yield 97%; mp. 293–295°C; IR (KBr, cm^{-1}): 3237 (NH), 1679 (C=N), 1570, 1513 (C=C); ^1H NMR (DMSO- d_6): δ = 1.52 (s, 4H, 2CH₂), 1.62 (d, 2H, J = 4.4 Hz, CH₂), 3.78 (s, 4H, 2CH₂N), 7.49 (d, 4H, J = 8.0 Hz, Ar), 7.68 (d, 4H, J = 8.4 Hz, Ar), 8.13 (s, 1H, CH), 11.20 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 24.2, 25.5, 43.7, 128.1, 128.8, 132.6, 135.9, 141.3, 160, 164.0 ppm; Anal. Calcd for C₂₂H₂₂Cl₂N₈ (469.37): C, 56.30; H, 4.72; N, 23.87. Found: C, 56.55; H, 4.94; N, 24.03. (m/z) Calcd: 469.37; LC-MS [M]; Found: 469 (Supporting Information, Figure S52 in Supplementary Data).

(38) 2,4-Bis(2-(4-Fluorobenzylidene)hydrazinyl)-6-(piperidin-1-yl)-1,3,5-triazine (**7g**, Supporting Information, Figure S39 in Supplementary Data). White solid in yield 95%; mp. 275–277°C; IR (KBr, cm^{-1}): 3358 (NH), 1680 (C=N), 1603, 1501 (C=C); ^1H NMR (DMSO- d_6): δ = 1.56 (s, 4H, 2CH₂), 1.64 (s, 2H, CH₂), 3.82 (s, 4H, 2CH₂N), 7.32 (brs, 4H, Ar), 7.85 (brs, 4H, Ar), 8.23 (brs, 1H, CH), 12.50 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 24.4, 25.9, 44.7, 116.4, 129.7, 130.8, 144.2, 162.7, 164.2 ppm; Anal. Calcd for C₂₂H₂₂F₂N₈

(436.46): C, 60.54; H, 5.08; N, 25.67. Found: C, 60.51; H, 5.23; N, 25.38.

(39) 2-(Piperidin-1-yl)-4,6-bis(2-(1-(*p*-tolyl)ethylidene)hydrazinyl)-1,3,5-triazine (**7h**, Supporting Information, Figure S40 in Supplementary Data). White solid in yield 91%; mp. 222–224°C; IR (KBr, cm^{-1}): 3356 (NH), 1628 (C=N), 1562, 1437 (C=C); ^1H NMR (DMSO- d_6): δ = 1.52 (s, 4H, 2CH₂), 1.62 (s, 2H, CH₂), 2.28 (s, 6H, 2CH₃CN), 2.33 (s, 6H, 2CH₃), 3.81 (s, 4H, 2CH₂N), 7.19 (d, 4H, J = 8.0 Hz, Ar), 7.71 (d, 4H, J = 8.0 Hz, Ar), 9.70 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 13.9, 21.3, 24.9, 26.0, 43.9, 126.4, 129.3, 136.5, 138.4, 147.9, 164.8, 165.6 ppm; Anal. Calcd for C₂₆H₃₂N₈ (456.59): C, 68.39; H, 7.06; N, 24.54. Found: C, 68.65; H, 7.33; N, 24.23.

(40) 4,4'-((2,2'-(6-(Piperidin-1-yl)-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(ethan-1-yl-1-ylidene))diphenol (**7i**, Supporting Information, Figure S41 in Supplementary Data). Beige solid in yield 85%; mp. 288–290°C; IR (KBr, cm^{-1}): 3440 (OH), 3358 (NH), 1672 (C=N), 1597, 1500 (C=C); ^1H NMR (DMSO- d_6): δ = 1.55 (s, 4H, 2CH₂), 1.64 (s, 2H, CH₂), 2.31 (s, 6H, 2CH₃), 3.84 (s, 4H, 2CH₂N), 6.82 (brs, 4H, Ar), 7.70 (brs, 4H, Ar), 9.90 (brs, 2H, OH, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 14.4, 24.6, 26.0, 44.3, 115.6, 128.2, 131.1, 154.2, 159.1, 160.0 ppm; Anal. Calcd for C₂₄H₂₈N₈O₂ (460.53): C, 62.59; H, 6.13; N, 24.33. Found: C, 62.82; H, 6.42; N, 24.61. (m/z) Calcd: 460.53; LC-MS [M + 2H]; Found: 463.2 (Supporting Information, Figure S42 in Supplementary Data).

(41) 2,4-Bis(2-benzylidenehydrazinyl)-6-methoxy-1,3,5-triazine (**8a**, Supporting Information, Figure S43 in Supplementary Data). Beige solid in yield 70%; mp 292–294°C; IR (KBr, cm^{-1}): 3224 (NH), 1620 (C=N), 1586, 1547 (C=C); ^1H NMR (DMSO- d_6): δ = 3.91 (s, 3H, OCH₃), 7.36–7.47 (m, 6H, Ar), 7.67 (d, 4H, J = 7.2 Hz, Ar), 8.19 (s, 1H, CH), 11.35 (brs, 1H, NH), 11.51 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 54.0, 126.8, 128.8, 129.5, 134.7, 144.5, 161.4, 165.5, 171.0 ppm; Anal. Calcd for C₁₈H₁₇N₇O (347.38): C, 62.24; H, 4.93; N, 28.23. Found: C, 62.51; H, 5.04; N, 28.49.

(42) 2-Methoxy-4,6-bis(2-(4-methylbenzylidene)hydrazinyl)-1,3,5-triazine (**8b**, Supporting Information, Figure S44 in Supplementary Data). White solid in yield 84%; mp 298–300°C; IR (KBr, cm^{-1}): 3323 (NH), 1616 (C=N), 1586, 1540 (C=C); ^1H NMR (DMSO- d_6): δ = 2.33 (s, 6H, 2CH₃), 3.89 (s, 3H, OCH₃), 7.25 (d, 4H, J = 7.2 Hz, Ar), 7.56 (d, 4H, J = 7.2 Hz, Ar), 8.14 (s, 1H, CH), 11.28 (brs, 1H, NH), ppm; ^{13}C NMR (DMSO- d_6): δ = 21.5, 54.4, 127.1, 129.8, 132.4, 139.6, 144.3, 165.5, 166.2 ppm; Anal. Calcd for C₂₀H₂₁N₇O (375.44): C, 63.98; H, 5.64; N, 26.12. Found: C, 63.82; H, 5.73; N, 26.33.

(43) 2-Methoxy-4,6-bis(2-(4-methoxybenzylidene)hydrazinyl)-1,3,5-triazine (**8c**, Supporting Information, Figure S45 in Supplementary Data). White solid in yield 86%; mp 300–302°C; IR (KBr, cm^{-1}): 3277 (NH), 1612 (C=N), 1589, 1545 (C=C); ^1H NMR (DMSO- d_6): δ = 3.79 (s, 6H, 2OCH₃), 3.89 (s, 3H, OCH₃), 7.0 (d, 4H, J = 8.0 Hz, Ar), 7.61 (d, 4H, J = 7.2 Hz, Ar),

8.13 (s, 1H, CH), 11.22 (brs, 2H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 53.9, 55.3, 114.3, 127.2, 128.2, 144.0, 160.4, 164.7 ppm; Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{N}_7\text{O}_3$ (407.43): C, 58.96; H, 5.20; N, 24.06. Found: C, 58.77; H, 5.02; N, 24.33.

(44) 4,4'-(2,2'-(6-Methoxy-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene)bis(methanylylidene))diphenol (**8d**, Supporting Information, Figure S46 in Supplementary Data). Beige solid in yield 83%; mp 294–296°C; IR (KBr, cm^{-1}): 3522 (OH), 3268 (NH), 1618 (C=N), 1545, 1514 (C=C); ^1H NMR (DMSO- d_6): δ = 3.84 (s, 3H, OCH_3), 6.77 (d, 4H, J = 8.0 Hz, Ar), 7.45 (d, 4H, J = 7.2 Hz, Ar), 8.03 (s, 1H, CH), 9.83 (s, 1H, OH), 11.10–11.30 (brs, 2H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 53.9, 115.7, 125.8, 128.4, 144.0, 158.9, 165.8, 171.0 ppm; Anal. Calcd for $\text{C}_{18}\text{H}_{17}\text{N}_7\text{O}_3$ (379.37): C, 56.99; H, 4.52; N, 25.84. Found: C, 56.76; H, 4.76; N, 26.07.

(45) 2,4-Bis(2-(4-bromobenzylidene)hydrazinyl)-6-methoxy-1,3,5-triazine (**8e**, Supporting Information, Figure S47 in Supplementary Data). White solid in yield 87%; mp. 317–319°C; IR (KBr, cm^{-1}): 3318 (NH), 1678 (C=N), 1586, 1535 (C=C); ^1H NMR (DMSO- d_6): δ = 3.89 (s, 3H, OCH_3), 7.62 (brs, 8H, Ar), 8.15 (s, 1H, CH), 11.4 (brs, 1H, NH), 11.70 (brs, 1H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 54.0, 122.6, 128.5, 131.7, 133.9, 142.8, 163.8 ppm; Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{Br}_2\text{N}_7\text{O}$ (505.17): C, 42.80; H, 2.99; N, 19.41. Found: C, 43.01; H, 3.15; N, 19.66.

(46) 2,4-Bis(2-(4-chlorobenzylidene)hydrazinyl)-6-methoxy-1,3,5-triazine (**8f**, Supporting Information, Figure S48 in Supplementary Data). Beige solid in yield 86%; mp 308–310°C; IR (KBr, cm^{-1}): 3315 (NH), 1616 (C=N), 1586, 1534 (C=C); ^1H NMR (DMSO- d_6): δ = 3.91 (s, 3H, OCH_3), 7.50 (d, 4H, J = 7.2 Hz, Ar), 7.69 (d, 4H, J = 8.0 Hz, Ar), 8.17 (s, 1H, CH), 11.50 (brs, 2H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 53.9, 122.5, 128.4, 128.9, 131.7, 133.9, 141.4, 161.0 ppm; Anal. Calc. for $\text{C}_{18}\text{H}_{15}\text{Cl}_2\text{N}_7\text{O}$ (416.26): C, 51.94; H, 3.63; N, 23.55. Found: C, 51.86; H, 3.88; N, 23.79.

(47) 2,4-Bis(2-(4-fluorobenzylidene)hydrazinyl)-6-methoxy-1,3,5-triazine (**8g**, Supporting Information, Figure S49 in Supplementary Data). Beige solid in yield 69%; mp 323–325°C; IR (KBr, cm^{-1}): 3217 (NH), 1603 (C=N), 1547, 1504 (C=C); ^1H NMR (DMSO- d_6): δ = 3.90 (s, 3H, OCH_3), 7.26–7.30 (m, 4H, Ar), 7.71–7.74 (m, 4H, Ar), 8.17 (s, 1H, CH), 11.40 (brs, 2H, NH) ppm; ^{13}C NMR (DMSO- d_6): δ = 54.0, 116.0, 128.6, 131.2, 143.2, 161.4, 164.2 ppm; Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{F}_2\text{N}_7\text{O}$ (383.35): C, 56.39; H, 3.94; N, 25.58. Found: C, 56.65; H, 4.05; N, 25.32.

2.2. Antimicrobial Activity. The antibacterial activity of the prepared compounds was assessed against two selected bacteria groups: Gram-positive bacteria, namely, *Streptococcus mutans* (wild strain) and *Staphylococcus aureus* (ATCC 29213), and Gram-negative bacteria, namely, *Escherichia coli* (ATCC 25922) and *Salmonella typhimurium* (ATCC 14028), maintained in Brain Heart Infusion medium at -20°C . One mL of each targeted grown bacterial culture was added to 100 mL of Brain Heart Infusion broth and incubated at $37^\circ\text{C} \pm 1^\circ\text{C}$ for subculture of all targeted bacterial cultures. Using

of sterile physiological solution, concentration of bacterial inoculum was determined to be 10^8 CFU/mL (0.5 McFarland turbidity standards) for antimicrobial tests. In case of fungus *Candida albicans*, the PDA was used as a medium in antagonistic activity against the tested fungi.

2.2.1. Primary Screening of Synthetic Organic Compounds for Bioactivity by Agar Diffusion Disc. The tested compounds were dissolved in DMSO and prepared at concentration 2500 $\mu\text{g/mL}$. The sterilized Mueller Hinton agar plates seeded with pathogenic bacteria were prepared; 6 mm paper discs with 100 μg of the tested compounds were placed in the plates seeded with tested pathogenic bacteria. The inoculated plates were kept at 37°C for 24 h for bacteria while, in case of fungi, the used medium in antagonistic activity against tested fungi is PDA. Antimicrobial activity was determined by the inhibition zone [46].

2.2.2. Minimum Inhibitory Concentration (MIC) Determination. The bioactivity of the tested compounds was determined by using a microdilution method, using Mueller–Hinton broth. The inoculum was prepared as described previously and the tested compounds were stocked in 0.25% of DMSO (concentration of each compound 1.25 mg/mL). After that, serial dilutions were done by addition of 100 μL of culture broth which was poured in microplate and 25 μL of each concentration of tested compound in the first line of microplate to reach concentrations ranging from 125 to 0.973 $\mu\text{g/mL}$ in presence of the positive and negative control. The microplates were vaccinated with 5 μL of a bacterial inoculum (10^8 CFU/mL) and were kept for 24 h at 37°C in triplicate tests. After incubation period, we added 20 μL INT (0.5 mg/mL) to achieve bacterial growth. The INT containing microplates were incubated at 37°C for 30 min and the collected data were expressed in micrograms per milliliters [47].

3. Results and Discussion

3.1. Chemistry. The first chlorine atom of cyanuric chloride **1** was replaced by benzylamine, *N*-methylbenzylamine, morpholine, piperidine, or methoxy, while two hydrazine groups replaced the second and third chlorine atoms. Accordingly, 2,4-dihydrazino-6-substituted-1,3,5-triazine derivatives **3a–e** were prepared by displacement of the two chlorine atoms by two hydrazine groups.

Cyanuric chloride **1** was first reacted with different amines such as benzylamine, *N*-methylbenzylamine, morpholine, piperidine, or methanol at 0°C for 1–2 h to afford the products **2a–e** (Scheme 1); the spectral data agreed with the reported data [44, 45]. The dichloro derivatives **2a–e** were reacted with hydrazine hydrate (80%) under reflux using acetonitrile as a solvent to render the products of 2,4-dihydrazino-6-substituted-1,3,5-triazine derivatives **3a–e** which were used directly in the next step without further purification (Scheme 1).

The products **4–8** were prepared by condensation of the hydrazine derivatives **3a–e** with different aldehydes or

SCHEME 1: Synthesis of 6-substituted-s-triazine-Schiff base derivative.

substituted acetophenone in ethanol as solvent and in the presence of drops of acetic acid (Scheme 1).

The ^1H NMR spectrum of **4b** (Figure 1) as a prototype for the **4a-m** series displayed two doublet peaks at δ 4.54 (J = 6.4 Hz) and 7.04 ppm (J = 8.0), attributed to resonances of protons of CH_2N and 2H-6 respectively, doublet of doublet at δ 6.77 ppm which integrated two protons for 2H-4, two triplets at δ 7.22 (J = 8.0 Hz) and 7.32 ppm (J = 7.6) for H-4', 2H-5, and H-3', H-5', and four singlets at δ 7.08, 7.38, 8.06,

and 9.54 ppm for 2H-2, H-2' combined H-6', CH, and OH, respectively, while the NH proton appeared as abroad singlet at δ 10.82 ppm. The ^{13}C NMR spectrum also reveals the symmetry of **4b**, as indicated by the appearance of fourteen distinct carbon peaks, among which are four peaks at δ 112.9, 116.8, 118.3, 126.9, 127.6, 128.6, 130.1, 136.8, 140.7, and 157.9 ppm related to the aromatic carbons, in addition to four peaks at δ 43.8, 143.0, 164.6, and 166.6 ppm belonging to (CH_2NH), and ($\text{C}=\text{N}$), respectively.

FIGURE 1: Structure of **4b**.

The ^1H NMR spectrum of **5b** as a prototype for the **5a-i** series exhibited three singlets at δ 2.33, 3.10, and 4.87 ppm, which were assigned to protons of 2CH_3 , CH_3N , and CH_2N , respectively. This spectrum showed three signals for aromatic rings at δ 7.21–7.26 (m, 4H, 2H-3, and 2H-5), 7.31–7.38 (m, 5H, H-2', H-3', H-4', H-5', and H-6'), and 7.53 (d, 4H, J = 8.0 Hz, 2H-2, and 2H-6) ppm. This spectrum also showed a sharp singlet at δ 8.11 ppm integrating one proton, which was assigned to CH, in addition the broad singlet at δ 11.90 ppm for NH. The ^{13}C NMR spectrum of **5b** showed fourteen individual carbon signals. The three signals in the sp^3 region appeared at δ 21.4, 34.1, and 51.1 ppm for (2CH_3), (CH_3N), and (CH_2N), respectively, in addition to signals at δ 142.8, 164.7, and 165.9 ppm for ($\text{C}=\text{N}$). Assignments of resonances of six carbons of the benzyl were at δ 127.4, 128.1, 128.8, and 138.7 ppm. The signals of the remaining aromatic carbons appeared at δ 126.8, 129.8, 132.8, and 139.1 ppm.

The ^1H NMR spectrum of **6a** as a prototype for the **6a-i** series showed two sharp singlets at δ 3.66 and 3.78 ppm for 2 OCH_2 - and 2 NCH_2 - related to the morpholine, respectively, singlet at δ 8.14 ppm for CH, and a broad singlet at δ 11.0 ppm for NH. Three signals for aromatic protons appeared at δ 7.42 (t, 4H, J = 7.2 Hz, 2H-3, and 2H-5) and 7.65 (d, 4H, J = 7.6 Hz, 2H-2, and 2H-6). The ^{13}C NMR spectrum also confirms the structure for compound **6a**, as indicated by the appearance of two signals at δ 43.8 and 66.5 ppm for (2 NCH_2 -) and (2 OCH_2 -), respectively, for the morpholino moiety, while the other three signals appeared at δ 142.8, 164.8, and 165.3 ppm for ($\text{C}=\text{N}$), in addition to the observation of the aromatic ring carbons appearing at δ 126.9, 129.2, 129.6, and 135.5 ppm, respectively. The LC-MS of compound **6a** (Supporting Information, Figure S24 in Supplementary Data) the exact molecular mass [$\text{M} + \text{H}$] 403 (m/z calcd. 402.45) with R_t 15.3 min, using buffer A: 0.1% formic acid in H_2O ; and buffer B: 0.1% formic acid in CH_3CN in 30 min.

The ^1H NMR spectrum of **7i** as a prototype for **7a-i** series showed three singlet peaks at δ 1.55, 1.64, and 3.84 ppm for the 2CH_2 , CH_2 , and 2NCH_2 (piperidine moiety), while it showed a singlet at δ 2.31 ppm for the methyl group of the acetophenone moiety and two broad singlets at δ 9.90 and 11.0 ppm for OH and NH, respectively, beside two doublets forming AB system at δ 6.82 (J = 8.0 Hz) and 7.70 ppm (J = 7.2 Hz) for the aromatic proton H-3, H-5 and H-2, H-6, respectively. The ^{13}C NMR spectrum of **7i** exhibited absorption peak at δ 14.4 related to methyl group, three peaks for the piperidine residue at δ 24.6, 26.0, and 44.3 ppm

related to (CH_2), (2CH_2), and ($2\text{CH}_2\text{N}$), respectively, and absorption peaks at δ 154.2 and 160.0 ppm related to ($\text{C}=\text{N}$), besides four peaks at δ 115.6, 128.2, 131.1, and 159.1 ppm for the aromatic carbons. The LC-MS of compound **7i** (Supporting Information, Figure S42 in Supplementary Data) using buffer A: 0.1% formic acid in H_2O and buffer B: 0.1% formic acid in CH_3CN in 30 min showed one peak at R_t 14.08 min with the expected mass [$\text{M} + 2\text{H}$] 463.2 (m/z calcd. 460.53).

The ^1H NMR spectrum of **8b** as a prototype for the **8a-g** series revealed the resonances of protons methyl group at δ 2.33 ppm (s, 2CH_3) and methoxy group at δ 3.89 ppm (s, OCH_3), in addition to the two doublets that appeared at δ 7.25 (d, 4H, J = 7.2 Hz) and 7.56 ppm (d, 4H, J = 7.2 Hz) representing of 2H-2, 2H-3, and 2H-5, 2H-6 (AB system) while singlet (integrated for one proton) appearing at δ 8.14 ppm was attributed to CH. Finally, the proton of NH group appeared as broad singlet at δ 11.28 ppm. The ^{13}C NMR spectrum of **8b** showed signal at δ 21.5 for the methyl carbon while methoxy carbon appeared at δ 54.4 ppm and the aromatic carbons signals at δ 127.1, 129.8, 132.4, and 139.6 ppm. The three signals at δ 144.3, 165.5, and 166.2 ppm in the later spectrum were assigned to ($\text{C}=\text{N}$).

3.2. Biology

3.2.1. Bioactivity of Tested Compounds against the Pathogenic Bacteria and Fungus *Candida albicans*. In the present study, twenty-two compounds from the above series **4-8** were tested for bioactivities against the selected bacteria and fungi; only eight compounds have antiactivities of target pathogenic bacteria. The results in Table 1 showed that the tested compounds, namely, **4f**, **4k**, **4l**, **4m**, **5d**, **5g**, **6i**, and **6h**, have bioactivity against the target pathogenic bacteria as antimicrobial agents and do not have antagonistic effect against fungus *Candida albicans*. The present study also showed that compounds **4k** and **5g** have wide-range effect presently in Gram-positive and Gram-negative bacteria, while compounds **6i** and **6h** were specific in the effect against the Gram-negative and Gram-positive bacteria, respectively. Table 1 also showed that compounds **4f** and **4l** showed activity against *Streptococcus mutans*, while compounds **4m** and **5d** showed activity against *Staphylococcus aureus*.

A lowest concentration of chemical compound that inhibits the bacterial growth is the minimum inhibitory concentration (MIC, $\mu\text{g/mL}$). The MIC of compound **4k** that inhibited all the tested pathogenic bacteria was 31.25 $\mu\text{g/mL}$, while the MIC of compound **4f** that inhibited the *Streptococcus mutans* was 62.5 $\mu\text{g/mL}$, the MIC of compound **4l** that inhibited *Streptococcus mutans* was 100 $\mu\text{g/mL}$, and the MIC of compound **4m** that inhibited *Staphylococcus aureus* was 62.50 $\mu\text{g/mL}$. The MIC of compound **5d** that inhibited *Staphylococcus aureus* was 31.25 $\mu\text{g/mL}$, the MIC of compound **5g** that inhibited *Staphylococcus aureus* and *Salmonella typhimurium* was 100 $\mu\text{g/mL}$, the MIC of compound **6i** that inhibited *Escherichia coli* and *Salmonella typhimurium* was 62.50 $\mu\text{g/mL}$, and the MIC of compound **6h** that inhibited *Streptococcus mutans* and *Staphylococcus aureus* was 31.25 and 62.50 $\mu\text{g/mL}$, respectively (Table 2).

TABLE 1: Bioactivities of the synthetic organic compounds against some pathogenic bacteria and unicellular fungus *Candida albicans* at 100 μ g/paper disc (6 mm).

Compd. number	<i>Streptococcus mutans</i>	<i>Staphylococcus aureus</i>	<i>Escherichia coli</i>	<i>Salmonella typhimurium</i>	<i>Candida albicans</i>
Control	–	–	–	–	–
4k	+	+	+	+	–
4j	–	–	–	–	–
4f	+	–	–	–	–
4e	–	–	–	–	–
4i	–	–	–	–	–
4l	+	–	–	–	–
4m	–	+	–	–	–
5c	–	–	–	–	–
5d	–	+	–	–	–
5e	–	–	–	–	–
5f	–	–	–	–	–
5g	–	+	–	+	–
5i	–	–	–	–	–
6d	–	–	–	–	–
6f	–	–	–	–	–
6g	–	–	–	–	–
6i	–	–	+	+	–
6h	+	+	–	–	–
8c	–	–	–	–	–
8d	–	–	–	–	–
8f	–	–	–	–	–
8g	–	–	–	–	–

TABLE 2: Minimum inhibitory concentration (MIC, μ g/mL) of bioactive synthetic compounds against target pathogenic bacteria.

Compd. number	<i>Streptococcus mutans</i> wild strain	<i>Staphylococcus aureus</i>	<i>Escherichia coli</i>	<i>Salmonella typhimurium</i>
4k	31.25	31.25	31.25	31.25
4f	62.50	–	–	–
4l	100.00	–	–	–
4m	–	62.50	–	–
5d	–	31.25	–	–
5g	–	100.00	–	100.00
6i	–	–	62.50	62.50
6h	31.25	62.50	–	–

It is obvious from Table 1 that small structural variations in s-triazine ring may induce an effect on antibacterial activity, such as in series of benzyl derivatives (**4** series) and the *N*-methylbenzyl derivatives (**5** series); the benzyl derivatives showed more activity than the *N*-benzyl derivatives. The same was observed when the benzyl was substituted by methoxy (**8** series); the benzyl derivatives showed more activity. Also the substituent effect on the benzylidene moiety has a great effect on the antibacterial activity as shown in case of **4** series and **5** series, and it is interesting to note that in most of the cases the fluorine derivatives showed higher activity than the chlorine and bromine derivatives as in case of **5** series. On the other hand, not all the tested compounds showed any antifungal activity; this observation agreed with the reported data for s-triazine derivatives [48, 49].

4. Conclusion

The present work represents the synthesis, characterization, and biological activity of novel series of the s-triazine *bis*-Schiff base. Only twenty-two compounds from the above series **4a–m**, **5a–i**, **6a–i**, **7a–i**, and **8a–g** were tested against Gram-positive and Gram-negative bacteria according to their solubility in aqueous DMSO. Only eight compounds **4k**, **4f**, **4l**, **4m**, **5d**, **5g**, **6i**, and **6h** showed bioactivity against the target pathogenic bacteria as antimicrobial agents and did not show antagonistic effect against fungus *Candida albicans*. Two compounds **4k** and **5g** have wide-range effect presently in Gram-positive and Gram-negative bacteria, while compounds **6i** and **6h** have specific effect against the Gram-negative and Gram-positive bacteria, respectively.

In addition, compounds **4f** and **4l** showed activity against *Streptococcus mutans* while compounds **4m** and **5d** showed activity against *Staphylococcus aureus*.

It is clear that the substituent in the benzylidene as well as the triazine ring may have great effect on the antimicrobial activity. Further investigation is being run in our lab to get a clear depiction on the mode of action and the relation between the biological activity and substituent effect on the s-triazine moiety.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for funding this work through Research Group no. RGP-234, Saudi Arabia.

Supplementary Materials

Figures S1–S52: ^1H NMR, ^{13}C NMR, and LC-MS spectra. (Supplementary Materials)

References

- [1] A. Sousa-Pedrares, N. Camiña, J. Romero, M. L. Durán, J. A. García-Vázquez, and A. Sousa, "Electrochemical synthesis and crystal structure of cobalt(II), nickel(II), copper(II), zinc(II) and cadmium(II) complexes with 2-pyridinecarbaldehyde-(2I-aminosulfonylbenzoyl)hydrazone," *Polyhedron*, vol. 27, no. 16, pp. 3391–3397, 2008.
- [2] M. Sönmez, M. Çelebi, and I. Berber, "Synthesis, spectroscopic and biological studies on the new symmetric Schiff base derived from 2,6-diformyl-4-methylphenol with N-aminopyrimidine," *European Journal of Medicinal Chemistry*, vol. 45, no. 5, pp. 1935–1940, 2010.
- [3] S.-G. Teoh, G.-Y. Yeap, C.-C. Loh, L.-W. Foong, S.-B. Teo, and H.-K. Fun, "Inner coordination sphere tin(IV) complexes with some O,N,N-terdentate N-(2-hydroxybenzaldehyde)-1-amino-2-phenyleneimine and N-(2-hydroxy-1-naphthaldehyde)-1-amino-2-phenyleneimine and O,N,N,O-quadridentate N,N'-bis(2-hydroxybenzaldehyde)-1,2-phenylenediimine and N,N'-bis(2-hydroxy-1-naphthaldehyde)-1,2-phenylenediimine Schiff bases," *Polyhedron*, vol. 16, no. 13, pp. 2213–2221, 1997.
- [4] G. P. Pokhariyal, B. K. Lal, and V. Rastogi, "Synthesis, spectral and thermal studies of some divalent metal complexes with 8-acetyl-7-hydroxy-4-methyl coumarin oxime-hdrazone," *Asian Journal of Chemistry*, vol. 9, pp. 541–543, 1997.
- [5] P. Barbazán, R. Carballo, B. Covelo, C. Lodeiro, J. C. Lima, and E. M. Vázquez-López, "Synthesis, characterization, and photo-physical properties of 2-hydroxybenzaldehyde [(1E)-1-pyridin-2-ylethylidene]hydrazone and its rhenium(I) complexes," *European Journal of Inorganic Chemistry*, no. 17, pp. 2713–2720, 2008.
- [6] S. Banerjee, S. Mondal, W. Chakraborty et al., "Interaction of boric acid with salicyl derivatives as an anchor group of boron-selective adsorbents," *European Journal of Organic Chemistry*, pp. 2785–2790, 2008.
- [7] N. Ghavtadze, R. Fröhlich, and E.-U. Würthwein, "2H-pyrrole derivatives from an aza-nazarov reaction cascade involving indole as the neutral leaving group," *European Journal of Organic Chemistry*, no. 21, pp. 3656–3667, 2008.
- [8] J. P. Cornelissen, J. H. Van Diemen, L. R. Groeneveld, J. G. Haasnoot, A. L. Spek, and J. Reedijk, "Synthesis and properties of isostructural transition-metal (copper, nickel, cobalt, and iron) compounds with 7,7',8,8'-tetracyanoquinodimethanide(1-) in an unusual monodentate coordination mode. Crystal structure of bis(3,5-bis(pyridin-2-yl)-4-amino-1,2,4-triazole)bis(7,7',8,8'-tetracyanoquinodimethanido)copper(II)," *Inorganic Chemistry*, vol. 31, no. 2, pp. 198–202, 1992.
- [9] F. Kandil, M. K. Chebani, and W. Al Zoubi, "Synthesis of Macrocyclic Bis-Hydrazone and Their Use in Metal Cations Extraction," *ISRN Organic Chemistry*, vol. 2012, pp. 1–8, 2012.
- [10] N. Nawar, M. A. Khattab, and N. M. Hosny, "Some metal(II) complexes of o-aminoacetophenone benzoylhydrazone (AABH): Their preparation, characterization and antimicrobial activity," *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*, vol. 29, no. 8, pp. 1365–1384, 1999.
- [11] M.-D. Gui, R.-H. Zhang, H. Yan, H.-M. Wang, X.-B. Chen, and S.-J. Zhu, "Synthesis and magnetic and catalytic properties of manganous isonicotinoyl hydrazone complexes," *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*, vol. 28, no. 8, pp. 1381–1391, 1998.
- [12] Z. H. Chohan and A. Rauf, "Some biologically active mixed ligand complexes of Co(II), Cu(II) and Ni(II) with ONO, NNO and SNO donor nicotinoylhydrazine-derived ligands," *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*, vol. 26, no. 4, pp. 591–604, 1996.
- [13] S. K. Agarwal, J. Jain, and S. Chand, "Synthetic and antimicrobial studies of hexacoordinated ternary complexes of Mn(II) and Cu(II)," *Asian Journal of Chemistry*, vol. 14, no. 1, pp. 489–492, 2002.
- [14] X. Zhu, C. Wang, W. Dong et al., "Synthesis, structure, and biological activities of the O,O'-diethylphosphorohydrazonothionate ligand and complexes with nickel(II), copper(II), and zinc(II)," *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*, vol. 32, no. 3, pp. 475–487, 2002.
- [15] K. A. El-Manakhly, H. A. Bayoumi, M. M. Ezz Eldin, and H. A. Hammad, "Physicochemical and antimicrobial studies of some metal complexes," *Journal of the Indian Dental Association*, vol. 76, no. 2, pp. 63–64, 1999.
- [16] K. P. Deepa, K. K. Aravindakshan, and F. Suhara, "Synthesis, characterization and antifungal studies of metal complexes of benzoyl- and salicylhydrazones of N-methylacetoacetanilide," *Asian Journal of Chemistry*, vol. 13, no. 2, pp. 513–523, 2001.
- [17] A. K. Panda, D. C. Dash, P. Mishra, and H. Mohanty, "Metallo ligands: Part V - Homo and bimetallic trinuclear complexes of copper(II), nickel(II) and cobalt(II) with 1,8-di-(2I-hydroxyphenyl)-4,5-diphenyl-2,3,6,7-tetrazaoctan-1,3,5,7-tetraene," *Indian Journal of Chemistry - Section A Inorganic, Physical, Theoretical and Analytical Chemistry*, vol. 35, no. 10, pp. 848–851, 1996.
- [18] M. F. Iskander, L. El-Sayed, N. M. H. Salem, W. Haase, H. J. Linder, and S. Foro, "Synthesis, characterization and magnetochemical studies of some copper(II) complexes derived from n-salicylidene-n-alkanoylhydrazins: X-ray crystal and molecular structure of bis [monochloro-(μ-n-salicylidenemyristoylhydrazine) ono(-1)] dicopper(II)," *Polyhedron*, vol. 23, no. 1, pp. 23–31, 2004.

- [19] C. Li, L. Wang, X. Meng, and H. Zhao, "Synthesis, characterization and antitumor activity of benzaldehyde nitrogen mustard picolinoyl hydrazone complexes," *Transition Metal Chemistry*, vol. 24, no. 2, pp. 206–209, 1999.
- [20] Q. Zhang, C. He, P. Yin, and J. M. Shreeve, "Insensitive nitrogen-rich materials incorporating the nitroguanidyl functionality," *Chemistry - An Asian Journal*, vol. 9, no. 1, pp. 212–217, 2014.
- [21] A. El-Faham, S. M. Osman, H. A. Al-Lohedan, and G. A. El-Mahdy, "Hydrazino-methoxy-1,3,5-triazine derivatives' excellent corrosion organic inhibitors of steel in acidic chloride solution," *Molecules*, vol. 21, no. 6, article no. 714, 2016.
- [22] G. A. El-Mahdy, H. H. Al-Rasheed, M. A. Alshaikh, H. A. Al-Lohedan, and A. El-Faham, "2,4-Dihydrazino-6-Morpholino-1,3,5-Triazine (DHMT) and 2,4-Dihydrazino-6-Piperidino-1,3,5-Triazine (DHPT) as promising corrosion inhibitors of steel in acidic media," *International Journal of Electrochemical Science*, vol. 11, no. 7, pp. 5459–5472, 2016.
- [23] P. Gamez and J. Reedijk, "1,3,5-Triazine-based synthons in supramolecular chemistry," *European Journal of Inorganic Chemistry*, no. 1, pp. 29–42, 2006.
- [24] S. M. Soliman, A. El-Faham, and J. H. Albering, "Synthesis, X-ray crystal structure and DFT studies of two octahedral cobalt(II) complexes with N,N,N-tridentate triazine-type ligand," *Journal of Coordination Chemistry*, vol. 70, no. 13, pp. 2261–2279, 2017.
- [25] S. Demeshko, S. Dechert, and F. Meyer, "Anion- π Interactions in a Carousel Copper(II)-Triazine Complex," *Journal of the American Chemical Society*, vol. 126, no. 14, pp. 4508–4509, 2004.
- [26] J. A. Zerkowski, C. T. Seto, and G. M. Whitesides, "Solid-state structures of 'rosette' and 'crinkled tape' motifs derived from the cyanuric acid-melamine lattice," *Journal of the American Chemical Society*, vol. 114, no. 13, pp. 5473–5475, 1992.
- [27] J. Ramírez, A.-M. Stadler, J. M. Harrowfield et al., "Coordination architectures of large heavy metal cations (Hg_2^+ and Pb_2^+) with bis-tridentate ligands: Solution and solid-state studies," *Zeitschrift für anorganische und allgemeine Chemie*, vol. 633, no. 13–14, pp. 2435–2444, 2007.
- [28] J. Ramírez, A.-M. Stadler, L. Brelot, and J.-M. Lehn, "Coordinative, conformational and motional behaviour of triazine-based ligand strands on binding of $\text{Pb}(\text{II})$ cations," *Tetrahedron*, vol. 64, no. 36, pp. 8402–8410, 2008.
- [29] A. Das, S. Demeshko, S. Dechert, and F. Meyer, "A new triazine-based tricompartamental ligand for stepwise assembly of mononuclear, dinuclear, and 1D-polymeric heptacoordinate manganese(II)/azido complexes," *European Journal of Inorganic Chemistry*, no. 8, pp. 1240–1248, 2011.
- [30] F. Yaoting, L. Gang, L. Zifeng, H. Hongwei, and M. Hairong, "Synthesis, structure and third-order nonlinear optical properties of 1,3,5-triazine-based $\text{Zn}(\text{II})$ three-dimensional supramolecule," *Journal of Molecular Structure*, vol. 693, no. 1–3, pp. 217–224, 2004.
- [31] L. Pavelek, V. Ladányi, M. Nečas, S. Vallová, and K. Wichterle, "Dioxouranium complexes with pentadentate s-triazine Schiff base ligands: Synthesis, crystal structure and optical properties," *Polyhedron*, vol. 107, pp. 89–96, 2016.
- [32] L. Pavelek, V. Ladányi, M. Nečas, Z. Moravec, and K. Wichterle, "Synthesis and characterization of lanthanide complexes with a pentadentate triazine-based ligand," *Polyhedron*, vol. 119, pp. 134–141, 2016.
- [33] L. Tei, Z. Baranyai, E. Brücher et al., "Dramatic increase of selectivity for heavy lanthanide(III) cations by tuning the flexibility of polydentate chelators," *Inorganic Chemistry*, vol. 49, no. 2, pp. 616–625, 2010.
- [34] F. W. Lewis, L. M. Harwood, M. J. Hudson et al., "Highly efficient separation of actinides from lanthanides by a phenanthroline-derived bis-triazine ligand," *Journal of the American Chemical Society*, vol. 133, no. 33, pp. 13093–13102, 2011.
- [35] M. J. Hudson, L. M. Harwood, D. M. Laventine, and F. W. Lewis, "Use of soft heterocyclic N-donor ligands to separate actinides and lanthanides," *Inorganic Chemistry*, vol. 52, no. 7, pp. 3414–3428, 2013.
- [36] M. G. B. Drew, C. Hill, M. J. Hudson, P. B. Iveson, C. Madic, and T. G. A. Youngs, "Solvent extraction and lanthanide complexation studies with new terdentate ligands containing two 1,3,5-triazine moieties," *Dalton Transactions*, no. 2, pp. 244–251, 2004.
- [37] T. Vilaivan, N. Saesaengseerung, D. Jarprung, S. Kamchonwongpaisan, W. Sirawaraporn, and Y. Yuthavong, "Synthesis of solution-phase combinatorial library of 4,6-diamino-1,2-dihydro-1,3,5-triazine and identification of new leads against $\text{Al6V}+\text{S108T}$ mutant dihydrofolate reductase of *Plasmodium falciparum*," *Bioorganic & Medicinal Chemistry*, vol. 11, no. 2, pp. 217–224, 2003.
- [38] J. N. Sangshetti and D. B. Shinde, "One pot synthesis and SAR of some novel 3-substituted 5,6-diphenyl-1,2,4-triazines as antifungal agents," *Bioorganic & Medicinal Chemistry Letters*, vol. 20, no. 2, pp. 742–745, 2010.
- [39] W. Lv, B. Banerjee, K. L. Molland et al., "Synthesis of 3-(3-aryl-pyrrolidin-1-yl)-5-aryl-1,2,4-triazines that have antibacterial activity and also inhibit inorganic pyrophosphatase," *Bioorganic & Medicinal Chemistry*, vol. 22, no. 1, pp. 406–418, 2014.
- [40] K. K. Bansal, D. Kakde, U. Gupta, and N. K. Jain, "Development and characterization of triazine based dendrimers for delivery of antitumor agent," *Journal of Nanoscience and Nanotechnology*, vol. 10, no. 12, pp. 8395–8404, 2010.
- [41] N. C. Desai, A. H. Makwana, and R. D. Senta, "Synthesis, characterization and antimicrobial activity of some novel 4-(4-(arylamino)-6-(piperidin-1-yl)-1,3,5-triazine-2-ylamino)-N-(pyrimidin-2-yl)benzenesulfonamides," *Journal of Saudi Chemical Society*, vol. 20, no. 6, pp. 686–694, 2016.
- [42] G. Blotny, "Recent applications of 2,4,6-trichloro-1,3,5-triazine and its derivatives in organic synthesis," *Tetrahedron*, vol. 62, no. 41, pp. 9507–9522, 2006.
- [43] C. A. M. Afonso, N. M. T. Lourenço, and A. A. de Rosatella, "Synthesis of 2,4,6-tri-substituted-1,3,5-triazines," *Molecules*, vol. 11, no. 1, pp. 81–102, 2006.
- [44] V. V. Bakharev, A. A. Gidasov, V. E. Parfenov et al., "Synthesis of 4-amino-6-chloro-1,3,5-triazin-2(1H)-ones," *Russian Chemical Bulletin*, vol. 61, no. 1, pp. 99–112, 2012.
- [45] W. Huang, W. Zheng, D. J. Urban et al., "N4-Phenyl modifications of N2-(2-hydroxy)ethyl-6-(pyrrolidin-1-yl)-1,3,5-triazine-2,4-diamines enhance glucocerebrosidase inhibition by small molecules with potential as chemical chaperones for Gaucher disease," *Bioorganic & Medicinal Chemistry Letters*, vol. 17, no. 21, pp. 5783–5789, 2007.
- [46] D. A. Vanden and A. J. Berghe, "Screening methods for antibacterial and antiviral agents from higher plants," in *Methods in Plant Biochemistry*, P. M. Dey and J. D. Harbone, Eds., pp. 47–69, Academic Press, London, 1991.
- [47] S. T. Asundaria, S. A. Patel, K. M. Mehta, and K. C. Patel, "Synthesis, characterization and antimicrobial studies of some novel 1,3,4-thiadiazolium-2-thiolate derivatives," *South African Journal of Chemistry*, vol. 63, pp. 141–144, 2015.

- [48] K. Srinivas, U. Srinivas, K. Bhanuprakash, K. Harakishore, U. S. N. Murthy, and V. J. Rao, "Synthesis and antibacterial activity of various substituted s-triazines?" *European Journal of Medicinal Chemistry*, vol. 41, pp. 1240–1246, 2006.
- [49] H. H. Al-Rasheed, M. Al Alshaikh, J. M. Khaled, N. S. Alharbi, and A. El-Faham, "Ultrasonic irradiation: synthesis, characterization, and preliminary antimicrobial activity of novel series of 4,6-disubstituted-1,3,5-triazine containing hydrazone derivatives," *Journal of Chemistry*, vol. 2016, Article ID 3464758, 9 pages, 2016.