Research Article

Evaluation of Glycemic and Lipid Profile of Offspring of Diabetic Wistar Rats Treated with *Malpighia emarginata* Juice

Sandra M. Barbalho,¹ Débora C. Damasceno,² Ana Paula Machado Spada,³ Miréia Palhares,⁴ Karla Aparecida Martuchi,⁵ Marie Oshiiwa,⁶ Viviane Sazaki,⁴ and Vanessa Sellis da Silva⁴

¹ Medical School of Medicine of Marília (UNIMAR), Faculty of Technology and Foods of Marília, Avenida Higino Muzzi Filho, 1001, Marília, SP 17525902, Brazil

² Laboratory of Experimental Research on Gynecology and Obstetrics, Botucatu Medical School, UNESP, Distrito Rubião Junior, S/N, Botucatu, SP 18618000, Brazil

³ Biochemistry Department, Faculty at UNINOVE and SENAC, USP/São Paulo, SP 01041000, Brazil

⁴ Nutrition Department, Methodist University of Piracicaba, UNIMEP, Rua Tenente Florencio Pupo Neto, 300, Lins, SP 16400680, Brazil

⁵ Nutrition Department, Frigoríficos Bertin S/A, Rodovia de Acesso Lins-Getulina s/n, Parque Industrial, Lins, SP 16400000, Brazil

⁶ Faculty of Technology and Foods of Marília, FATEC, Avenida Castro Alves, 62, Marília, SP 17506000, Brazil

Correspondence should be addressed to Sandra M. Barbalho, smbarbalho@terra.com.br

Received 10 August 2010; Revised 6 December 2010; Accepted 3 January 2011

Academic Editor: I. Raz

Copyright © 2011 Sandra M. Barbalho et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Knowing that maternal diabetes is related to hyperglycemia and fetal hyperinsulinemia, which affect the lipid metabolism, the aim of this study was to evaluate the effects of *Malpighia emarginata* (acerola) juice on the glycemic and lipid profile of offspring of diabetic and nondiabetic Wistar rats. The adult offspring of non-diabetic dams and of dams with severe streptozotocin-induced diabetes were divided into groups: G1, offspring (of control dams) treated with water, G2, offspring (of diabetic dams) treated with water, G3, male offspring (of control dams) treated with acerola juice, and G4, male offspring (of diabetic dams) treated with acerola juice. The offspring of diabetic dams treated with acerola juice showed significantly decreased levels of glucose, cholesterol, triglycerides, and increased HDL-c. The use of acerola juice is a potential strategy to aid in the prevention of DM and dyslipidemia and its complications or to act as an auxiliary in the treatment of these diseases.

1. Introduction

Diabetes mellitus is a disease characterized by chronic hyperglycemia because of a total or relative lack of insulin [1].

Among the many groups of people who suffer from this disease are gestating women. Gestational diabetes leads to modifications in the metabolism of the mother and her offspring caused by the mother's hyperinsulinemia and hyperglycemia. These conditions affect the fetal metabolism during the gestational period and extend throughout the life of the offspring [2, 3].

Maternal hyperinsulinemia and hyperglycemia give rise to complications such as macrosomia, spontaneous

termination of pregnancy, premature birth, and metabolic and respiratory complications in the newborn [4, 5].

The presence of obesity and diabetes is more frequent in childhood and adolescence when there is a history of maternal or gestational diabetes. In these conditions, metabolic disorders can affect the growth and metabolism of descendants and subsequent generations [6, 7].

There are many allopathic medications for controlling glycemia and dyslipidemia but they are usually expensive and have been implicated in congenital anomalies [8]. Due to the high cost of medications, pregnant women turn to alternative remedies to treat diabetes and other diseases, motivating the search for alternatives that are normally easier

to find and inexpensive. In view of this tendency, there is a need for studies to determine the effects of medicinal plants and phytotherapeutic products and to establish their ideal therapeutic schemes to take advantage of their benefits and reduce the occurrence of adverse effects. In Brazil and many parts of the world, medicinal plants are sold in open street markets and grown in the backyards of private homes [9, 10].

The acerola tree (*Malpighia emarginata*) is a very popular plant in South America, and its fruit is known to be rich in ascorbic acid and polyphenols. Hanamura et al. [11] studied the antihyperglycemic effects of polyphenols from acerola fruit and found positive effects. However, there are no reports in the literature about the effects of *Malpighia emarginata* on the glycemic and lipid profile in the offspring of streptozotocin-induced diabetic Wistar rats.

Knowing that maternal diabetes is related to hyperglycemia and fetal hyperinsulinemia, which affect the lipid metabolism, the aim of this study was to evaluate the effects of *Malpighia emarginata* (acerola) juice on the glycemic and lipid profile of the offspring of diabetic and nondiabetic Wistar rats.

2. Material and Methods

2.1. Parental Generation. The animals included in this study were treated according to the “Guide to the care and use of experimental animals”, which delineates the principles by the Canadian Council for the care to laboratory animals. The study was initiated after its approval by the Ethics Committee under registration number 2500000764/2007-47. Wistar rats were kept in the vivarium of our university under controlled conditions (12/12-hour light/dark cycle, ambient temperature of $22 \pm 2^\circ\text{C}$, relative humidity of $60 \pm 5\%$, and water and chow ad libitum).

2.2. Diabetes Induction. Nondiabetic female rats ($n = 20$) weighing approximately 250 g underwent a seven-day adaptation period in the room where the experiment was designed. After such period, randomly selected rats received the intravenous administration (caudal vein) of 40 mg of streptozotocin/Kg (STZ—SIGMA Chemical Company, St. Louis, MO, USA) diluted in citrate buffer (0.1 M; pH 4.5). Nondiabetic animals received a similar volume of citrate buffer by the same route of administration of the diabetogenic drug [12]. For criteria of inclusion, the rats given citrate buffer and presenting with glycemia $<120\text{ mg/dL}$ at day 0 of pregnancy were included in the nondiabetic groups. Rats from the STZ group presenting with glycemia 300 mg/dL at day 0 of pregnancy were included in the diabetic groups.

2.3. Mating Period. All female rats were mated overnight to nondiabetic male rats untreated with acerola juice. The mornings that spermatozoa were found in the vaginal smear were designated gestational day 0. The mating procedure continued for 15 consecutive days, which comprises approximately three oestral cycles, and until replicate numbers were obtained for each group ($n = 15$ animals/group). Nonmated female rats in this period were considered infertile and discounted from the study [13].

2.4. Pregnancy Period. Glycaemic level (food ad libitum overnight) was monitored during pregnancy in the morning using the glucose oxidase method [13] at days 0, 7, 14, and 21 of pregnancy in all experimental groups. Approximately at the day 21 of pregnancy, after birth of the offspring, the dams remained in individual cages with their offspring until weaning (21 days). Food intake was addressed every day and body weight once a week.

2.5. Offspring. After weaning, the offspring (60 males) were kept in collective cages until they reached adulthood. The adult animals, weighing approximately 250 g, were divided into 4 experimental groups: G1 ($n = 15$), male offspring (from control dams) treated with vehicle (water), G2 ($n = 15$), male offspring (from diabetic dams) treated with vehicle (water), G3 ($n = 15$), male offspring (from control dams) treated with *Malpighia emarginata* (acerola) juice, and G4 ($n = 15$), male offspring (from diabetic dams) treated with *M. emarginata* (acerola) juice.

The animals from groups G3 and G4 received *Malpighia emarginata* (acerola) juice at a dose of 0.58 g/kg once a day (at early morning) for 30 consecutive days. The dose administered to the animals was based on 200 g/L, which corresponds to the daily intake of 200 mL of juice by an adult man weighing 70.0 Kg (this consumption was based on popular consultation to prepare the juice).

Food intake was addressed every day and body weight once a week.

2.6. Acerola Juice Preparation. Acerola fruits (obtained from Cooperativa Agrícola de Lins-SP (COALINS)) were washed, weighed (200 g/L), and triturated with water in a blender for 2 minutes. The juice was filtered and administered daily. It was prepared every day 15 minutes before its administration.

2.7. Blood Collection and Biochemical Profile Determination. At the end of the treatment (30 days), the offspring from diabetic and non diabetic dams (60 male rats) were anesthetized with sodium pentobarbital (150 mg/kg) and killed. Blood samples were collected in order to determine the biochemical profile (total cholesterol, HDL-c, LDL-c, triglycerides, and glucose). Tests were performed according to the methodology proposed by commercial kits: LABTEST (Lagoa Santa, Belo Horizonte, MG) for glycemia, total cholesterol and HDL-c, and triglycerides and WIENER LAB (São Paulo, SP) for LDL-c. The results were interpreted according to the criteria established by the American Diabetes Association [1].

2.8. Statistical Analysis. Data analysis was performed by using Students *t*-test, and the level of significance adopted was 5%.

3. Results

The following 4 tables show the results of the biochemical profile of the offspring from nondiabetic and diabetic dams.

TABLE 1: Biochemical profile of offspring treated with water (from nondiabetic dams: G1 and from diabetic dams: G2).

	Groups		P value
	G1	G2	
Glucose	67.7 ± 11.9	88.9 ± 11.6	.0002*
Triglycerides	119.9 ± 72.4	139.0 ± 92.5	.0000*
Cholesterol	76.8 ± 67.3	112.7 ± 24.7	.0131*
HDL-c	34.4 ± 3.3	30.4 ± 4.1	.0175*
LDL-c	65.0 ± 19.5	84.5 ± 37.1	.2676

Values presented as mean ± standard deviation (SD)/*significant values.

TABLE 2: Biochemical profile of offspring (from nondiabetic dams) treated with vehicle (G1) and offspring (from nondiabetic dams) treated with acerola juice (G3).

	Groups		P value
	G1	G3	
Glucose	67.7 ± 11.9	68.68 ± 23.75	.0771
Triglycerides	119.9 ± 72.4	83.00 ± 48.41	.2354
Cholesterol	76.8 ± 67.3	98.00 ± 22.70	.0610
HDL-c	34.4 ± 3.3	45.25 ± 19.98	.0000*
LDL-c	65.0 ± 19.5	57.36 ± 35.19	.0620

Values presented as mean ± standard deviation (SD)/*significant values.

TABLE 3: Biochemical profile of offspring treated with acerola juice (from nondiabetic dams: G3 and from diabetic dams: G4).

	Groups		P value
	G3	G4	
Glucose	68.68 ± 23.75	72.69 ± 16.09	.0637
Triglycerides	83.00 ± 48.41	87.47 ± 53.60	.3891
Cholesterol	98.00 ± 22.70	94.85 ± 16.73	.0604
HDL-c	45.25 ± 19.98	60.68 ± 13.43	.1047
LDL-c	57.36 ± 35.19	51.85 ± 13.6	.0556

Values presented as mean ± standard deviation (SD).

Table 1 compares G1 (offspring of nondiabetic dams treated with water) and G2 (offspring of diabetic dams treated with water). Male offspring of diabetic dams (G2) presented significantly higher levels of glycemia, triglycerides and total cholesterol and significantly lower HDL-c levels than G1.

Table 2 compares G1 (offspring of control dams), that received water, and G3 (offspring of control dams), treated with acerola juice. As can be seen, the male offspring treated with acerola juice showed significantly higher HDL-c levels.

Table 3 shows that there are no differences in the biochemical profile of G3 (nondiabetic offspring treated with acerola juice) and G4 (diabetic offspring treated with acerola juice).

A comparison of G2 (offspring of diabetic dams), treated with water, and G4 (offspring of diabetic dams), treated with acerola juice, suggests that the use of acerola juice reduced glycemia, triglycerides, total cholesterol, and LDL-c and increased HDL-c levels (Table 4).

TABLE 4: Biochemical profile of offspring treated with vehicle (G2) and offspring treated with acerola juice (G4).

	Groups		P value
	G2	G4	
Glucose	88.9 ± 11.6	72.69 ± 16.09	.0000*
Triglycerides	139.0 ± 92.5	87.47 ± 53.60	.0000*
Cholesterol	112.7 ± 24.7	94.85 ± 16.73	.0335*
HDL-c	30.4 ± 4.1	60.68 ± 13.43	.0000*
LDL-c	84.5 ± 37.1	51.85 ± 13.6	.0011*

Values presented as mean ± standard deviation (SD)/*significant values.

No differences were found in food intake and body weight among the groups at the beginning and end of the treatment (Table 5).

Table 6 shows that the food intake of nondiabetic and diabetic dams on the first day of pregnancy and on the day preceding the birth of their offspring did not differ significantly among the groups. The mothers showed a significant weight increase at the beginning and end of gestation, but no significant differences in weight were found between diabetic and nondiabetic mothers.

4. Discussion

The results of this study reveal metabolic modifications in the offspring of diabetic dams when compared to offspring of nondiabetic dams. The animals (offspring of both diabetic and nondiabetic dams) treated with acerola juice presented higher levels of HDL-c and lower values of LDL-c. Modifications in levels of blood glucose and plasma fats are risk factors for metabolic syndrome and its future complications as vascular diseases [14–18]. The alterations resulting from maternal diabetes are related to hyperglycemia and fetal hyperinsulinemia, which affect lipid and protein synthesis [2, 4]. Furthermore, maternal hyperglycemia stimulates fetal growth due to the greater availability of glucose in the blood flow and the regulation of growth factors [4, 19]. In this work, the animals in the diabetic and nondiabetic groups did not show significant differences in body weight.

The results of this work indicate that the use of acerola juice can be beneficial in preventing and helpful in treating hyperglycemia and dyslipidemias in the offspring of diabetic dams. Numerous studies in the literature demonstrate that plants can also be used for this purpose in the offspring of diabetic dams [20–22]. Many other authors have reported the beneficial effects of plants on glycemic and lipid profiles. Babiloni et al. [23] showed the antihyperglycemic activity of stem bark extract of *Vatairea macrocarpa* in the treatment of diabetic rats. Badole et al. [24] found significant antihyperglycemic effects in alloxan-induced diabetic mice treated with *Pleurotus pulmonarius* and observed that it has potent synergistic antihyperglycemic effects when used in association with glyburide. Umar et al. [25] found that *Tetracera scandens* has antidiabetic efficacy in diabetic rats. Another study [26] observed a reduction in blood glucose and in the lipid profile of streptozotocin-induced diabetic rats treated with *Cassia glauca*. Bera et al. [22] observed that

TABLE 5: Food intake values before (time 1) and after (time 2) treatment with water (G1 and G2) and acerola juice (G3 and G4).

	Groups				P value
	G1	G2	G3	G4	
Food intake (time 1)	120.0 ± 58.2	139.5 ± 15.5	115.5 ± 11.1	127.0 ± 39.1	.0732
Food intake (time 2)	112.8 ± 38.8	133.2 ± 11.6	118.7 ± 5.9	123.8 ± 27.2	.0670
P value	.0661	.1927	.0603	.1003	
Body weigh (time 1)	248.6 ± 49.9	255.5 ± 59.7	268.6 ± 61.3	282.12 ± 56.6	.0501
Body weigh (time 2)	305.5 ± 89.2	326.2 ± 48.2	329.6 ± 61.9	338.26 ± 76.2	.0578
P value	.0781	.0689	.0789	.0798	

Values presented as mean ± standard deviation (SD).

TABLE 6: Food intake values on the first day of pregnancy (time 1) and on the day preceding the birth of their offspring (time 2) in the non diabetic (NDM) and in diabetic dams (DM).

	Groups		P value
	Dam NDM	Dam DM	
Food intake (time1)	120.0 ± 37.1	139.5 ± 25.3	.0732
Food intake (time 2)	142.8 ± 39.4	153.2 ± 59.5	.0670
P value	.0661	.1927	
Body weigh (time 1)	306.6 ± 59.4	280.5 ± 39.7	.0652
Body weigh (time 2)	429.5 ± 45.7	430.9 ± 54.2	.0668
P value	.0381*	.0299*	

Values presented as mean ± standard deviation (SD)/* significant values.

the effects of a polyherbal formulation (of eight medicinal plants) for the management of streptozotocin-induced diabetes in rats were comparable to those of the antidiabetic drug (Glibenclamide). Decreased lipids and glycemia were reported by Daisy et al. [27] using *Hunteria umbellata* and by Adeneye and Adeyemi [28] using isolated components of *Gymnema sylvestre*. Many other studies have demonstrated the beneficial effects of plants such as *Cynodon dactylon*, *Coriandrum sativum*, *Mentha piperita*, and *Amorphophallus konjac* on the glycemic and lipid profiles of diabetic rats (streptozotocin or alloxan-induced) [29–32]. Lin et al. [33] also showed that *Siraitia grosvenorii* polysaccharides can be helpful in reducing glucose and lipid levels in rabbits.

Acerola fruit is rich in vitamin C, calcium, iron, and polyphenols such as anthocyanins (cyanidin-3- α -O-rhamnoside and pelargonidin-3- α -O-rhamnoside), quercetin (quercetin-3- α -O-rhamnoside), chlorogenic and caffeic acids, and kaempferol [11, 34–36]. Kawaguchi et al. [37] described the flavonoid leucocyanidin-3-O-beta-D-glucoside and gave it the trivial name “aceronidin”, which possesses stronger antioxidative activity than alphatocopherol. Acerola also contains carotenoids that associated with bioflavonoids provide important nutrients and have a potential use as antioxidant. Brazil’s climate and soil are very favorable for cultivating acerola, making this country the world’s main producer of this fruit. Acerola is commercialized in different forms such as juices, jams, ice-creams, sweets, and liqueurs [38]. The presence of vitamins and antioxidants is known to reduce blood glucose, cholesterol, triglycerides, and LDL-c levels, and to elevate HDL-c levels

[39–41]. Bhandari et al. [20] studied the antioxidant effects of *Embelia ribes* on streptozotocin-induced diabetes in Wistar rats and found that it has important antioxidant activity in combating free radicals produced in hyperglycemic conditions, thus protecting against loss of beta cells.

Flavonoids may act in distinct ways on various plasma lipid components [42], but the literature still does not explain the exact mechanisms whereby phenolic acids, flavonoids, and terpenoids exert these beneficial effects. The presence of diabetes may hasten the development of endothelial lesions and atherosclerosis, which may be minimized by the presence of antioxidants [43, 44].

Polyphenolic substances extracted from acerola can have inhibitory effects on maltase and α -glucosidase and reduce glucose absorption, which ends up in reducing glycemia [11]. Polyphenols also participate in preventing the formation of AGEs (advanced glycation products) in diabetics, which are implicated in many micro and macrovascular complications of diabetes mellitus [45]. The antioxidant properties of polyphenols prevent oxidative stress caused by the scavenging activity of free radicals and also show anti-inflammatory activities [46–48]. The results of this work indicate that acerola juice can be helpful in preventing future damage caused by diabetes and its complications, such as vascular diseases, which, worldwide, are among the main causes of death.

5. Conclusion

The use of acerola juice is a potential strategy to aid in the prevention of DM and dyslipidemia and its complications or to act as an auxiliary in the treatment of these diseases.

Although the results of the use of acerola juice are promising, further studies are essential to evaluate its effects on human beings and to determine the ideal dosages to be used.

References

- [1] American Diabetes Association (ADA), 2010, http://care.diabetesjournals.org/content/33/Supplement_1/S3.full.pdf+html.
- [2] P. D. C. Padilha, C. Saunders, R. C. M. Machado et al., “Association between pre-gestational nutritional status and prediction of the risk of adverse pregnancy outcome,” *Revista Brasileira de Ginecologia e Obstetricia*, vol. 29, no. 10, pp. 511–518, 2007.

- [3] L. A. George, A. B. Uthlaut, N. M. Long et al., "Different levels of overnutrition and weight gain during pregnancy have differential effects on fetal growth and organ development," *Reproductive Biology and Endocrinology*, vol. 8, article 75, 2010.
- [4] A. Maayan-Metzger, D. Lubin, and J. Kuint, "Hypoglycemia rates in the first days of life among term infants born to diabetic mothers," *Neonatology*, vol. 96, no. 2, pp. 80–85, 2009.
- [5] C. Y. Li, F. C. Sung, P. C. Hsieh, M. D. Lee, T. H. Lu, and H. F. Chen, "Offspring birth 8 weight and risk of mortality from diabetes in mothers," *Journal of Epidemiology & Community Health*. In press.
- [6] E. F. Buzinaro, C. B. Berchieri, A. L. M. Haddad, C. R. Padovani, and W. De Paula Pimenta, "Overweight in adolescent offspring of women with hyperglycemia during pregnancy," *Arquivos Brasileiros de Endocrinologia e Metabologia*, vol. 52, no. 1, pp. 85–92, 2008.
- [7] U. Simeoni and D. J. Barker, "Offspring of diabetic pregnancy: long-term outcomes," *Seminars in Fetal and Neonatal Medicine*, vol. 14, no. 2, pp. 119–124, 2009.
- [8] T. D. R. Brandolt, C. C. Ferrão, S. M. N. Nunes, and G. M. B. Silva, "Effects of the extract of the *Plectranthus barbatus* (Andr.) Benth in the reproductive performance of *Rattus norvegicus*," *Biotemas*, vol. 20, no. 2, pp. 49–58, 2007.
- [9] R. Ananthan, M. Latha, K. M. Ramkumar, L. Pari, C. Baskar, and V. Narmatha Bai, "Effect of *Gymnema montanum* leaves on serum and tissue lipids in alloxan diabetic rats," *Experimental Diabetes Research*, vol. 4, no. 3, pp. 183–189, 2003.
- [10] S. Dev, "Impact of natural products in modern drug development," *Indian Journal of Experimental Biology*, vol. 48, no. 3, pp. 191–198, 2010.
- [11] T. Hanamura, C. Mayama, H. Aoki, Y. Hirayama, and M. Shimizu, "Antihyperglycemic effect of polyphenols from acerola (*Malpighia emarginata* DC.) fruit," *Bioscience, Biotechnology and Biochemistry*, vol. 70, no. 8, pp. 1813–1820, 2006.
- [12] I. M. P. Calderon, M. V. C. Rudge, and M. A. M. Brasil, "Diabetes and experimental pregnancy in rats: I. Diabetes induction, pregnancy obtention and evolution," *Acta Cirurgica Brasileira*, vol. 7, pp. 9–14, 2002.
- [13] D. C. Damasceno, W. G. Kempinas, G. T. Volpato, M. Consoni, M. V. C. Rudge, and F. J. R. Paumgartten, *Anomalias Congénitas: Estudos Experimentais*, Coopmed, Belo Horizonte, Brazil, 1st edition, 2008.
- [14] K. Paletas, E. Athanasiadou, M. Sarigianni et al., "The protective role of the mediterranean diet on the prevalence of metabolic syndrome in a population of greek obese subjects," *Journal of the American College of Nutrition*, vol. 29, no. 1, pp. 41–45, 2010.
- [15] A. Basu, M. Du, K. Sanchez et al., "Green tea minimally affects biomarkers of inflammation in obese subjects with metabolic syndrome," *Nutrition*, vol. 27, no. 2, pp. 206–213, 2011.
- [16] H. O. El Mesallamy, E. El-Demerdash, L. N. Hammad, and H. M. El Magdoub, "Effect of taurine supplementation on hyperhomocysteinemia and markers of oxidative stress in high fructose diet induced insulin resistance," *Diabetology and Metabolic Syndrome*, vol. 2, no. 1, article 46, 2010.
- [17] L. P. M. Oliveira, A. M. O. Assis, M. D. A. C. da Silva et al., "Factors associated with overweight and abdominal fat in adults in Salvador, Bahia State, Brazil," *Cadernos de Saude Publica*, vol. 25, no. 3, pp. 570–582, 2009.
- [18] E. S. Ford, C. Li, G. Zhao, W. S. Pearson, and A. H. Mokdad, "Hypertriglyceridemia and Its pharmacologic treatment among US adults," *Archives of Internal Medicine*, vol. 169, no. 6, pp. 572–578, 2009.
- [19] R. M. Y. Nomura, N. V. Costa, K. Sakamoto, C. A. Maguanha, S. Miyadahira, and M. Zugaib, "Computerized cardiotocography in pregnancies complicated by pregestational diabetes mellitus: heart rate patterns in large for gestational age fetuses," *Revista Brasileira de Ginecologia e Obstetrícia*, vol. 27, no. 12, pp. 712–718, 2005.
- [20] U. Bhandari, N. Jain, and K. K. Pillai, "Further studies on antioxidant potential and protection of pancreatic β -cells by *Embelia ribes* in experimental diabetes," *Experimental Diabetes Research*, vol. 2007, Article ID 15803, 2007.
- [21] G. T. Volpato, F. L. Vieira, D. C. Damasceno, F. L. A. Câmara, L. C. Di Stasi, and I. P. Lemonica, "Effects of leaves of *Polytmia sonchifolia* (yacon) aqueous extract in diabetic rats," *Revista Brasileira de Plantas Mediciniais*, vol. 9, no. 2, pp. 88–93, 2007.
- [22] T. K. Bera, D. De, K. Chatterjee, K. M. Ali, and D. Ghosh, "Effect of Diashis, a polyherbal formulation, in streptozotocin-induced diabetic male albino rats," *International Journal of Ayurveda Research*, vol. 1, no. 1, pp. 18–24, 2010.
- [23] P. D. Babiloni, M. P. dos Santos, G. M. Aiko et al., "Mechanism of anti-hyperglycemic action of *Vatairea macrocarpa* (Leguminosae): investigation in peripheral tissues," *Journal of Ethnopharmacology*, vol. 131, pp. 135–139, 2010.
- [24] S. L. Badole, N. M. Patel, P. A. Thakurdesai, and S. L. Bodhankar, "Interaction of aqueous extract of *Pleurotus pulmonarius* (Fr.) Quel-Champ. with glyburide in alloxan induced diabetic mice," *Evidence-based Complementary and Alternative Medicine*, vol. 5, no. 2, pp. 159–164, 2008.
- [25] A. Umar, Q. U. Ahmed, B. Y. Muhammad, B. B. S. Dogarai, and S. Z. B. T. M. Soad, "Anti-hyperglycemic activity of the leaves of *Tetracera scandens* Linn. Merr. (Dilleniaceae) in alloxan induced diabetic rats," *Journal of Ethnopharmacology*, vol. 131, no. 1, pp. 140–145, 2010.
- [26] M. D. Salahuddin, S. S. Jalalpure, and N. B. Gadge, "Antidiabetic activity of aqueous bark extract of *Cassia glauca* in streptozotocin-induced diabetic rats," *Canadian Journal of Physiology and Pharmacology*, vol. 88, no. 2, pp. 153–160, 2010.
- [27] P. Daisy, J. Eliza, and K. A. M. Mohamed Farook, "A novel dihydroxy gymnemic triacetate isolated from *Gymnema sylvestre* possessing normoglycemic and hypolipidemic activity on STZ-induced diabetic rats," *Journal of Ethnopharmacology*, vol. 126, no. 2, pp. 339–344, 2009.
- [28] A. A. Adeneye and O. O. Adeyemi, "Further evaluation of antihyperglycemic activity of *Hunteria umbellata* (K. Schum) Hallier f. seed extract in experimental diabetes," *Journal of Ethnopharmacology*, vol. 126, no. 2, pp. 238–243, 2009.
- [29] S. K. Singh, P. K. Rai, D. Jaiswal, and G. Watal, "Evidence-based critical evaluation of glycemic potential of *Cynodon dactylon*," *Evidence-Based Complementary and Alternative Medicine*, vol. 5, no. 4, pp. 415–420, 2008.
- [30] P. Dhanapakiam, J. M. Joseph, V. K. Ramaswamy, M. Moorthi, and A. S. Kumar, "The cholesterol lowering property of coriander seeds (*Coriandrum sativum*): mechanism of action," *Journal of Environmental Biology*, vol. 29, no. 1, pp. 53–56, 2008.
- [31] S. M. Barbalho, A. P. M. Spada, E. P. de Oliveira et al., "Mentha piperita effects on wistar rats plasma lipids," *Brazilian Archives of Biology and Technology*, vol. 52, no. 5, pp. 1137–1143, 2009.
- [32] M. Chua, T. C. Baldwin, T. J. Hocking, and K. Chan, "Traditional uses and potential health benefits of *Amorphophallus konjac* K. Koch ex N.E.Br," *Journal of Ethnopharmacology*, vol. 128, no. 2, pp. 268–278, 2010.

- [33] G. P. Lin, T. Jiang, X. B. Hu, X. H. Qiao, and Q. H. Tuo, "Effect of *Siraitia grosvenorii* polysaccharide on glucose and lipid of diabetic rabbits induced by feeding high fat/high sucrose chow," *Experimental diabetes research*, vol. 2007, Article ID 67435, 2007.
- [34] A. L. A. Vendramini and L. C. Trugo, "Phenolic compounds in acerola fruit (*Malpighia puniceifolia*, L.)," *Journal of the Brazilian Chemical Society*, vol. 15, no. 5, pp. 664–668, 2004.
- [35] N. Motohashi, H. Wakabayashi, T. Kurihara et al., "Biological activity of Barbados cherry (Acerola fruits, fruit of *Malpighia emarginata* DC) extracts and fractions," *Phytotherapy Research*, vol. 18, no. 3, pp. 212–223, 2004.
- [36] M. E. Schreckinger, J. Lotton, M. A. Lila, and E. G. de Mejia, "Berries from South America: a comprehensive review on chemistry, health potential, and commercialization," *Journal of medicinal food*, vol. 13, no. 2, pp. 233–246, 2010.
- [37] M. Kawaguchi, H. Tanabe, and K. Nagamine, "Isolation and characterization of a novel flavonoid possessing a 4,2''-glycosidic linkage from green mature acerola (*Malpighia emarginata* DC.) fruit," *Bioscience, Biotechnology and Biochemistry*, vol. 71, no. 5, pp. 1130–1135, 2007.
- [38] R. P. Choudhury, A. Kumar, and A. N. Garg, "Analysis of Indian mint (*Mentha spicata*) for essential, trace and toxic elements and its antioxidant behaviour," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 41, no. 3, pp. 825–832, 2006.
- [39] W. R. Leifert and M. Y. Abeywardena, "Cardioprotective actions of grape polyphenols," *Nutrition Research*, vol. 28, no. 11, pp. 729–737, 2008.
- [40] J. T. Doyama, H. G. Rodrigues, E. L. B. Novelli, E. Cereda, and W. Vilegas, "Chemical investigation and effects of the tea of *Passiflora alata* on biochemical parameters in rats," *Journal of Ethnopharmacology*, vol. 96, no. 3, pp. 371–374, 2005.
- [41] A. N. Pan, O. H. Franco, J. Ye et al., "Soy protein intake has sex-specific effects on the risk of metabolic syndrome in middle-aged and elderly Chinese," *Journal of Nutrition*, vol. 138, no. 12, pp. 2413–2421, 2008.
- [42] T. Hanamura, T. Hagiwara, and H. Kawagishi, "Structural and functional characterization of polyphenols isolated from acerola (*Malpighia emarginata* DC.) fruit," *Bioscience, Biotechnology and Biochemistry*, vol. 69, no. 2, pp. 280–286, 2005.
- [43] B. Özdemir, A. Ekbül, N. B. Topal et al., "Effects of *Origanum onites* on endothelial function and serum biochemical markers in hyperlipidaemic patients," *Journal of International Medical Research*, vol. 36, no. 6, pp. 1326–1334, 2008.
- [44] L. A. Maile, L. B. Allen, U. Veluvolu et al., "Identification of compounds that inhibit IGF-I signaling in hyperglycemia," *Experimental Diabetes Research*, vol. 2009, Article ID 267107, 2009.
- [45] S. I. Andreea, C. Marieta, and D. Anca, "AGEs and glucose levels modulate type I and III procollagen mRNA synthesis in dermal fibroblasts cells culture," *Experimental Diabetes Research*, vol. 2008, Article ID 473603, 2008.
- [46] T. Hanamura, E. Uchida, and H. Aoki, "Skin-lightening effect of a polyphenol extract from acerola (*Malpighia emarginata* DC.) fruit on UV-induced pigmentation," *Bioscience, Biotechnology and Biochemistry*, vol. 72, no. 12, pp. 3211–3218, 2008.
- [47] M. Kawaguchi, H. Tanabe, and K. Nagamine, "Isolation and characterization of a novel flavonoid possessing a 4,2''-glycosidic linkage from green mature acerola (*Malpighia emarginata* DC.) fruit," *Bioscience, Biotechnology and Biochemistry*, vol. 71, no. 5, pp. 1130–1135, 2007.
- [48] T. Hanamura and H. Aoki, "Toxicological evaluation of polyphenol extract from acerola (*Malpighia emarginata* DC.) fruit," *Journal of Food Science*, vol. 73, no. 4, pp. T55–T61, 2008.