Research Article

T-Cell Cytokine Gene Polymorphisms and Vitamin D Pathway Gene Polymorphisms in End-Stage Renal Disease due to Type 2 Diabetes Mellitus Nephropathy: Comparisons with Health Status and Other Main Causes of End-Stage Renal Disease

Alicja E. Grzegorzewska,¹ Grzegorz Ostromecki,² Paulina Zielińska,³
Adrianna Mostowska,⁴ and Paweł P. Jagodziński⁴

¹Department of Nephrology, Transplantology and Internal Diseases, Poznań University of Medical Sciences (PUMS),
49 Przybyszewskiego Boulevard, 60-355 Poznań, Poland

²DaVita Clinic Piła Dialysis Center, Wojska Polskiego 43, 64-420 Piła, Poland

³Student Nephrology Research Group, Department of Nephrology, Transplantology and Internal Diseases, PUMS,
Przybyszewskiego 49, 60-355 Poznań, Poland

⁴Department of Biochemistry and Molecular Biology, PUMS, Święcickiego 6, 60-781 Poznań, Poland

Correspondence should be addressed to Alicja E. Grzegorzewska; alicja_grzegorzewska@yahoo.com

Received 17 July 2014; Revised 22 September 2014; Accepted 22 September 2014; Published 22 December 2014

Academic Editor: Salwa Ibrahim

Copyright © 2014 Alicja E. Grzegorzewska et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. T-cell cytokine gene polymorphisms and vitamin D pathway gene polymorphisms were evaluated as possibly associated with end-stage renal disease (ESRD) resulting from type 2 diabetes mellitus (DM) nephropathy. **Methods.** Studies were conducted among hemodialysis (HD) patients with ESRD due to type 2 DM nephropathy, chronic glomerulonephritis, chronic infective tubulointerstitial nephritis, and hypertensive nephropathy as well as in healthy subjects. A frequency distribution of T-cell-related interleukin (IL) genes (*IL18* rs360719, *IL12A* rs568408, *IL12B* rs3212227, *IL4R* rs1805015, *IL13* rs20541, *IL28B* rs8099917, *IL28B*, and rs12979860) and vitamin D pathway genes (GC genes: rs2298849, rs7041, and rs1155563; VDR genes: rs2228570, rs1544410; and RXRA genes: rs10776909, rs10881578, and rs749759) was compared between groups. **Results.** No significant differences in a frequency distribution of tested polymorphisms were shown between type 2 DM nephropathy patients and controls. A difference was found in *IL18* rs360719 polymorphic distribution between the former group and chronic infective tubulointerstitial nephritic patients ($P_{\text{trend}} = 0.033$), which also differed in this polymorphism from controls ($P_{\text{trend}} = 0.005$). **Conclusion.** T-cell cytokine and vitamin D pathway gene polymorphisms are not associated with ESRD due to type 2 DM nephropathy in Polish HD patients. *IL18* rs360719 is probably associated with the pathogenesis of chronic infective tubulointerstitial nephritis.

1. Introduction

Diabetes mellitus (DM) is the most common cause of end-stage renal disease (ESRD) in many hemodialysis (HD) centers. In Australia and New Zealand, the incident ESRD population (1991–2005) who began renal replacement therapy (RRT) included 30.0% type 2 DM and 4.5% type 1 DM subjects [1]. In the HEMODIALYSIS (HEMO) study, the group of HD patients comprised approximately 45% of DM subjects [2].

Diabetic ESRD patients compared to nondiabetic ESRD subjects show higher both mortality rate [3] and prevalence of coronary artery disease (CAD) [4], are more prone to severe infections [5] and worse response to hepatitis B vaccination [6], and more often suffer from adynamic bone disease associated with low serum parathyroid hormone (PTH) levels [7]. In this paper we will focus on ESRD due to type 2 DM nephropathy. Together with altered glucose metabolism and insulin resistance, deficiency of vitamin D [8] and aberrant T-cell cytokine balance [9] were found to be associated with

TABLE 1: HRM and RFLP conditions for the identification of polymorphisms genotyped in the vitamin D pathway related genes.

Gene symbol	rs number	Alleles	Primers for PCR amplification (5'-3')	Annealing temp. (°C)	PCR product length (bp)	HRM ^a analysis Melting temp. range (°C)	Restriction enzyme	RFLP ^b analysis Restriction fragment length (bp)
GC	rs7041	G/T	F: GGAGGTGAGTTTATGGAACAGC R: GGCATTAAGCTGGTATGAGGTC	66.3	493		HaeIII	T = 493 G = 414 + 79
	rs1155563	C/T	F: GGTATTCTAAGACTGTGCTCTTGC R: ATGTGTTCTCACTGTTTCGACATCC	63.0	116	71-78		
	rs2298849	C/T	F: TCCACTGGCAAAACACATTAC R: GGGACATCTGCATTATCCTG	60.6	118	73-83		
	rs10881578	A/G	F: TCTTGAGCAATGCCAGCAG R: CCACAGCTCACACATCCAATC	60.6	75	80-90		
RXRA	rs10776909	C/T	F: CAGCCTGTGGCCTGCTCA R: AACCTCCGGCCCTTGGAG	60.6	95	82-92		
	rs749759	A/G	F: ATAGGGCTTGCCTGCCTAGA R: CTCCACCATAGCCCAAGTGA	62.6	382		BstXI	A = 382 G = 243 + 139
VDR	rs1544410	A/G	F: GGAGACACAGATAAGGAAATAC R: CCGCAAGAAACCTCAAATAACA	60.6	248		FspI	A (B) = 248 G (b) = 175 + 73
	rs2228570	C/T	F: GCACTGACTCTGGCTCTGAC R: ACCCTCCTGCTCCTGTGGCT	72.5	341		FokI	C (F) = 341 T (f) = 282 + 59

^aHRM analysis: high resolution melt analysis.^bRFLP analysis: restriction fragment length polymorphism analysis.

TABLE 2: Characteristics of hemodialysis patients ($n = 893$).

Parameter	Type 2 DM nephropathy	Other causes of ESRD	<i>P</i> value
Demographic data	$n = 366$	$n = 527$	
Male sex, n (% of all)	201 (54.9)	307 (58.3)	0.337 ^b
Age at RRT beginning, years	62.9 ± 14.1	57.2 ± 17.2	<0.0001^c
RRT duration, years	3.29 (0.06–28.0)	4.42 (0.12–28.2)	<0.0001^c
Death rate, cases per 100 patient-years	0.48	0.42	
Death rate, cases per 100 RRT-years	7.97	4.63	
Clinical data	$n = 332$	$n = 527$	
Coronary artery disease, n (% of all)	174 (52.4)	168 (31.9)	<0.0001^b
Myocardial infarction, n (% of all)	98 (29.5)	101 (19.2)	0.009^b
Parathyroidectomy, n (% of all)	2 (0.60)	21 (3.98)	0.0009^b
Treatment with cinacalcet hydrochloride	24 (7.2)	98 (18.6)	<0.0001^b
Laboratory data	$n = 366$	$n = 527$	
Anti-HBc positive, n (% of all)	95 (26.0)	126 (23.9)	0.528 ^b
HBsAg positive, n (% of all anti-HBc positive)	7 (7.4)	11 (8.7)	0.807 ^b
Anti-HCV positive, n (% of all)	26 (7.1)	57 (10.8)	0.062 ^b
HCV RNA positive, n (% of all anti-HCV positive)	14 (53.8)	39 (68.4)	0.225 ^b
Responders to hepatitis B vaccine, n (% of all)	202 (55.2)	315 (59.8)	0.191 ^b
25(OH)D (ng/mL) ^a	13.3 ± 3.9	14.5 ± 5.6	0.182 ^{a,d}
Total calcium (mg/dL)	8.83 ± 0.67	8.91 ± 0.82	0.264 ^d
Phosphates (mg/dL)	5.03 ± 1.44	5.25 ± 1.49	0.054 ^d
PTH (pg/mL)	296 (12.9–3,757)	463 (12.7–3,741)	<0.0001^c
Total alkaline phosphatase (U/L)	98.2 (25.8–1,353)	97.1 (40.5–1,684)	0.528 ^c

25(OH)D: 25-hydroxycholecalciferol, anti-HBc: antibodies to core antigen of hepatitis B virus, anti-HCV: antibodies to hepatitis C virus, HBsAg: surface antigen of hepatitis B virus, DM: diabetes mellitus, ESRD: end-stage renal disease, HCV RNA: ribonucleic acid of hepatitis C virus, PTH: parathyroid hormone, and RRT: renal replacement therapy.

A significant difference is indicated using bold font.

^a $n = 66$ for type 2 DM nephropathy; $n = 96$ for other renal diseases.

^bFisher's exact test.

^cMann-Whitney test.

^dUnpaired *t*-test, Welch corrected.

this severe complication of type 2 DM. There is a link between vitamin D and T-cell functional balance: active form of vitamin D [1,25(OH)₂D] has the inhibitory effect on the T helper (Th) 17 and Th1 response [10].

Abnormalities in T-cell cytokine equilibrium [11–13] and plasma vitamin D concentrations [14–16] are related to cardiovascular events [13, 16] and immunononcompetence during infections [11, 14] and vaccinations [12, 15]. Serum PTH levels are dependent on serum vitamin D concentrations [17], and T cells are implicated in the mechanism of PTH action in bone [18].

Vitamin D activity may be adequately expressed if vitamin D pathway components (vitamin D binding protein, also referred to as group-specific component (GC), vitamin D receptor (VDR), and retinoid X receptors (RXRs)) are properly structured and regulated. The recent study by Zhang et al. [19] has shown that VDR BsmI polymorphism correlates with type 2 DM nephropathy and may be susceptible for early onset of this nephropathy. Among T-cell-related cytokine gene polymorphisms, promoter polymorphic variants of *IL10* [20, 21] and *IL6* [22] were already associated with the risk of type 2 DM nephropathy. Monocyte chemoattractant protein 1 (MCP-1) has been reported to participate in the

pathogenesis of early type 2 DM nephropathy [23], but *MCPI* polymorphism in the promoter region was not differentially distributed between ESRD patients with type 2 DM nephropathy and healthy controls [24, 25].

To our knowledge, there are scarce data, if any, on ESRD due to type 2 DM nephropathy showing a frequency distribution of single nucleotide polymorphisms (SNPs) of T-cell-related IL genes: *IL18* rs360719, *IL12A* rs568408, *IL12B* rs3212227, *IL4R* rs1805015, *IL13* rs20541, *IL28B* rs8099917, and *IL28B* rs12979860 as well as vitamin D pathway genes: GC genes (*GC* rs2298849, rs7041, and rs1155563), VDR genes (*VDR* rs2228570, rs1544410), and RXR α genes (*RXRA* rs10776909, rs10881578, and rs749759). The aim of our study was to determine the potential association between aforementioned polymorphisms of T-cell-related cytokine genes and vitamin D pathway genes and ESRD due to type 2 DM nephropathy. For comparisons, aforementioned genotype frequencies of healthy controls as well as ESRD patients with other main causes of ESRD were used. Polymorphism related associations, if exist, could contribute to explanation of susceptibility to ESRD due to type 2 DM nephropathy and phenotype differences between ESRD patients with type 2 DM nephropathy and other causes of ESRD.

TABLE 3: Characteristics of hemodialysis patients grouped by a cause of ESRD.

Parameter	Type 2 DM nephropathy (1)	Chronic glomerulonephritis (2)	Chronic tubulointerstitial nephritis (3)	Hypertensive nephropathy (4)	<i>P</i> value
Demographic data	<i>n</i> = 366	<i>n</i> = 178	<i>n</i> = 118	<i>n</i> = 231	
Male sex, <i>n</i> (% of all)	201 (54.9)	110 (61.8)	63 (53.4)	134 (58.0)	0.386 ^b
Age at RRT beginning, years	62.9 ± 14.1	47.4 ± 17.6	59.9 ± 16.6	63.3 ± 13.6	<0.0001 ^c 1 versus 2: <0.001 ^c 2 versus 3: <0.001 ^c 2 versus 4: <0.001 ^c
RRT duration, years	3.29 (0.06–28.0)	5.73 (0.16–28.2)	4.82 (0.33–26.5)	3.82 (0.12–20.4)	<0.0001 ^c 1 versus 2: <0.001 ^c 1 versus 3: <0.01 ^c 2 versus 4: <0.001 ^c
Death rate, cases per 100 patient-years	0.48	0.41	0.44	0.42	
Death rate, cases per 100 dialysis-years	7.97	2.87	5.28	6.70	
Clinical data	<i>n</i> = 332	<i>n</i> = 178	<i>n</i> = 118	<i>n</i> = 231	
Coronary artery disease, <i>n</i> (% of all)	174 (52.4)	43 (24.2)	29 (24.6)	96 (41.5)	<0.0001 ^b 1 versus 2: <0.0001 ^c 1 versus 3: <0.0001 ^c 1 versus 4: 0.013 ^c 2 versus 4: 0.0002 ^c 3 versus 4: 0.002 ^c
Myocardial infarction, <i>n</i> (% of all)	98 (29.5)	25 (14.0)	17 (14.4)	59 (25.5)	<0.0001 ^b 1 versus 2: <0.0001 ^c 1 versus 3: <0.0001 ^c 1 versus 4: <0.0001 ^c 2 versus 4: 0.005 ^c 3 versus 4: 0.02 ^e
PTX, <i>n</i> (% of all)	2 (0.60)	14 (7.9)	5 (4.2)	2 (0.87)	<0.0001 ^b 1 versus 2: <0.0001 ^c 1 versus 3: 0.015 ^c 2 versus 4: 0.0004 ^c 3 versus 4: 0.046 ^c
Treatment with cinacalcet hydrochloride	24 (7.2)	48 (27.0)	21 (17.8)	29 (12.6)	<0.0001 ^b 1 versus 2: <0.0001 ^c 1 versus 3: 0.0008 ^c 1 versus 4: 0.017 ^c 2 versus 4: 0.0003 ^c
Laboratory data	<i>n</i> = 366	<i>n</i> = 178	<i>n</i> = 118	<i>n</i> = 231	
Anti-HBc positive, <i>n</i> (% of all)	95 (26.0)	53 (29.8)	25 (21.2)	48 (20.8)	0.233 ^b
HBsAg positive, <i>n</i> (% of all anti-HBc positive)	7 (7.4)	10 (18.9)	0 (0.0)	1 (2.08)	0.0007 ^b 1 versus 2: 0.032 ^c 2 versus 3: 0.007 ^c 2 versus 4: 0.001 ^c
Anti-HCV positive, <i>n</i> (% of all)	26 (7.1)	33 (18.5)	11 (9.3)	13 (5.6)	<0.0001 ^b 1 versus 2: 0.0004 ^c 2 versus 3: 0.031 ^c 2 versus 4: <0.0001 ^c
HCV RNA positive, <i>n</i> (% of all anti-HCV positive)	14 (53.8)	27 (81.8)	4 (36.4)	8 (61.5)	<0.0001 ^b 1 versus 2: <0.0001 ^c 2 versus 3: 0.0004 ^c 2 versus 4: <0.0001 ^c

TABLE 3: Continued.

Parameter	Type 2 DM nephropathy (1)	Chronic glomerulonephritis (2)	Chronic tubulointerstitial nephritis (3)	Hypertensive nephropathy (4)	P value
Responders to hepatitis B vaccine, <i>n</i> (% of all)	202 (55.2)	107 (60.1)	70 (59.3)	138 (59.7)	0.598 ^b
25(OH)D (ng/mL) ^a	13.3 ± 3.9	14.2 ± 7.3	15.7 ± 4.3	14.1 ± 3.9	0.453 ^d
Total calcium (mg/dL)	8.83 ± 0.67	8.85 ± 0.85	9.04 ± 0.61	8.88 ± 0.87	0.239 ^d
Phosphates (mg/dL)	5.03 ± 1.44	5.63 ± 1.59	4.92 ± 1.29	5.15 ± 1.47	0.0007^d 1 versus 2: <0.001^c 2 versus 3: <0.01^c 2 versus 4: <0.05^c
PTH (pg/mL)	296 (12.9–3,757)	632 (12.7–3,118)	426 (45.8–3,741)	364 (19.5–2,351)	<0.0001^c 1 versus 2: <0.001^c 1 versus 3: <0.05^c 1 versus 4: <0.05^c 2 versus 4: <0.001^c
Total ALP (U/L)	98.2 (25.8–1,353)	113 (44.5–860)	89.0 (40.5–1,684)	90.9 (41.0–1,110)	0.010^c 2 versus 4: <0.05^c

25(OH)D: 25-hydroxycholecalciferol, anti-HBc: antibodies to core antigen of hepatitis B virus, anti-HCV: antibodies to hepatitis C virus, HBsAg: surface antigen of hepatitis B virus, DM: diabetes mellitus, ESRD: end-stage renal disease, HCV RNA: ribonucleic acid of hepatitis C virus, PTH: parathyroid hormone, and RRT: renal replacement therapy.

^a*n* = 66 for type 2 DM nephropathy, *n* = 40 for chronic glomerulonephritis, *n* = 13 for chronic interstitial nephritis, and *n* = 43 for hypertensive nephropathy.

^bChi squared test.

^cKruskal-Wallis test.

^dANOVA test.

^eFisher's exact test.

2. Material and Methods

2.1. Patients and Controls. Blood samples for genotype analyses are collected since 2009 from ESRD patients (estimated glomerular filtration rate (eGFR) category G5 in accordance with KDIGO recommendations [26]). All subjects were treated with HD on enrolment. Controls were recruited from blood donors and healthy volunteers unrelated to patients. All enrolled individuals live/lived in the Greater Poland region of Poland.

Genotyping of *IL18* rs360719, *IL12A* rs568408, *IL12B* rs3212227, *IL4R* rs1805015, and *IL13* rs20541 polymorphisms was performed in 2009–2012 using currently available material. Results had been analyzed in our previous studies in the context of responsiveness to the surface antigen of hepatitis B virus (HBsAg) using data of all (not segregated) patients [27–30]. For this study, we used results of controls and patients with type 2 DM nephropathy, chronic glomerulonephritis, chronic infective tubulointerstitial nephritis, and hypertensive nephropathy.

IL28B rs8099917, *IL28B* rs12979860, *GC* rs2298849, *GC* rs7041, *GC* rs1155563, *VDR* rs2228570, *VDR* rs1544410, *RXRA* rs10776909, *RXRA* rs10881578, and *RXRA* rs749759 polymorphisms were analyzed in winter 2013/2014 among HD patients with ESRD (*n* = 893) due to type 2 DM nephropathy (*n* = 366), chronic glomerulonephritis (*n* = 178), chronic infective tubulointerstitial nephritis (*n* = 118), and hypertensive nephropathy (*n* = 231) as well as healthy controls (*n* = 378).

DM was not diagnosed in patients having renal diseases other than type 2 DM nephropathy.

Healthy individuals and HD patients with other renal diseases as cause of ESRD served as reference groups for a frequency distribution of tested polymorphic variants. All examined subjects were of Caucasian race.

Basic clinical and laboratory data were collected on enrolment and they are updated every year.

2.2. Genotyping. Genomic DNA for genotype analysis was isolated from peripheral blood lymphocytes by salt-out extraction procedure.

Genotyping of *IL18* rs360719, *IL12A* rs568408, *IL12B* rs3212227, *IL4R* rs1805015, and *IL13* rs20541 polymorphisms was performed as previously described [27–30].

IL28B rs8099917 and *IL28B* rs12979860 polymorphisms were genotyped using high-resolution melting curve analysis (HRM) on the LightCycler 480 system (Roche Diagnostics, Mannheim, Germany) with the use of 5x HOT FIREPol EvaGreen HRM Mix (Solis BioDyne, Tartu, Estonia). The PCR program consisted of an initial step at 95°C for 15 min to activate HOT FIREPol DNA polymerase, followed by 50 amplification cycles of denaturation at 95°C for 10 s, annealing at 61°C for 10 s, and elongation at 72°C for 15 s. Amplified DNA fragments were then subjected to HRM with 0.1°C increments in temperatures ranging from 76 to 96°C. Primers used for PCR with subsequent HRM analysis were as follows: rs8099917F 5' TTTGTCACTGTTCTCCTTTTG3', rs8099917R 5' AAGACATAAAAAGCCAGCTACCA3',

TABLE 4: Comparison of the distribution of polymorphic variants of tested genes between ESRD patients treated with hemodialysis due to type 2 DM nephropathy and healthy subjects.

Parameter	Type 2 DM nephropathy (frequency)	Healthy subjects (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
<i>IL18</i> rs360719	<i>n</i> = 248	<i>n</i> = 240			
TT	133 (0.54)	121 (0.50)	Referent		0.233
CT	102 (0.41)	98 (0.41)	0.947 (0.654–1.372)	0.777	
CC	13 (0.05)	21 (0.09)	0.563 (0.270–1.174)	0.145	
CT + CC	115 (0.46)	119 (0.50)	0.879 (0.616–1.254)	0.526	
MAF	128 (0.26)	140 (0.29)	0.845 (0.638–1.119)	0.268	
<i>IL12A</i> rs568408	<i>n</i> = 234	<i>n</i> = 240			
GG	173 (0.74)	171 (0.71)	Referent		0.782
AG	52 (0.22)	63 (0.26)	0.816 (0.534–1.246)	0.389	
AA	9 (0.04)	6 (0.03)	1.483 (0.517–4.256)	0.600	
AG + AA	61 (0.26)	69 (0.29)	0.874 (0.583–1.309)	0.538	
MAF	70 (0.15)	75 (0.16)	0.976 (0.684–1.393)	0.965	
<i>IL12B</i> rs3212227	<i>n</i> = 247	<i>n</i> = 240			
AA	156 (0.63)	151 (0.63)	Referent		0.639
AC	84 (0.34)	77 (0.32)	1.056 (0.721–1.547)	0.846	
CC	7 (0.03)	12 (0.05)	0.563 (0.217–1.473)	0.345	
AC + CC	91 (0.37)	89 (0.37)	0.990 (0.685–1.430)	1.000	
MAF	98 (0.20)	101 (0.21)	0.927 (0.680–1.268)	0.699	
<i>IL4R</i> rs1805015	<i>n</i> = 303	<i>n</i> = 225			
TT	205 (0.68)	162 (0.72)	Referent		0.304
CT	82 (0.27)	53 (0.24)	1.223 (0.818–1.828)	0.360	
CC	16 (0.05)	10 (0.04)	1.264 (0.559–2.861)	0.684	
CT + CC	98 (0.32)	63 (0.28)	1.229 (0.843–1.793)	0.295	
MAF	114 (0.19)	73 (0.16)	1.197 (0.866–1.653)	0.313	
<i>IL13</i> rs20541	<i>n</i> = 303	<i>n</i> = 230			
CC	168 (0.55)	124 (0.54)	Referent		0.457
CT	114 (0.38)	84 (0.36)	1.002 (0.695–1.443)	1.000	
TT	21 (0.07)	22 (0.10)	0.705 (0.371–1.338)	0.324	
CT + TT	135 (0.45)	106 (0.46)	0.940 (0.666–1.326)	0.726	
MAF	156 (0.26)	128 (0.28)	0.899 (0.684–1.182)	0.489	
<i>IL28B</i> rs8099917	<i>n</i> = 339	<i>n</i> = 375			
TT	219 (0.65)	245 (0.65)	Referent		0.504
GT	107 (0.31)	123 (0.33)	0.973 (0.709–1.336)	0.872	
GG	13 (0.04)	7 (0.02)	2.078 (0.814–5.302)	0.169	
GT + GG	120 (0.35)	130 (0.35)	1.033 (0.759–1.405)	0.875	
MAF	133 (0.20)	137 (0.18)	1.092 (0.837–1.423)	0.560	
<i>IL28B</i> rs12979860	<i>n</i> = 336	<i>n</i> = 372			
CC	141 (0.42)	164 (0.44)	Referent		0.669
CT	157 (0.47)	166 (0.45)	1.100 (0.804–1.505)	0.576	
TT	38 (0.11)	42 (0.11)	1.052 (0.643–1.723)	0.900	
CT + TT	195 (0.56)	208 (0.56)	1.090 (0.809–1.469)	0.595	
MAF	116 (0.29)	250 (0.34)	1.049 (0.842–1.307)	0.713	
<i>GC</i> rs2298849	<i>n</i> = 364 ^a	<i>n</i> = 375			
TT	226 (0.62)	237 (0.63)	Referent		0.250
CT	110 (0.30)	124 (0.33)	0.930 (0.679–1.274)	0.688	
CC	28 (0.08)	14 (0.04)	2.097 (1.077–4.086)	0.035	
CT + CC	138 (0.38)	138 (0.37)	1.049 (0.778–1.413)	0.762	
MAF	166 (0.23)	152 (0.20)	1.162 (0.907–1.490)	0.262	

TABLE 4: Continued.

Parameter	Type 2 DM nephropathy (frequency)	Healthy subjects (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
GC rs7041	<i>n</i> = 343	<i>n</i> = 361			
GG	112 (0.33)	116 (0.32)	Referent		0.572
GT	163 (0.47)	186 (0.52)	0.908 (0.650–1.268)	0.609	
TT	68 (0.20)	59 (0.16)	1.194 (0.773–1.844)	0.440	
GT + TT	231 (0.67)	245 (0.68)	0.977 (0.712–1.339)	0.936	
MAF	299 (0.44)	304 (0.42)	1.062 (0.860–1.312)	0.612	
GC rs1155563	<i>n</i> = 362	<i>n</i> = 377			
TT	180 (0.50)	189 (0.50)	Referent		0.541
CT	141 (0.39)	155 (0.41)	0.955 (0.703–1.297)	0.815	
CC	41 (0.11)	33 (0.09)	1.305 (0.789–2.155)	0.311	
CT + CC	182 (0.50)	188 (0.50)	1.017 (0.762–1.356)	0.941	
MAF	223 (0.31)	221 (0.29)	1.074 (0.859–1.341)	0.567	
VDR rs2228570	<i>n</i> = 345	<i>n</i> = 371			
CC	101 (0.29)	103 (0.28)	Referent		0.401
CT	175 (0.51)	183 (0.49)	0.975 (0.691–1.376)	0.930	
TT	69 (0.20)	85 (0.23)	0.828 (0.544–1.260)	0.394	
CT + TT	244 (0.71)	268 (0.72)	0.929 (0.671–1.285)	0.679	
MAF	313 (0.45)	353 (0.48)	0.915 (0.743–1.126)	0.432	
VDR rs1544410	<i>n</i> = 359	<i>n</i> = 372			
GG	137 (0.38)	148 (0.40)	Referent		0.753
AG	165 (0.46)	165 (0.44)	1.080 (0.787–1.483)	0.686	
AA	57 (0.16)	59 (0.16)	1.044 (0.678–1.607)	0.912	
AG + AA	222 (0.62)	224 (0.60)	1.071 (0.795–1.442)	0.705	
MAF	279 (0.39)	283 (0.38)	1.035 (0.839–1.278)	0.788	
RXRA rs10776909	<i>n</i> = 364	<i>n</i> = 378			
CC	233 (0.64)	250 (0.66)	Referent		0.426
CT	111 (0.30)	112 (0.30)	1.063 (0.774–1.461)	0.746	
TT	20 (0.05)	16 (0.04)	1.341 (0.679–2.651)	0.490	
CT + TT	131 (0.36)	128 (0.34)	1.098 (0.812–1.485)	0.590	
MAF	151 (0.21)	144 (0.19)	1.112 (0.862–1.435)	0.452	
RXRA rs10881578	<i>n</i> = 365	<i>n</i> = 377			
AA	197 (0.54)	183 (0.48)	Referent		0.168
AG	134 (0.37)	154 (0.41)	0.808 (0.775–1.046)	0.185	
GG	34 (0.09)	40 (0.11)	0.790 (0.479–1.301)	0.376	
AG + GG	168 (0.46)	194 (0.51)	0.804 (0.603–1.073)	0.143	
MAF	202 (0.28)	234 (0.31)	0.850 (0.680–1.063)	0.172	
RXRA rs749759	<i>n</i> = 355	<i>n</i> = 370			
GG	207 (0.58)	221 (0.60)	Referent		0.850
AG	125 (0.35)	123 (0.33)	1.085 (0.794–1.216)	0.632	
AA	23 (0.06)	26 (0.07)	0.944 (0.522–1.708)	0.881	
AG + AA	148 (0.42)	149 (0.40)	1.061 (0.789–1.426)	0.706	
MAF	171 (0.24)	175 (0.24)	1.024 (0.804–1.304)	0.894	

ESRD: end-stage renal disease, DM: diabetes mellitus, and MAF: minor allele frequency.

^aNot consistent with Hardy-Weinberg equilibrium.

rs12979860F 5'CGTGCCCTGTCGTGTACTGAA3', and rs12979860R 5'AGGCTCAGGGTCAATCACAG3'.

Genotyping of the GC rs1155563, GC rs2298849, RXRA rs10881578, and RXRA rs10776909 polymorphisms was carried out by HRM on the Bio-Rad CFX96 Real Time PCR system (Bio-Rad, Hercules, CA). DNA fragments amplified

with the use of specific primers were subjected to HRM with 0.1°C increments in temperatures ranging from 71 to 92°C. Genotyping of the GC rs7041, RXRA rs749759, VDR rs1544410, and VDR rs2228570 was performed using the polymerase chain reaction and restriction fragment length polymorphism (PCR-RFLP) method according to the

TABLE 5: Comparison of the distribution of polymorphic variants of tested genes between ESRD patients treated with hemodialysis due to type 2 DM nephropathy and the most common causes of ESRD other than type 2 DM nephropathy (chronic glomerulonephritis, chronic tubulointerstitial nephritis, and hypertensive nephritis).

Genotype	Type 2 DM nephropathy (frequency)	Other causes of ESRD (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
<i>IL18</i> rs360719	<i>n</i> = 248	<i>n</i> = 353			
TT	133 (0.54)	186 (0.53)	Referent	—	0.362
CT	102 (0.41)	135 (0.38)	1.057 (0.752–1.485)	0.795	
CC	13 (0.05)	32 (0.09)	0.568 (0.287–1.124)	0.107	
CT + CC	115 (0.46)	167 (0.47)	0.963 (0.696–1.334)	0.868	
MAF	128 (0.26)	199 (0.28)	0.886 (0.684–1.149)	0.370	
<i>IL12A</i> rs568408	<i>n</i> = 234	<i>n</i> = 337			
GG	173 (0.74)	234 (0.69)	Referent	—	0.303
AG	52 (0.22)	89 (0.26)	0.790 (0.533–1.060)	0.275	
AA	9 (0.04)	14 (0.04)	0.870 (0.368–2.055)	0.831	
AG + AA	61 (0.26)	103 (0.31)	0.801 (0.552–1.163)	0.260	
MAF	70 (0.15)	117 (0.17)	0.837 (0.606–1.157)	0.319	
<i>IL12B</i> rs3212227	<i>n</i> = 247	<i>n</i> = 352			
AA	156 (0.63)	205 (0.58)	Referent	—	0.176
AC	84 (0.34)	132 (0.38)	0.836 (0.593–1.068)	0.337	
CC	7 (0.03)	15 (0.04)	0.613 (0.244–1.540)	0.376	
AC + CC	91 (0.37)	147 (0.42)	0.814 (0.582–1.136)	0.236	
MAF	98 (0.20)	162 (0.23)	0.828 (0.624–1.098)	0.215	
<i>IL4R</i> rs1805015	<i>n</i> = 303	<i>n</i> = 436			
TT	205 (0.68)	295 (0.68)	Referent	—	0.871
CT	82 (0.27)	121 (0.28)	0.975 (0.700–2.360)	0.933	
CC	16 (0.05)	20 (0.05)	1.151 (0.583–2.275)	0.728	
CT + CC	98 (0.32)	141 (0.32)	1.000 (0.731–1.368)	1.000	
MAF	114 (0.19)	161 (0.18)	1.023 (0.784–1.335)	0.919	
<i>IL13</i> rs20541	<i>n</i> = 303	<i>n</i> = 436			
CC	168 (0.55)	242 (0.56)	Referent	—	0.902
CT	114 (0.38)	166 (0.38)	0.989 (0.726–1.348)	1.000	
TT	21 (0.07)	28 (0.06)	1.080 (0.594–1.967)	0.878	
CT + TT	135 (0.45)	194 (0.44)	1.002 (0.746–1.346)	1.000	
MAF	156 (0.26)	222 (0.25)	1.015 (0.800–1.287)	0.950	
<i>IL28B</i> rs8099917	<i>n</i> = 339	<i>n</i> = 493			
TT	219 (0.65)	317 (0.64)	Referent	—	0.858
GT	107 (0.31)	162 (0.33)	0.956 (0.709–1.289)	0.820	
GG	13 (0.04)	14 (0.03)	1.344 (0.620–2.916)	0.549	
GT + GG	120 (0.35)	176 (0.36)	0.987 (0.739–1.318)	0.941	
MAF	133 (0.20)	190 (0.19)	1.022 (0.799–1.309)	0.910	
<i>IL28B</i> rs12979860	<i>n</i> = 336	<i>n</i> = 488			
CC	141 (0.42)	209 (0.43)	Referent	—	0.952
CT	157 (0.47)	221 (0.45)	1.053 (0.783–1.415)	0.763	
TT	38 (0.11)	58 (0.12)	0.971 (0.612–1.541)	0.907	
CT + TT	195 (0.56)	279 (0.57)	1.036 (0.782–1.373)	0.830	
MAF	116 (0.29)	337 (0.35)	1.006 (0.819–1.237)	0.994	
<i>GC</i> rs2298849	<i>n</i> = 364 ^a	<i>n</i> = 524			
TT	226 (0.62)	339 (0.65)	Referent	—	0.109
CT	110 (0.30)	165 (0.31)	1.000 (0.745–1.342)	1.000	
CC	28 (0.08)	20 (0.04)	2.100 (1.155–3.819)	0.014	
CT + CC	138 (0.38)	185 (0.35)	1.119 (0.848–1.477)	0.436	
MAF	166 (0.23)	205 (0.20)	1.215 (0.964–1.530)	0.111	

TABLE 5: Continued.

Genotype	Type 2 DM nephropathy (frequency)	Other causes of ESRD (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
GC rs7041	<i>n</i> = 343	<i>n</i> = 506			
GG	112 (0.33)	182 (0.36)	Referent	—	0.247
GT	163 (0.47)	236 (0.47)	1.122 (0.824–1.528)	0.480	
TT	68 (0.20)	88 (0.17)	1.256 (0.846–1.863)	0.267	
GT + TT	231 (0.67)	324 (0.64)	1.159 (0.867–1.548)	0.340	
MAF	299 (0.44)	412 (0.41)	1.125 (0.925–1.369)	0.259	
GC rs1155563	<i>n</i> = 362	<i>n</i> = 527			
TT	180 (0.50)	252 (0.48)	Referent	—	0.614
CT	141 (0.39)	213 (0.40)	0.927 (0.696–1.234)	0.610	
CC	41 (0.11)	62 (0.12)	0.926 (0.597–1.435)	0.740	
CT + CC	182 (0.50)	275 (0.52)	0.927 (0.709–1.211)	0.585	
MAF	223 (0.31)	337 (0.32)	0.947 (0.772–1.161)	0.638	
VDR rs2228570	<i>n</i> = 345	<i>n</i> = 503			
CC	101 (0.29)	130 (0.26)	Referent	—	0.541
CT	175 (0.51)	275 (0.55)	0.819 (0.594–1.130)	0.249	
TT	69 (0.20)	98 (0.19)	0.906 (0.606–1.356)	0.682	
CT + TT	244 (0.71)	373 (0.74)	0.842 (0.620–1.143)	0.273	
MAF	313 (0.45)	471 (0.47)	0.943 (0.776–1.145)	0.588	
VDR rs1544410	<i>n</i> = 359	<i>n</i> = 512			
GG	137 (0.38)	189 (0.37)	Referent	—	0.598
AG	165 (0.46)	235 (0.46)	0.969 (0.720–1.303)	0.880	
AA	57 (0.16)	88 (0.17)	0.894 (0.599–1.332)	0.613	
AG + AA	222 (0.62)	323 (0.63)	0.948 (0.718–1.253)	0.722	
MAF	279 (0.39)	411 (0.40)	0.948 (0.778–1.152)	0.626	
RXRA rs10776909	<i>n</i> = 364	<i>n</i> = 526			
CC	233 (0.64)	308 (0.59)	Referent	—	0.298
CT	111 (0.30)	196 (0.37)	0.749 (0.561–0.999)	0.050	
TT	20 (0.05)	22 (0.04)	1.202 (0.641–2.254)	0.629	
CT + TT	131 (0.36)	218 (0.41)	0.794 (0.603–1.046)	0.108	
MAF	151 (0.21)	240 (0.23)	0.883 (0.702–1.112)	0.317	
RXRA rs10881578	<i>n</i> = 365	<i>n</i> = 525			
AA	197 (0.54)	252 (0.48)	Referent	—	0.134
AG	134 (0.37)	220 (0.42)	0.779 (0.586–1.035)	0.096	
GG	34 (0.09)	53 (0.10)	0.821 (0.513–1.312)	0.478	
AG + GG	168 (0.46)	273 (0.52)	0.787 (0.602–1.029)	0.088	
MAF	202 (0.28)	326 (0.31)	0.850 (0.690–1.046)	0.139	
RXRA rs749759	<i>n</i> = 355	<i>n</i> = 514			
GG	207 (0.58)	265 (0.52)	Referent	—	0.082
AG	125 (0.35)	212 (0.41)	0.755 (0.567–1.005)	0.059	
AA	23 (0.06)	37 (0.07)	0.796 (0.459–1.381)	0.490	
AG + AA	148 (0.42)	249 (0.48)	0.761 (0.579–1.000)	0.053	
MAF	171 (0.24)	286 (0.28)	0.823 (0.661–1.025)	0.092	

ESRD: end-stage renal disease, DM: diabetes mellitus, and MAF: minor allele frequency.

^aNot consistent with Hardy-Weinberg equilibrium.

manufacturer's instructions (Fermentas, Vilnius, Lithuania). Primer sequences and conditions for HRM and PCR-RFLP analyses are presented in Table 1.

For quality control, the genotyping analysis was blinded to the subject's case-control status. In addition, approximately 10% of the randomly chosen samples were regenotyped. Samples that failed the genotyping were excluded from further statistical analyses.

2.3. 25(OH)D Testing. Plasma 25(OH)D was determined in blindly selected 162 HD patients in the winter season of the year to avoid differences in sunlight exposure between patients who used to sunbathe and those who did not. Plasma 25(OH)D concentration was measured in HD patients who had not been treated with vitamin D or had stopped such a treatment for at least 3 weeks to obtain the so-called basic vitamin D concentrations. Under these conditions, there were

TABLE 6: Selected comparisons of the polymorphic variants distribution of tested genes between type 2 DM nephropathy patients, chronic infective tubulointerstitial nephritic patients, and healthy subjects.

Genotype	Genotype frequencies		Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
Type 2 DM nephropathy versus chronic infective tubulointerstitial nephritis					
<i>IL18</i> rs360719	<i>n</i> = 248	<i>n</i> = 77			
TT	133 (0.54)	54 (0.70)	Referent		0.033
CT	102 (0.41)	19 (0.25)	2.180 (1.217–3.905)	0.009^a	
CC	13 (0.05)	4 (0.05)	1.320 (0.412–4.228)	0.783	
CT + CC	115 (0.46)	23 (0.30)	2.030 (1.173–3.512)	0.012^a	
MAF	128 (0.26)	27 (0.18)	1.636 (1.031–2.596)	0.046	
Chronic infective tubulointerstitial nephritis versus healthy controls					
<i>IL18</i> rs360719	<i>n</i> = 77	<i>n</i> = 240			
TT	54 (0.70)	121 (0.50)	Referent		0.005
CT	19 (0.25)	98 (0.41)	0.434 (0.242–0.781)	0.006^a	
CC	4 (0.05)	21 (0.09)	0.427 (0.140–1.303)	0.160	
CT + CC	23 (0.30)	119 (0.50)	0.433 (0.250–0.750)	0.004^a	
MAF	27 (0.18)	140 (0.29)	0.516 (0.326–0.818)	0.006	

DM: diabetes mellitus; MAF: minor allele frequency.

Significant differences are indicated using bold font.

^aSignificant after the Bonferroni correction ($P < 0.017$).

no patients showing optimal plasma 25(OH)D levels (35–80 ng/mL for adults). To examine plasma 25(OH)D levels, a chemiluminescent microparticle immunoassay (CMIA) was used according to the manufacturer's instructions (Abbott Diagnostics ARCHITECT 25-OH VITAMIN D CMIA).

2.4. Statistical Methods. Results are presented as percentage for categorical variables, as mean with one standard deviation for normally distributed continuous variables or as median with range for not normally distributed continuous variables as tested by the Shapiro-Wilk test. Statistical tests used for comparison of data obtained in selected groups are indicated at *P* values.

Hardy-Weinberg equilibrium (HWE) was tested to compare the observed genotype frequencies to the expected ones using Chi-square test. Distributions of tested polymorphisms were consistent with HWE with three exceptions which are indicated in tables showing analysis of genotype and allele distributions. The Fisher exact probability test or Chi-square test was used to evaluate differences in genotype and allele prevalence between the examined groups. Homozygotes for the major allele were the reference group. The odds ratio (OR) with *P* value and 95% confidence intervals (95% CI) value were calculated. All probabilities were two-tailed. Polymorphisms were tested for association using the Chi-square test for trend (*P*_{trend}). Power analysis was performed by Fisher's exact test.

Values of $P < 0.05$ were judged to be significant. However, associations were reported only if the following conditions were fulfilled.

- (1) A genotype distribution was consistent with HWE in a tested group and a referent group.
- (2) *P*_{trend} was below 0.05.

- (3) Odds ratio remained significant after the Bonferroni correction applied for multiple testing, if appropriate.

Aforementioned statistical calculations were performed using GraphPad InStat 3.10, 32 bit for Windows, created on July 9, 2009 (GraphPad Software, Inc., La Jolla, USA), Cytel-Studio version 10.0, created on January 16, 2013 (CytelStudio Software Corporation, Cambridge, USA), and Statistica version 10, 2011 (StatSoft, Inc., Tulsa, USA).

3. Results

Characteristics of the examined HD patients are presented in Tables 2 and 3. ESRD patients due to type 2 DM nephropathy compared to non-DM ESRD patients showed older age at RRT onset, shorter treatment with RRT, higher death rate on RRT, higher prevalence of CAD and myocardial infarction, lower serum PTH level, and lower frequency of parathyroidectomy and treatment with cinacalcet.

In respect of the examined parameters, type 2 DM nephropathy patients differed the most significantly from chronic glomerulonephritic subjects, the least significantly from hypertensive nephropathy patients.

There were no differences in frequency distributions of tested genotypes between type 2 DM nephropathy patients and healthy subjects (Table 4) as well as other ESRD patients analyzed together (Table 5) which could be judged as significant associations.

Comparisons of genotype and allele frequencies between type 2 DM nephropathy patients and other ESRD groups revealed associations only with chronic infective tubulointerstitial nephritic patients in respect of *IL18* rs360719 (Table 6, no significant results are shown). Frequency of *IL18* rs360719 allele C carriers was higher in type 2 DM nephropathy patients than in those with chronic infective tubulointerstitial

TABLE 7: Comparison of the distribution of polymorphic variants of tested genes between ESRD patients treated with hemodialysis due to type 2 DM nephropathy grouped by diagnosis of CAD.

Parameter	Type 2 DM nephropathy with CAD (frequency)	Type 2 DM nephropathy without CAD (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
<i>IL18</i> rs360719	<i>n</i> = 124	<i>n</i> = 109			
TT	68 (0.55)	53 (0.49)	Referent		0.269
CT	51 (0.41)	49 (0.45)	1.128 (0.725–1.754)	0.653	
CC	5 (0.04)	7 (0.06)	0.628 (0.194–2.036)	0.557	
CT + CC	56 (0.45)	56 (0.51)	0.879 (0.560–1.380)	0.645	
MAF	61 (0.25)	63 (0.29)	0.803 (0.532–1.211)	0.345	
<i>IL12A</i> rs568408	<i>n</i> = 117	<i>n</i> = 102			
GG	83 (0.71)	77 (0.63)	Referent		0.361
AG	28 (0.24)	22 (0.22)	1.181 (0.623–2.236)	0.630	
AA	6 (0.05)	3 (0.03)	1.855 (0.448–7.678)	0.502	
AG + AA	34 (0.29)	25 (0.25)	1.262 (0.691–2.304)	0.542	
MAF	40 (0.17)	28 (0.14)	1.311 (0.776–2.214)	0.378	
<i>IL12B</i> rs3212227	<i>n</i> = 124	<i>n</i> = 109			
AA	78 (0.63)	69 (0.63)	Referent		0.906
AC	43 (0.35)	36 (0.33)	1.057 (0.611–1.829)	0.889	
CC	3 (0.02)	4 (0.04)	0.664 (0.143–3.069)	0.708	
AC + CC	46 (0.37)	40 (0.37)	1.017 (0.597–1.734)	1.000	
MAF	49 (0.20)	44 (0.20)	0.974 (0.618–1.535)	0.909	
<i>IL4R</i> rs1805015	<i>n</i> = 144	<i>n</i> = 127			
TT	95 (0.66)	86 (0.68)	Referent		0.947
CT	42 (0.29)	32 (0.25)	1.188 (0.689–2.048)	0.581	
CC	7 (0.05)	9 (0.07)	0.704 (0.251–1.972)	0.605	
CT + CC	49 (0.34)	41 (0.32)	1.082 (0.652–1.797)	0.797	
MAF	56 (0.19)	50 (0.20)	0.985 (0.644–1.504)	0.944	
<i>IL13</i> rs20541	<i>n</i> = 144	<i>n</i> = 127			
CC	80 (0.56)	71 (0.56)	Referent		0.867
CT	55 (0.38)	46 (0.36)	1.061 (0.640–1.759)	0.898	
TT	9 (0.06)	10 (0.08)	0.799 (0.307–2.077)	0.808	
CT + TT	64 (0.44)	56 (0.44)	1.014 (0.627–1.640)	1.000	
MAF	73 (0.25)	92 (0.26)	0.967 (0.657–1.423)	0.944	
<i>IL28B</i> rs8099917	<i>n</i> = 163	<i>n</i> = 145			
TT	105 (0.64)	97 (0.67)	Referent		0.752
GT	52 (0.32)	42 (0.29)	1.144 (0.700–1.870)	0.618	
GG	6 (0.04)	6 (0.04)	0.924 (0.288–2.961)	1.000	
GT + GG	58 (0.36)	48 (0.33)	1.116 (0.697–1.189)	0.719	
MAF	64 (0.20)	54 (0.19)	1.068 (0.714–1.597)	0.829	
<i>IL28B</i> rs12979860	<i>n</i> = 163	<i>n</i> = 142			
CC	69 (0.42)	66 (0.46)	Referent		0.352
CT	73 (0.45)	62 (0.44)	1.126 (0.698–1.816)	0.715	
TT	21 (0.13)	14 (0.10)	1.435 (0.674–3.055)	0.448	
CT + TT	94 (0.58)	76 (0.54)	1.183 (0.752–1.861)	0.490	
MAF	115 (0.35)	90 (0.32)	1.175 (0.838–1.647)	0.396	
<i>GC</i> rs2298849	<i>n</i> = 172	<i>n</i> = 158 ^a			
TT	99 (0.58)	106 (0.67)	Referent		0.173
CT	60 (0.35)	40 (0.25)	1.606 (0.989–2.608)	0.067	
CC	13 (0.07)	12 (0.08)	1.160 (0.505–2.663)	0.833	
CT + CC	73 (0.42)	52 (0.33)	1.503 (0.959–2.355)	0.088	
MAF	166 (0.25)	64 (0.20)	1.313 (0.909–1.895)	0.174	

TABLE 7: Continued.

Parameter	Type 2 DM nephropathy with CAD (frequency)	Type 2 DM nephropathy without CAD (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
<i>GC rs7041</i>	<i>n</i> = 161	<i>n</i> = 151			
GG	57 (0.35)	46 (0.30)	Referent		0.844
GT	69 (0.43)	82 (0.54)	1.327 (0.825–2.134)	0.277	
TT	35 (0.22)	23 (0.15)	1.629 (0.900–2.949)	0.136	
GT + TT	104 (0.65)	105 (0.70)	1.061 (0.721–1.559)	0.769	
MAF	139 (0.43)	128 (0.42)	1.025 (0.746–1.409)	0.943	
<i>GC rs1155563</i>	<i>n</i> = 172	<i>n</i> = 157			
TT	82 (0.48)	79 (0.50)	Referent		0.645
CT	70 (0.41)	61 (0.39)	1.106 (0.697–1.755)	0.724	
CC	20 (0.12)	17 (0.11)	1.133 (0.554–2.321)	0.856	
CT + CC	90 (0.52)	78 (0.50)	1.112 (0.721–1.714)	0.660	
MAF	110 (0.32)	95 (0.30)	1.084 (0.779–1.508)	0.695	
<i>VDR rs2228570</i>	<i>n</i> = 162	<i>n</i> = 152			
CC	43 (0.27)	44 (0.29)	Referent		0.316
CT	93 (0.57)	68 (0.45)	1.400 (0.829–2.363)	0.230	
TT	26 (0.16)	40 (0.26)	0.665 (0.348–1.272)	0.252	
CT + TT	119 (0.73)	108 (0.71)	1.128 (0.688–1.849)	0.705	
MAF	145 (0.45)	148 (0.49)	0.854 (0.624–1.169)	0.365	
<i>VDR rs1544410</i>	<i>n</i> = 170	<i>n</i> = 155			
GG	65 (0.38)	61 (0.39)	Referent		0.772
AG	79 (0.46)	72 (0.46)	1.030 (0.641–1.653)	0.905	
AA	26 (0.15)	22 (0.14)	1.109 (0.569–2.160)	0.865	
AG + AA	105 (0.62)	94 (0.61)	1.048 (0.671–1.639)	0.909	
MAF	131 (0.39)	116 (0.37)	1.048 (0.763–1.440)	0.833	
<i>RXRA rs10776909</i>	<i>n</i> = 172	<i>n</i> = 158			
CC	112 (0.65)	104 (0.66)	Referent		0.621
CT	48 (0.28)	47 (0.30)	0.948 (0.585–1.537)	0.902	
TT	12 (0.07)	7 (0.04)	1.592 (0.604–4.198)	0.473	
CT + TT	60 (0.35)	54 (0.34)	1.032 (0.655–1.625)	0.908	
MAF	72 (0.21)	61 (0.19)	1.107 (0.756–1.621)	0.672	
<i>RXRA rs10881578</i>	<i>n</i> = 173	<i>n</i> = 158			
AA	89 (0.51)	92 (0.58)	Referent		0.192
AG	65 (0.38)	53 (0.34)	1.268 (0.796–2.019)	0.345	
GG	19 (0.11)	13 (0.08)	1.511 (0.704–3.241)	0.340	
AG + GG	84 (0.49)	66 (0.42)	1.316 (0.852–2.032)	0.226	
MAF	103 (0.30)	79 (0.25)	1.272 (0.902–1.793)	0.199	
<i>RXRA rs749759</i>	<i>n</i> = 169	<i>n</i> = 153			
GG	100 (0.59)	89 (0.58)	Referent		0.812
AG	59 (0.35)	54 (0.35)	0.972 (0.610–1.551)	1.000	
AA	10 (0.06)	10 (0.07)	0.890 (0.354–2.238)	0.818	
AG + AA	69 (0.41)	64 (0.42)	0.960 (0.615–1.496)	0.910	
MAF	79 (0.23)	74 (0.24)	0.956 (0.665–1.375)	0.882	

CAD: coronary artery disease, ESRD: end-stage renal disease, DM: diabetes mellitus, and MAF: minor allele frequency.

^aNot consistent with Hardy-Weinberg equilibrium.

nephritis. The latter group showed lower frequency of *IL18* rs360719 allele C carriers compared to healthy controls (Table 6).

Type 2 DM nephropathy patients with diagnosed CAD differed in tested genotype frequencies neither from type 2 DM nephropathy subjects without CAD (Table 7) nor from healthy controls (Table 8).

4. Discussion

Genetic studies involving DM nephropathy and related complications are not consistent in many aspects [31–34]. Some polymorphisms tested in this study were reported as being associated with type 1 DM (*IL12B* rs3212227 [35], *IL4R* [36, 37], *IL13* [37], *VDR* rs1544410 [38, 39], and *VDR*

TABLE 8: Comparison of the distribution of polymorphic variants of tested genes between type 2 DM nephropathy patients with diagnosis of CAD and healthy controls.

Parameter	Type 2 DM nephropathy with CAD (frequency)	Healthy controls (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
<i>IL18</i> rs360719	<i>n</i> = 124	<i>n</i> = 240			
TT	68 (0.55)	121 (0.50)	Referent		0.186
CT	51 (0.41)	98 (0.41)	0.926 (0.590–1.453)	0.819	
CC	5 (0.04)	21 (0.09)	0.424 (0.153–1.174)	0.122	
CT + CC	56 (0.45)	119 (0.50)	0.837 (0.542–1.294)	0.440	
MAF	61 (0.25)	140 (0.29)	0.792 (0.558–1.124)	0.223	
<i>IL12A</i> rs568408	<i>n</i> = 117	<i>n</i> = 240			
GG	83 (0.71)	171 (0.71)	Referent		0.626
AG	28 (0.24)	63 (0.26)	0.916 (0.546–1.535)	0.794	
AA	6 (0.05)	6 (0.03)	2.060 (0.645–6.583)	0.348	
AG + AA	34 (0.29)	69 (0.29)	1.015 (0.624–1.653)	1.000	
MAF	40 (0.17)	75 (0.16)	1.113 (0.731–1.695)	0.695	
<i>IL12B</i> rs3212227	<i>n</i> = 124	<i>n</i> = 240			
AA	78 (0.63)	151 (0.63)	Referent		0.475
AC	43 (0.35)	77 (0.32)	1.081 (0.681–1.717)	0.813	
CC	3 (0.02)	12 (0.05)	0.484 (0.133–1.766)	0.397	
AC + CC	46 (0.37)	89 (0.37)	1.001 (0.639–1.567)	1.000	
MAF	49 (0.20)	101 (0.21)	0.924 (0.631–1.354)	0.757	
<i>IL4R</i> rs1805015	<i>n</i> = 144	<i>n</i> = 225			
TT	95 (0.66)	162 (0.72)	Referent		0.285
CT	42 (0.29)	53 (0.24)	1.351 (0.838–2.179)	0.221	
CC	7 (0.05)	10 (0.04)	1.194 (0.440–3.240)	0.798	
CT + CC	49 (0.34)	63 (0.28)	1.326 (0.845–2.083)	0.246	
MAF	56 (0.19)	73 (0.16)	1.247 (0.848–1.832)	0.305	
<i>IL13</i> rs20541	<i>n</i> = 144	<i>n</i> = 230			
CC	80 (0.56)	124 (0.54)	Referent		0.469
CT	55 (0.38)	84 (0.36)	1.015 (0.653–1.578)	1.000	
TT	9 (0.06)	22 (0.10)	0.634 (0.278–1.447)	0.324	
CT + TT	64 (0.44)	106 (0.46)	0.936 (0.616–1.422)	0.831	
MAF	73 (0.25)	128 (0.28)	0.881 (0.630–1.231)	0.510	
<i>IL28B</i> rs8099917	<i>n</i> = 163	<i>n</i> = 375			
TT	105 (0.64)	245 (0.65)	Referent		0.584
GT	52 (0.32)	123 (0.33)	0.986 (0.663–1.467)	1.000	
GG	6 (0.04)	7 (0.02)	2.000 (0.656–6.094)	0.229	
GT + GG	58 (0.36)	130 (0.35)	1.041 (0.709–1.530)	0.845	
MAF	64 (0.20)	137 (0.18)	1.093 (0.786–1.521)	0.658	
<i>IL28B</i> rs12979860	<i>n</i> = 163	<i>n</i> = 372			
CC	69 (0.42)	164 (0.44)	Referent		0.281
CT	73 (0.45)	166 (0.45)	1.045 (0.705–1.549)	0.841	
TT	21 (0.13)	42 (0.11)	1.188 (0.656–2.154)	0.644	
CT + TT	94 (0.58)	208 (0.56)	1.074 (0.740–1.558)	0.776	
MAF	115 (0.35)	250 (0.34)	1.077 (0.819–1.416)	0.644	
<i>GC</i> rs2298849	<i>n</i> = 172	<i>n</i> = 375			
TT	99 (0.58)	237 (0.63)	Referent		0.080
CT	60 (0.35)	124 (0.33)	1.158 (0.786–1.706)	0.486	
CC	13 (0.07)	14 (0.04)	2.223 (1.008–4.901)	0.052	
CT + CC	73 (0.42)	138 (0.37)	1.266 (0.876–1.830)	0.220	
MAF	166 (0.25)	152 (0.20)	1.311 (0.969–1.774)	0.092	

TABLE 8: Continued.

Parameter	Type 2 DM nephropathy with CAD (frequency)	Healthy controls (frequency)	Odds ratio (95% CI)	Two-tailed <i>P</i>	<i>P</i> _{trend}
<i>GC rs7041</i>	<i>n</i> = 161	<i>n</i> = 361			
GG	57 (0.35)	116 (0.32)	Referent		0.748
GT	69 (0.43)	186 (0.52)	0.755 (0.496–1.150)	0.196	
TT	35 (0.22)	59 (0.16)	1.207 (0.714–2.040)	0.502	
GT + TT	104 (0.65)	245 (0.68)	0.864 (0.584–1.278)	0.482	
MAF	139 (0.43)	304 (0.42)	1.044 (0.801–1.362)	0.800	
<i>GC rs1155563</i>	<i>n</i> = 172	<i>n</i> = 377			
TT	82 (0.48)	189 (0.50)	Referent		0.378
CT	70 (0.41)	155 (0.41)	1.041 (0.710–1.527)	0.845	
CC	20 (0.12)	33 (0.09)	1.397 (0.757–2.578)	0.332	
CT + CC	90 (0.52)	188 (0.50)	1.103 (0.769–1.583)	0.646	
MAF	110 (0.32)	221 (0.29)	1.134 (0.861–1.494)	0.411	
<i>VDR rs2228570</i>	<i>n</i> = 162	<i>n</i> = 371			
CC	43 (0.27)	103 (0.28)	Referent		0.386
CT	93 (0.57)	183 (0.49)	1.217 (0.788–1.880)	0.384	
TT	26 (0.16)	85 (0.23)	0.733 (0.416–1.290)	0.321	
CT + TT	119 (0.73)	268 (0.72)	1.064 (0.702–1.613)	0.833	
MAF	145 (0.45)	353 (0.48)	0.893 (0.687–1.160)	0.434	
<i>VDR rs1544410</i>	<i>n</i> = 170	<i>n</i> = 372			
GG	65 (0.38)	148 (0.40)	Referent		0.880
AG	79 (0.46)	165 (0.44)	1.090 (0.734–1.620)	0.687	
AA	26 (0.15)	59 (0.16)	1.003 (0.581–1.732)	1.000	
AG + AA	105 (0.62)	224 (0.60)	1.067 (0.735–1.549)	0.776	
MAF	131 (0.39)	283 (0.38)	1.021 (0.784–1.329)	0.931	
<i>RXRA rs10776909</i>	<i>n</i> = 172	<i>n</i> = 378			
CC	112 (0.65)	250 (0.66)	Referent		0.483
CT	48 (0.28)	112 (0.30)	0.957 (0.638–1.434)	0.838	
TT	12 (0.07)	16 (0.04)	1.674 (0.767–3.656)	0.209	
CT + TT	60 (0.35)	128 (0.34)	1.046 (0.716–1.529)	0.846	
MAF	72 (0.21)	144 (0.19)	1.125 (0.819–1.545)	0.518	
<i>RXRA rs10881578</i>	<i>n</i> = 173	<i>n</i> = 377			
AA	89 (0.51)	183 (0.48)	Referent		0.682
AG	65 (0.38)	154 (0.41)	0.868 (0.591–1.275)	0.494	
GG	19 (0.11)	40 (0.11)	0.977 (0.535–1.783)	1.000	
AG + GG	84 (0.49)	194 (0.51)	0.890 (0.621–1.276)	0.582	
MAF	103 (0.30)	234 (0.31)	0.942 (0.714–1.243)	0.725	
<i>RXRA rs749759</i>	<i>n</i> = 169	<i>n</i> = 370			
GG	100 (0.59)	221 (0.60)	Referent		0.924
AG	59 (0.35)	123 (0.33)	1.060 (0.718–1.566)	0.842	
AA	10 (0.06)	26 (0.07)	0.850 (0.395–1.830)	0.710	
AG + AA	69 (0.41)	149 (0.40)	1.023 (0.707–1.482)	0.925	
MAF	79 (0.23)	175 (0.24)	0.985 (0.727–1.334)	0.983	

CAD: coronary artery disease, DM: diabetes mellitus, and MAF: minor allele frequency.

rs2228570 [38]), type 2 DM susceptibility (*VDR* rs2228570 [40], *VDR* rs1544410 [41]), and phenotype of type 2 DM (*VDR* rs2228570 [42], *VDR* rs1544410 [41, 43]). *VDR* rs2228570 and *IL4* polymorphisms were also related to the risk of chronic kidney disease [44, 45]. On the other hand, there are also data indicating no major effect of *IL12B* on type 1 DM susceptibility in the entire study group [46], no association of *IL4R* with type 1 DM [47], no evident causal relationship between vitamin D pathway genes and type 2 DM, myocardial infarction or mortality [48], similar distribution of genotypes, allele and haplotypes of *VDR* rs2228570 and *VDR* rs731236 between type 2 DM patients and controls [49], no contribution of *VDR* rs1544410 to type 1 DM susceptibility [50], and no association of *VDR* rs1544410 with chronic kidney disease susceptibility [51].

In this study we were not able to show significant differences in the frequency distribution of tested polymorphic variants of T-cell-related cytokine genes or vitamin D pathway genes between HD patients with ESRD due to type 2 DM nephropathy and controls as well as HD patients with other causes of ESRD analyzed together. This lack of association was present although the examined type 2 DM nephropathy patients showed clinical complications more frequently than HD patients with other renal diseases: higher dialysis related mortality rate [3], higher prevalence of CAD including myocardial infarction [4], lower serum PTH, and lower frequency of parathyroidectomy and treatment with cinacalcet, all of them predictive for higher tendency to adynamic bone disease [7]. Type 2 DM nephropathy patients with or without diagnosis of CAD also did not differ in tested genotype distributions.

Development of ESRD substantially ameliorates interpatient clinical variability related to underlying renal impairment and exposes uremia-related signs and symptoms. Comparisons of type 2 DM nephropathy patients in respect of tested genotype frequencies with subjects showing other common causes of ESRD revealed that the former group has a higher *IL18* rs360719 minor allele frequency than chronic infective tubulointerstitial nephritic group. In this case, lower *IL18* rs360719 minor allele frequency in tubulointerstitial nephritic patients was observed also when their results were compared to those of healthy subjects. Sánchez et al. [52] have found a significant increase in the relative expression of *IL-18* mRNA in individuals carrying the rs360719 minor allele. *IL-18* is IFN- γ inducing factor. Infective tubulointerstitial nephritic patients are known to have diminished ability of blood leukocytes to produce IFN- γ [53]. Our study indicates that this may be related to lower frequency of *IL18* rs360719 minor allele in this group compared to controls and type 2 DM nephropathy patients. In type 2 DM patients with overt nephropathy, positive correlations between plasma IFN- γ , proteinuria, and eGFR were found [54].

Due to limited financial support, we did not perform any functional studies regarding T-cell-related interleukin and vitamin D pathway genes, especially that multiple influences independent or dependent on genetic profile need to be taken into account in such studies conducted in the uremic milieu. Although the examined patients showing ESRD due to type 2 DM nephropathy were well-defined group, they obviously

were not consistent in HLA DRB1 alleles. The latter could be important in modulating susceptibility to advanced type 2 DM nephropathy and related complications, like it was shown for type 1 DM [55] or type 2 DM [41], regardless of their complications.

5. Summary

Distributions of tested T-cell cytokine gene polymorphisms or vitamin D pathway gene polymorphisms are not significantly different among patients with ESRD due to type 2 DM nephropathy and healthy individuals. Subjects with ESRD due to type 2 DM nephropathy differ in clinical manifestation from patients with other nephropathies leading to dialysis dependency, but differences in tested genotype distributions were found only in *IL18* rs360719 compared with chronic tubulointerstitial nephritic patients. This difference probably arose from the fact that pathology of chronic infective tubulointerstitial nephritis might have been associated with this specific polymorphism.

6. Conclusions

In Polish HD patients, T-cell cytokine gene polymorphisms and vitamin D pathway gene polymorphisms are not associated with ESRD due to type 2 DM nephropathy. *IL18* polymorphism is worthy to be further investigated in chronic infective tubulointerstitial nephritic patients as being possibly associated with this disease.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] E. Villar, H. C. Sean, and S. P. McDonald, "Incidences, treatments, outcomes, and sex effect on survival in patients with end-stage renal disease by diabetes status in Australia and New Zealand (1991–2005)," *Diabetes Care*, vol. 30, no. 12, pp. 3070–3076, 2007.
- [2] A. Sattar, C. Argyropoulos, L. Weissfeld et al., "All-cause and cause-specific mortality associated with diabetes in prevalent hemodialysis patients," *BMC Nephrology*, vol. 13, article 130, 2012.
- [3] F. Chantrel, I. Enache, M. Bouiller et al., "Abysmal prognosis of patients with type 2 diabetes entering dialysis," *Nephrology Dialysis Transplantation*, vol. 14, no. 1, pp. 129–136, 1999.
- [4] H. Al-Thani, A. Shabana, A. Hussein et al., "Cardiovascular complications in diabetic patients undergoing regular hemodialysis: a 5-year observational study," *Angiology*, 2014.
- [5] M. J. Sarnak and B. L. Jaber, "Mortality caused by sepsis in patients with end-stage renal disease compared with the general population," *Kidney International*, vol. 58, no. 4, pp. 1758–1764, 2000.
- [6] S.-M. Alavian and S. V. Tabatabaei, "The effect of diabetes mellitus on immunological response to hepatitis B virus vaccine in individuals with chronic kidney disease: a meta-analysis of current literature," *Vaccine*, vol. 28, no. 22, pp. 3773–3777, 2010.

- [7] D. Zayour, M. Daouk, W. Medawar, M. Salamoun, and G. El-Hajj Fuleihan, "Predictors of bone mineral density in patients on hemodialysis," *Transplantation Proceedings*, vol. 36, no. 5, pp. 1297–1301, 2004.
- [8] D. M. Li, Y. Zhang, B. Ding et al., "The association between vitamin D deficiency and diabetic nephropathy in type 2 diabetic patients," *Zhonghua Nei Ke Za Zhi*, vol. 52, no. 11, pp. 970–974, 2013 (Chinese).
- [9] C.-C. Wu, H.-K. Sytwu, K.-C. Lu, and Y.-F. Lin, "Role of T cells in type 2 diabetic nephropathy," *Experimental Diabetes Research*, vol. 2011, Article ID 514738, 9 pages, 2011.
- [10] Y. Tian, C. Wang, Z. Ye, X. Xiao, A. Kijlstra, and P. Yang, "Effect of 1,25-Dihydroxyvitamin D3 on Th17 and Th1 response in patients with Behçet's disease," *Investigative Ophthalmology and Visual Science*, vol. 53, no. 10, pp. 6434–6441, 2012.
- [11] J. Stachowski, "Hepatitis C virus infection in renal diseases: state of knowledge, therapeutic problems and perspectives," *Polski Merkuriusz Lekarski*, vol. 8, no. 46, pp. 303–306, 2000 (Polish).
- [12] B. D. Livingston, J. Alexander, C. Crimi et al., "Altered helper T lymphocyte function associated with chronic hepatitis B virus infection and its role in response to therapeutic vaccination in humans," *Journal of Immunology*, vol. 162, no. 5, pp. 3088–3095, 1999.
- [13] J. Zhang, G. Hua, X. Zhang, R. Tong, X. Du, and Z. Li, "Regulatory T cells/T-helper cell 17 functional imbalance in uraemic patients on maintenance haemodialysis: a pivotal link between microinflammation and adverse cardiovascular events," *Nephrology*, vol. 15, no. 1, pp. 33–41, 2010.
- [14] E. Borella, G. Neshet, E. Israeli, and Y. Shoenfeld, "Vitamin D: a new anti-infective agent?" *Annals of the New York Academy of Sciences*, vol. 1317, no. 1, pp. 76–83, 2014.
- [15] E. Zitt, H. Sprenger-Mähr, F. Knoll, U. Neyer, and K. Lhotta, "Vitamin D deficiency is associated with poor response to active hepatitis B immunisation in patients with chronic kidney disease," *Vaccine*, vol. 30, no. 5, pp. 931–935, 2012.
- [16] T. Shoji and Y. Nishizawa, "Vitamin D and survival of hemodialysis patients," *Clinical calcium*, vol. 14, no. 9, pp. 64–68, 2004 (Japanese).
- [17] L. Steingrimsdottir, O. Gunnarsson, O. S. Indridason, L. Franzson, and G. Sigurdsson, "Relationship between serum parathyroid hormone levels, vitamin D sufficiency, and calcium intake," *The Journal of the American Medical Association*, vol. 294, no. 18, pp. 2336–2341, 2005.
- [18] R. Pacifici, "Role of T cells in the modulation of PTH action: physiological and clinical significance," *Endocrine*, vol. 44, no. 3, pp. 576–582, 2012.
- [19] H. Zhang, J. Wang, B. Yi et al., "BsmI polymorphisms in vitamin D receptor gene are associated with diabetic nephropathy in type 2 diabetes in the Han Chinese population," *Gene*, vol. 495, no. 2, pp. 183–188, 2012.
- [20] N. Mtiraoui, I. Ezzidi, M. Kacem et al., "Predictive value of interleukin-10 promoter genotypes and haplotypes in determining the susceptibility to nephropathy in type 2 diabetes patients," *Diabetes/Metabolism Research and Reviews*, vol. 25, no. 1, pp. 57–63, 2009.
- [21] I. Ezzidi, N. Mtiraoui, M. Kacem et al., "Interleukin-10-592C/A, -819C/T and -1082A/G promoter variants affect the susceptibility to nephropathy in Tunisian type 2 diabetes (T2DM) patients," *Clinical Endocrinology*, vol. 70, no. 3, pp. 401–407, 2009.
- [22] M. Karadeniz, M. Erdogan, A. Berdeli, and C. Yilmaz, "Association of interleukin-6 -174 G>C promoter polymorphism with increased risk of type 2 diabetes mellitus patients with diabetic nephropathy in Turkey," *Genetic Testing and Molecular Biomarkers*, vol. 18, no. 1, pp. 62–65, 2014.
- [23] M. Murea, T. C. Register, J. Divers et al., "Relationships between serum MCP-1 and subclinical kidney disease: African American-Diabetes Heart Study," *BMC Nephrology*, vol. 13, no. 1, article 148, 2012.
- [24] A. E. Grzegorzewska, D. Pajzderski, A. Sowińska, and P. P. Jagodziński, "Polymorphism of monocyte chemoattractant protein 1 (MCP-1-2518 A/G) and responsiveness to hepatitis B vaccination in hemodialysis patients," *Polskie Archiwum Medycyny Wewnętrznej*, vol. 124, no. 1-2, pp. 10–18, 2014.
- [25] A. E. Grzegorzewska, D. Pajzderski, A. Sowińska, and P. P. Jagodziński, "Monocyte chemoattractant protein-1 gene (MCP-1-2518 A/G) polymorphism and serological markers of hepatitis B virus infection in hemodialysis patients," *Medical Science Monitor*, vol. 20, pp. 1101–1116, 2014.
- [26] Group KDIGO, "KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease," *Kidney International Supplements*, vol. 3, pp. 1–150, 2013.
- [27] A. E. Grzegorzewska, P. Wobszal, and P. P. Jagodziński, "Interleukin-18 promoter polymorphism and development of antibodies to surface antigen of hepatitis B virus in hemodialysis patients," *Kidney and Blood Pressure Research*, vol. 35, no. 1, pp. 1–8, 2012.
- [28] A. E. Grzegorzewska, P. M. Wobszal, A. Sowińska, A. Mostowska, and P. P. Jagodziński, "Association of the interleukin-12 polymorphic variants with the development of antibodies to surface antigen of hepatitis B virus in hemodialysis patients in response to vaccination or infection," *Molecular Biology Reports*, vol. 40, no. 12, pp. 6899–6911, 2013.
- [29] A. E. Grzegorzewska, P. M. Wobszal, A. Mostowska, and P. P. Jagodziński, "Antibodies to hepatitis B virus surface antigen and interleukin 12 and interleukin 18 gene polymorphisms in hemodialysis patients," *BMC Nephrology*, vol. 13, no. 1, article 75, 2012.
- [30] A. E. Grzegorzewska, D. Pajzderski, A. Sowińska, A. Mostowska, and P. P. Jagodziński, "IL4R and IL13 polymorphic variants and development of antibodies to surface antigen of hepatitis B virus in hemodialysis patients in response to HBV vaccination or infection," *Vaccine*, vol. 31, no. 14, pp. 1766–1770, 2013.
- [31] S. S. Rich, "Genetics of diabetes and its complications," *Journal of the American Society of Nephrology*, vol. 17, no. 2, pp. 353–360, 2006.
- [32] N. D. Palmer and B. I. Freedman, "Insights into the genetic architecture of diabetic nephropathy," *Current Diabetes Reports*, vol. 12, no. 4, pp. 423–431, 2012.
- [33] N. Franceschini, N. M. Shara, H. Wang et al., "The association of genetic variants of type 2 diabetes with kidney function," *Kidney International*, vol. 82, no. 2, pp. 220–225, 2012.
- [34] N. D. Palmer, C. W. McDonough, P. J. Hicks et al., "A genome-wide association search for type 2 diabetes genes in african americans," *PLoS ONE*, vol. 7, no. 1, Article ID e29202, 2012.
- [35] A. Davoodi-Semiromi, J. J. Yang, and J.-X. She, "IL-12p40 is associated with type 1 diabetes in Caucasian-American families," *Diabetes*, vol. 51, no. 7, pp. 2334–2336, 2002.
- [36] D. B. Mirel, A. M. Valdes, L. C. Lazzaroni, R. L. Reynolds, H. A. Erlich, and J. A. Noble, "Association of IL4R haplotypes with type 1 diabetes," *Diabetes*, vol. 51, no. 11, pp. 3336–3341, 2002.
- [37] T. L. Bugawan, D. B. Mirel, A. M. Valdes, A. Panelo, P. Pozzilli, and H. A. Erlich, "Association and interaction of the IL4R,

- IL4, and IL13 loci with type 1 diabetes among filipinos," *The American Journal of Human Genetics*, vol. 72, no. 6, pp. 1505–1514, 2003.
- [38] B. Frederiksen, E. Liu, J. Romanos et al., "Investigation of the vitamin D receptor gene (VDR) and its interaction with protein tyrosine phosphatase, non-receptor type 2 gene (PTPN2) on risk of islet autoimmunity and type 1 diabetes: the Diabetes Autoimmunity Study in the Young (DAISY)," *The Journal of Steroid Biochemistry and Molecular Biology*, vol. 133, no. 1, pp. 51–57, 2013.
- [39] C. Panierakis, G. Goulielmos, D. Mamoulakis, E. Petraki, E. Papavasiliou, and E. Galanakis, "Vitamin D receptor gene polymorphisms and susceptibility to type 1 diabetes in Crete, Greece," *Clinical Immunology*, vol. 133, no. 2, pp. 276–281, 2009.
- [40] Q. Wang, B. Xi, K. H. Reilly, M. Liu, and M. Fu, "Quantitative assessment of the associations between four polymorphisms (FokI, ApaI, BsmI, TaqI) of vitamin D receptor gene and risk of diabetes mellitus," *Molecular Biology Reports*, vol. 39, no. 10, pp. 9405–9414, 2012.
- [41] N. M. Al-Daghri, O. Al-Attas, M. S. Alokail et al., "Vitamin D receptor gene polymorphisms and HLA DRB1*04 cosegregation in Saudi type 2 diabetes patients," *Journal of Immunology*, vol. 188, no. 3, pp. 1325–1332, 2012.
- [42] F.-L. Vélazoudom-Céphise, L. Larifla, J.-P. Donnet et al., "Vitamin D deficiency, vitamin D receptor gene polymorphisms and cardiovascular risk factors in Caribbean patients with type 2 diabetes," *Diabetes & Metabolism*, vol. 37, no. 6, pp. 540–545, 2011.
- [43] D. A. F. Ferrarezi, N. Bellili-Muñoz, D. Dubois-Laforgue et al., "Allelic variations of the vitamin D receptor (VDR) gene are associated with increased risk of coronary artery disease in type 2 diabetics: the DIABHYCAR prospective study," *Diabetes and Metabolism*, vol. 39, no. 3, pp. 263–270, 2013.
- [44] T. B. Zhou, Z. P. Jiang, M. F. Huang, and N. Su, "Association of vitamin D receptor FokI (rs2228570), TaqI (rs731236) and ApaI (rs7975232) gene polymorphism with the risk of chronic kidney disease," *Journal of Receptors and Signal Transduction Research*, vol. 5, pp. 1–5, 2014.
- [45] R. D. Mittal and P. K. Manchanda, "Association of interleukin (IL)-4 intron-3 and IL-6 -174 G/C gene polymorphism with susceptibility to end-stage renal disease," *Immunogenetics*, vol. 59, no. 2, pp. 159–165, 2007.
- [46] G. Morahan, E. McKinnon, J. Berry et al., "Evaluation of IL12B as a candidate type I diabetes susceptibility gene using data from the Type I Diabetes Genetics Consortium," *Genes and Immunity*, vol. 10, supplement 1, pp. S64–S68, 2009.
- [47] H. A. Erlich, K. Lohman, S. J. MacK et al., "Association analysis of SNPs in the IL4R locus with type i diabetes," *Genes and Immunity*, vol. 10, supplement 1, pp. S33–S41, 2009.
- [48] R. Jorde, H. Schirmer, T. Wilsgaard et al., "Polymorphisms related to the serum 25-Hydroxyvitamin D level and risk of Myocardial infarction, diabetes, cancer and mortality. The Tromsø study," *PLoS ONE*, vol. 7, no. 5, Article ID e37295, 2012.
- [49] H. Bid, R. Konwar, C. Aggarwal et al., "Vitamin D receptor (FokI, BsmI and TaqI) gene polymorphisms and type 2 diabetes mellitus: a North Indian study," *Indian Journal of Medical Sciences*, vol. 63, no. 5, pp. 187–194, 2009.
- [50] M. C. Lemos, A. Fagulha, E. Coutinho et al., "Lack of association of vitamin D receptor gene polymorphisms with susceptibility to type 1 diabetes mellitus in the Portuguese population," *Human Immunology*, vol. 69, no. 2, pp. 134–138, 2008.
- [51] T. B. Zhou, Z. P. Jiang, and M. F. Huang, "Association of vitamin D receptor BsmI (rs1544410) gene polymorphism with the chronic kidney disease susceptibility," *Journal of Receptors and Signal Transduction Research*, 2014.
- [52] E. Sánchez, R. J. Palomino-Morales, N. Ortego-Centeno et al., "Identification of a new putative functional IL18 gene variant through an association study in systemic lupus erythematosus," *Human Molecular Genetics*, vol. 18, no. 19, pp. 3739–3748, 2009.
- [53] I. P. Kudriashova, T. P. Ospel'nikova, and F. I. Ershov, "Cycloferon administration in chronic pyelonephritis: changes in interferon status," *Terapevticheskiĭ Arkhiv*, vol. 83, no. 6, pp. 33–35, 2011 (Russian).
- [54] C.-C. Wu, J.-S. Chen, K.-C. Lu et al., "Aberrant cytokines/chemokines production correlate with proteinuria in patients with overt diabetic nephropathy," *Clinica Chimica Acta*, vol. 411, no. 9–10, pp. 700–704, 2010.
- [55] N. Israni, R. Goswami, A. Kumar, and R. Rani, "Interaction of Vitamin D receptor with HLA DRB1*0301 in Type 1 diabetes patients from North India," *PLoS ONE*, vol. 4, no. 12, Article ID e8023, 2009.