Review Article

Nonhuman Primate Models of Type 1 Diabetes Mellitus for Islet Transplantation

Haitao Zhu,¹ Liang Yu,¹ Yayi He,² and Bo Wang¹

¹ Department of Hepatobiliary Surgery, First Affiliated Hospital, Medical College, Xi'an Jiaotong University, Xi'an 710061, China

² Department of Endocrinology, First Affiliated Hospital, Medical College, Xi'an Jiaotong University, Xi'an 710061, China

Correspondence should be addressed to Bo Wang; bobwang75@yeah.net

Received 13 August 2014; Revised 2 October 2014; Accepted 7 October 2014; Published 20 October 2014

Academic Editor: Subrata Chakrabarti

Copyright © 2014 Haitao Zhu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Islet transplantation is an attractive treatment of type 1 diabetes mellitus (T1DM). Animal models of diabetes mellitus (DM) contribute a lot to the experimental studies of islet transplantation and to evaluations of isolated islet grafts for future clinical applications. Diabetic nonhuman primates (NHPs) represent the suitable models of DMs to better evaluate the effectiveness of islet transplantation, to assess new strategies for controlling blood glucose (BG), relieving immune rejection, or prolonging islet survival, and eventually to translate the preclinical data into tangible clinical practice. This review introduces some NHP models of DM, clarifies why and how the models should be used, and elucidates the usefulness and limitations of the models in islet transplantation.

1. Introduction

Type 1 diabetes mellitus (T1DM), once known as insulin-dependent diabetes mellitus (IDDM), is an autoimmune disorder caused by progressive destruction of insulin-producing pancreatic β -cell that results in hyperglycemia [1, 2]. Curing patients with T1DM requires both ablation of β -cell-specific autoimmune reaction and β -cell replacement therapy [3]. At present, pancreatic islet transplantation is a promising and minimally invasive treatment that has the great potential to restore normoglycemia and achieve complete independence from exogenous insulin in T1DM patients [3–6]. Since Ballinger's and colleagues first demonstrated that islet cells isolated from rat pancreata could be transplanted into a diabetic rat model to reverse hyperglycemia, scholars pursued efforts to develop this treatment for clinical application [7, 8]. In the year 2000, by using the "Edmonton Protocol" based on a glucocorticoid-free immunosuppressive regimen, Shaprio et al. achieved insulin independence with an excellent metabolic control in diabetic patients after allotransplantation of islet mass [9]. However, due to shortage of human donors, another islet source such as from xenogenic pig or stem-cell derived functional islet emerges as an alternative

strategy for transplantation [10]. Recently, several trials demonstrated the possibility of achieving sustained survival of xenogenic islet grafts in diabetic rodents or primates [11–16]; and stem-cell therapy also promised a reproducible and nearly unlimited supply of transplantable islets [17–19]. All these steady and great progresses made in the islet transplantation field likely depend on improvements in techniques for high-quality islet preparation [20–23], strategies for immune suppression or tolerance [5, 24, 25], and protocols for differentiation of stem-cell into functional β -cell [26]. However, animal models of diabetes mellitus (DM) also contribute a lot to islet research and to evaluation of isolated islet grafts for clinical applications [27, 28]. Moreover, to better investigate the achieved findings and to assess new protocols for relieving immune rejection and drugs/devices for prolonging islet survival, experimental research in appropriate animal models is essential. Hence, rigorous trial test of the islet grafts from various sources on preclinical large animal models of DM represents an important step to better verify the effectiveness of islet transplantation and develop novel approaches for T1DM treatment.

In large animals, diabetic nonhuman primates (NHPs) are considered as the good models for translating obtained

islet transplant research safely into tangible clinical applications due to advantages including (1) evolutionary proximity to human beings, with approximately 95% of homology at the nucleotide level [29]; (2) similar clinical features of diabetes between NHPs and humans [30–33]; (3) the fact that large body size and long lifespan of NHPs make it possible and convenient to perform longitudinal studies and numerous procedures (e.g., biopsy of pancreatic tissues, extraction of large volumes of blood samples, or surgical implantation of catheters) [34]; and the fact that (4) immune system of NHPs closely resembles that of humans, making it possible and useful to study immunological aspects related with islet transplantation, as well as to develop and verify strategies to improve islet graft survival for future clinical practice. Besides, another important requirement for preclinical test in NHPs is based on the indefinite translation of rodent protocols to outbred NHPs. Even though numerous protocols are effectively applied to induce immune tolerance in rodent models, only a small amount of strategies can prolong rejection-free interval (RFI) survival without sustained severe immunosuppressive treatment in NHP models [35].

Currently, the commonly used NHP animals are baboons and macaques (rhesus and cynomolgus monkeys). Nevertheless, there are persistent hurdles to the advancement and large-scale application of these models in islet transplantation. In this review, we introduce some NHP models of DM, clarify why and how the models should be used, and elucidate the usefulness and limitations of the models in islet transplantation.

2. Housing and Feeding of NHPs

For promoting animal welfare and improving reliability and quality of research data in islet transplantation, academics recommend adaptations of housing and nursing care of NHP models of DM in the experimental animal center or laboratory [36]. Prior to experimental process, NHPs should be socially kept (if possible, that is, in a naturally composed breeding group, or in a peer group or same-sex group, or as a pair) in enriched laboratory conditions with (1) enhanced housing (a cage with a perch and bedding, with minimal height of 1.50 m for monkeys and 2.50 m for chimpanzees), (2) controlled temperature and light-dark cycle, (3) regular standard chow (at least twice a day) and supplementary feed (e.g., fresh vegetables and fruits; daily or weekly), (4) documented health evaluation, and (5) positive reinforcement training [37–39]. Through training, the diabetic NHPs may be able to offer a body part for blood glucose monitoring, subcutaneous or intramuscular injection, and blood collection; some primates may cooperate with intragastric administration of immunosuppressants after islet transplantation.

3. Species Selection

Macaque monkeys, especially cynomolgus monkey (*Macaca fascicularis*) and rhesus macaque (*Macaca mulatta*), have become the major species of choice for DM induction in

the field of preclinical islet transplantation [25, 40–42] with several superior characteristics of relatively easy upkeep in captivity, wide availability, polymorphism of the major histocompatibility complex (MHC), anatomical and physiological similarity of humans, and resemblance in immune systems [34, 43–45]. More importantly, these species are usually smaller than baboons and chimpanzees, only 5–9 kilograms of males and 3–6 kilograms of females [37], and therefore have a clear superiority of much less expensive housing, care, feeding, and charges for immunosuppressive regimens. Besides that, fewer islet grafts are required to be implanted to achieve insulin independence in these diabetic recipients. Baboons and chimpanzees, although used less often [46], are likely more robust than macaques, potentially rendering them less prone to surgical and other complications [47].

4. Induction of DM in NHPs for Islet Transplantation

Animal experiments have enormously contributed to the study of DM, as well as islet research and transplantation. Generally, diabetic animals are classified as spontaneous and secondary (pancreatized, chemical-induced, or gene transfection) models. However, up to now, no exact recommendation has been reached for a standardized method of DM induction.

Because spontaneous T1DM occurs in an extremely low incidence in NHPs [30], in the area of islet transplantation, DMs are usually induced in NHPs by pancreatectomy or/and by systemic administration of streptozotocin (STZ) [25, 36, 46]. Although alloxan has been reported to chemically induce a diabetic state in NHPs [34], it is generally acknowledged that this diabetogenic agent works selectively on β -cells of rodents and is therefore not an available and potent inducer of DM for NHPs [48–50]. Additionally, genetically induced models of DM are mainly performed in the rodent animals, such as Ins2Akita mice [51], human islet amyloid polypeptide (hIAPP) mice [52], and humanized mice with aspects of human immune system [53]. By contrast, the application of transgenic NHPs in biomedical research is just at infancy stage [54]. Currently, Huntington's disease (HD) monkey is the first and only reported transgenic NHP model of human disease [55, 56].

In NHPs, the various modeling methods exhibit different characters in pathophysiological mechanism, progression, and complications of the DM. In particular, the severity, stability, and extent of metabolic and endocrine abnormalities vary according to the induction protocols, primate species, and individual differences in monkeys.

4.1. Surgical Pancreatectomy. One of the most straightforward means of examining the effect of hyperglycemia in animal is to remove the pancreas, either partially or totally [57]. Pancreatized animals, which have more than 100 y of history, are often thought of as the earliest models of DM. In the 1880s, in order to observe the intestinal absorptive function, von Mering and his colleagues excised the pancreas of a dog, and then this animal developed symptoms

of polyuria and polydipsia and was eventually diagnosed as DM [57]. The extensive studies on pancreatectomized NHPs were first conducted by Gillman et al. in the 1960s [58, 59]. After pancreatectomy, hyperglycemia (hallmark of DM), polydipsia, lipaemia, and ketonaemia occurred in the baboons (*Papio ursinus*). In general, the pancreatectomized model is considered to be and utilized as a T1DM model on account of full depletion of the islet β cells.

Total pancreatectomy (TP) is a commonly used procedure to induce IDDM in NHPs (e.g., cynomolgus monkey, rhesus monkey, and baboon) in the preclinical studies of islet autotransplantation, as well as allotransplantation [60–65] and xenotransplantation [66, 67]. The resected pancreas can be utilized for islet cell isolation and preparation. Compared with STZ administration, TP induces IDDM more effectively, reliably, and permanently [68]. However, the complete deficiency of both pancreatic endocrine and exocrine functions also results in high probability of hypoglycemia (blood glucose < 2.8 mmol/L), unstable glycemic control, and loss of other islet hormones [69]. In view of the pancreatic function, compensatory response, and regeneration of islet mass, more than 90% of pancreas should be removed to induce stable DM in a rhesus monkey [64, 70]. Therefore, this invasive surgical procedure is more complex and requires precise manipulation. For successfully performing TP to induce DM in NHP and enabling the recipient monkey to undergo islet transplantation safely, special attention should be paid to the following points. (1) Vital signs should be monitored and maintained in a safe range during the operations: heart rate (80–160/min), systolic blood pressure (60–120 mmHg), oxygen saturation (>96%), and body temperature (35–37°C) [62]. (2) Splenic artery and vein, portal and superior mesenteric vein, arterial arcades around the duodenum, inferior mesenteric and middle colic veins, and common bile duct must be carefully preserved [64, 68, 71]. (3) Appropriate glycemic control should be performed after TP. Generally, blood glucose (BG) levels are controlled in the 100–300 mg/dL range with exogenous insulin administration before islet implantation [60–62]. (4) Controlling the fasting BG at 5–10 mmol/L and the glycosylated hemoglobin (HbA_{1c}) at 3%–6.5% is a suitable strategy to induce a lower probability of hypoglycemia [69]. (5) Intensive postoperative care and management must be carried out timely to avoid complications. (6) TP and islet implantation should not be conducted concurrently but consecutively as a 2-stage surgery. Fourteen days after TP is recommended as the optimal time for islet infusion [62].

4.2. STZ Administration. STZ, a naturally occurring chemical (molecular formula: C₈H₁₅N₃O₇, molecular weight: 265.2 g/mol) derived from the soil microbe *Streptomyces achromogenes*, has been used clinically for treating metastatic pancreatic islet cell carcinomas and used investigatively for inducing experimental DM in animals [27, 72–74]. This diabetogenic drug can inhibit insulin secretion and induce a state of IDDM through its ability to result in a selective death of pancreatic β -cells. The cell toxicity of STZ is mainly dependent on its methylation activities leading to DNA damages. After injection, STZ is selectively taken up by islet cells via the

glucose transporter 2 (GLUT2). And then, this glucosamine-nitrosourea compound leads to DNA breaks and subsequently activates poly-ADP-ribose polymerase (PARP) that result in depletion of cellular nicotinamide adenine dinucleotide (NAD⁺) and decline of subphysiological adenosine triphosphate (ATP) [72, 75]. The loss of cellular energy stores eventually causes cell death. In addition, other cells expressing the transporter GLUT2 such as hepatocytes and renal tubular cells are also susceptible to STZ; thus, the major disadvantages of STZ are hepatotoxicity and nephrotoxicity, particularly at high dose.

Compared with TP, pharmacologically induced IDDM using STZ is far more common, more convenient, and less invasive. At present, this nonsurgical approach is widely used to induce a reproducible form of DM in NHPs [36]. This is illustrated by the reports of Emory Transplant Center, in which TP in previous studies [66, 76] was replaced by STZ injection in recent studies [13, 77]. Following STZ administration, monkeys developed IDDM within 24 h, with elevated fasting BG levels (>400 mg/dL), decreased serum C-P levels (0.01–0.6 ng/mL), and loss of body weight. Even more importantly, the diabetic monkeys could remain the diabetic state up to 1 y with persistent low C-P levels (<0.6 ng/mL), no C-P release during intravenous glucose tolerance test (IVGTT), and pancreatic histology that revealed only small numbers of insulin-staining cells [78]. Although Bottino et al. reported that, in cynomolgus monkeys, pancreas could regain endogenous C-P production after complete β -cell destruction by high-dose STZ injection (125–150 mg/kg body weight), it seemed to be a nongeneralizable and incidental event with only 2 out of 11 diabetic monkeys recovering β -cell function [79].

Usually, for successful STZ administration in experimental NHPs and for optimum results, some recommendations should be followed: (1) NHPs must be fasted overnight (8–12 h) before STZ injection. (2) Owing to STZ's instability in solution, STZ should be freshly prepared, dissolved in citrate buffer (pH 4.4–4.5), stored on ice, and administered immediately within 10 min [75, 80] (Table 1). (3) The NHPs can receive antiemetic prophylaxis about 30–60 min prior to STZ injection to suppress postinfusion nausea and vomiting [81]. (4) Since obesity is a relevant risk factor for adverse events, it is suggested to preferentially select young monkeys or adult monkeys with lower girth-to-height ratio (GHtR, obesity indicator) for DM induction. In adult monkeys with a higher than normal GHtR, researchers should lower the STZ dosage by as much as 20% [81]. (5) After STZ injection, aggressive fluid administration (normal saline, dose: 20–30 mL/kg, rate: 1.0 mL/min) is helpful to reduce morbidity and eliminate mortality in diabetic monkeys [81, 82]. (6) Exogenous insulin is administered to maintain BG level < 300 mg/dL (maximum) in diabetic monkeys before transplant and following islet graft rejection [82–86]. (7) Since cyclosporine can facilitate renal dysfunction in STZ-induced diabetic monkeys [87], an intensive and careful adverse event monitoring (AEM) should be conducted as soon as immunosuppressive treatment starts in the islet transplant recipients.

TABLE 1: DM induction in NHPs by STZ administration.

Author and publication year	Animals tested (gender, age, and BW)	STZ dose	Solvent	Route of administration	Results
Theriault et al., 1999 [82]	Cyno (6–8 months, 1.0–2.0 kg), $n = 11$	150 mg/kg	0.9% NS	via SVAP	All persistently hyperglycemic (BG > 200 mg/dL; 9/11 serum C-P < 0.5 ng/mL, 60 days; 2/11 C-P at 0.9 ng/mL, 159 days); all without any observable clinical complications except transient nausea and vomiting
Shibata et al., 2002 [92]	Rhesus (female, 2.5–7.5 years, 4.2–6.8 kg)	STZ (Upjohn) G1: 100 mg/kg ($n = 2$); G2: 125 mg/kg ($n = 4$); G3: 150 mg/kg ($n = 7$)	Citric acid (pH 4.5)	via CVC	G1: none diabetic (unchanged basal insulin, N_{IPC} , in islets was 77% of normal values); G2: all diabetic (IDDM) without mortality and hepatic/renal injury (N_{IPC} was 0.3% of normal values); G3: 2/7 dead (28.6% mortality), 5/7 diabetic (up to 3 weeks, no IPCs)
Koulmanda et al., 2003 [78]	Cyno (male, 3–6 years, 3.1–9.0 kg)	G1: 100 mg/kg ($n = 4$); G2: 55 mg/kg ($n = 20$)	0.9% NS	i.v.	All groups diabetic (IDDM, BG > 400 mg/dL, Serum C-P < 0.6 ng/mL, loss of BW, decreased N_{IPC}); G1: impaired liver/kidney functions, severe hepatic steatosis, and acute renal tubular injury; G2: normal liver/kidney functions, no histological changes
Tal et al., 2004 [95]	Cyno (4–6 years, $n = 6$); Rhesus (8–12 years, $n = 3$); Pigtail (2 years, $n = 5$)	STZ (Zanosar) 50–70 mg/kg	5% dextrose	Selective arterial injection	All diabetic (BG > 200 mg/dL, complete obliteration of β -cell, 20–270 days of duration); 1/14 dead (gastric dilatation); no abnormalities in liver and kidney
Rood et al., 2006 [91]	Cyno (male, 2–3 years, 2.7–4.1 kg)	G1: STZ (Sigma), 1250 mg/m ² ($n = 4$); G2: STZ (Sigma), 60 mg/kg ($n = 1$); G3: STZ (Zanosar), 150 mg/kg ($n = 5$)	NM	i.v.	G1: all completely diabetic (BG > 200 mg/dL, C-P < 0.5 ng/mL, low N_{IPC}), with several AEs (vomiting, protein-losing nephropathy, and low ALB levels); G2: failed to lose complete β -cell function (C-P at 1.86 ng/mL), no AEs; G3: all completely diabetic (BG > 200 mg/dL, C-P < 0.5 ng/mL, low or no N_{IPC}), with only transient vomiting
He et al., 2011 [69]	Rhesus (3–6 years)	G1: single STZ ad. (80 mg/kg), $n = 7$; G2: multiple STZ ad. (25 mg/kg for 5 consecutive days), $n = 9$	NM	i.v.	All groups diabetic; G1: lower C-P levels (<0.1 nmol/L) and probability of hypoglycemia (0.41); G2: relatively higher C-P levels (<0.5 nmol/L) and probability of hypoglycemia (0.80)
Zou et al., 2012 [96]	Cyno (male, 2.4–4.7 kg)	STZ (Sigma) G1: 100 mg/kg ($n = 7$); G2: 68 mg/kg ($n = 6$); G3: 60 mg/kg ($n = 2$)	Citric acid (pH 4.5)	i.v. (via saphenous vein)	G1 and G2: all diabetic (mean fasting BG > 22 mmol/L, low N_{IPC} , >8 months of duration), without any marked hepatic/renal injury; G3: one diabetic (fasting BG at 26.4 mmol/L, C-P at 28 pmol/L); the other nondiabetic (fasting BG at 5.7 mmol/L, C-P at 413 pmol/L)

Overview of published studies of STZ-induced diabetic models in NHPs.

NM: not mentioned; DM: diabetes mellitus; NHP: nonhuman primate; STZ: streptozotocin; BW: body weight; G: group; Cyno: cynomolgus monkey; Rhesus: rhesus monkey; Pigtail: pigtail macaque; NS: normal saline; SVAP: subcutaneous vascular access ports; CVC: central venous catheter; BG: blood glucose; C-P: c-peptide; IDDM: insulin-dependent diabetes mellitus; N_{IPC} : number of insulin-positive cells; i.v.: intravenous; ALB: albumin; AEs: adverse events; and ad.: administration.

Most importantly, it is necessary to establish the optimal dose of STZ that is required for inducing irreversible and stable DM with less adverse effects before evaluating treatment strategies in islet transplantation. The diabetogenic dose depends on the animal species, age, body weight, route of administration, and nutritional status [27, 75, 88]. Presently, there is still an ongoing debate with regard to the proper dose of STZ in NHPs, and STZ administration for DM induction has plenty of potential variables (Table 1). Administration of a low-dose STZ (20–50 mg/kg body weight) was not sufficient and reliable to consistently induce complete DM (C-P negative DM) in cynomolgus monkeys [89, 90]. Higher doses of STZ (80–150 mg/kg body weight) were found to be effective and sufficient [13, 81, 84, 87, 91–93] but were associated with more systemic side effects (e.g., transient vomiting, severe hypoglycemia) and serious complications (e.g., hepatic and renal function/tissue injury), as well as higher morbidity and mortality (approximately 28.6%–100%) [69, 87, 92–94]. Koulmanda et al. [78], Tal et al. [95], Dufrane et al. [94], and Zou et al. [96] demonstrated that STZ dose of 50–70 mg/kg could induce stable IDDM in all macaques for up to 0.5–1 y without any evidence of regeneration of β -cell, as well as liver and kidney toxicity. On the contrary, Theriault and colleagues reported the safe and consistent induction of IDDM with a single high-dose of STZ (150 mg/kg body weight) in young cynomolgus monkeys (6–8-month-old). Moreover, Wijkstrom et al. [87] recommended that the STZ dose should depend on body surface area (BSA) (1250 mg/m²) rather than body weight (BW). Zou et al. [96] further demonstrated that a dose > 760 mg/m² of STZ might reliably induce IDDM in juvenile cynomolgus monkeys (2–3-year-old), whereas a dose < 700 mg/m² was not effective. One reason for these discrepancies is that the β -cells of younger monkeys may be more resistant to the toxic effect of STZ than that of older ones, and this may account for the varying STZ dosages to induce DM. Another variable that may influence the dosage is the way of STZ administration. Since STZ has a short half-life (5–15 min), higher dose of STZ is necessary if the agents cannot be administered immediately as a bolus. In addition, the individual genetic and physiological differences, variability in handling the STZ compound (i.e., scale difference, different solvent, and degradation due to preparation time and stock condition), and environmental factors (e.g., hydration protocols and subsequent regimens) also contribute to the wide variance in dose level among different research centers.

Nowadays, in preclinical studies of islet transplantation, DM is frequently induced by Zanosar STZ (Teva Pharmaceuticals) with a relatively high dosage (125–150 mg/kg or 1250–1600 mg/m², intravenous injection) in recipient macaques (weight range, 3–5 kg) [13, 77, 83–86, 97]. Zanosar STZ is a clinically used drug with greater purity and less variability and therefore has fewer adverse effects than other preparations of STZ in DM induction [91].

4.3. Partial Pancreatectomy with STZ Administration. Since TP causes relatively high surgical morbidity/mortality [69, 71] and high-dose (100–120 mg/kg body weight) STZ can

induce hepatotoxicity (e.g., hepatic steatosis) and nephrotoxicity (e.g., severe proteinuria, glomerular damage, and acute tubular injury) in diabetic NHPs [78, 81, 91, 93], partial pancreatectomy (PP, approximately 75% pancreatic tissues) combined with low-dose STZ injection (15 mg/kg body weight) (PP-STZ) is considered as an alternative method for IDDM induction [71, 93]. The procedure for PP is similar to that of TP, except that the pancreatic head and uncinate process are retained in order to prevent surgical damage to important blood vessels and bile duct system. The remaining β -cells are then disrupted by low-dose STZ administration to avoid liver and kidney injury in NHPs caused by high-dose STZ. Although TP can lead to much lower concentration of C-peptide (C-P) (<0.1 nmol/L) than PP-STZ, there is no significant difference between the two modeling methods in terms of BG level, HbA_{1c}, and exogenous insulin dosage [69, 71]. Six months after successful induction of DM, the C-P levels in the diabetic monkeys were still <0.5 mmol/L, suggesting that the DM model induced by PP-STZ was stable and credible [71]. Moreover, the PP-STZ has other advantages including stable glycemic control, less STZ-induced organ lesion, low incidence of hypoglycemia, relatively easy surgical operation, reduced complications, favorable postoperative recovery, and no mortality. All these make PP-STZ a better modeling strategy than TP and other methods (high-dose STZ administration).

5. Limitations of NHP Models of DM in Islet Research

On account of similar autoimmune responses that cause islet damage/loss between spontaneous type 1 diabetic animals and human T1DM, these diabetic animals are regarded as useful and reliable models for clearly elucidating the pathogenesis of T1DM, for comprehensively analyzing the induction of immune suppression/tolerance in islet transplantation, and for effectively translating the experimental findings into clinical islet transplantation in diabetic patients. Nevertheless, there is very low incidence of spontaneous T1DM in NHPs (also lower than the incidence of spontaneous type 2 diabetes), and the time course of diabetes progression is extensive too [30]. Nowadays, the most commonly used spontaneous type 1 diabetic animals are biobreeding (BB) rat and nonobese diabetic (NOD) mice [98, 99]. The autoimmunity of NOD mice is characterized by several abnormalities in the immune system including the presence of pancreas-specific autoantibodies, autoreactive CD4⁺ and CD8⁺ T cells, and defects in T regulatory cells [100, 101], which also appear in T1DM patients and rarely exist in surgical/STZ-induced NHP models of IDDM. In order to successfully perform clinical islet transplantation, it is essential to control the allo-, xeno-, and autoimmunity. The NOD mice represent the suitable models for dissecting tolerance mechanisms and assessing influences of different types of immunity in islet transplantation [102–105]. And obtained experimental results are the important improvements and supplements to the preclinical findings in diabetic NHPs. Finally, the combined data will roundly evaluate the immunosuppressive strategies

and further promote immune tolerance in the future clinical islet transplantation.

Additionally, in order to clarify and improve the translational value of preclinical trials of islet transplantation, other intrinsic limitations in the NHP models of DM should also be considered, especially in pig-to-NHPs islet xenotransplantation [106]. (1) Monkeys maintain a lower set point for normoglycemia but higher insulin and C-P secretion than pigs and humans. Given the differences in glucose metabolism, it is harder to achieve stable glucose homeostasis in pig-to-NHP islet xenotransplantation [107–109]. (2) The therapeutic window of some of the commonly used immunosuppressants (e.g., cyclosporine, tacrolimus, and rapamycin) is much smaller in NHPs than in humans [109], possibly resulting in more serious side effects and rendering the NHP models less predictive for the clinical condition. (3) Severe and progressive body weight loss occurs frequently because of gastrointestinal disturbance and diarrhea, particularly if TP is carried out to induce a stable IDDM. The nutritional deficiencies can result in susceptibility to infectious complications, as well as misconception of BG and C-P levels regarding islet graft function [109]. Addressing the problems of diabetic NHP model will eventually lead to optimal experimental designs and proper use of diabetic animals from experimental islet transplantation to translational studies.

6. Conclusions

In summary, NHPs represent useful and suitable diabetic animal models for evaluating the effects of therapeutic interventions on the outcome of preclinical islet transplantation. Although NHP models of DM can be induced by several methods, the revulsive mechanisms are greatly different. Therefore, in islet transplantation, it is necessary to choose a proper NHP model on the basis of the aim and translational value of the preclinical study. Moreover, further exploring and clarifying the characteristics and limitations of diabetic NHP models will significantly draw the obtained preclinical data closer to clinical trials.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

This work was supported by the National Natural Science Foundation of China (no. 30700772).

References

- [1] J. H. Kim, S.-M. Jin, H. S. Kim, K.-A. Kim, and M.-S. Lee, "Immunotherapeutic treatment of autoimmune diabetes," *Critical Reviews in Immunology*, vol. 33, no. 3, pp. 245–281, 2013.
- [2] Å. Lernmark and H. E. Larsson, "Immune therapy in type 1 diabetes mellitus," *Nature Reviews Endocrinology*, vol. 9, no. 2, pp. 92–103, 2013.
- [3] X. Clemente-Casares, S. Tsai, C. Huang, and P. Santamaria, "Antigen-specific therapeutic approaches in type 1 diabetes," *Cold Spring Harbor Perspectives in Medicine*, vol. 2, no. 2, Article ID a007773, 2012.
- [4] P. R. V. Johnson and K. E. Jones, "Pancreatic islet transplantation," *Seminars in Pediatric Surgery*, vol. 21, no. 3, pp. 272–280, 2012.
- [5] M. McCall and A. M. J. Shapiro, "Update on islet transplantation," *Cold Spring Harbor Perspectives in Medicine*, vol. 2, no. 7, Article ID a007823, 2012.
- [6] E. A. Ryan, J. R. T. Lakey, R. V. Rajotte et al., "Clinical outcomes and insulin secretion after islet transplantation with the edmonton protocol," *Diabetes*, vol. 50, no. 4, pp. 710–719, 2001.
- [7] W. F. Ballinger and P. E. Lacy, "Transplantation of intact pancreatic islets in rats," *Surgery*, vol. 72, no. 2, pp. 175–186, 1972.
- [8] C. B. Kemp, M. J. Knight, D. W. Scharp, P. E. Lacy, and W. F. Ballinger, "Transplantation of isolated pancreatic islets into the portal vein of diabetic rats," *Nature*, vol. 244, no. 5416, p. 447, 1973.
- [9] A. M. J. Shapiro, J. R. T. Lakey, E. A. Ryan et al., "Islet transplantation in seven patients with type 1 diabetes mellitus using a glucocorticoid-free immunosuppressive regimen," *The New England Journal of Medicine*, vol. 343, no. 4, pp. 230–238, 2000.
- [10] D. D. Lee, E. Grossman, and A. S. Chong, "Cellular therapies for type 1 diabetes," *Hormone and Metabolic Research*, vol. 40, no. 2, pp. 147–154, 2008.
- [11] D. Dufrane, R.-M. Goebbels, and P. Gianello, "Alginate macroencapsulation of pig islets allows correction of streptozotocin-induced diabetes in primates up to 6 months without immunosuppression," *Transplantation*, vol. 90, no. 10, pp. 1054–1062, 2010.
- [12] W. Wang, Z. Mo, B. Ye, P. Hu, S. Liu, and S. Yi, "A clinical trial of xenotransplantation of neonatal pig islets for diabetic patients," *Zhong Nan Da Xue Xue Bao Yi Xue Ban*, vol. 36, no. 12, pp. 1134–1140, 2011.
- [13] P. Thompson, I. R. Badell, M. Lowe et al., "Alternative immunomodulatory strategies for xenotransplantation: CD40/154 pathway-sparing regimens promote xenograft survival," *The American Journal of Transplantation*, vol. 12, no. 7, pp. 1765–1775, 2012.
- [14] E. Ashkenazi, B. M. Baranovski, G. Shahaf, and E. C. Lewis, "Pancreatic islet xenograft survival in mice is extended by a combination of alpha-1-antitrypsin and single-dose anti-CD4/CD8 therapy," *PLoS ONE*, vol. 8, no. 5, Article ID e63625, 2013.
- [15] H. Arefanian, E. B. Tredget, R. V. Rajotte, R. G. Gill, G. S. Korbitt, and G. R. Rayat, "Short-term administrations of a combination of anti-LFA-1 and anti-CD154 monoclonal antibodies induce tolerance to neonatal porcine islet xenografts in mice," *Diabetes*, vol. 59, no. 4, pp. 958–966, 2010.
- [16] M. Tian, Y. Lv, C. Zhai, H. Zhu, L. Yu, and B. Wang, "Alternative immunomodulatory strategies for xenotransplantation: CD80/CD86-CTLA4 pathway-modified immature dendritic cells promote xenograft survival," *PLoS ONE*, vol. 8, no. 7, Article ID e69640, 2013.
- [17] J. Stanekzai, E. R. Isenovic, and S. A. Mousa, "Treatment options for diabetes: potential role of stem cells," *Diabetes Research and Clinical Practice*, vol. 98, no. 3, pp. 361–368, 2012.

- [18] N. Lumelsky, O. Blondel, P. Laeng, I. Velasco, R. Ravin, and R. McKay, "Differentiation of embryonic stem cells to insulin-secreting structures similar to pancreatic islets," *Science*, vol. 292, no. 5520, pp. 1389–1394, 2001.
- [19] S. D. Dave, A. V. Vanikar, H. L. Trivedi, U. G. Thakkar, S. C. Gopal, and T. Chandra, "Novel therapy for insulin-dependent diabetes mellitus: infusion of in vitro-generated insulin-secreting cells," *Clinical and Experimental Medicine*, pp. 1–5, 2013.
- [20] H. Zhu, L. Yu, and B. Wang, "Progress in isolation and purification of porcine islets," *Chinese Journal of Reparative and Reconstructive Surgery*, vol. 26, no. 8, pp. 1012–1016, 2012.
- [21] J. D. Carter, S. B. Dula, K. L. Corbin, R. Wu, and C. S. Nunemaker, "A practical guide to rodent islet isolation and assessment," *Biological Procedures Online*, vol. 11, no. 1, pp. 3–31, 2009.
- [22] S. Matsumoto, H. Noguchi, B. Naziruddin et al., "Improvement of pancreatic islet cell isolation for transplantation," *Proceedings (Baylor University Medical Center)*, vol. 20, no. 4, pp. 357–362, 2007.
- [23] H. T. Zhu, L. Yu, Y. Lyu, and B. Wang, "Optimal pig donor selection in islet xenotransplantation: current status and future perspectives," *Journal of Zhejiang University Science B*, vol. 15, no. 8, pp. 681–691, 2014.
- [24] P. Chhabra and K. L. Brayman, "Current status of immunomodulatory and cellular therapies in preclinical and clinical islet transplantation," *Journal of Transplantation*, vol. 2011, Article ID 637692, 24 pages, 2011.
- [25] Q. H. Hu, Z. W. Liu, and H. T. Zhu, "Pig islets for islet xenotransplantation: current status and future perspectives," *Chinese Medical Journal*, vol. 127, no. 2, pp. 370–377, 2014.
- [26] C. Aguayo-Mazzucato and S. Bonner-Weir, "Stem cell therapy for type 1 diabetes mellitus," *Nature Reviews Endocrinology*, vol. 6, no. 3, pp. 139–148, 2010.
- [27] N. Sakata, G. Yoshimatsu, H. Tsuchiya, S. Egawa, and M. Unno, "Animal models of diabetes mellitus for islet transplantation," *Experimental diabetes research*, vol. 2012, Article ID 256707, 11 pages, 2012.
- [28] J.-P. Dehoux and P. Gianello, "The importance of large animal models in transplantation," *Frontiers in Bioscience*, vol. 12, no. 13, pp. 4864–4880, 2007.
- [29] E. M. Weerts, W. E. Fantegrossi, and A. K. Goodwin, "The value of nonhuman primates in drug abuse research," *Experimental and Clinical Psychopharmacology*, vol. 15, no. 4, pp. 309–327, 2007.
- [30] J. D. Wagner, J. M. Cline, M. K. Shadoan, B. C. Bullock, S. E. Rankin, and W. T. Cefalu, "Naturally occurring and experimental diabetes in cynomolgus monkeys: a comparison of carbohydrate and lipid metabolism and islet pathology," *Toxicologic Pathology*, vol. 29, no. 1, pp. 142–148, 2001.
- [31] J. A. Bluestone, K. Herold, and G. Eisenbarth, "Genetics, pathogenesis and clinical interventions in type 1 diabetes," *Nature*, vol. 464, no. 7293, pp. 1293–1300, 2010.
- [32] H. J. Harwood Jr., P. Listrani, and J. D. Wagner, "Nonhuman primates and other animal models in diabetes research," *Journal of Diabetes Science and Technology*, vol. 6, no. 3, pp. 503–514, 2012.
- [33] J. D. Wagner, K. Kavanagh, G. M. Ward, B. J. Auerbach, H. J. Harwood Jr., and J. R. Kaplan, "Old world nonhuman primate models of type 2 diabetes mellitus," *ILAR Journal*, vol. 47, no. 3, pp. 259–271, 2006.
- [34] C. F. Howard Jr., "Nonhuman primates as models for the study of human diabetes mellitus," *Diabetes*, vol. 31, no. 1, part 2, pp. 37–44, 1982.
- [35] A. D. Kirk, "Transplantation tolerance: a look at the nonhuman primate literature in the light of modern tolerance theories," *Critical Reviews in Immunology*, vol. 19, no. 5–6, pp. 349–388, 1999.
- [36] S. He, D. Wang, and L. Wei, "Practical and critical instruction for nonhuman primate diabetic models," *Transplantation Proceedings*, vol. 45, no. 5, pp. 1856–1865, 2013.
- [37] E. L. Röder and P. J. A. Timmermans, "Housing and care of monkeys and apes in laboratories: adaptations allowing essential species-specific behaviour," *Laboratory Animals*, vol. 36, no. 3, pp. 221–242, 2002.
- [38] G. Laule and M. Whittaker, "Enhancing nonhuman primate care and welfare through the use of positive reinforcement training," *Journal of Applied Animal Welfare Science*, vol. 10, no. 1, pp. 31–38, 2007.
- [39] S. J. Schapiro and S. P. Lambeth, "Control, choice, and assessments of the value of behavioral management to nonhuman primates in captivity," *Journal of Applied Animal Welfare Science*, vol. 10, no. 1, pp. 39–47, 2007.
- [40] M. Koulmanda, A. Qipo, Z. Fan et al., "Prolonged survival of allogeneic islets in cynomolgus monkeys after short-term triple therapy," *American Journal of Transplantation*, vol. 12, no. 5, pp. 1296–1302, 2012.
- [41] M. Marigliano, A. Casu, S. Bertera, M. Trucco, and R. Bottino, "Hemoglobin A1C percentage in nonhuman primates: a useful tool to monitor diabetes before and after porcine pancreatic islet xenotransplantation," *Journal of Transplantation*, vol. 2011, Article ID 965605, 8 pages, 2011.
- [42] B. Hirshberg, S. Mog, N. Patterson, J. Leconte, and D. M. Harlan, "Histopathological study of intrahepatic islets transplanted in the nonhuman primate model using edmonton protocol immunosuppression," *Journal of Clinical Endocrinology and Metabolism*, vol. 87, no. 12, pp. 5424–5429, 2002.
- [43] E. Pennisi, "Genomicists tackle the primate tree," *Science*, vol. 316, no. 5822, pp. 218–221, 2007.
- [44] M. Vaccari and G. Franchini, "Memory T cells in rhesus macaques," *Advances in Experimental Medicine and Biology*, vol. 684, pp. 126–144, 2010.
- [45] A. Blancher, A. Aarnink, K. Tanaka et al., "Study of cynomolgus monkey (*Macaca fascicularis*) Mhc DRB gene polymorphism in four populations," *Immunogenetics*, vol. 64, no. 8, pp. 605–614, 2012.
- [46] P. P. M. Rood, L. H. Buhler, R. Bottino, M. Trucco, and D. K. C. Cooper, "Pig-to-nonhuman primate islet xenotransplantation: a review of current problems," *Cell Transplantation*, vol. 15, no. 2, pp. 89–104, 2006.
- [47] C. R. Abec, K. Mansfield, S. D. Tardif, and T. Morris, *Nonhuman Primates in Biomedical Research: Diseases*, Elsevier Science, Amsterdam, The Netherlands, 2012.
- [48] D. Cheța, "Animal models of type I (insulin-dependent) diabetes mellitus," *Journal of Pediatric Endocrinology and Metabolism*, vol. 11, no. 1, pp. 11–19, 1998.
- [49] B. Tyrberg, A. Andersson, and L. A. Borg, "Species differences in susceptibility of transplanted and cultured pancreatic islets to the β -cell toxin alloxan," *General and Comparative Endocrinology*, vol. 122, no. 3, pp. 238–251, 2001.
- [50] T. Szkudelski, "The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas," *Physiological Research*, vol. 50, no. 6, pp. 537–546, 2001.

- [51] C. E. Mathews, S. H. Langley, and E. H. Leiter, "New mouse model to study islet transplantation in insulin-dependent diabetes mellitus," *Transplantation*, vol. 73, no. 8, pp. 1333–1336, 2002.
- [52] A. V. Matveyenko and P. C. Butler, "Islet amyloid polypeptide (IAPP) transgenic rodents as models for type 2 diabetes," *ILAR Journal*, vol. 47, no. 3, pp. 225–233, 2006.
- [53] A. A. Viehmann Milam, S. E. Maher, J. A. Gibson et al., "A humanized mouse model of autoimmune insulinitis," *Diabetes*, vol. 63, no. 5, pp. 1712–1724, 2014.
- [54] A. W. S. Chan, "Progress and prospects for genetic modification of nonhuman primate models in biomedical research," *ILAR Journal*, vol. 54, no. 2, Article ID ilt035, pp. 211–223, 2013.
- [55] S.-H. Yang, P.-H. Cheng, H. Banta et al., "Towards a transgenic model of Huntington's disease in a non-human primate," *Nature*, vol. 453, no. 7197, pp. 921–924, 2008.
- [56] Y. Niu, Y. Yu, A. Bernat et al., "Transgenic rhesus monkeys produced by gene transfer into early-cleavage-stage embryos using a simian immunodeficiency virus-based vector," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 41, pp. 17663–17667, 2010.
- [57] D. A. Rees and J. C. Alcolado, "Animal models of diabetes mellitus," *Diabetic Medicine*, vol. 22, no. 4, pp. 359–370, 2005.
- [58] J. Gillman, C. Gilbert, and J. C. Allan, "The relationship of hyperglycaemia to hyperlipaemia and ketonaemia in depancreatized baboons (*Papio ursinus*)," *Journal of Endocrinology*, vol. 17, no. 4, pp. 349–362, 1958.
- [59] J. Gillman, C. Gilbert, E. Epstein, and J. C. Allan, "Endocrine control of blood sugar, lipaemia, and ketonaemia in diabetic," *British Medical Journal*, vol. 2, no. 5107, pp. 1260–1263, 1958.
- [60] J. L. Buss, E. Essig, K. Osei, S. Brodsky, G. Hadley, and A. Rajab, "Steroid-free maintenance of islet allografts using mycophenolate mofetil and cyclosporine in the nonhuman primate," *Annals of Transplantation*, vol. 16, no. 2, pp. 88–97, 2011.
- [61] M. Watanabe, K. Yamashita, T. Suzuki et al., "ASKP1240, a fully human anti-CD40 monoclonal antibody, prolongs pancreatic islet allograft survival in nonhuman primates," *American Journal of Transplantation*, vol. 13, no. 8, pp. 1976–1988, 2013.
- [62] T. Yoshida, T. Suzuki, M. Watanabe et al., "Induction of insulin-dependent diabetes mellitus by total pancreatectomy for pancreatic islet transplantation in cynomolgus monkeys," *Journal of Hepato-Biliary-Pancreatic Sciences*, vol. 19, no. 6, pp. 661–666, 2012.
- [63] N. S. Kenyon, L. A. Fernandez, R. Lehmann et al., "Long-term survival and function of intrahepatic islet allografts in baboons treated with humanized anti-CD154," *Diabetes*, vol. 48, no. 7, pp. 1473–1481, 1999.
- [64] N. S. Kenyon, M. Chatzipetrou, M. Masetti et al., "Long-term survival and function of intrahepatic islet allografts in rhesus monkeys treated with humanized anti-CD154," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 14, pp. 8132–8137, 1999.
- [65] D. W. R. Gray, G. L. Warnock, R. Sutton, M. Peters, P. McShane, and P. J. Morris, "Successful autotransplantation of isolated islets of Langerhans in the cynomolgus monkey," *British Journal of Surgery*, vol. 73, no. 10, pp. 850–853, 1986.
- [66] K. Cardona, Z. Milas, E. Strobert et al., "Engraftment of adult porcine islet xenografts in diabetic nonhuman primates through targeting of costimulation pathways," *The American Journal of Transplantation*, vol. 7, no. 10, pp. 2260–2268, 2007.
- [67] P. Thompson, K. Cardona, M. Russell et al., "CD40-specific costimulation blockade enhances neonatal porcine islet survival in nonhuman primates," *The American Journal of Transplantation*, vol. 11, no. 5, pp. 947–957, 2011.
- [68] B.-G. Ericzon, R. M. H. Wijnen, K. Kubota, A. V. D. Bogaard, and G. Kootstra, "Diabetes induction and pancreatic transplantation in the cynomolgus monkey: methodological considerations," *Transplant International*, vol. 4, no. 2, pp. 103–109, 1991.
- [69] S. He, Y. Chen, L. Wei et al., "Treatment and risk factor analysis of hypoglycemia in diabetic rhesus monkeys," *Experimental Biology and Medicine*, vol. 236, no. 2, pp. 212–218, 2011.
- [70] Y. Zhang, L. Fu, Y.-R. Lu et al., "Pancreas anatomy and surgical procedure for pancreatectomy in rhesus monkeys," *Journal of Medical Primatology*, vol. 40, no. 6, pp. 376–382, 2011.
- [71] C. F. Qiao, B. L. Tian, G. Mai et al., "Induction of diabetes in rhesus monkeys and establishment of insulin administration strategy," *Transplantation Proceedings*, vol. 41, no. 1, pp. 413–417, 2009.
- [72] S. Lenzen, "The mechanisms of alloxan- and streptozotocin-induced diabetes," *Diabetologia*, vol. 51, no. 2, pp. 216–226, 2008.
- [73] J. Frame, D. Kelsen, N. Kemeny et al., "A phase II trial of streptozotocin and adriamycin in advanced APUD tumors," *American Journal of Clinical Oncology*, vol. 11, no. 4, pp. 490–495, 1988.
- [74] J. Naidoo, D. O'Toole, M. J. Kennedy, J. V. Reynolds, M. O'Connor, and K. O'Byrne, "A single institution experience of streptozotocin/fluorouracil combination chemotherapy: a case series," *Irish Journal of Medical Science*, vol. 181, no. 2, pp. 211–214, 2012.
- [75] C. O. Eleazu, K. C. Eleazu, S. Chukwuma, and U. N. Essien, "Review of the mechanism of cell death resulting from streptozotocin challenge in experimental animals, its practical use and potential risk to humans," *Journal of Diabetes & Metabolic Disorders*, vol. 12, article 60, 2013.
- [76] K. Cardona, G. S. Korbitt, Z. Milas et al., "Long-term survival of neonatal porcine islets in nonhuman primates by targeting costimulation pathways," *Nature Medicine*, vol. 12, no. 3, pp. 304–306, 2006.
- [77] P. Thompson, I. R. Badell, M. Lowe et al., "Islet xenotransplantation using gal-deficient neonatal donors improves engraftment and function," *American Journal of Transplantation*, vol. 11, no. 12, pp. 2593–2602, 2011.
- [78] M. Koulmanda, A. Qipo, S. Chebrolu, J. O'Neil, H. Auchincloss Jr., and R. N. Smith, "The effect of low versus high dose of streptozotocin in cynomolgus monkeys (*Macaca fascicularis*)," *American Journal of Transplantation*, vol. 3, no. 3, pp. 267–272, 2003.
- [79] R. Bottino, A. Criscimanna, A. Casu et al., "Recovery of endogenous β -cell function in nonhuman primates after chemical diabetes induction and islet transplantation," *Diabetes*, vol. 58, no. 2, pp. 442–447, 2009.
- [80] M. C. Deeds, J. M. Anderson, A. S. Armstrong et al., "Single dose streptozotocin-induced diabetes: considerations for study design in islet transplantation models," *Laboratory Animals*, vol. 45, no. 3, pp. 131–140, 2011.
- [81] M. L. Graham, L. A. Mutch, E. F. Rieke et al., "Refining the high-dose streptozotocin-induced diabetic non-human primate model: an evaluation of risk factors and outcomes," *Experimental Biology and Medicine*, vol. 236, no. 10, pp. 1218–1230, 2011.

- [82] B. R. Theriault, J. R. Thistlethwaite Jr., M. G. Levisetti et al., "Induction, maintenance, and reversal, of streptozotocin-induced insulin-dependent diabetes mellitus in the juvenile cynomolgus monkey (*Macaca fascicularis*)," *Transplantation*, vol. 68, no. 3, pp. 331–337, 1999.
- [83] M. Lowe, I. R. Badell, P. Thompson et al., "A novel monoclonal antibody to CD40 prolongs islet allograft survival," *The American Journal of Transplantation*, vol. 12, no. 8, pp. 2079–2087, 2012.
- [84] I. R. Badell, M. C. Russell, K. Cardona et al., "CTLA4Ig prevents alloantibody formation following nonhuman primate islet transplantation using the CD40-specific antibody 3A8," *American Journal of Transplantation*, vol. 12, no. 7, pp. 1918–1923, 2012.
- [85] D. J. Van Der Windt, R. Bottino, A. Casu et al., "Long-term controlled normoglycemia in diabetic non-human primates after transplantation with hCD46 transgenic porcine islets," *The American Journal of Transplantation*, vol. 9, no. 12, pp. 2716–2726, 2009.
- [86] G. Hecht, S. Eventov-Friedman, C. Rosen et al., "Embryonic pig pancreatic tissue for the treatment of diabetes in a nonhuman primate model," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 21, pp. 8659–8664, 2009.
- [87] M. Wijkstrom, N. Kirchhof, M. Graham et al., "Cyclosporine toxicity in immunosuppressed streptozotocin-diabetic nonhuman primates," *Toxicology*, vol. 207, no. 1, pp. 117–127, 2005.
- [88] K. Hayashi, R. Kojima, and M. Ito, "Strain differences in the diabetogenic activity of streptozotocin in mice," *Biological and Pharmaceutical Bulletin*, vol. 29, no. 6, pp. 1110–1119, 2006.
- [89] R. M. Pitkin and W. A. Reynolds, "Diabetogenic effects of streptozotocin in rhesus monkeys," *Diabetes*, vol. 19, no. 2, pp. 85–90, 1970.
- [90] K. N. Litwak, W. T. Cefalu, and J. D. Wagner, "Streptozotocin-induced diabetes mellitus in cynomolgus monkeys: changes in carbohydrate metabolism, skin glycation, and pancreatic islets," *Laboratory Animal Science*, vol. 48, no. 2, pp. 172–178, 1998.
- [91] P. P. M. Rood, R. Bottino, A. N. Balamurugan et al., "Induction of diabetes in cynomolgus monkeys with high-dose streptozotocin: adverse effects and early responses," *Pancreas*, vol. 33, no. 3, pp. 287–292, 2006.
- [92] S. Shibata, N. Kirchhof, S. Matsumoto et al., "High-dose streptozotocin for diabetes induction in adult rhesus monkeys," *Transplantation Proceedings*, vol. 34, no. 4, pp. 1341–1344, 2002.
- [93] X. Jin, L. Zeng, S. He et al., "Comparison of single high-dose streptozotocin with partial pancreatectomy combined with low-dose streptozotocin for diabetes induction in rhesus monkeys," *Experimental Biology and Medicine*, vol. 235, no. 7, pp. 877–885, 2010.
- [94] D. Dufrane, M. Van Steenberghe, Y. Guiot, R.-M. Goebbels, A. Saliez, and P. Gianello, "Streptozotocin-induced diabetes in large animals (pigs/primates): role of GLUT2 transporter and β -cell plasticity," *Transplantation*, vol. 81, no. 1, pp. 36–45, 2006.
- [95] M. G. Tal, B. Hirshberg, Z. Neeman et al., "Induction of diabetes in nonhuman primates by means of temporary arterial embolization and selective arterial injection of streptozotocin," *Radiology*, vol. 230, no. 1, pp. 163–168, 2004.
- [96] C. L. Zou, J. Y. Wang, S. Y. Wang et al., "Characterizing the induction of diabetes in juvenile cynomolgus monkeys with different doses of streptozotocin," *Science China Life Sciences*, vol. 55, no. 3, pp. 210–218, 2012.
- [97] I. R. Badell, M. C. Russell, P. W. Thompson et al., "LFA-1-Specific therapy prolongs allograft survival in rhesus macaques," *Journal of Clinical Investigation*, vol. 120, no. 12, pp. 4520–4531, 2010.
- [98] C. Polychronakos, "Animal models of spontaneous autoimmune diabetes: notes on their relevance to the human disease," *Current Diabetes Reports*, vol. 4, no. 2, pp. 151–154, 2004.
- [99] B. J. Whalen, J. P. Mordes, and A. A. Rossini, "The BB rat as a model of human insulin-dependent diabetes mellitus," in *Current Protocols in Immunology*, chapter 15, unit 15.3, 2001.
- [100] M. S. Anderson and J. A. Bluestone, "The NOD mouse: a model of immune dysregulation," *Annual Review of Immunology*, vol. 23, pp. 447–485, 2005.
- [101] S. M. Cabrera, M. R. Rigby, and R. G. Mirmira, "Targeting regulatory T cells in the treatment of type 1 diabetes mellitus," *Current Molecular Medicine*, vol. 12, no. 10, pp. 1261–1272, 2012.
- [102] M. Wang, J. Racine, M. Zhang et al., "MHC-mismatched chimerism is required for induction of transplantation tolerance in autoimmune nonobese diabetic recipients," *The Journal of Immunology*, vol. 193, no. 4, pp. 2005–2015, 2014.
- [103] R. Deng, M. Khattar, A. Xie et al., "Anti-TCR mab induces peripheral tolerance to alloantigens and delays islet allograft rejection in autoimmune diabetic nod mice," *Transplantation*, vol. 97, no. 12, pp. 1216–1224, 2014.
- [104] Q. Shi, D. Wang, G. A. Hadley, A. W. Bingaman, S. T. Bartlett, and D. L. Farber, "Long-term islet graft survival in NOD mice by abrogation of recurrent autoimmunity," *Diabetes*, vol. 53, no. 9, pp. 2338–2345, 2004.
- [105] F. T. Thomas, K. Pittman, and J. M. Thomas, "Induction of functional xenograft tolerance of pig islets in the autoimmune NOD mouse," *Transplantation Proceedings*, vol. 27, no. 6, pp. 3323–3325, 1995.
- [106] M. Wijkstrom, R. Bottino, and D. K. C. Cooper, "Limitations of the pig-to-non-human primate islet transplantation model," *Xenotransplantation*, vol. 20, no. 1, pp. 2–4, 2013.
- [107] A. Casu, R. Bottino, A. N. Balamurugan et al., "Metabolic aspects of pig-to-monkey (*Macaca fascicularis*) islet transplantation: implications for translation into clinical practice," *Diabetologia*, vol. 51, no. 1, pp. 120–129, 2008.
- [108] M. L. Graham, M. D. Bellin, K. K. Papas, B. J. Hering, and H.-J. Schuurman, "Species incompatibilities in the pig-to-macaque islet xenotransplant model affect transplant outcome: a comparison with allotransplantation," *Xenotransplantation*, vol. 18, no. 6, pp. 328–342, 2011.
- [109] M. L. Graham and H.-J. Schuurman, "The usefulness and limitations of the diabetic macaque model in evaluating long-term porcine islet xenograft survival," *Xenotransplantation*, vol. 20, no. 1, pp. 5–17, 2013.