Review Article

Updating Experimental Models of Diabetic Cardiomyopathy

**J. Fuentes-Antrás,¹ B. Picatoste,^{1,2} A. Gómez-Hernández,^{2,3} J. Egido,^{1,2}
J. Tuñón,¹ and Ó. Lorenzo^{1,2}**

¹IIS-Fundación Jiménez Díaz, Autónoma University, 28040 Madrid, Spain

²Spanish Biomedical Research Centre in Diabetes and Associated Metabolic Disorders (CIBERDEM) Network, 28040 Madrid, Spain

³Biochemistry and Molecular Biology Department, School of Pharmacy, Complutense University of Madrid, 28040 Madrid, Spain

Correspondence should be addressed to Ó. Lorenzo; olorenzo@fjd.es

Received 14 January 2015; Revised 26 March 2015; Accepted 29 March 2015

Academic Editor: Geoff Werstuck

Copyright © 2015 J. Fuentes-Antrás et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Diabetic cardiomyopathy entails a serious cardiac dysfunction induced by alterations in structure and contractility of the myocardium. This pathology is initiated by changes in energy substrates and occurs in the absence of atherosclerosis, hypertension, or other cardiomyopathies. Inflammation, hypertrophy, fibrosis, steatosis, and apoptosis in the myocardium have been studied in numerous diabetic experimental models in animals, mostly rodents. Type I and type II diabetes were induced by genetic manipulation, pancreatic toxins, and fat and sweet diets, and animals recapitulate the main features of human diabetes and related cardiomyopathy. In this review we update and discuss the main experimental models of diabetic cardiomyopathy, analysing the associated metabolic, structural, and functional abnormalities, and including current tools for detection of these responses. Also, novel experimental models based on genetic modifications of specific related genes have been discussed. The study of specific pathways or factors responsible for cardiac failures may be useful to design new pharmacological strategies for diabetic patients.

1. Main Models of Diabetic Cardiomyopathy (DCM)

Experimental models of both type I and type II diabetes (T1DM and T2DM) consistently exhibit alterations in the circulating levels of glucose and in the lipid profile (Table 1). Main T1DM and T2DM animals show both hyperglycemia (early after pancreas-toxin or fat/sweet-diet administration, resp., and later after genetic mutations) and hyperlipidemia, represented by elevated levels of TAG, cholesterol, and lipoproteins. More interestingly, these models also exhibit functional, structural, and metabolic abnormalities that recapitulate the human DCM pathology.

Dysfunction in Cardiac Mechanics. In patients, early stages of DCM are marked by a deterioration of longitudinal systolic function, a compensative elevated radial function, and diastolic dysfunction [1]. DCM progression is characterized by reduced ejection fraction (EF) and ventricular dilatation and, later, mimicking dilative cardiomyopathy. Experimental

T1DM and T2DM animals are prone to develop diastolic and/or systolic dysfunction, as demonstrated in numerous *in vivo* studies using echocardiography, magnetic resonance imaging (MRI), and hemodynamic measurements [2, 3]. Diastolic dysfunction usually precedes the alteration of cardiac contractility. As recently updated [4], a broad analysis of diastolic performance should include several Doppler indexes: ratio of peak velocity of early to late filling of mitral inflow (E/A) and deceleration time of early filling of mitral inflow, as well as the isovolumetric relaxation/contraction time. Given the subjectivity of visual echocardiographic estimations and the frequent coexistence of different Doppler severity patterns, measurement of left ventricular (LV) diastolic pressures by catheterization is also highly valuable. The main systolic parameters include LV septum thicknesses, LV internal dimensions, posterior wall (PW) thicknesses at diastole and systole, LV mass, LVEF, LV fractional shortening (FS), and ventricular contractility assessment (dP/dt). Regardless of rather inconstant particularities mainly displayed by Akita mice, all T1DM models have been reported

TABLE 1: Main experimental models of T1DM and T2DM and related plasma and cardiac characteristics.

	Type I diabetes				Type II diabetes							DIO																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
	Toxics	Genetics		Genetics	 

TABLE I: Continued.

	Type I diabetes				Type II diabetes							DIO		
	Toxics	Genetics			Genetics			Genetics				Fat	Sweet	
	STZ (Antibiotic toxicity)	Alloxan (Uric acid derivative toxicity)	OVE26 (Calmodulin overexpression)	Akita (<i>Ins2</i> ^{+/-})	NOD (Nonobese diabetic mouse)	BB rat (Nonobese diabetic rat)	ob/ob (<i>Lep</i> ^{-/-})	db/db (<i>Lep</i> ^{-/-})	ZDF (<i>Lep</i> ^{-/-})	KK A ^y (Polygenic obese + A ^y)	OLETF (Polygenic obese)	GK (Polygenic nonobese)		
Metabolic alterations														
Glucose oxidation	↓[70]			↓[24, 71]			↓[5, 64]	↓[26]	↓[10, 72]			↓[31, 80]	↓[74]	
FA oxidation	↑[70]		↑[62]	↑[24, 71]			↑[5, 64]	↑[63]	↑[10, 86]			↑[31]	~ [74]/↑[73]	↑[53]
Mitoch. function	↓[70]		↓[62]	↓[71]			↓[81]	↓[152]	↓[10]	↓[151]	↓[83]	↓[84]	↓[73]	
Oxidative stress	↑[70, 76]	↑[153]	↑[78, 79]	~[71]			↑[92]	↑[152]	↑[10, 86]		↑[47, 83]	↑[84, 85]	↑[34]	↑[53]
Ca ²⁺ mobilization	↓[70]		↓[90]	↓[24]		↓[91]	↓[25]	↓[93]	~[72]			↓[94]	↓[13, 95]	↓[14]

T1DM was induced by toxins [streptozotocin (STZ) or alloxan] or genetic mutations (OVE26, calmodulin transgenic; Akita, insulin-2 deficient; NOD, nonobese diabetic or BB, BioBreeding diabetes-prone mice). T2DM models were produced by genetic alterations [*ob/ob*, leptin deficient mice; *db/db*, leptin receptor deficient mice; ZDF, Zucker Diabetic Fatty rats; KK A^y, yellow obese gene transgenic KK mice; OLETF, Otsuka Long-Evans Tokushima fatty rats; GK, Goto-Kakizaki rats or DIO, diet-induced (fat, sweet) obesity].

to meet most of these criteria (Table 1). Cardiac performance has also been extensively studied in T2DM experimental models. Genetically obese mice displayed severe diastolic dysfunction, as evidenced by diminished E/A velocities in *ob/ob* and *db/db* mice [5, 6]. Contractile properties are yet marginally affected in *ob/ob* mice [5]. On the contrary, *db/db* mice exhibited a reduced FS and velocity of circumferential fibre shortening at age 12 weeks [6]. Van den Bergh et al. also reported in *db/db* a decreased preload recruitable stroke work, end-systolic elastance, and dP/dt from age 24 weeks onwards but found preserved cardiac output, EF, and dP/dt [7] (Table 1). Other T2DM models presented heterogeneous data regarding cardiac performance. In ZDF rats, impaired diastolic function has been extensively described [8–10]. However, Zhou's work initially reported a decrease of FS at age 20 weeks [8], and we and others found no evidence of systolic dysfunction by both echocardiographic and MRI determinations at ages 16 and 44 weeks, respectively [9, 10]. Similarly, OLETF and GK rats preserved contractile function while displaying abnormalities of ventricular relaxation, as suggested by a prolonged deceleration time, decreased peak velocity, and reduced LV diameters associated with increased LVPW thickness and LV mass [11, 12]. In contrast, diet-induced obesity appears to invariably associate both diastolic and systolic dysfunctions. High-fat diet increased LV mass and reduced dP/dt and FS after 6 weeks [13]. Alike, sucrose fed rats exhibited early abnormalities in LV fillings, as demonstrated by reduced E/A ratios, together with depressed FS and EF after 10 weeks [14]. Consistently, echocardiographic findings in other high-fat and high-sucrose fed mice revealed decreased FS, EF, and velocity of circumferential fibre shortening, as well as dramatically impaired parameters of diastolic function after 16 weeks [15].

Thus, cardiac dysfunction in T1DM patients could be roughly reproduced in all conventional models (induced by toxins and genetic alterations). In T2DM, diet-induced models may represent the human pathology more appropriately, at least in the advanced states of the disease.

(1) Alterations of Cardiac Structure. Maladaptive structural modifications underlie both the diastolic and systolic impairments that eventually lead to heart failure. Although the progression of these events has not been fully established, the hallmark of changes includes cell hypertrophy, local inflammation, and interstitial fibrosis promoted by steatosis and cell-death processes in the injured myocardium.

(a) Hypertrophy. Undiagnosed cardiac hypertrophy is prevalent in asymptomatic T2DM patients, reaching up to 56% in recent reports [16]. Myocyte hypertrophy is common in biopsy of diabetic hearts and is induced by long-standing metabolic imbalances and microcirculation anomalies. However, the contribution of cardiomyocyte hypertrophy to the development of ventricular hypertrophy is still unclear [17]. Most models of DCM display progressive concentric or eccentric hypertrophy, which is directly associated with diastolic and systolic derangement, respectively. Indicative parameters include LV mass and heart-to-body weight (or femur length) ratio, LVPW and septum thicknesses, LV

internal diameters, and cardiomyocyte size. In addition, complementary plasma levels of brain natriuretic peptide (BNP), atrial natriuretic peptide (ANP), and β -myosin heavy chain (β -MHC) have traditionally been considered supportive [18]. However, a recent systematic analysis of hypertrophic biomarkers in diabetic rodents revealed that most commonly measured genes are confounded by diabetogenic methods and do not correlate with cardiac hypertrophy [19]. In STZ-induced models, reduced wall thicknesses and increased LV internal diameters are prominent features, accounting for extensive ventricular dilatational remodeling [20] (Table 1). We observed cardiomyocyte size enlargement in long-term T1DM (22 weeks), reproduced to a lesser extent in short-term counterparts (6 weeks) [21]. Similar histological findings have been reported in OVE26 mice together with the upregulation of ANP and β -MHC [22, 23]. Conversely, no evidence of myocardial hypertrophy has been found in Akita mice [24]. In T2DM mice models, though systolic dysfunction is developed as a late event, myocardial mass may increase earlier. A higher heart weight has been reported in *ob/ob* and *db/db* mice at ages 12 and 9 weeks, respectively [25, 26]. A comprehensive MRI analysis of *db/db* hearts by Yue et al. revealed a significant increase in LV mass and LVPW, and septum thicknesses, as soon as age 13 weeks [26]. However, early architectural changes are subtle and not consistently identified. In this line, echocardiographic and cell dimension assessments did not find hypertrophy in 15-week-old *db/db* mice, despite evidence of systolic compromise [27]. In contrast, ZDF and GK rats usually exhibited a significant increase of the septum and LVPW thicknesses, reduced LV diastolic and systolic diameters, enlarged cardiomyocyte volume, and higher expression rates of ANP, in spite of the fact that contractile function was frequently preserved [10, 12, 28–31]. Finally, the limited body of evidence on diet-induced T2DM rather pointed hypertrophy as the only morphological alteration constantly present, independently of its high-fat or high-sugar diet origin [13, 32–34]. However, a recent work reported no difference in heart weight and cardiomyocyte size but overexpression of ANP, BNP, and β -MHC after a 16-week-long high-fat diet [35].

(b) Inflammation. Chronic low-grade inflammation has been recently added to the features of DCM in human patients [36]. Inflammatory signaling in cardiomyocytes usually occurs as an early response to myocardial injury and entails an activation of the proinflammatory nuclear transcription factor- κ B (NF- κ B) and the related expression of cytokines (i.e., tumour necrosis factor- α (TNF α), interleukins (IL-1 β , IL-6), and chemokines (i.e., MCP-1)), and adhesion molecules (i.e., intracellular and vascular cell adhesion molecule-1 (ICAM-1, VCAM-1, resp.)) [37, 38]. Toll-like receptors and inflammasome signalling platforms may also be key participants in DCM-associated inflammation [39]. In STZ-treated hearts, both macrophage and lymphocyte infiltration and increased expression of proinflammatory cytokines and adhesion molecules were detected at 22 weeks, but not 6 weeks, after injection [21] (Table 1). Likewise, Li's paper reported an augmented NF- κ B activity in OVE26 mice [22]. A comprehensive approach to myocardial inflammation

in Akita mice was recently available, indicating the upregulation of TNF α and the attenuation of anti-inflammatory IL-10 [40]. However, data from T2DM models is more variable. *Ob/ob* and *db/db* mice displayed inflammation and increased myocardial proinflammatory factors [41], whereas we are not aware of any work addressing inflammation in the heart of GK and OLETF rats or KK A^y mice. ZDF myocardia showed either absence of inflammatory cell infiltrates at ages 9 and 13 months or higher levels of proinflammatory cytokines at age 22 weeks [42]. Inflammation is also present in diet-induced models of obesity and diabetes. Several studies on high-fat fed mice showed upregulation of proinflammatory factors, together with lower measurements of anti-inflammatory adiponectin and IFN γ [35, 43, 44]. In turn, high-sucrose fed mice did not present inflammation [32].

(c) *Fibrosis*. Interstitial fibrosis is the histologic hallmark of human DCM as a result of both replacement of focal myocyte death and response to inflammatory infiltrate. Accelerated extracellular matrix (ECM) deposition leads to ventricular stiffness in the diabetic heart, which also stimulates a local increase of cytokines and NF- κ B [45]. Collagen type I and type III fibres accumulate in the epicardial and perivascular domains, whereas type IV is mostly found in the endocardium. Collagen can further undergo glycation by advanced glycation end-products (AGEs) and impair its degradation, leading to fibrosis, myocardial stiffness, and decreased cardiac relaxation. Quantification of interstitial fibrosis may be evaluated by histological staining (Masson's trichrome, Azan, and Sirius red) and gene/protein expression of ECM components. Valuable insight into the molecular substrate of ECM deposition is provided by the analysis of TGF β -Smad-API axis, poly(ADP-ribose) polymerase-1 (PARP-1), and matrix metalloproteinases (MMP) activity (MMP/TIMP (tissue inhibitor metalloproteinase) ratio). We reported that the myocardium of both short-term and long-term STZ-treated rats undergoes intense fibrotic remodeling, mainly at the expense of interstitial and perivascular ECM deposits and TGF β axis [21] (Table 1). OVE26 mice and BB rats also displayed myocardial fibrosis, as determined by increases in collagen accumulation and reduced MMP activity [22, 46]. As with hypertrophy, histological study of Akita mice showed no evidence of fibrosis and elevated levels of ECM proteins [24]. In T2DM, most models exhibited extensive myocardial collagen deposits. In ZDF rats, we and others described an enlarged interstitial space mostly confined to the subendocardial domain [10, 29]. Similarly, we recently reported increased ECM deposition and upregulated expression profibrotic factors in the GK myocardium [12]. Other models in which fibrosis has been clearly detected include *db/db* and *ob/ob* mice and OLETF rats [18, 26, 47]. On the contrary, several groups pointed the absence of increased collagen deposition and ventricular stiffness in *ob/ob* despite strong evidence of other DCM traits such as hypertrophy and apoptosis [5, 25]. Much less has been done so far to assess fibrosis in diet-induced obesity models, in which a complete histological examination is lacked. High-fat fed mice presented overexpression of TGF β axis and downregulation of antifibrotic p-Smad1/5 and bone morphogenetic protein-2

(BMP-2) [34, 43]. However, Nunes et al. found no differences in perivascular and interstitial fibrosis in high-sucrose fed mice compared to controls [32].

(d) *Steatosis*. Accumulation of fat in nonadipose tissues in human diabetes may be a protective response to provide a store of fuel for subsequent oxidation and to prevent exposure to toxic lipid metabolites such as ceramides [48]. However, a chronic imbalance of lipid storage versus lipid oxidation may lead to mechanical dysfunction. Insulin resistance and high levels of intramyocardial triacylglycerol (TAG), long-chain fatty acid (FA), TAG derivatives (i.e., diacylglycerols (DAG) and phospholipids), and ceramides are main triggers of heart disease in obesity and T2DM [49]. Also, ceramide and DAG can reduce insulin-stimulated signalling through the activation of PKC and NF- κ B. Surprisingly, myocardial steatosis is a common feature among T1DM models (Table 1). STZ-treated mice showed a 50% increase in cardiac content of FA [50], and both FA and ceramide deposits have been reported in NOD and Akita mice already at age 3 months [24, 51]. An association between increased myocardial lipid content and cardiac dysfunction has also been observed in rodent models of obesity and T2DM. Interestingly, Zhou's lab described that cardiac steatosis was evident in ZDF at 6 weeks even before the reduction of FS [8]. Also, *ob/ob* and *db/db* showed significantly elevated contents of phospholipids, whereas ceramides, cardiolipin, and unesterified cholesterol remained unaltered [5, 41, 52]. As expected, both high-fat and high-fructose-induced obesity models also displayed extensive myocardial steatosis [15, 53].

(e) *Apoptosis*. Myocardial cell death is recognized as a major event in the progression of DCM. Biopsies from diabetic patients revealed that cardiomyocytes are more susceptible to diabetes-induced apoptosis than endothelial cells or fibroblasts [54]. Standard methods to study apoptosis include TUNEL staining, phosphatidylserine exposure in cell membrane, and analysis of caspase-3 and other components of the extrinsic (i.e., Fas, FasL) and intrinsic (i.e., Bax/Bcl-2 ratio, cytochrome-C) apoptotic pathways. Of note, the DNA repairing enzyme PARP-1, which is also recognized as an apoptotic marker, has been recently shown to be central in a genuine type of cell death termed parthanatos [55], and its potential role in DCM has already been stated [56]. STZ-treated hearts displayed a significant increase in cardiomyocyte apoptosis already at 3 days after injection. A progressive decrease of apoptosis was reported from 4 to 24 weeks after the induction of diabetes [57] (Table 1). In this line, we showed that apoptosis is more severe in long-standing STZ-treated hearts than in short-term ones [21]. Similarly, apoptosis was stimulated in OVE26 [58]. T2DM hearts also augmented rates of cell death. Comprehensive studies on *ob/ob* and *db/db* mice reported marked pro-apoptotic molecules and apoptosis [59, 60]. Moreover, cell death became more accentuated with aging [46, 61–63]. Likewise, we and others have described spread apoptosis in ZDF and GK myocardia [10, 12, 64]. Finally, most of the works on high-fat and high-sugar fed mice reported increased myocardial apoptosis [13, 43, 61, 65].

FIGURE 1: The diabetic milieu at the myocardium. Thickened lines represent activated pathways, and dotted lines denote reduced pathways. The absence of insulin response by defect in its secretion (T1DM) or sensibility (T2DM) promotes a lack of glucose assimilation for energy demand in the cardiomyocytes and extracellular glucose accumulation. Thus, FA may be the unique energetic substrate but β -oxidation may saturate and lipid can accumulate and cause steatosis and lipotoxicity by DAG, ceramide, and ROS formation. These metabolites together with glucose derivatives (i.e., polyols, AGEs) damage the mitochondria and endoplasmic reticulum (ER) and regulate the expression of proinflammatory, hypertrophy, fibrotic, and apoptotic genes. RAGE, receptor for advanced glycation end products; UR, unspecific receptors; IR, insulin receptor; FAR, fatty acid receptors. Ch: cholesterol; Cyt. C: cytochrome-C.

However, no change in apoptosis markers was identified in a more recent contribution in high-sucrose fed mice [66].

In summary, inflammation in human DCM usually occurs in early stages of the disease. Main established T1DM and T2DM models could emulate this response when analysed soon after the onset of the pathology. Also, most of them (but Akita) can exhibit myocardial hypertrophy, fibrosis, steatosis and apoptosis, preferentially at late stages of the disease. Thus, these models may be used to reproduce specific human responses to DCM, in particular when obesity coexists in the patients.

(2) *Alterations in Cardiac Metabolism and Calcium Handling.* The structural alterations in DCM may be originated by disruptions in energetic metabolism, which almost completely relies on FA and requires higher rates of O₂ consumption. This scenario induces a progressive loss of cardiac efficiency by mitochondrial uncoupling and endoplasmic reticulum stress, and subsequent apoptosis and cardiac dysfunction [67] (Figure 1).

(a) *Energetic Imbalance.* In human DCM, FA constitute principally the unique energy source of fuel due to the insulin

inefficiency and reduction of glucose transporters and glucose oxidation enzymes [68]. As a consequence, PPAR(α , β , and γ) transcription factors are stimulated to increase gene expression of those proteins involved in FA-uptake, -transport, and -oxidation (FAO) and ceramide production, in both peroxisomes and mitochondria [69]. Main models of T1DM [62, 70, 71] exhibited a reduced expression of GLUT1 and GLUT4, and higher levels of the glucose oxidation inhibitor pyruvate dehydrogenase kinase 4 (PDK4) (Table 1). Upregulation of PPAR α and FAO proteins has also been extensively described [62]. Likewise, Akita mice hearts showed lower glucose degradation and greater palmitate oxidation and expression of FA-transporters [24]. Nevertheless, most of the experimental data regarding metabolic abnormalities of the diabetic heart derive from T2DM models. Insulin resistance and higher FA utilization have been observed in *ob/ob* mice. As a result, myocardial O₂ consumption rose, and cardiac performance substantially declined [64]. In the same model, a consistent decrease of key regulators of glucose metabolism such as phosphofructokinase (PFK) and GLUT-4 and increased expression of a set of PPAR α and PPAR γ independent genes involved in FA metabolism was reported [5]. In like fashion, *db/db* mice displayed a 40% decrease

in glucose uptake, together with a considerable reduction in carbohydrate oxidation from age 10 weeks onwards. At this age, palmitate oxidation was shown to contribute over 90% to the cardiac FA production [26, 63]. In addition, we and others demonstrated the abnormal lipid metabolism of ZDF hearts, which include upregulation of genes involved in FA uptake, lipid reesterification, and β -oxidation [8, 10, 72]. Several studies ranging from gene expression assessment to quantitative positron emission tomography imaging have recently uncovered similar alterations in OLETF and GK rats [11, 31]. Diet-induced models have also disclosed similar disturbances. A recent work described an excessive FA metabolism in high-fat fed mice, supported also by upregulated PPAR α [15]. In the same model, Cole et al. found increased FAO rates [73], though Ussher's group observed no association between lipid accumulation and insulin sensitivity [74].

(b) Mitochondrial Dysfunction and Oxidative Stress. Energetic imbalance triggers defects of mitochondrial respiration and ROS overload in human diabetes. In both T1DM and T2DM, metabolic maladaptation contributes to ROS imbalance by affecting multiple enzymatic systems including NADH oxidase (NOX), endothelial NO synthase, and, mainly, the mitochondrial respiratory chain [68, 75]. In STZ-treated rats, mitochondrial damage was evidenced by loss of membrane potential, increases in ROS production, and reduction in antioxidant glutathione [76] (Table 1). Further proteomic analysis revealed marked alterations in the expression of 24 different cardiac proteins implicated in metabolic derangement and ROS excess, half of them located in the mitochondria [77]. Likewise, OVE26 mice showed altered mitochondrial function, reduced glutathione, and increased mitochondrial biogenesis [62, 78, 79]. A comprehensive study pointed that mitochondria of Akita mice also displayed respiratory defects, reduced cristae density, and increased volume at the mitochondria. However, despite morphological aberrations, mitochondrial number and antioxidant defences remained unchanged, and the absence of mitochondrial uncoupling was evidenced by unaffected ATP-to-O₂ ratio [71]. In T2DM, *db/db* mice exhibited increased O₂ consumption, mitochondrial ROS generation, and lipid peroxidation. Mitochondrial uncoupling was associated with upregulated FAO genes and electron transfer flavoproteins [80]. *Ob/ob* mice and OLETF maintain unchanged levels of uncoupled proteins despite signal of mitochondrial dysfunction [81, 82]. Higher mitochondrial ROS production and lipid peroxidation rates have been also described in ZDF, OLETF, and GK rats [10, 83–85]. In addition, an elevation in antioxidant levels was demonstrated in ZDF rats [86]. Interestingly, ZDF rats neither displayed increased myocardial O₂ consumption nor showed a significant reduction of mitochondrial biogenesis-related factors [10, 87]. Finally, mitochondrial uncoupling appears to be invariably present in cardiomyocytes from diet-induced obese mice. High-fat intake is associated with an increased burden of oxidized proteins, NOX and uncoupling-related factors, and antioxidant responses [43, 65, 73].

(c) Altered Ca²⁺ Mobilization. A progressive dysregulation of Ca²⁺ handling underlies the relaxation-contraction defects

in human DCM. Impaired Ca²⁺ signaling may be the consequence of altered sarcolemmal or sarcoplasmic Ca²⁺ pumps and channels and ryanodine receptors [88]. Cardiac mitochondria from STZ-treated rats decreased the capacity to stimulate ATP synthesis via stimulation of Ca²⁺ pumps, and, consequently, it abridged the relaxation during work load [89] (Table 1). Impaired contractility and reuptake of Ca²⁺ in cardiomyocytes from OVE26 mice were associated with reduced Ca²⁺ ATPase-60A (SERCA2a) levels and cytoplasmic Ca²⁺ clearance [90]. BB rats and Akita mice reported altered cardiac excitation-contraction coupling as well [24, 91]. In experimental T2DM models, the mechanisms explaining decreased Ca²⁺ uptake are not yet fully resolved. In *ob/ob*, Li et al. noted that SERCA2a was markedly modified by oxidation [92], whereas Van Den Bergh et al. showed increased SERCA2a levels with less affinity for Ca²⁺, which reduced contractile capacity [25]. In *db/db* mice, Belke's lab did not report significant variance in ryanodine receptor levels but lessened SERCA2a [93]. However, unchanged expression of SERCA2a has been described in ZDF [72] and intracellular Ca²⁺ transient was unaltered in GK cardiomyocytes [94]. In addition, cardiomyocytes from high-fat fed mice displayed a significantly elevated baseline intracellular Ca²⁺ and reduced Ca²⁺ decay rate, together with a marked decrease in SERCA2a expression [13, 65]. Further, sucrose fed rats showed reduced Ca²⁺ uptake [14], and fructose fed mice exhibited an attenuation of diastolic Ca²⁺ and Ca²⁺ transient amplitude, as well as SERCA2a expression [95].

Thus, the main T1DM and T2DM experimental models could replicate the metabolic alterations occurring in human hearts, including the energetic imbalance and mitochondrial and Ca²⁺ defects.

2. Cardiac Responses of DCM-Like Models Induced by Genetic Modification

Specific target genes have been modulated to create DCM-like models in rodents (Table 2). As detected in T1DM patients, cardiac dysfunction, hypertrophy, and fibrosis were demonstrated by upregulation of PKC β [96] or by downregulation of GLUT4 [97–99], phosphoinositide dependent kinase-1 (PDK1) [100], phosphoinositide-3 kinase (PI3K) [101], or glucokinase (GCK) [102] genes (Table 2). Apoptosis and inflammation were less common (only in PKC β and/or PI3K), and some metabolic controversies were described. For example, the lack of GLUT4 or the cardiac-specific insulin receptor (CIRKO) [103, 104] decreased both FA and glucose metabolisms, but it intriguingly elevated oxidative stress without reducing mitochondrial function. On the other hand, as observed in T2DM patients, a decline of cardiac function and increased myocardial hypertrophy, fibrosis, and steatosis were predominantly found in PPAR α [105, 106], PPAR γ [107, 108], long-chain acyl-CoA synthetase-1 (LCACSI) [109, 110], lipoprotein lipase (LPL) [111, 112], and fatty acid transport protein-1 (FATP1) [113, 114] overexpressed mice. Apoptosis was also presented in PPAR γ , LCACSI, and, perhaps, LPL null mice. Interestingly, all these responses were exhibited by adipose triglyceride lipase (ATGL) deletion [115, 116]. Also,

TABLE 2: Genetic models of DCM and related cardiac responses.

	Upregulation					Downregulation						
	PPAR α	PPAR γ	LCACSI	LPL	FATP1	PKC β	UCPDTA	CIRKO	GLUT4	ATGL	PDK1	GCK
Heart failure markers	$\uparrow$ ANP, BNP	$\uparrow$ ANP, BNP	$\uparrow$ ANP, BNP	$\uparrow$ ANP, BNP					$\uparrow$ ANP, BNP			
Cardiac abnormalities												
Functional												
Diastolic function	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow/\sim$	$\downarrow$	$\downarrow$	$\downarrow$
Systolic function	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\sim$	$\downarrow$	$\sim$	$\downarrow$	$\downarrow/\sim$	$\downarrow$	$\sim$	
Structural												
Hypertrophy	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\downarrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$
Inflammation						$\uparrow$				$\sim$		
Fibrosis	$\uparrow$		$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\sim$	$\uparrow/\sim$	$\uparrow$	$\uparrow$	$\uparrow$
Steatosis	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$			$\sim$	$\sim$	$\uparrow$		
Apoptosis	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow/\sim$		$\uparrow$				$\uparrow$	$\sim$	$\uparrow$
Metabolic alterations												
Glucose oxidation	$\downarrow$	$\sim$		$\downarrow$				$\downarrow$	$\downarrow/\sim$	$\uparrow$	$\downarrow$	$\downarrow$
FA oxidation	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$	$\uparrow$		$\uparrow$	$\downarrow$	$\downarrow$	$\downarrow$	$\uparrow$	$\uparrow$
Mitochondrial function	$\uparrow$	$\uparrow$	$\uparrow$				$\uparrow$	$\uparrow$		$\downarrow$		$\uparrow$
Oxidative stress	$\uparrow$	$\uparrow$	$\uparrow$					$\uparrow$	$\uparrow$		$\uparrow$	$\uparrow$
Ca ²⁺ mobilization				$\sim$	$\sim$				$\downarrow$		$\sim$	
References	[105, 106]	[107, 108]	[109, 110]	[111, 112]	[113, 114]	[96]	[117, 118]	[103, 104]	[97–99]	[115, 116]	[100]	[101] [102]

Genetic overexpression of PPAR α , PPAR γ , long-chain acyl-CoA synthetase-1 (LCACSI), lipoprotein lipase (LPL), fatty acid transport protein-1 (FATP1), PKC β and uncoupling protein-diphtheria toxin A (UCPDTA), or ablation of cardiac-specific insulin receptor (CIRKO), GLUT4, adipose triglyceride lipase (ATGL), phosphoinositide dependent kinase-1 (PDK1), phosphoinositide-3 kinase (PI3K), and glucokinase (GCK) were forced in rodents for DCM. The associated cardiac effects and the levels of natriuretic peptides in plasma are also highlighted for each model. $\uparrow$, $\downarrow$ and $\sim$ stand for increased, decreased, or not modified effect, respectively.

by ablation of the brown adipose tissue in the transgenic model of uncoupling protein-diphtheria toxin A (UCPDTA) [117, 118], mice exhibited a T2DM-like phenotype though a potential rise of related cardiac steatosis and apoptosis was not assessed. However, metabolic changes in all these models were principally represented by an increase of FA oxidation. The glucose metabolism, mitochondrial function, and Ca^{2+} mobilization were unpredictably changed or were not analysed.

3. Major Animal Models Recapitulating the Human DCM Pathology

3.1. T1DM Associated DCM-Like Models. The STZ-administration could recapitulate the most human DCM features associated with T1DM (Table 1). The injection of this β -cell killer can induce cardiac dysfunction in parallel with typical myocardial alterations in the structure and metabolism. However, STZ may also injure nonpancreatic tissues such as brain, and the rapid and accentuated onset of T1DM (i.e., robust hyperglycemia, lipotoxicity, and cardiac apoptosis after only 2-3 days of injection) can diverge from the human nature. In this regard, an upregulation of calmodulin in OVE26 mice may develop a slower and more humanized T1DM-associated DCM, though the energetic imbalance should be further examined. However, this point mutation disagrees with the patient aetiology, in which an autoimmune response is usually the origin of the disease. In addition, the lack of insulin production in Akita leads to some uncommon responses in the myocardium such as the absence of hypertrophy and fibrosis and unchanged levels of oxidation in spite of mitochondrial dysfunction. Nonconventional T1DM models may also emulate the human disorder (Table 2). Aside CIRKO and GLUT4 metabolic controversies, the absence of other mediators for glucose assimilation as GCK may mirror the T1DM phenotype in patients. Also, the lack of PDK1 might reduce the activation of PI3K-dependent insulin signaling [119], leading to a useful T1DM-associated DCM model.

3.2. T2DM Associated DCM-Like Models. Traditional models of T2DM are those with an abolished leptin-system activity (Table 1). Considering leptin activities other than satiety control, these models develop obesity and an average onset of T2DM pathology, with increased hyperglycemia, insulin resistance, and hyperlipidemia, in parallel with cardiac dysfunction. They mostly show expected structural alterations as in the human disorder. However, a polygenic model could be more representative. In this regard, GK rats trigger similar responses but the lack of obesity and steatosis is not usually observed in clinic. More valid models of T2DM-associated DCM can be the DIO. In particular, high-fat diet emulates responses of the leptin-system defect, including obesity and steatosis, but not as a consequence of a single mutation. These effects can be additionally enhanced by coadministration of STZ [120], though tissue toxicity may be increased. Other nonconventional approaches may mimic human DCM in T2DM (Table 2). The upregulation of PPAR α

or PPAR γ unexpectedly leads to similar actions to the lack of leptin stimulation. In spite of increased FA-transporters and FAO enzymes, the myocardium accumulates lipids and ROS. Likely, the level of other PPAR-regulated genes such as reesterification enzymes, glucose transporters, and proinflammatory cytokines could also explain those actions [121]. Moreover, ATGL, LPL, and FATP1 are involved in FA uptake and accumulation in the cardiomyocytes, and LCACS1 catalyses the first step of lipid biosynthesis. In this regard, an excess of FA and lipid content could increase lipotoxicity and subsequent cardiac failure, as occurs in T2DM patients.

4. Conclusions and Future Remarks

Since human trials focused on identifying the cardiovascular outcomes of tight glycemic control failed to show significant benefit [122], the prevention and treatment of DCM may not be limited to the restoration of systemic parameters. It should also include the preservation of cardiac structure and function along with the local metabolic balances (Figure 1). Thus, the knowledge and manipulation of specific key pathways and mediators in appropriate experimental models may be useful for studying new approaches for diabetic patients. Although the experimental milieu (genetic background, diets, and animal location) must be obviously considered, we may have valuable armamentarium to investigate key factors for DCM pathology. In relation with current data, the most appropriate T1DM-model induced by pancreas toxicity is achieved by STZ. However, unique (nontoxic) mutations on glucose-related genes (not on insulin) have demonstrated a DCM-like T1DM phenotype, as occurs with GCK. In addition, due to its possible origin as an autoimmune failure, T1DM could be ideally produced by triggers of specific autoimmune reactions or β -cell death (i.e., virus, modified genes). For T2DM-linked DCM, we may have more working options, including both T2DM-obesity and T2DM aetiologies. Since human T2DM is usually acquired with an unhealthy lifestyle and/or unknown polygenic mutations, high-fat fed models can be closer to that point mutations on leptin system or in lipid storage. Polygenetic mutations of obesity, such as KK A y , and OLETF, should be further characterized, and nonobese polygenetic mutated GK rats could facilitate the study of T2DM *per se*. However, we probably need diet-conditioned models of polygenetic mutations of T2DM to exemplify a *bona fide* approach to a related DCM.

Abbreviations

AGEs:	Advanced glycation end products
ANP:	Atrial natriuretic peptide
AP-1:	Activating protein-1
ATGL:	Adipose triglyceride lipase
Bax:	Bcl-2-associated X protein
Bcl2:	B-cell lymphoma 2
BMP-2:	Bone morphogenetic protein-2
BNP:	Brain natriuretic peptide
DCM:	Diabetic cardiomyopathy
ECM:	Extracellular matrix

EF:	Ejection fraction
FA:	Fatty acid
FAO:	Fatty acid oxidation
FasL:	Fas ligand
FATP1:	Fatty acid transport protein-1
FS:	Fractional shortening
GCK:	Glucokinase
GLUT:	Glucose transporter
ICAM-1:	Intercellular adhesion molecule-1
IFN γ :	Interferon- γ
IL:	Interleukin
LCACSl:	Long-chain acyl-CoA synthetase-1
LV:	Left ventricular
MCP-1:	Monocyte chemoattractant protein-1
MMP:	Matrix metalloproteinase
MRI:	Magnetic resonance imaging
NF κ B:	Nuclear factor of κ -light polypeptide gene enhancer in B-cells
NOX:	NADH oxidase
PARP-1:	Poly(ADP-ribose) polymerase 1
PDK:	Phosphoinositide dependent kinase
PI3K:	Phosphatidylinositol 3-kinase
PKC:	Protein kinase C
PPAR:	Peroxisome proliferator-activated receptor
PW:	Posterior wall
ROS:	Reactive oxygen species
SERCA2a:	Sarco(endo)plasmic reticulum calcium ATPase
TAG:	Triacylglyceride
TGF β :	Transforming growth factor β
TIMP:	Tissue inhibitor of metalloprotease
TNF α :	Tumor necrosis factor α
TUNEL:	Terminal deoxynucleotidyl transferase-(TdT-) mediated dUTP nick end labelling
UCPDTA:	Uncoupling protein- (UCP-) promoter-driven diphtheria toxin A
VCAM-1:	Vascular cell adhesion molecule-1
β -MHC:	β -Myosin heavy chain.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This work was supported by national grants from Ministerio de Educación y Ciencia (SAF2009-08367) and Comunidad de Madrid (CCG10-UAM/BIO-5289).

References

- [1] M. Niemann, S. Herrmann, G. Ertl, and F. Weidemann, "Echocardiography in diabetic cardiomyopathy," *Herz*, vol. 38, no. 1, pp. 42–47, 2013.
- [2] J. N. Khan, E. G. Wilmot, M. Leggate et al., "Subclinical diastolic dysfunction in young adults with Type 2 diabetes mellitus: a multiparametric contrast-enhanced cardiovascular magnetic resonance pilot study assessing potential mechanisms," *European Heart Journal—Cardiovascular Imaging*, vol. 15, no. 11, pp. 1263–1269, 2014.
- [3] C. Jellis, J. Wright, D. Kennedy et al., "Association of imaging markers of myocardial fibrosis with metabolic and functional disturbances in early diabetic cardiomyopathy," *Circulation: Cardiovascular Imaging*, vol. 4, no. 6, pp. 693–702, 2011.
- [4] S. Gao, D. Ho, D. E. Vatner, and S. F. Vatner, "Echocardiography in mice," *Current Protocols in Mouse Biology*, vol. 1, pp. 71–83, 2011.
- [5] C. Christoffersen, E. Bollano, M. L. S. Lindegaard et al., "Cardiac lipid accumulation associated with diastolic dysfunction in obese mice," *Endocrinology*, vol. 144, no. 8, pp. 3483–3490, 2003.
- [6] L. M. Semeniuk, A. J. Kryski, and D. L. Severson, "Echocardiographic assessment of cardiac function in diabetic *db/db* and transgenic *db/db*-hGLUT4 mice," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 283, no. 3, pp. H976–H982, 2002.
- [7] A. van den Bergh, W. Flameng, and P. Herijgers, "Type II diabetic mice exhibit contractile dysfunction but maintain cardiac output by favourable loading conditions," *European Journal of Heart Failure*, vol. 8, no. 8, pp. 777–783, 2006.
- [8] Y.-T. Zhou, P. Grayburn, A. Karim et al., "Lipotoxic heart disease in obese rats: implications for human obesity," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 4, pp. 1784–1789, 2000.
- [9] A. Daniels, D. Linz, M. van Bilsen et al., "Long-term severe diabetes only leads to mild cardiac diastolic dysfunction in Zucker diabetic fatty rats," *European Journal of Heart Failure*, vol. 14, no. 2, pp. 193–201, 2012.
- [10] E. Ramírez, M. Klett-Mingo, S. Ares-Carrasco et al., "Eplerenone attenuated cardiac steatosis, apoptosis and diastolic dysfunction in experimental type-II diabetes," *Cardiovascular Diabetology*, vol. 12, no. 1, article 172, 2013.
- [11] K. Mizushige, L. Yao, T. Noma et al., "Alteration in left ventricular diastolic filling and accumulation of myocardial collagen at insulin-resistant prediabetic stage of a type II diabetic rat model," *Circulation*, vol. 101, no. 8, pp. 899–907, 2000.
- [12] B. Picatoste, E. Ramírez, A. Caro-Vadillo et al., "Sitagliptin reduces cardiac apoptosis, hypertrophy and fibrosis primarily by insulin-Dependent mechanisms in experimental type-II diabetes. Potential roles of GLP-1 isoforms," *PLoS ONE*, vol. 8, no. 10, Article ID e78330, 2013.
- [13] C. X. Fang, F. Dong, D. P. Thomas, H. Ma, L. He, and J. Ren, "Hypertrophic cardiomyopathy in high-fat diet-induced obesity: role of suppression of forkhead transcription factor and atrophy gene transcription," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 295, no. 3, pp. H1206–H1215, 2008.
- [14] Z. Vasanji, E. J. F. Cantor, D. Juric, M. Moyen, and T. Netticadan, "Alterations in cardiac contractile performance and sarcoplasmic reticulum function in sucrose-fed rats is associated with insulin resistance," *The American Journal of Physiology—Cell Physiology*, vol. 291, no. 4, pp. C772–C780, 2006.
- [15] T. Pulinilkunnil, P. C. Kienesberger, J. Nagendran, N. Sharma, M. E. Young, and J. R. B. Dyck, "Cardiac-specific adipose triglyceride lipase overexpression protects from cardiac steatosis and dilated cardiomyopathy following diet-induced obesity," *International Journal of Obesity*, vol. 38, no. 2, pp. 205–215, 2014.
- [16] J. B. Somaratne, G. A. Whalley, K. K. Poppe et al., "Screening for left ventricular hypertrophy in patients with type 2 diabetes

- mellitus in the community," *Cardiovascular Diabetology*, vol. 10, article 29, 2011.
- [17] A. Dei Cas, V. Spigoni, V. Ridolfi, and M. Metra, "Diabetes and chronic heart failure: from diabetic cardiomyopathy to therapeutic approach," *Endocrine, Metabolic and Immune Disorders—Drug Targets*, vol. 13, no. 1, pp. 38–50, 2013.
 - [18] S. Zibadi, F. Cordova, E. H. Slack, R. R. Watson, and D. F. Larson, "Leptin's regulation of obesity-induced cardiac extracellular matrix remodeling," *Cardiovascular Toxicology*, vol. 11, no. 4, pp. 325–333, 2011.
 - [19] E. J. Cox and S. A. Marsh, "A systematic review of fetal genes as biomarkers of cardiac hypertrophy in rodent models of diabetes," *PLoS ONE*, vol. 9, no. 3, Article ID e92903, 2014.
 - [20] O. Nemoto, M. Kawaguchi, H. Yaoita, K. Miyake, K. Maehara, and Y. Maruyama, "Left ventricular dysfunction and remodeling in streptozotocin-induced diabetic rats," *Circulation Journal*, vol. 70, no. 3, pp. 327–334, 2006.
 - [21] S. Ares-Carrasco, B. Picatoste, A. Benito-Martín et al., "Myocardial fibrosis and apoptosis, but not inflammation, are present in long-term experimental diabetes," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 297, no. 6, pp. H2109–H2119, 2009.
 - [22] Y. Li, J. Ma, H. Zhu et al., "Targeted inhibition of calpain reduces myocardial hypertrophy and fibrosis in mouse models of type 1 diabetes," *Diabetes*, vol. 60, no. 11, pp. 2985–2994, 2011.
 - [23] Y. Wang, W. Sun, B. Du et al., "Therapeutic effect of MG-132 on diabetic cardiomyopathy is associated with its suppression of proteasomal activities: roles of Nrf2 and NF- κ B," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 304, no. 4, pp. H567–H578, 2013.
 - [24] R. Basu, G. Y. Oudit, X. Wang et al., "Type 1 diabetic cardiomyopathy in the Akita (Ins2WT/IC96Y) mouse model is characterized by lipotoxicity and diastolic dysfunction with preserved systolic function," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 297, no. 6, pp. H2096–H2108, 2009.
 - [25] A. Van Den Bergh, A. Vanderper, P. Vangheluwe et al., "Dyslipidaemia in type II diabetic mice does not aggravate contractile impairment but increases ventricular stiffness," *Cardiovascular Research*, vol. 77, no. 2, pp. 371–379, 2008.
 - [26] P. Yue, T. Arai, M. Terashima et al., "Magnetic resonance imaging of progressive cardiomyopathic changes in the db/db mouse," *American Journal of Physiology: Heart and Circulatory Physiology*, vol. 292, no. 5, pp. H2106–H2118, 2007.
 - [27] L. Pereira, J. Matthes, I. Schuster et al., "Mechanisms of $[Ca^{2+}]_i$ transient decrease in cardiomyopathy of db/db type 2 diabetic mice," *Diabetes*, vol. 55, no. 3, pp. 608–615, 2006.
 - [28] S. Fredersdorf, C. Thumann, C. Ulucan et al., "Myocardial hypertrophy and enhanced left ventricular contractility in Zucker diabetic fatty rats," *Cardiovascular Pathology*, vol. 13, no. 1, pp. 11–19, 2004.
 - [29] M. Conti, I. M. Renaud, B. Poirier et al., "High levels of myocardial antioxidant defense in aging nondiabetic normotensive Zucker obese rats," *The American Journal of Physiology—Regulatory Integrative and Comparative Physiology*, vol. 286, no. 4, pp. R793–R800, 2004.
 - [30] M. Desrois, R. J. Sidell, D. Gauguier, C. L. Davey, G. K. Radda, and K. Clarke, "Gender differences in hypertrophy, insulin resistance and ischemic injury in the aging type 2 diabetic rat heart," *Journal of Molecular and Cellular Cardiology*, vol. 37, no. 2, pp. 547–555, 2004.
 - [31] S. Devanathan, S. T. Nemanich, A. Kovacs, N. Fettig, R. J. Gropler, and K. I. Shoghi, "Genomic and metabolic disposition of non-obese type 2 diabetic rats to increased myocardial Fatty Acid metabolism," *PLoS ONE*, vol. 8, no. 10, Article ID e78477, 2013.
 - [32] S. Nunes, E. Soares, J. Fernandes et al., "Early cardiac changes in a rat model of prediabetes: brain natriuretic peptide overexpression seems to be the best marker," *Cardiovascular Diabetology*, vol. 12, article 44, 2013.
 - [33] L. N. Axelsen, J. B. Lademann, J. S. Petersen et al., "Cardiac and metabolic changes in long-term high fructose-fat fed rats with severe obesity and extensive intramyocardial lipid accumulation," *The American Journal of Physiology—Regulatory Integrative and Comparative Physiology*, vol. 298, no. 6, pp. R1560–R1570, 2010.
 - [34] A. K. M. Tarikuz Zaman, D. L. McLean, and B. E. Sobel, "The efficacy and tolerability of azilsartan in obese insulin-resistant mice with left ventricular pressure overload," *Journal of Cardiovascular Pharmacology*, vol. 62, no. 4, pp. 381–387, 2013.
 - [35] M. H. Noyan-Ashraf, E. A. Shikatani, I. Schuiki et al., "A glucagon-like peptide-1 analog reverses the molecular pathology and cardiac dysfunction of a mouse model of obesity," *Circulation*, vol. 127, no. 1, pp. 74–85, 2013.
 - [36] J. D. Schilling and D. L. Mann, "Diabetic cardiomyopathy: bench to bedside," *Heart Failure Clinics*, vol. 8, no. 4, pp. 619–631, 2012.
 - [37] D. L. Mann, "Inflammatory mediators and the failing heart: past, present, and the foreseeable future," *Circulation Research*, vol. 91, no. 11, pp. 988–998, 2002.
 - [38] W. Huang and C. K. Glass, "Nuclear receptors and inflammation control: molecular mechanisms and pathophysiological relevance," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 30, no. 8, pp. 1542–1549, 2010.
 - [39] J. Fuentes-Antrás, A. M. Ioan, J. Tuñón, J. Egido, and Ó. Lorenzo, "Activation of toll-like receptors and inflammasome complexes in the diabetic cardiomyopathy-associated inflammation," *International Journal of Endocrinology*, vol. 2014, Article ID 847827, 10 pages, 2014.
 - [40] V. Chavali, S. C. Tyagi, and P. K. Mishra, "Differential expression of dicer, miRNAs, and inflammatory markers in diabetic Ins2 $^{+/-}$ akita hearts," *Cell Biochemistry and Biophysics*, vol. 68, no. 1, pp. 25–35, 2014.
 - [41] J. Mori, V. B. Patel, O. Abo Alrob et al., "Angiotensin 1-7 ameliorates diabetic cardiomyopathy and diastolic dysfunction in db/db mice by reducing lipotoxicity and inflammation," *Circulation. Heart Failure*, vol. 7, no. 2, pp. 327–339, 2014.
 - [42] A. Jadhav, S. Tiwari, P. Lee, and J. F. Ndisang, "The heme oxygenase system selectively enhances the anti-inflammatory macrophage-m2 phenotype, reduces pericardial adiposity, and ameliorated cardiac injury in diabetic cardiomyopathy in Zucker diabetic fatty rats," *Journal of Pharmacology and Experimental Therapeutics*, vol. 345, no. 2, pp. 239–249, 2013.
 - [43] H.-T. Wang, C.-F. Liu, T.-H. Tsai et al., "Effect of obesity reduction on preservation of heart function and attenuation of left ventricular remodeling, oxidative stress and inflammation in obese mice," *Journal of Translational Medicine*, vol. 10, no. 1, article 145, 2012.
 - [44] D. Álvarez-Guardia, X. Palomer, T. Coll et al., "PPAR β/δ activation blocks lipid-induced inflammatory pathways in mouse heart and human cardiac cells," *Biochimica et Biophysica Acta*, vol. 1811, no. 2, pp. 59–67, 2011.

- [45] K. J. Way, K. Isshiki, K. Suzuma et al., "Expression of connective tissue growth factor is increased in injured myocardium associated with protein kinase C $\beta 2$ activation and diabetes," *Diabetes*, vol. 51, no. 9, pp. 2709–2718, 2002.
- [46] S. Katayama, M. Abe, K. Negishi, K. Takahashi, J. Ishii, and K. Komeda, "Reciprocal changes in left ventricular collagen $\alpha 1$ chain gene expression between types I and IV in spontaneously diabetic rats," *Diabetes Research and Clinical Practice*, vol. 26, no. 3, pp. 163–169, 1994.
- [47] J. E. Lee, C.-O. Yi, B. T. Jeon et al., " α -lipoic acid attenuates cardiac fibrosis in Otsuka Long-Evans Tokushima Fatty rats," *Cardiovascular Diabetology*, vol. 11, article 111, 2012.
- [48] G. F. Lewis, A. Carpentier, K. Adeli, and A. Giacca, "Disordered fat storage and mobilization in the pathogenesis of insulin resistance and type 2 diabetes," *Endocrine Reviews*, vol. 23, no. 2, pp. 201–229, 2002.
- [49] G. Korosoglou, P. M. Humpert, J. Ahrens et al., "Left ventricular diastolic function in type 2 diabetes mellitus is associated with myocardial triglyceride content but not with impaired myocardial perfusion reserve," *Journal of Magnetic Resonance Imaging*, vol. 35, no. 4, pp. 804–811, 2012.
- [50] M. Ueno, J. Suzuki, Y. Zenimaru et al., "Cardiac overexpression of hormone-sensitive lipase inhibits myocardial steatosis and fibrosis in streptozotocin diabetic mice," *The American Journal of Physiology—Endocrinology and Metabolism*, vol. 294, no. 6, pp. E1109–E1118, 2008.
- [51] B. Dong, D. Qi, L. Yang et al., "TLR4 regulates cardiac lipid accumulation and diabetic heart disease in the nonobese diabetic mouse model of type 1 diabetes," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 303, no. 6, pp. H732–H742, 2012.
- [52] L. A. Barouch, D. Gao, L. Chen et al., "Cardiac myocyte apoptosis is associated with increased DNA damage and decreased survival in murine models of obesity," *Circulation Research*, vol. 98, no. 1, pp. 119–124, 2006.
- [53] Y. S. Diniz, P. P. Santos, H. B. Assalin et al., "Conjugated linoleic acid and cardiac health: oxidative stress and energetic metabolism in standard and sucrose-rich diets," *European Journal of Pharmacology*, vol. 579, no. 1–3, pp. 318–325, 2008.
- [54] A. Frustaci, J. Kajstura, C. Chimenti et al., "Myocardial cell death in human diabetes," *Circulation Research*, vol. 87, no. 12, pp. 1123–1132, 2000.
- [55] S. A. Andrabi, T. M. Dawson, and V. L. Dawson, "Mitochondrial and nuclear cross talk in cell death: parthanatos," *Annals of the New York Academy of Sciences*, vol. 1147, pp. 233–241, 2008.
- [56] P. Puthanveetil, D. Zhang, Y. Wang et al., "Diabetes triggers a PARP1 mediated death pathway in the heart through participation of FoxO1," *Journal of Molecular and Cellular Cardiology*, vol. 53, no. 5, pp. 677–686, 2012.
- [57] F. Fiordaliso, B. Li, R. Latini et al., "Myocyte death in streptozotocin-induced diabetes in rats is angiotensin II-dependent," *Laboratory Investigation*, vol. 80, no. 4, pp. 513–527, 2000.
- [58] Z. Xie, K. Lau, B. Eby et al., "Improvement of cardiac functions by chronic metformin treatment is associated with enhanced cardiac autophagy in diabetic OVE26 mice," *Diabetes*, vol. 60, no. 6, pp. 1770–1778, 2011.
- [59] E. Shen, Y. Li, L. Shan et al., "Rac1 is required for cardiomyocyte apoptosis during hyperglycemia," *Diabetes*, vol. 58, no. 10, pp. 2386–2395, 2009.
- [60] R. S. Guleria, R. Choudhary, T. Tanaka, K. M. Baker, and J. Pan, "Retinoic acid receptor-mediated signaling protects cardiomyocytes from hyperglycemia induced apoptosis: role of the renin-angiotensin system," *Journal of Cellular Physiology*, vol. 226, no. 5, pp. 1292–1307, 2011.
- [61] H. Matsui, T. Yokoyama, K. Sekiguchi et al., "Stearoyl-CoA desaturase-1 (SCD1) augments saturated fatty acid-induced lipid accumulation and inhibits apoptosis in cardiac myocytes," *PLoS ONE*, vol. 7, no. 3, Article ID e33283, 2012.
- [62] X. Shen, S. Zheng, V. Thongboonkerd et al., "Cardiac mitochondrial damage and biogenesis in a chronic model of type 1 diabetes," *American Journal of Physiology—Endocrinology and Metabolism*, vol. 287, no. 5, pp. E896–E905, 2004.
- [63] E. Aasum, A. D. Hafstad, D. L. Severson, and T. S. Larsen, "Age-dependent changes in metabolism, contractile function, and ischemic sensitivity in hearts from db/db mice," *Diabetes*, vol. 52, no. 2, pp. 434–441, 2003.
- [64] P. K. Mazumder, B. T. O'Neill, M. W. Roberts et al., "Impaired cardiac efficiency and increased fatty acid oxidation in insulin-resistant ob/ob mouse hearts," *Diabetes*, vol. 53, no. 9, pp. 2366–2374, 2004.
- [65] K. Ballal, C. R. Wilson, R. Harmancey, and H. Taegtmeier, "Obesogenic high fat western diet induces oxidative stress and apoptosis in rat heart," *Molecular and Cellular Biochemistry*, vol. 344, no. 1–2, pp. 221–230, 2010.
- [66] S. Nunes, E. Soares, J. Fernandes et al., "Early cardiac changes in a rat model of prediabetes: brain natriuretic peptide overexpression seems to be the best marker," *Cardiovascular Diabetology*, vol. 12, no. 1, article 44, 2013.
- [67] Z. Y. Fang, J. B. Prins, and T. H. Marwick, "Diabetic cardiomyopathy: evidence, mechanisms, and therapeutic implications," *Endocrine Reviews*, vol. 25, no. 4, pp. 543–567, 2004.
- [68] H. Bugger and E. D. Abel, "Mitochondria in the diabetic heart," *Cardiovascular Research*, vol. 88, no. 2, pp. 229–240, 2010.
- [69] T.-I. Lee, Y.-H. Kao, Y.-C. Chen, J.-H. Huang, F.-C. Hsiao, and Y.-J. Chen, "Peroxisome proliferator-activated receptors modulate cardiac dysfunction in diabetic cardiomyopathy," *Diabetes Research and Clinical Practice*, vol. 100, no. 3, pp. 330–339, 2013.
- [70] C. E. Flarsheim, I. L. Grupp, and M. A. Matlib, "Mitochondrial dysfunction accompanies diastolic dysfunction in diabetic rat heart," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 271, no. 1, part 2, pp. H192–H202, 1996.
- [71] H. Bugger, S. Boudina, X. X. Hu et al., "Type 1 diabetic akita mouse hearts are insulin sensitive but manifest structurally abnormal mitochondria that remain coupled despite increased uncoupling protein 3," *Diabetes*, vol. 57, no. 11, pp. 2924–2932, 2008.
- [72] L. S. Golfman, C. R. Wilson, S. Sharma et al., "Activation of PPAR γ enhances myocardial glucose oxidation and improves contractile function in isolated working hearts of ZDF rats," *American Journal of Physiology—Endocrinology and Metabolism*, vol. 289, no. 2, pp. E328–E336, 2005.
- [73] M. A. Cole, A. J. Murray, L. E. Cochlin et al., "A high fat diet increases mitochondrial fatty acid oxidation and uncoupling to decrease efficiency in rat heart," *Basic Research in Cardiology*, vol. 106, no. 3, pp. 447–457, 2011.
- [74] J. R. Ussher, T. R. Koves, J. S. Jaswal et al., "Insulin-stimulated cardiac glucose oxidation is increased in high-fat diet-induced obese mice lacking malonyl CoA decarboxylase," *Diabetes*, vol. 58, pp. 1766–1775, 2009.
- [75] O. Ilkun and S. Boudina, "Cardiac dysfunction and oxidative stress in the metabolic syndrome: an update on antioxidant

- therapies," *Current Pharmaceutical Design*, vol. 19, no. 27, pp. 4806–4817, 2013.
- [76] S. Ghosh, T. Pulinilkunnil, G. Yuen et al., "Cardiomyocyte apoptosis induced by short-term diabetes requires mitochondrial GSH depletion," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 289, no. 2, pp. H768–H776, 2005.
- [77] M. Hamblin, D. B. Friedman, S. Hill, R. M. Caprioli, H. M. Smith, and M. F. Hill, "Alterations in the diabetic myocardial proteome coupled with increased myocardial oxidative stress underlies diabetic cardiomyopathy," *Journal of Molecular and Cellular Cardiology*, vol. 42, no. 4, pp. 884–895, 2007.
- [78] Y. Song, Y. Du, S. D. Prabhu, and P. N. Epstein, "Diabetic cardiomyopathy in OVE26 mice shows mitochondrial ROS production and divergence between in vivo and in vitro contractility," *Review of Diabetic Studies*, vol. 4, no. 3, pp. 159–168, 2007.
- [79] Q. Liang, E. C. Carlson, R. V. Donthi, P. M. Kralik, X. Shen, and P. N. Epstein, "Overexpression of metallothionein reduces diabetic cardiomyopathy," *Diabetes*, vol. 51, no. 1, pp. 174–181, 2002.
- [80] S. Boudina, S. Sena, H. Theobald et al., "Mitochondrial energetics in the heart in obesity-related diabetes: direct evidence for increased uncoupled respiration and activation of uncoupling proteins," *Diabetes*, vol. 56, no. 10, pp. 2457–2466, 2007.
- [81] S. Boudina, S. Sena, B. T. O'Neill, P. Tathireddy, M. E. Young, and E. D. Abel, "Reduced mitochondrial oxidative capacity and increased mitochondrial uncoupling impair myocardial energetics in obesity," *Circulation*, vol. 112, no. 17, pp. 2686–2695, 2005.
- [82] S. Boudina and E. D. Abel, "Diabetic cardiomyopathy revisited," *Circulation*, vol. 115, no. 25, pp. 3213–3223, 2007.
- [83] J. P. Vázquez-Medina, I. Popovich, M. A. Thorwald et al., "Angiotensin receptor-mediated oxidative stress is associated with impaired cardiac redox signaling and mitochondrial function in insulin-resistant rats," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 305, no. 4, pp. H599–H607, 2013.
- [84] D. L. Santos, C. M. Palmeira, R. Seica et al., "Diabetes and mitochondrial oxidative stress: a study using heart mitochondria from the diabetic Goto-Kakizaki rat," *Molecular and Cellular Biochemistry*, vol. 246, no. 1–2, pp. 163–170, 2003.
- [85] J. Grijalva, S. Hicks, X. Zhao et al., "Exercise training enhanced myocardial endothelial nitric oxide synthase (eNOS) function in diabetic Goto-Kakizaki (GK) rats," *Cardiovascular Diabetology*, vol. 7, article 34, 2008.
- [86] H. K. Vincent, S. K. Powers, A. J. Dirks, and P. J. Scarpace, "Mechanism for obesity-induced increase in myocardial lipid peroxidation," *International Journal of Obesity*, vol. 25, no. 3, pp. 378–388, 2001.
- [87] P. Wang, S. G. Lloyd, H. Zeng, A. Bonen, and J. C. Chatham, "Impact of altered substrate utilization on cardiac function in isolated hearts from Zucker diabetic fatty rats," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 288, no. 5, pp. H2102–H2110, 2005.
- [88] B. Turan and G. Vassort, "Ryanodine receptor: a new therapeutic target to control diabetic cardiomyopathy," *Antioxidants and Redox Signaling*, vol. 15, no. 7, pp. 1847–1861, 2011.
- [89] C. E. Flarsheim, I. L. Grupp, and M. A. Matlib, "Mitochondrial dysfunction accompanies diastolic dysfunction in diabetic rat heart," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 271, no. 1, pp. H192–H202, 1996.
- [90] P. M. Kralik, G. Ye, N. S. Metreveli, X. Shen, and P. N. Epstein, "Cardiomyocyte dysfunction in models of type 1 and type 2 diabetes," *Cardiovascular Toxicology*, vol. 5, no. 3, pp. 285–292, 2005.
- [91] J. Ren and A. M. Bode, "Altered cardiac excitation-contraction coupling in ventricular myocytes from spontaneously diabetic BB rats," *American Journal of Physiology: Heart and Circulatory Physiology*, vol. 279, no. 1, pp. H238–H244, 2000.
- [92] S.-Y. Li, X. Yang, A. F. Ceylan-Isik, M. Du, N. Sreejayan, and J. Ren, "Cardiac contractile dysfunction in *Lep/Lep* obesity is accompanied by NADPH oxidase activation, oxidative modification of sarco(endo)plasmic reticulum Ca^{2+} -ATPase and myosin heavy chain isozyme switch," *Diabetologia*, vol. 49, no. 6, pp. 1434–1446, 2006.
- [93] D. D. Belke, E. A. Swanson, and W. H. Dillmann, "Decreased sarcoplasmic reticulum activity and contractility in diabetic db/db mouse heart," *Diabetes*, vol. 53, no. 12, pp. 3201–3208, 2004.
- [94] K. A. Salem, T. E. Adrian, M. A. Qureshi, K. Parekh, M. Oz, and F. C. Howarth, "Shortening and intracellular Ca^{2+} in ventricular myocytes and expression of genes encoding cardiac muscle proteins in early onset type 2 diabetic Goto-Kakizaki rats," *Experimental Physiology*, vol. 97, no. 12, pp. 1281–1291, 2012.
- [95] K. M. Mellor, I. R. Wendt, R. H. Ritchie, and L. M. D. Delbridge, "Fructose diet treatment in mice induces fundamental disturbance of cardiomyocyte Ca^{2+} handling and myofilament responsiveness," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 302, no. 4, pp. H964–H972, 2012.
- [96] H. Wakasaki, D. Koya, F. J. Schoen et al., "Targeted overexpression of protein kinase C $\beta 2$ isoform in myocardium causes cardiomyopathy," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 17, pp. 9320–9325, 1997.
- [97] E. D. Abel, H. C. Kaulbach, R. Tian et al., "Cardiac hypertrophy with preserved contractile function after selective deletion of GLUT4 from the heart," *The Journal of Clinical Investigation*, vol. 104, no. 12, pp. 1703–1714, 1999.
- [98] A. A. Domenighetti, V. R. Danes, C. L. Curl, J. M. Favaloro, J. Proietto, and L. M. D. Delbridge, "Targeted GLUT-4 deficiency in the heart induces cardiomyocyte hypertrophy and impaired contractility linked with Ca^{2+} and proton flux dysregulation," *Journal of Molecular and Cellular Cardiology*, vol. 48, no. 4, pp. 663–672, 2010.
- [99] Y. Li, A. R. Wende, O. Nunthakungwan et al., "Cytosolic, but not mitochondrial, oxidative stress is a likely contributor to cardiac hypertrophy resulting from cardiac specific GLUT4 deletion in mice," *FEBS Journal*, vol. 279, no. 4, pp. 599–611, 2012.
- [100] A. Mora, A. M. Davies, L. Bertrand et al., "Deficiency of PDK1 in cardiac muscle results in heart failure and increased sensitivity to hypoxia," *The EMBO Journal*, vol. 22, no. 18, pp. 4666–4676, 2003.
- [101] R. H. Ritchie, J. E. Love, K. Huynh et al., "Enhanced phosphoinositide 3-kinase($\text{pI}10\alpha$) activity prevents diabetes-induced cardiomyopathy and superoxide generation in a mouse model of diabetes," *Diabetologia*, vol. 55, no. 12, pp. 3369–3381, 2012.
- [102] H. Li, X. Wang, Y. Mao et al., "Long term liver specific glucokinase gene defect induced diabetic cardiomyopathy by up regulating NADPH oxidase and down regulating insulin receptor and p-AMPK," *Cardiovascular Diabetology*, vol. 13, no. 1, article 24, 2014.

- [103] D. D. Belke, S. Betuing, M. J. Tuttle et al., "Insulin signaling coordinately regulates cardiac size, metabolism, and contractile protein isoform expression," *Journal of Clinical Investigation*, vol. 109, no. 5, pp. 629–639, 2002.
- [104] S. Boudina, H. Bugger, S. Sena et al., "Contribution of impaired myocardial insulin signaling to mitochondrial dysfunction and oxidative stress in the heart," *Circulation*, vol. 119, no. 9, pp. 1272–1283, 2009.
- [105] B. N. Finck, X. Han, M. Courtois et al., "A critical role for PPAR α -mediated lipotoxicity in the pathogenesis of diabetic cardiomyopathy: modulation by dietary fat content," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 3, pp. 1226–1231, 2003.
- [106] B. N. Finck, J. J. Lehman, T. C. Leone et al., "The cardiac phenotype induced by PPAR α overexpression mimics that caused by diabetes mellitus," *Journal of Clinical Investigation*, vol. 109, no. 1, pp. 121–130, 2002.
- [107] N.-H. Son, T.-S. Park, H. Yamashita et al., "Cardiomyocyte expression of PPAR γ leads to cardiac dysfunction in mice," *Journal of Clinical Investigation*, vol. 117, no. 10, pp. 2791–2801, 2007.
- [108] N.-H. Son, S. Yu, J. Tuinei et al., "PPAR γ -induced cardiolipotoxicity in mice is ameliorated by PPAR α deficiency despite increases in fatty acid oxidation," *Journal of Clinical Investigation*, vol. 120, no. 10, pp. 3443–3454, 2010.
- [109] H.-C. Chiu, A. Kovacs, D. A. Ford et al., "A novel mouse model of lipotoxic cardiomyopathy," *Journal of Clinical Investigation*, vol. 107, no. 7, pp. 813–822, 2001.
- [110] L. Liu, X. J. Shi, K. G. Bharadwaj et al., "DGAT1 expression increases heart triglyceride content but ameliorates lipotoxicity," *The Journal of Biological Chemistry*, vol. 284, no. 52, pp. 36312–36323, 2009.
- [111] H. Yagyu, G. Chen, M. Yokoyama et al., "Lipoprotein lipase (LpL) on the surface of cardiomyocytes increases lipid uptake and produces a cardiomyopathy," *Journal of Clinical Investigation*, vol. 111, no. 3, pp. 419–426, 2003.
- [112] M. Yokoyama, H. Yagyu, Y. Hu et al., "Apolipoprotein B production reduces lipotoxic cardiomyopathy. Studies in heart-specific lipoprotein lipase transgenic mouse," *The Journal of Biological Chemistry*, vol. 279, no. 6, pp. 4204–4211, 2004.
- [113] H.-C. Chiu, A. Kovacs, R. M. Blanton et al., "Transgenic expression of fatty acid transport protein 1 in the heart causes lipotoxic cardiomyopathy," *Circulation Research*, vol. 96, no. 2, pp. 225–233, 2005.
- [114] T. P. Flagg, O. Cazorla, M. S. Remedi et al., "Ca²⁺-independent alterations in diastolic sarcomere length and relaxation kinetics in a mouse model of lipotoxic diabetic cardiomyopathy," *Circulation Research*, vol. 104, no. 1, pp. 95–103, 2009.
- [115] G. Haemmerle, A. Lass, R. Zimmermann et al., "Defective lipolysis and altered energy metabolism in mice lacking adipose triglyceride lipase," *Science*, vol. 312, no. 5774, pp. 734–737, 2006.
- [116] P. C. Kienesberger, T. Pulini, J. Nagendran et al., "Early structural and metabolic cardiac remodeling in response to inducible adipose triglyceride lipase ablation," *Cardiovascular Research*, vol. 99, no. 3, pp. 442–451, 2013.
- [117] A. Cittadini, C. S. Mantzoros, T. G. Hampton et al., "Cardiovascular abnormalities in transgenic mice with reduced brown fat: an animal model of human obesity," *Circulation*, vol. 100, no. 21, pp. 2177–2183, 1999.
- [118] J. G. Duncan, J. L. Fong, D. M. Medeiros, B. N. Finck, and D. P. Kelly, "Insulin-resistant heart exhibits a mitochondrial biogenic response driven by the peroxisome proliferator-activated receptor- α /PGC-1 α gene regulatory pathway," *Circulation*, vol. 115, no. 7, pp. 909–917, 2007.
- [119] J. Y. Cho and J. Park, "Contribution of natural inhibitors to the understanding of the PI3K/PDK1/PKB pathway in the insulin-mediated intracellular signaling cascade," *International Journal of Molecular Sciences*, vol. 9, no. 11, pp. 2217–2230, 2008.
- [120] S. A. Marsh, L. J. Dell'Italia, and J. C. Chatham, "Interaction of diet and diabetes on cardiovascular function in rats," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 296, no. 2, pp. H282–H292, 2009.
- [121] J. Fuentes-Antrás, B. Picatoste, E. Ramírez, J. Egido, J. Tuñón, and Ó. Lorenzo, "Targeting metabolic disturbance in the diabetic heart," *Cardiovascular Diabetology*, vol. 14, article 17, 2015.
- [122] T. Mazzone, "Intensive glucose lowering and cardiovascular disease prevention in diabetes: reconciling the recent clinical trial data," *Circulation*, vol. 122, no. 21, pp. 2201–2211, 2010.
- [123] S. Islas-Andrade, M. C. R. Monsalve, J. E. De La Peña, A. C. Polanco, M. A. Palomino, and A. F. Velasco, "Streptozotocin and alloxan in experimental diabetes: comparison of the two models in rats," *Acta Histochemica et Cytochemica*, vol. 33, no. 3, pp. 201–208, 2000.
- [124] G. D. Lopaschuk and H. Tsang, "Metabolism of palmitate in isolated working hearts from spontaneously diabetic 'BB' Wistar rats," *Circulation Research*, vol. 61, no. 6, pp. 853–858, 1987.
- [125] J. Buchanan, P. K. Mazumder, P. Hu et al., "Reduced cardiac efficiency and altered substrate metabolism precedes the onset of hyperglycemia and contractile dysfunction in two mouse models of insulin resistance and obesity," *Endocrinology*, vol. 146, no. 12, pp. 5341–5349, 2005.
- [126] F. Forcheron, A. Basset, P. Abdallah, P. D. Carmine, N. Gadot, and M. Beylot, "Diabetic cardiomyopathy: effects of fenofibrate and metformin in an experimental model—the Zucker diabetic rat," *Cardiovascular Diabetology*, vol. 8, article 16, 2009.
- [127] H. Iwatsuka, A. Shino, and Z. Suzuoki, "General survey of diabetic features of yellow KK mice," *Endocrinologia Japonica*, vol. 17, no. 1, pp. 23–35, 1970.
- [128] K. Kawano, T. Hirashima, S. Mori, Y. Saitoh, M. Kurosumi, and T. Natori, "Spontaneous long-term hyperglycemic rat with diabetic complications: Otsuka Long-Evans Tokushima Fatty (OLETF) strain," *Diabetes*, vol. 41, no. 11, pp. 1422–1428, 1992.
- [129] C. Tourrel, D. Bailbe, M. Lacorne, M.-J. Meile, M. Kergoat, and B. Portha, "Persistent improvement of type 2 diabetes in the Goto-Kakizaki rat model by expansion of the β -cell mass during the prediabetic period with glucagon-like peptide-1 or exendin-4," *Diabetes*, vol. 51, no. 5, pp. 1443–1452, 2002.
- [130] M. S. Winzell and B. Ahrén, "The high-fat diet-fed mouse: a model for studying mechanisms and treatment of impaired glucose tolerance and type 2 diabetes," *Diabetes*, vol. 53, supplement 3, pp. S215–S219, 2004.
- [131] H. Bar On, P. S. Roheim, and H. A. Eder, "Serum lipoproteins and apolipoproteins in rats with streptozotocin induced diabetes," *Journal of Clinical Investigation*, vol. 57, no. 3, pp. 714–721, 1976.
- [132] J. Altomonte, L. Cong, S. Harbaran et al., "Foxo1 mediates insulin action on apoC-III and triglyceride metabolism," *The Journal of Clinical Investigation*, vol. 114, no. 10, pp. 1493–1503, 2004.
- [133] C. Tang, J. E. Kanter, K. E. Bornfeldt, R. C. Leboeuf, and J. F. Oram, "Diabetes reduces the cholesterol exporter ABCA1 in mouse macrophages and kidneys," *Journal of Lipid Research*, vol. 51, no. 7, pp. 1719–1728, 2010.

- [134] P. Naveilhan, L. Svensson, S. Nyström, A. J. Ekstrand, and P. Ern-fors, "Attenuation of hypercholesterolemia and hyperglycemia in ob/ob mice by NPY Y2 receptor ablation," *Peptides*, vol. 23, no. 6, pp. 1087–1091, 2002.
- [135] K. Kobayashi, T. M. Forte, S. Taniguchi, B. Y. Ishida, K. Oka, and L. Chan, "The db/db mouse, a model for diabetic dyslipidemia: Molecular characterization and effects of western diet feeding," *Metabolism: Clinical and Experimental*, vol. 49, no. 1, pp. 22–31, 2000.
- [136] J. D. Sparks, T. L. Phung, M. Bolognino et al., "Lipoprotein alterations in 10- and 20-week-old Zucker diabetic fatty rats: hyperinsulinemic versus insulinopenic hyperglycemia," *Metabolism: Clinical and Experimental*, vol. 47, no. 11, pp. 1315–1324, 1998.
- [137] T. Shimizugawa, M. Ono, M. Shimamura et al., "ANGPTL3 decreases very low density lipoprotein triglyceride clearance by inhibition of lipoprotein lipase," *The Journal of Biological Chemistry*, vol. 277, no. 37, pp. 33742–33748, 2002.
- [138] S. D. Calligaris, M. Lecanda, F. Solis et al., "Mice long-term high-fat diet feeding recapitulates human cardiovascular alterations: an animal model to study the early phases of diabetic cardiomyopathy," *PLoS ONE*, vol. 8, no. 4, Article ID e60931, 2013.
- [139] S. A. Oliveira-Junior, P. F. Martinez, D. M. Guizoni et al., "AT1 receptor blockade attenuates insulin resistance and myocardial remodeling in rats with diet-induced obesity," *PLoS ONE*, vol. 9, no. 1, Article ID e86447, 2014.
- [140] I. I. Joffe, K. E. Travers, C. L. Perreault-Micale et al., "Abnormal cardiac function in the streptozotocin-induced, non-insulin-dependent diabetic rat: noninvasive assessment with Doppler echocardiography and contribution of the nitric oxide pathway," *Journal of the American College of Cardiology*, vol. 34, no. 7, pp. 2111–2119, 1999.
- [141] F. S. Fein, B. Miller-Green, and E. H. Sonnenblick, "Altered myocardial mechanics in diabetic rabbits," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 248, no. 5, part 2, pp. H729–H736, 1985.
- [142] L. M. Semeniuk, A. J. Kryski, and D. L. Severson, "Echocardiographic assessment of cardiac function in diabetic db/db and transgenic db/db-hGLUT4 mice," *American Journal of Physiology: Heart and Circulatory Physiology*, vol. 283, no. 3, pp. H976–H982, 2002.
- [143] B. Rodrigues and J. H. McNeill, "Cardiac dysfunction in isolated perfused hearts from spontaneously diabetic BB rats," *Canadian Journal of Physiology and Pharmacology*, vol. 68, no. 4, pp. 514–518, 1990.
- [144] T. Pulinilkunnil, P. C. Kienesberger, J. Nagendran, N. Sharma, M. E. Young, and J. R. B. Dyck, "Cardiac-specific adipose triglyceride lipase overexpression protects from cardiac steatosis and dilated cardiomyopathy following diet-induced obesity," *International Journal of Obesity*, vol. 38, no. 2, pp. 205–215, 2014.
- [145] Z. Lu, Y.-P. Jiang, X.-H. Xu, L. M. Ballou, I. S. Cohen, and R. Z. Lin, "Decreased L-type Ca^{2+} current in cardiac myocytes of type 1 diabetic akita mice due to reduced phosphatidylinositol 3-kinase signaling," *Diabetes*, vol. 56, no. 11, pp. 2780–2789, 2007.
- [146] P. Pacher, L. Liaudet, F. G. Soriano, J. G. Mabley, É. Szabó, and C. Szabó, "The role of poly(ADP-ribose) polymerase activation in the development of myocardial and endothelial dysfunction in diabetes," *Diabetes*, vol. 51, no. 2, pp. 514–521, 2002.
- [147] T. L. Broderick and A. K. Hutchison, "Cardiac dysfunction in the euglycemic diabetic-prone BB Wor rat," *Metabolism: Clinical and Experimental*, vol. 53, no. 11, pp. 1391–1394, 2004.
- [148] A. Van den Bergh, W. Flameng, and P. Herijgers, "Type II diabetic mice exhibit contractile dysfunction but maintain cardiac output by favourable loading conditions," *European Journal of Heart Failure*, vol. 8, no. 8, pp. 777–783, 2006.
- [149] Y.-C. Hsiao, K.-I. Suzuki, H. Abe, and T. Toyota, "Ultrastructural alterations in cardiac muscle of diabetic BB Wistar rats," *Virchows Archiv A*, vol. 411, no. 1, pp. 45–52, 1987.
- [150] L. Li, Y. Hua, M. Dong et al., "Short-term lenalidomide (Revlimid) administration ameliorates cardiomyocyte contractile dysfunction in ob/ob obese mice," *Obesity*, vol. 20, no. 11, pp. 2174–2185, 2012.
- [151] K. Saito, S. Nishi, T. Kashima, and H. Tanaka, "Histologic and ultrastructural studies on the myocardium in spontaneously diabetic KK mice: a new animal model of cardiomyopathy," *The American Journal of Cardiology*, vol. 53, no. 2, pp. 320–323, 1984.
- [152] S. Boudina, S. Sena, H. Theobald et al., "Mitochondrial energetics in the heart in obesity-related diabetes: direct evidence for increased uncoupled respiration and activation of uncoupling proteins," *Diabetes*, vol. 56, no. 10, pp. 2457–2466, 2007.
- [153] V. Akhileshwar, S. P. Patel, and S. S. Katyare, "Diabetic cardiomyopathy and reactive oxygen species (ROS) related parameters in male and female rats: a comparative study," *Indian Journal of Clinical Biochemistry*, vol. 22, no. 1, pp. 84–90, 2007.