Research Article

On Necessary Condition for the Variable Exponent Hardy Inequality

Aziz Harman

Faculty of Education, Dicle University, 21280 Diyarbakir, Turkey

Correspondence should be addressed to Aziz Harman, aharman@dicle.edu.tr

Received 2 January 2012; Revised 20 February 2012; Accepted 24 February 2012

Academic Editor: Carlo Sbordone

Copyright © 2012 Aziz Harman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We derive a necessary condition for exponent functions p, β such that the variable exponent Hardy inequality $\|x^{\beta(x)-1} \int_0^x f(t) dt\|_{L^{p(\cdot)}(0,l)} \leq C \|x^{\beta(x)} f\|_{L^{p(\cdot)}(0,l)}$ holds.

1. Introduction

A sufficient condition on measurable functions $p : (0, l) \rightarrow [1, \infty)$, $\beta : (0, l) \rightarrow (-\infty, \infty)$ for which the variable exponent Hardy inequality

$$\left\| |x|^{\beta(\cdot)-1} Hf \right\|_{L^{p(\cdot)}(0,l)} \leq C \left\| |x|^{\beta(\cdot)} f \right\|_{L^{p(\cdot)}(0,l)}, \quad Hf(x) = \int_0^x f(t) dt \quad (1.1)$$

holds for all $f \geq 0$ have been known (see [1–3]). According to mentioned works, if $\beta(0) < 1 - (1/p(0))$, $p^- := \inf\{p(x) : x \in (0, l)\} > 1$, then a sufficient condition is $p, \beta \in \Lambda$, where Λ means a class of measurable functions $g : (0, l) \rightarrow (-\infty, \infty)$ such that $\exists g(0), C > 0$

$$|g(x) - g(0)| \ln \frac{1}{|x|} \leq C, \quad 0 < x < \frac{1}{2}. \quad (1.2)$$

The purpose of this paper is to prove that a weaker continuity condition on $p(\cdot)$ and $\beta(\cdot)$ is necessary for the norm inequality to hold provided that $p(\cdot)$ and $\beta(\cdot)$ are monotone (see Theorems 2.1 and 2.2 for a precise statement), which is the following condition:

$$|g(2x) - g(x)| \ln \frac{1}{x} \leq C, \quad 0 < x < \frac{1}{2}. \quad (1.3)$$

Note that condition (1.3) is strictly weaker than (1.2). For example, it is satisfied by $p(x) = C/|\log(x)|^{1/2}$. This condition is new and somewhat surprising. Since in the corresponding theorem for the maximal operator, it is known that $p(\cdot)$ need not be continuous, and the problem of determining which exponent conditions are necessary and/or sufficient is an open one.

If the powers are not monotone, it follows from the results of the paper [2] that condition (1.2) is close to be sharp. Also in [2], the necessity of conditions $p^- > 1$ and $\beta(0) < 1 - 1/p(0)$ was proved. Recently, there have been quite a number of papers discussing the Hardy inequality in norms of the variable exponent Lebesgue spaces [3–11].

For problems of boundedness of classical integral operators in variable exponent Lebesgue spaces and regularity results for nonlinear equations with nonstandard growth condition, see monograph [12] and references therein.

2. Main Results

As to the basic properties of spaces $L^{p(\cdot)}$, we refer to [13]. Throughout this paper it is assumed that $p(x)$ is a measurable function in $(0, l)$ taking its values from the interval $[1, \infty)$ with $p^+ := \sup\{p(x) : x \in (0, l)\} < \infty$. The space of functions $L^{p(\cdot)}(0, l)$ is introduced as the class of measurable functions $f(x)$ in $(0, l)$, which have a finite $I_p(f) := \int_0^l |f(x)|^{p(x)} dx$ -modular. A norm in $L^{p(\cdot)}(0, l)$ is given in the form

$$\|f\|_{L^{p(\cdot)}(0, l)} = \inf \left\{ \lambda > 0 : I_p\left(\frac{f}{\lambda}\right) \leq 1 \right\}. \quad (2.1)$$

There exists a relation between modular and norm, which is expressed by the following inequalities:

$$\|f\|_{L^{p(\cdot)}(0, l)}^{p^+} \leq I_p(f) \leq \|f\|_{L^{p(\cdot)}(0, l)}^{p^-}, \quad 1 \geq \|f\|_{p(\cdot)}, \quad (2.2)$$

$$\|f\|_{L^{p(\cdot)}(0, l)}^{p^-} \leq I_p(f) \leq \|f\|_{L^{p(\cdot)}(0, l)}^{p^+}, \quad 1 \leq \|f\|_{p(\cdot)}. \quad (2.3)$$

Such estimates allow us to perform our estimates in terms of a modular. In the following two theorems, we show that if functions p, β are monotone, then condition (1.3) for them is necessary for inequality (1.1) to hold.

Theorem 2.1. *Let $\beta \in \mathbb{R}$ a function $p : (0, l) \rightarrow [1, \infty)$ be increasing on $(0, \varepsilon)$ and such that $p(0) = \lim_{x \rightarrow 0} p(x)$ exists, $\beta(0) < 1 - 1/p(0)$, $p^- > 1$. Then for inequality (1.1) to hold, it is necessary that for the function $p(\cdot)$ condition (1.3) is satisfied.*

Theorem 2.2. Let $p \in \mathbb{R}$, let $\beta : (0, l) \rightarrow [-\infty, \infty)$ be a function decreasing on $(0, \varepsilon)$ such that $\beta(0) = \lim_{x \rightarrow 0} \beta(x)$ exists, and let the conditions $\beta(0) < 1 - 1/p(0)$, $p^- > 1$ be satisfied. Then for inequality (1.1) to hold, it is necessary that for the function $\beta(\cdot)$ condition (1.3) is satisfied.

The following two theorems show that the logarithmic regularity conditions (1.2) for the functions p, β are essential for inequality (1.1) to hold.

Theorem 2.3. Let $\beta \in \mathbb{R}$, and $\delta_n = 4^{-n}, n \in \mathbb{N}$. There exist a sequence of functions $\{f_n\}$ and a function $p : (0, l) \rightarrow [1, \infty)$ satisfying the conditions $\beta < 1 - 1/p(0)$, $p^- > 1$ such that

$$\lim_{n \rightarrow \infty} |p(\delta_n) - p(0)| \ln \frac{1}{\delta_n} = \infty, \quad (2.4)$$

and inequality (1.1) is violated.

Theorem 2.4. Let $p > 1$, $\delta_n = 4^{-n}, n \in \mathbb{N}$. Then there exist a sequence of functions $\{f_n\}$ and a function $\beta : (0, l) \rightarrow (-\infty, \infty)$ satisfying the conditions $\beta(0) < 1 - 1/p$,

$$\lim_{n \rightarrow \infty} |\beta(\delta_n) - \beta(0)| \ln \frac{1}{\delta_n} = \infty, \quad (2.5)$$

such that inequality (1.1) is violated.

3. Proofs of Main Results

Proof of Theorem 2.1. Denote $I_{p(\cdot)}(f) = \int_0^l |f(t)|^{p(t)} dt$. By (2.2) note that the condition $I_p(f) \leq 1$ is equivalent to $\|f\|_{L^{p(\cdot)}(0, l)} \leq 1$.

Put $\delta_k = \varepsilon 4^{-k}, k \in \mathbb{N}$, and $f_k(x) = x^{-1/p(x) - \beta} \chi_{(\delta_k, 2\delta_k)}(x), x \in (0, l)$. Then for sufficiently large k ,

$$\begin{aligned} I_{p(\cdot)}(x^{\beta(x)} f_k(x)) &= \int_{\delta_k}^{2\delta_k} (t^\beta t^{-1/p(t) - \beta})^{p(t)} dt \\ &= \int_{\delta_k}^{2\delta_k} t^{-1} dt = \ln 2. \end{aligned} \quad (3.1)$$

Also

$$\begin{aligned} I_{p(\cdot)}(x^{\beta(x)-1} H(f_k(x))) &\geq \int_{3\delta_k}^{4\delta_k} \left(x^{(\beta-1)} \int_{\delta_k}^{2\delta_k} t^{-(1/p(t) - \beta)} dt \right)^{p(x)} dx \\ &\geq C \int_{3\delta_k}^{4\delta_k} \delta_k^{(1-1/p(2\delta_k) - \beta)} x^{(\beta-1)p(2\delta_k)} dx \\ &\geq C \delta_k^{1-p(3\delta_k)/p(2\delta_k)} = C e^{(1/p^+)[p(3\delta_k) - p(2\delta_k)] \ln(1/2\delta_k)}. \end{aligned} \quad (3.2)$$

Applying inequality (1.1), we have

$$|p(3\delta_k) - p(2\delta_k)| \ln \frac{1}{2\delta_k} \leq C, \quad k \in \mathbb{N}, \quad (3.3)$$

which by using of monotony of p and its boundedness implies (1.3).

This completes the proof of Theorem 2.1. $\square$

Proof of Theorem 2.2. Put $\delta_k = \varepsilon 4^{-k}$, $k \in \mathbb{N}$ and $f_k(x) = x^{-1/p-\beta(x)} \chi_{(\delta_k, 2\delta_k)}(x)$, $x \in (0, l)$. Then

$$\begin{aligned} I_{p(\cdot)}(x^{\beta(x)} f_k(x)) &= \int_{\delta_k}^{2\delta_k} \left(t^{\beta(t)} t^{-1/p-\beta(t)} \right)^{p(t)} dt \\ &= \int_{\delta_k}^{2\delta_k} t^{-1} dt = \ln 2. \end{aligned} \quad (3.4)$$

Also

$$\begin{aligned} I_{p(\cdot)}(x^{\beta(x)-1} H(f_k(x))) &\geq \int_{3\delta_k}^{4\delta_k} \left(x^{\beta(x)-1} \int_{\delta_k}^{2\delta_k} t^{-1/p-\beta(t)} dt \right)^{p(x)} dx \\ &\geq C \delta_k^{[\beta(3\delta_k)-\beta(2\delta_k)]p} \geq C e^{p[\beta(3\delta_k)-\beta(2\delta_k)] \ln(1/\delta_k)}. \end{aligned} \quad (3.5)$$

Applying inequality (1.1), we have

$$|\beta(2\delta_k) - \beta(3\delta_k)| \ln \frac{1}{\delta_k} \leq C, \quad k \in \mathbb{N} \quad (3.6)$$

which by using monotony of β implies (1.3).

This completes the proof of Theorem 2.2. $\square$

Proof of Theorem 2.3. Let us assume that $f_k(x) = x^{-1/p(x)-\beta} \chi_{(\delta_k, 2\delta_k)}(x)$, $x \in (0, l)$. Fix $k \in \mathbb{N}$. We define the step function

$$p(x) = \begin{cases} p_0 + \alpha_n & \text{if } x \in (2\delta_n, 4\delta_n), \\ p_0 & \text{if } x \in (\delta_n, 2\delta_n), \end{cases} \quad n \in \mathbb{N}. \quad (3.7)$$

Here $\{\alpha_n\}$ is a sequence of positive numbers that satisfies the condition

$$n\alpha_n \longrightarrow \infty \text{ as } n \longrightarrow \infty. \quad (3.8)$$

Then $\alpha_n \ln(1/\delta_n) \rightarrow \infty$ as $n \rightarrow \infty$; that is, condition (1.2) is not satisfied for the function $p(x)$. Also note that this function $p(\cdot)$ is not monotone. We have

$$\begin{aligned} I_{p(\cdot)}(x^\beta f_k(x)) &= \int_{\delta_k}^{2\delta_k} \left(t^\beta t^{-1/p(t)-\beta} \right)^{p(t)} dt \\ &= \int_{\delta_k}^{2\delta_k} t^{-1} dt = \ln 2, \end{aligned} \quad (3.9)$$

$$\begin{aligned} I_{p(\cdot)}(x^{\beta-1} H(f_k(x))) &\geq \int_{2\delta_k}^{4\delta_k} \left(\int_{\delta_k}^{2\delta_k} t^{-1/p(t)-\beta} dt \right)^{(p_0+\alpha_k)} x^{(\beta-1)(p_0+\alpha_k)} dx \\ &\geq C \int_{2\delta_k}^{4\delta_k} \delta_k^{(1-1/p_0-\beta)(p_0+\alpha_k)} x^{(\beta-1)(p_0+\alpha_k)} dx \\ &\geq C \delta_k^{-\alpha_k/p_0} = C e^{\alpha_k/p_0 \ln(1/\delta_k)} \rightarrow \infty \text{ as } k \rightarrow \infty. \end{aligned} \quad (3.10)$$

The last relation shows violating of inequality (1.1) for sufficiently large k . $\square$

Proof of Theorem 2.4. Let us assume that $f_k(x) = x^{-1/p-\beta(x)} \chi_{(\delta_n, 2\delta_n)}(x)$, $x \in (0, l)$, $n \in \mathbb{N}$. Fix $n \in \mathbb{N}$. We define the step function β as

$$\beta(x) = \begin{cases} \beta_0 + \alpha_n & \text{if } x \in (\delta_n, 2\delta_n), \\ \beta_0 & \text{if } x \in (2\delta_n, 4\delta_n), \end{cases} \quad n \in \mathbb{N}, \quad (3.11)$$

where $\alpha_n \ln(1/\delta_n) \rightarrow \infty$; that is, condition (1.2) is not satisfied for the function β . Note that this function β is not monotone.

We have

$$I_{p(\cdot)}(x^{\beta(x)-1} f_n(x)) \geq C \delta_n^{-p\alpha_n} \rightarrow \infty \text{ as } n \rightarrow \infty, \quad (3.12)$$

$$I_{p(\cdot)}(x^{\beta(x)} f_n(x)) \leq C_0 \ln 2. \quad (3.13)$$

The last relation contradicts to inequality (1.1) for sufficiently large k . $\square$

Acknowledgment

The author thanks Professor Mamedov F.I. for his valuable discussion of the subject. The author also expresses his acknowledgement to the referee for drawing attention to the log example and comments of stylistic character.

References

- [1] L. Diening and S. Samko, "Hardy inequality in variable exponent Lebesgue spaces," *Fractional Calculus & Applied Analysis*, vol. 10, no. 1, pp. 1–17, 2007.
- [2] A. Harman and F. I. Mamedov, "On boundedness of weighted Hardy operator in $L^{p(\cdot)}$ and regularity condition," *Journal of Inequalities and Applications*, vol. 2010, Article ID 837951, 14 pages, 2010.

- [3] R. A. Mashiyev, B. Çekiç, F. I. Mamedov, and S. Ogras, "Hardy's inequality in power-type weighted $L^{p(\cdot)}$ spaces," *Journal of Mathematical Analysis and Applications*, vol. 334, no. 1, pp. 289–298, 2007.
- [4] D. Cruz-Uribe and F. I. Mamedov, "On a general weighted Hardy type inequality in the variable exponent Lebesgue spaces," *Revista Matemática Complutense*. In press.
- [5] D. E. Edmunds, V. Kokilashvili, and A. Meskhi, "On the boundedness and compactness of the weighted Hardy operators in spaces," *Georgian Mathematical Journal*, vol. 12, no. 1, pp. 27–44, 2005.
- [6] P. Harjulehto, P. Hästö, and M. Koskenoja, "Hardy's inequality in a variable exponent Sobolev space," *Georgian Mathematical Journal*, vol. 12, no. 3, pp. 431–442, 2005.
- [7] F. I. Mamedov and A. Harman, "On a weighted inequality of Hardy type in spaces $L^{p(\cdot)}$," *Journal of Mathematical Analysis and Applications*, vol. 353, no. 2, pp. 521–530, 2009.
- [8] F. I. Mamedov and A. Harman, "On a Hardy type general weighted inequality in spaces $L^{p(\cdot)}$," *Integral Equations and Operator Theory*, vol. 66, no. 4, pp. 565–592, 2010.
- [9] F. I. Mamedov, "On Hardy type inequality in variable exponent Lebesgue space $L^{p(\cdot)}(0, l)$," *Azerbaijan Journal of Mathematics*, vol. 2, no. 1, pp. 90–99, 2012.
- [10] H. Rafeiro and S. Samko, "Hardy type inequality in variable Lebesgue spaces," *Annales Academiæ Scientiarum Fennicæ*, vol. 34, no. 1, pp. 279–289, 2009.
- [11] S. Samko, "Hardy inequality in the generalized Lebesgue spaces," *Fractional Calculus & Applied Analysis*, vol. 6, no. 4, pp. 355–362, 2003.
- [12] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička, *Lebesgue and Sobolev Spaces with Variable Exponents*, vol. 2017 of *Lecture Notes in Mathematics*, Springer, Heidelberg, Germany, 2011.
- [13] X. Fan and D. Zhao, "On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$," *Journal of Mathematical Analysis and Applications*, vol. 263, no. 2, pp. 424–446, 2001.