Research Article

Damped Algorithms for the Split Fixed Point and Equilibrium Problems

Li-Jun Zhu^{1,2} and Minglun Ren¹

¹ School of Management, Hefei University of Technology, Hefei 230009, China

² School of Mathematics and Information Science, Beifang University of Nationalities, Yinchuan 750021, China

Correspondence should be addressed to Minglun Ren; hfutren@sina.com.cn

Received 23 June 2013; Accepted 5 August 2013

Academic Editor: J. Liang

Copyright © 2013 L.-J. Zhu and M. Ren. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The main purpose of this paper is to study the split fixed point and equilibrium problems which includes fixed point problems, equilibrium problems, and variational inequality problems as special cases. A damped algorithm is presented for solving this split common problem. Strong convergence analysis is shown.

1. Introduction

Very recently, the split problems (e.g., the split feasibility problem, the split common fixed points problem, and the split variational inequality problem) have been studied extensively, see, for instance, [1–19]. Now we recall the related history. Let H_1 and H_2 be two Hilbert spaces and $C \subset H_1$ and $Q \subset H_2$ two nonempty closed convex subsets. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator. The split feasibility problem is to solve the inclusion:

$$x \in C \cap A^{-1}(Q) \quad (1)$$

which arise in the field of intensity-modulated radiation therapy and was presented in [1]. The iteration $\rho^{n+1} = \text{proj}_C(\rho^n - \varsigma A^*(I - P_Q)A\rho^n)$ is popular with $\varsigma \in (0, 2/\|A\|^2)$. Further, Xu [3] suggested a single step regularized method. Dang and Gao [4] developed a damped projection algorithm. If C and Q are the fixed point sets of mappings U and T , respectively, then (1) becomes a special case of the split common fixed point problem:

$$\text{Find } x \in \text{Fix}(U) \cap A^{-1}(\text{Fix}(T)). \quad (2)$$

Censor and Segal [5] invented a scheme below to solve (2):

$$\rho^{n+1} = U(\rho^n - \varsigma A^*(I - T)A\rho^n), \quad n \in \mathbb{N}. \quad (3)$$

Cui et al., [6] extended the damped projection algorithm to the split common fixed point problems. Let $\psi : C \times C \rightarrow \mathbb{R}$ be a bifunction. The equilibrium problem is to find $x^\dagger \in C$ such that

$$\psi(x^\dagger, x) \geq 0, \quad \forall x \in C. \quad (4)$$

We will indicate with $\text{EP}(\psi)$ the set of solutions of (4).

In the present paper, our main purpose is to study the following split fixed point and equilibrium problem.

$$\begin{aligned} &\text{Find a point } u^\S \in \text{Fix}(W) \cap \text{EP}(\psi) \\ &\text{such that } Au^\S \in \text{Fix}(S) \cap \text{EP}(\varphi), \end{aligned} \quad (5)$$

where $\text{Fix}(S)$ and $\text{Fix}(W)$ are the sets of fixed points of two nonlinear mappings S and W , respectively; $\text{EP}(\psi)$ and $\text{EP}(\varphi)$ are the solution sets of two equilibrium problems with bifunctions ψ and φ , respectively, and A is a bounded linear mapping. Denote the solution set of (5) by

$$\Theta = \{x \in \text{Fix}(W) \cap \text{EP}(\psi) : Ax \in \text{Fix}(S) \cap \text{EP}(\varphi)\}. \quad (6)$$

We develop a damped algorithm to solve this split fixed point and equilibrium problem. Strong convergence of the suggested damped algorithm is demonstrated.

2. Concepts and Lemmas

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, respectively. Let C be a nonempty closed convex subset of H . A mapping $W : C \rightarrow C$ is called nonexpansive if

$$\|Wx^\dagger - Wx^\ddagger\| \leq \|x^\dagger - x^\ddagger\|, \quad (7)$$

for all $x^\dagger, x^\ddagger \in C$. We call $\text{proj}_C : H \rightarrow C$ the metric projection if for each $x^b \in H$

$$\|x^b - \text{proj}_C(x^b)\| = \inf \{\|x^b - x^\dagger\| : x^\dagger \in C\}. \quad (8)$$

It is well known that the metric projection $\text{proj}_C : H \rightarrow C$ is firmly nonexpansive, that is,

$$\begin{aligned} & \|\text{proj}_C(x^\dagger) - \text{proj}_C(x^\ddagger)\|^2 \\ & \leq \langle x^\dagger - x^\ddagger, \text{proj}_C(x^\dagger) - \text{proj}_C(x^\ddagger) \rangle \end{aligned} \quad (9)$$

for all $x^\dagger, x^\ddagger \in H$. Hence proj_C is also nonexpansive.

Lemma 1 (see [20]). *Let C be a nonempty closed convex subset of a real Hilbert space H . Let $\psi : C \times C \rightarrow \mathbb{R}$ be a bifunction which satisfies the following conditions:*

- (H1) $\psi(x^\ddagger, x^\ddagger) = 0$ for all $x^\ddagger \in C$;
- (H2) ψ is monotone, that is, $\psi(x^\ddagger, x^\dagger) + \psi(x^\dagger, x^\ddagger) \leq 0$ for all $x^\ddagger, x^\dagger \in C$;
- (H3) for each $x^\dagger, x^\ddagger, x^\natural \in C$, $\lim_{t \downarrow 0} \psi(tx^\natural + (1-t)x^\dagger, x^\ddagger) \leq \psi(x^\dagger, x^\ddagger)$;
- (H4) for each $x^\dagger \in C$, $x^\ddagger \mapsto \psi(x^\dagger, x^\ddagger)$ is convex and lower semicontinuous.

Let $\omega > 0$ and $x^\dagger \in C$. Then, there exists $x^\natural \in C$ such that

$$\psi(x^\natural, x^\ddagger) + \frac{1}{\omega} \langle x^\ddagger - x^\natural, x^\natural - x^\dagger \rangle \geq 0, \quad \forall x^\ddagger \in C. \quad (10)$$

Further, if $U_\omega^\psi(x^\dagger) = \{x^\natural \in C : \psi(x^\natural, x^\ddagger) + (1/\omega)\langle x^\ddagger - x^\natural, x^\natural - x^\dagger \rangle \geq 0, \text{ for all } x^\ddagger \in C\}$, then the following hold:

- (i) U_ω^ψ is single-valued and U_ω^ψ is firmly nonexpansive, that is, for any $x^\dagger, x^\ddagger \in H$, $\|U_\omega^\psi x^\dagger - U_\omega^\psi x^\ddagger\|^2 \leq \langle U_\omega^\psi x^\dagger - U_\omega^\psi x^\ddagger, x^\dagger - x^\ddagger \rangle$;
- (ii) $\text{EP}(\psi)$ is closed and convex and $\text{EP}(\psi) = \text{Fix}(U_\omega^\psi)$.

Lemma 2 (see [21]). *Let H be a Hilbert space and $C \subset H$ a closed convex subset. Let $W : C \rightarrow C$ be a nonexpansive mapping. Then, the mapping $I - W$ is demiclosed. That is, if $\{\rho^n\}$ is a sequence in C such that $\rho^n \rightarrow v$ weakly and $(I - W)\rho^n \rightarrow u$ strongly, then $(I - W)v = u$.*

Lemma 3 (see [22]). *Assume that $\{\eta_n\}$ is a sequence of nonnegative real numbers such that*

$$\eta_{n+1} \leq (1 - \kappa_n)\eta_n + \varsigma_n, \quad n \in \mathbb{N}, \quad (11)$$

where $\{\kappa_n\}$ is a sequence in $(0, 1)$ and $\{\varsigma_n\}$ is a sequence such that

- (1) $\sum_{n=1}^\infty \kappa_n = \infty$;
- (2) $\limsup_{n \rightarrow \infty} (\varsigma_n / \kappa_n) \leq 0$ or $\sum_{n=1}^\infty |\varsigma_n| < \infty$.

Then $\lim_{n \rightarrow \infty} \eta_n = 0$.

3. Main Results

Let H_1 and H_2 be two Hilbert spaces and $C \subset H_1$ and $Q \subset H_2$ two nonempty closed convex subsets. Let $A : H_1 \rightarrow H_2$ be a bounded linear operator with its adjoint A^* . Let $\psi : C \times C \rightarrow \mathbb{R}$ and let $\varphi : D \times D \rightarrow \mathbb{R}$ be two bifunctions satisfying the conditions (H1)–(H4) in Lemma 1. Let $S : D \rightarrow D$ and $W : C \rightarrow C$ be two nonexpansive mappings.

Algorithm 4. Let $x_0 \in H_1$. Define a sequence $\{x_n\}$ as follows:

$$\begin{aligned} \rho^{n+1} &= WU_i^\psi[(1 - \zeta_n) \\ &\quad \times (\rho^n + \varsigma A^*(SU_\kappa^\varphi - I)A\rho^n)], \quad \forall n \in \mathbb{N}, \end{aligned} \quad (12)$$

where ι, κ , and ς are three constants satisfying $\iota \in (0, \infty)$, $\kappa \in (0, \infty)$, $\varsigma \in (0, 1/\|A\|^2)$, and $\{\zeta_n\}$ is a real number sequence in $(0, 1)$.

In the sequel, we assume that

$$\Theta = \{x \in \text{Fix}(W) \cap \text{EP}(\psi) : Ax \in \text{Fix}(S) \cap \text{EP}(\varphi)\} \neq \emptyset. \quad (13)$$

Theorem 5. *If $\{\zeta_n\}$ satisfies $\lim_{n \rightarrow \infty} \zeta_n = 0$, $\sum_{n=1}^\infty \zeta_n = \infty$ and $\lim_{n \rightarrow \infty} \zeta_{n+1}/\zeta_n = 1$, then $\{\rho^n\}$ generated by algorithm (12) converges strongly to $\text{proj}_\Theta(0)$ which is the minimum-norm element in Θ .*

Proof. Let $p = \text{proj}_\Theta(0)$. Then, $p \in \text{Fix}(W) \cap \text{EP}(\psi)$ and $Ap \in \text{Fix}(S) \cap \text{EP}(\varphi)$. Set $z^n = U_\kappa^\varphi A\rho^n$, $y^n = (1 - \zeta_n)(\rho^n + \varsigma A^*(SU_\kappa^\varphi - I)A\rho^n)$ and $u^n = U_i^\psi[(1 - \zeta_n)(\rho^n + \varsigma A^*(SU_\kappa^\varphi - I)A\rho^n)]$ for all $n \in \mathbb{N}$. Then $u^n = U_i^\psi y^n$. From Lemma 1, we know that U_i^ψ and U_κ^φ are firmly nonexpansive. Thus, we have

$$\|z^n - Ap\| = \|U_\kappa^\varphi A\rho^n - Ap\| \leq \|A\rho^n - Ap\|, \quad (14)$$

$$\|u^n - p\| = \|U_i^\psi y^n - p\| \leq \|y^n - p\|, \quad (15)$$

$$\begin{aligned} \|SU_\kappa^\varphi A\rho^n - Ap\|^2 &= \|SU_\kappa^\varphi A\rho^n - SU_\kappa^\varphi Ap\|^2 \\ &\leq \|U_\kappa^\varphi A\rho^n - U_\kappa^\varphi Ap\|^2 \\ &\leq \|A\rho^n - Ap\|^2 - \|U_\kappa^\varphi A\rho^n - A\rho^n\|^2. \end{aligned} \quad (16)$$

Note that

$$\begin{aligned}\|u^{n+1} - u^n\| &= \|U_i^\psi y^{n+1} - U_i^\psi y^n\| \\ &\leq \|y^{n+1} - y^n\|,\end{aligned}\quad (17)$$

$$\begin{aligned}\|z^{n+1} - z^n\| &= \|U_\kappa^\phi A\rho^{n+1} - U_\kappa^\phi A\rho^n\| \\ &\leq \|A\rho^{n+1} - A\rho^n\|.\end{aligned}\quad (18)$$

From (12) and (15), we have

$$\|\rho^{n+1} - p\| = \|Wu^n - p\| \leq \|u^n - p\| \leq \|y^n - p\|. \quad (19)$$

Observe that

$$\begin{aligned}\|y^n - p\|^2 &= \|(1 - \zeta_n) \\ &\quad \times (\rho^n - p + \varsigma A^* (Sz^n - A\rho^n)) - \zeta_n p\|^2 \\ &\leq (1 - \zeta_n) \|(\rho^n - p + \varsigma A^* (Sz^n - A\rho^n))\|^2 \\ &\quad + \zeta_n \|p\|^2 \\ &= (1 - \zeta_n) \left[\|\rho^n - p\|^2 + 2\varsigma \right. \\ &\quad \times \langle \rho^n - p, A^* (Sz^n - A\rho^n) \rangle \\ &\quad \left. + \varsigma^2 \|A^* (Sz^n - A\rho^n)\|^2 \right] \\ &\quad + \zeta_n \|p\|^2.\end{aligned}\quad (20)$$

Since A^* is the adjoint of A , we have

$$\begin{aligned}\langle \rho^n - p, A^* (Sz^n - A\rho^n) \rangle &= \langle A(\rho^n - p), Sz^n - A\rho^n \rangle \\ &= \langle A\rho^n - Ap + Sz^n - A\rho^n \\ &\quad - (Sz^n - A\rho^n), Sz^n - A\rho^n \rangle \\ &= \langle Sz^n - Ap, Sz^n - A\rho^n \rangle - \|Sz^n - A\rho^n\|^2.\end{aligned}\quad (21)$$

Using parallelogram law, we obtain

$$\begin{aligned}\langle Sz^n - Ap, Sz^n - A\rho^n \rangle &= \frac{1}{2} (\|Sz^n - Ap\|^2 + \|Sz^n - A\rho^n\|^2 \\ &\quad - \|A\rho^n - Ap\|^2).\end{aligned}\quad (22)$$

From (16), (21) and (22), we have

$$\begin{aligned}\langle \rho^n - p, A^* (Sz^n - A\rho^n) \rangle &= \frac{1}{2} (\|Sz^n - Ap\|^2 + \|Sz^n - A\rho^n\|^2 \\ &\quad - \|A\rho^n - Ap\|^2) \\ &\quad - \|Sz^n - A\rho^n\|^2 \\ &\leq \frac{1}{2} (\|A\rho^n - Ap\|^2 - \|z^n - A\rho^n\|^2 \\ &\quad + \|Sz^n - A\rho^n\|^2 - \|A\rho^n - Ap\|^2) \\ &\quad - \|Sz^n - A\rho^n\|^2 \\ &= -\frac{1}{2} \|z^n - A\rho^n\|^2 \\ &\quad - \frac{1}{2} \|Sz^n - A\rho^n\|^2.\end{aligned}\quad (23)$$

By (20) and (23), we deduce

$$\begin{aligned}\|y^n - p\|^2 &\leq (1 - \zeta_n) \left[\|\rho^n - p\|^2 + \varsigma^2 \|A\|^2 \|Sz^n - A\rho^n\|^2 \right. \\ &\quad \left. + 2\varsigma \left(-\frac{1}{2} \|z^n - A\rho^n\|^2 \right. \right. \\ &\quad \left. \left. - \frac{1}{2} \|Sz^n - A\rho^n\|^2 \right) \right] + \zeta_n \|p\|^2 \\ &= (1 - \zeta_n) \\ &\quad \times \left[\|\rho^n - p\|^2 + (\varsigma^2 \|A\|^2 - \varsigma) \|Sz^n - A\rho^n\|^2 \right. \\ &\quad \left. - \varsigma \|z^n - A\rho^n\|^2 \right] + \zeta_n \|p\|^2 \\ &\leq (1 - \zeta_n) \|\rho^n - p\|^2 + \zeta_n \|p\|^2.\end{aligned}\quad (24)$$

It follows from (19), we get

$$\begin{aligned}\|\rho^{n+1} - p\|^2 &\leq \|y^n - p\|^2 \\ &\leq (1 - \zeta_n) \|\rho^n - p\|^2 + \zeta_n \|p\|^2 \\ &\leq \max \{ \|\rho^n - p\|^2, \|p\|^2 \}.\end{aligned}\quad (25)$$

The boundedness of the sequence $\{\rho^n\}$ yields.

Set $v^n = \rho^n + \varsigma A^*(SU_\kappa^\varphi - I)A\rho^n$. Then, we have

$$\begin{aligned}
\|v^{n+1} - v^n\|^2 &= \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + \varsigma \left[A^* (Sz^{n+1} - A\rho^{n+1}) - A^* (Sz^n - A\rho^n) \right]^2 \\
&= \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + 2\varsigma \langle \rho^{n+1} - \rho^n, \\
&\quad \quad A^* [(Sz^{n+1} - A\rho^{n+1}) - (Sz^n - A\rho^n)] \rangle \\
&\quad + \varsigma^2 \|A^* (Sz^{n+1} - A\rho^{n+1}) - A^* (Sz^n - A\rho^n)\|^2 \\
&\leq \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + 2\varsigma \langle A\rho^{n+1} - A\rho^n, \\
&\quad \quad Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n) \rangle \\
&\quad + \varsigma^2 \|A\|^2 \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&= \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + \varsigma^2 \|A\|^2 \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&\quad + 2\varsigma \langle Sz^{n+1} - Sz^n, \\
&\quad \quad Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n) \rangle \\
&\quad - 2\varsigma \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&= \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + \varsigma^2 \|A\|^2 \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&\quad + \varsigma \left(\|Sz^{n+1} - Sz^n\|^2 \right. \\
&\quad \quad + \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&\quad \quad \left. - \|A\rho^{n+1} - A\rho^n\|^2 \right) \\
&\quad - 2\varsigma \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&= \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + (\varsigma^2 \|A\|^2 - \varsigma) \\
&\quad \quad \times \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&\quad + \varsigma \left(\|Sz^{n+1} - Sz^n\|^2 - \|(A\rho^{n+1} - A\rho^n)\|^2 \right)
\end{aligned}$$

$$\begin{aligned}
&\leq \|\rho^{n+1} - \rho^n\|^2 \\
&\quad + (\varsigma^2 \|A\|^2 - \varsigma) \\
&\quad \quad \times \|Sz^{n+1} - Sz^n - (A\rho^{n+1} - A\rho^n)\|^2 \\
&\quad + \varsigma \left(\|Sz^{n+1} - Sz^n\|^2 - \|(A\rho^{n+1} - A\rho^n)\|^2 \right).
\end{aligned} \tag{26}$$

Since $\varsigma \in (0, 1/\|A\|^2)$, we derive by virtue of (18) and (26) that

$$\|v^{n+1} - v^n\| \leq \|\rho^{n+1} - \rho^n\|. \tag{27}$$

According to (17) and (27), we have

$$\begin{aligned}
\|\rho^{n+1} - \rho^n\| &= \|Wu^{n+1} - Wu^n\| \\
&\leq \|u^{n+1} - u^n\| \\
&\leq \|y^{n+1} - y^n\| \\
&= \|(1 - \zeta_{n+1})v^{n+1} - (1 - \zeta_n)v^n\| \\
&= \|(1 - \zeta_{n+1})(v^{n+1} - v^n) + (\zeta_n - \zeta_{n+1})v^n\| \\
&\leq (1 - \zeta_{n+1})\|v^{n+1} - v^n\| + |\zeta_{n+1} - \zeta_n|\|v^n\| \\
&\leq (1 - \zeta_{n+1})\|\rho^{n+1} - \rho^n\| + |\zeta_{n+1} - \zeta_n|\|v^n\|.
\end{aligned} \tag{28}$$

It follows that

$$\|\rho^{n+1} - \rho^n\| \leq \frac{|\zeta_{n+1} - \zeta_n|}{\zeta_{n+1}} \|v^n\|. \tag{29}$$

Since $\{\rho^n\}$ is bounded, we can deduce $\{v^n\}$ is also bounded. From (29), we have

$$\lim_{n \rightarrow \infty} \|\rho^{n+1} - \rho^n\| = 0. \tag{30}$$

Hence,

$$\lim_{n \rightarrow \infty} \|\rho^n - Wu^n\| = 0. \tag{31}$$

Using the firmly-nonexpansiveness of U_i^ψ , we have

$$\begin{aligned}
\|u^n - p\|^2 &= \|U_i^\psi y^n - p\|^2 \\
&\leq \|y^n - p\|^2 - \|U_i^\psi y^n - y^n\|^2 \\
&= \|y^n - p\|^2 - \|u^n - y^n\|^2.
\end{aligned} \tag{32}$$

Thus, we get

$$\begin{aligned}
\|\rho^{n+1} - p\|^2 &\leq \|u^n - p\|^2 \\
&\leq \|y^n - p\|^2 - \|u^n - y^n\|^2 \\
&\leq (1 - \zeta_n)\|\rho^n - p\|^2 + \zeta_n\|p\|^2 - \|u^n - y^n\|^2.
\end{aligned} \tag{33}$$

It follows that

$$\begin{aligned}
\|u^n - y^n\|^2 &\leq \|\rho^n - p\|^2 - \|\rho^{n+1} - p\|^2 + \zeta_n\|p\|^2 \\
&\leq (\|\rho^n - p\| + \|\rho^{n+1} - p\|) \|\rho^{n+1} - p\| + \zeta_n\|p\|^2.
\end{aligned} \tag{34}$$

This together with (30) and (C1) implies that

$$\lim_{n \rightarrow \infty} \|u^n - y^n\| = 0. \quad (35)$$

Note that

$$\begin{aligned} \|\rho^{n+1} - p\|^2 &\leq \|y^n - p\|^2 \\ &\leq (1 - \zeta_n) \|\rho^n - p\|^2 \\ &\quad + (1 - \zeta_n) (\zeta^2 \|A\|^2 - \varsigma) \|Sz^n - A\rho^n\|^2 \\ &\quad - (1 - \zeta_n) \varsigma \|z^n - A\rho^n\|^2 + \zeta_n \|p\|^2. \end{aligned} \quad (36)$$

Hence,

$$\begin{aligned} (1 - \zeta_n) (\varsigma - \zeta^2 \|A\|^2) \|Sz^n - A\rho^n\|^2 \\ + (1 - \zeta_n) \varsigma \|z^n - A\rho^n\|^2 \\ \leq \|\rho^n - p\|^2 - \|\rho^{n+1} - p\|^2 + \zeta_n \|p\|^2 \\ \leq (\|\rho^n - p\| + \|\rho^{n+1} - p\|) \|\rho^{n+1} - \rho^n\| \\ + \zeta_n \|p\|^2, \end{aligned} \quad (37)$$

which implies that

$$\lim_{n \rightarrow \infty} \|Sz^n - A\rho^n\| = \lim_{n \rightarrow \infty} \|z^n - A\rho^n\| = 0. \quad (38)$$

So, we get

$$\lim_{n \rightarrow \infty} \|Sz^n - z^n\| = 0. \quad (39)$$

Since

$$\begin{aligned} \|y^n - p^n\| &= \|\varsigma A^* (SU_\kappa^\varphi - I) A\rho^n + \zeta_n v^n\| \\ &\leq \varsigma \|A\| \|Sz^n - A\rho^n\| + \zeta_n \|v^n\|, \end{aligned} \quad (40)$$

we get

$$\lim_{n \rightarrow \infty} \|\rho^n - y^n\| = 0. \quad (41)$$

From (31), (35), and (41), we get

$$\lim_{n \rightarrow \infty} \|\rho^n - W\rho^n\| = 0. \quad (42)$$

Now, we show that

$$\limsup_{n \rightarrow \infty} \langle p, y^n - p \rangle \geq 0. \quad (43)$$

Choose a subsequence $\{y^{n_i}\}$ of $\{y^n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle p, y^n - p \rangle = \lim_{i \rightarrow \infty} \langle p, y^{n_i} - p \rangle. \quad (44)$$

Notice that $\{y^{n_i}\}$ is bounded, we can choose $\{y^{n_{i_j}}\}$ of $\{y^{n_i}\}$ such that $y^{n_{i_j}} \rightharpoonup z$. Without loss of generality, we assume that $y^{n_i} \rightharpoonup z$. From the above conclusions, we derive that

$$\begin{aligned} \rho^{n_i} &\rightharpoonup z, & u_{n_i} &\rightharpoonup z, \\ A\rho^{n_i} &\rightharpoonup Az, & z^{n_i} &\rightharpoonup Az. \end{aligned} \quad (45)$$

By Lemma 2, (39), and (41), we deduce $z \in \text{Fix}(W)$ and $Az \in \text{Fix}(S)$.

Next, we show that $z \in \text{EP}(\psi)$. Since $u^n = U_t^\psi y^n$, we have

$$\psi(u^n, x^\dagger) + \frac{1}{t} \langle x^\dagger - u^n, u^n - y^n \rangle \geq 0, \quad \forall x^\dagger \in C. \quad (46)$$

By the monotonicity of ψ , we have

$$\frac{1}{t} \langle x^\dagger - u^n, u^n - y^n \rangle \geq \psi(x^\dagger, u^n), \quad (47)$$

and so

$$\left\langle x^\dagger - u_{n_i}, \frac{u_{n_i} - y^{n_i}}{t} \right\rangle \geq \psi(x^\dagger, u_{n_i}). \quad (48)$$

Since $\|u^n - y^n\| \rightarrow 0$, $u_{n_i} \rightarrow z$, we obtain $(u_{n_i} - y^{n_i})/t \rightarrow 0$. Thus, $0 \geq \psi(x^\dagger, z)$. For t with $0 < t \leq 1$ and $x^\dagger \in C$, let $y^t = tx^\dagger + (1-t)z \in C$. We obtain $\psi(y^t, z) \leq 0$. Hence,

$$0 = \psi(y^t, y^t) \leq t\psi(y^t, x^\dagger) + (1-t)\psi(y^t, z) \leq t\psi(y^t, x^\dagger). \quad (49)$$

So, $0 \leq \psi(y^t, x^\dagger)$. And, thus, $0 \leq \psi(z, x^\dagger)$. This implies that $z \in \text{EP}(\psi)$. Similarity, we can prove that $Az \in \text{EP}(\varphi)$. To this end, we deduce $z \in \text{Fix}(W) \cap \text{EP}(\psi)$ and $Az \in \text{Fix}(S) \cap \text{EP}(\varphi)$. That is to say, $z \in \Theta$. Therefore,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle p, y^n - p \rangle &= \lim_{i \rightarrow \infty} \langle p, y^{n_i} - p \rangle \\ &= \lim_{i \rightarrow \infty} \langle p, z - p \rangle \\ &\geq 0. \end{aligned} \quad (50)$$

Finally, we prove $\rho^n \rightarrow p$. From (12), we have

$$\begin{aligned} \|\rho^{n+1} - p\|^2 &\leq \|y^n - p\|^2 \\ &= \|(1 - \zeta_n)(v^n - p) - \zeta_n p\|^2 \\ &\leq (1 - \zeta_n) \|v^n - p\|^2 - 2\zeta_n \langle p, y^n - p \rangle \\ &\leq (1 - \zeta_n) \|\rho^n - p\|^2 - 2\zeta_n \langle p, y^n - p \rangle. \end{aligned} \quad (51)$$

Applying Lemma 3 and (50) to (51), we deduce $\rho^n \rightarrow p$. The proof is completed. $\square$

Algorithm 6. Let $\rho^0 \in H_1$ arbitrarily define a sequence $\{\rho^n\}$ by the following:

$$\rho^{n+1} = W((1 - \zeta_n)(\rho^n + \varsigma A^*(S - I)A\rho^n)), \quad (52)$$

for all $n \in \mathbb{N}$, where $\varsigma \in (0, 1/\|A\|^2)$ and $\{\zeta_n\}$ is a real number sequence in $(0, 1)$.

Corollary 7. Suppose $\Theta_1 = \{x \in \text{Fix}(W) : Ax \in \text{Fix}(S)\} \neq \emptyset$. If $\{\zeta_n\}$ satisfies $\lim_{n \rightarrow \infty} \zeta_n = 0$, $\sum_{n=1}^\infty \zeta_n = \infty$, and $\lim_{n \rightarrow \infty} \zeta_{n+1}/\zeta_n = 1$, then the sequence $\{\rho^n\}$ generated by algorithm (52) converges strongly to $p = \text{proj}_{\Theta_1}(0)$ which is the mum-norm element in Θ_1 .

Algorithm 8. Let $\rho^0 \in H_1$ arbitrarily define a sequence $\{\rho^n\}$ by the following:

$$\rho^{n+1} = U_t^\psi((1 - \zeta_n)(\rho^n + \varsigma A^*(U_\kappa^\varphi - I)A\rho^n)), \quad (53)$$

for all $n \in \mathbb{N}$, where ι, κ , and ς are three constants satisfying $\iota \in (0, \infty)$, $\kappa \in (0, \infty)$, $\varsigma \in (0, 1/\|A\|^2)$, and $\{\zeta_n\}$ is a real number sequence in $(0, 1)$.

Corollary 9. Suppose $\Theta_2 = \{x \in \text{EP}(\psi) : Ax \in \text{EP}(\varphi)\} \neq \emptyset$. If $\{\zeta_n\}$ satisfies $\lim_{n \rightarrow \infty} \zeta_n = 0$, $\sum_{n=1}^{\infty} \zeta_n = \infty$, and $\lim_{n \rightarrow \infty} \zeta_{n+1}/\zeta_n = 1$, then the sequence $\{\rho^n\}$ generated by algorithm (53) converges strongly to $p = \text{proj}_{\Theta_2}(0)$ which is the mum-norm element in Θ_2 .

Algorithm 10. Let $\rho^0 \in H_1$ arbitrarily define a sequence $\{\rho^n\}$ by the following:

$$\rho^{n+1} = \text{proj}_C((1 - \zeta_n)(\rho^n + \varsigma A^*(\text{proj}_Q - I)A\rho^n)), \quad (54)$$

for all $n \in \mathbb{N}$, where $\varsigma \in (0, 1/\|A\|^2)$ and $\{\zeta_n\}$ is a real number sequence in $(0, 1)$.

Corollary 11. Suppose $\Theta_3 = \{x \in C : Ax \in Q\} \neq \emptyset$. If $\{\zeta_n\}$ satisfies $\lim_{n \rightarrow \infty} \zeta_n = 0$, $\sum_{n=1}^{\infty} \zeta_n = \infty$, and $\lim_{n \rightarrow \infty} \zeta_{n+1}/\zeta_n = 1$, then the sequence $\{\rho^n\}$ generated by algorithm (54) converges strongly to $p = \text{proj}_{\Theta_3}(0)$ which is the mum-norm element in Θ_3 .

Acknowledgment

Li-Jun Zhu was supported in part by NNSF of China (61362033), NZ13087, NGY2012097 and 2013xyz023.

References

- [1] Y. Censor and T. Elfving, "A multiprojection algorithm using Bregman projections in a product space," *Numerical Algorithms*, vol. 8, no. 2–4, pp. 221–239, 1994.
- [2] C. Byrne, "Iterative oblique projection onto convex sets and the split feasibility problem," *Inverse Problems*, vol. 18, no. 2, pp. 441–453, 2002.
- [3] H.-K. Xu, "Iterative methods for the split feasibility problem in infinite-dimensional Hilbert spaces," *Inverse Problems*, vol. 26, no. 10, Article ID 105018, 17 pages, 2010.
- [4] Y. Dang and Y. Gao, "The strong convergence of a KM-CQ-like algorithm for a split feasibility problem," *Inverse Problems*, vol. 27, no. 1, Article ID 015007, 9 pages, 2011.
- [5] Y. Censor and A. Segal, "The split common fixed point problem for directed operators," *Journal of Convex Analysis*, vol. 16, no. 2, pp. 587–600, 2009.
- [6] H. Cui, M. Su, and F. Wang, "Damped projection method for split common fixed point problems," *Journal of Inequalities and Applications*, vol. 2013, article 123, 2013.
- [7] L.-C. Ceng, Q. H. Ansari, and J.-C. Yao, "An extragradient method for solving split feasibility and fixed point problems," *Computers & Mathematics with Applications*, vol. 64, no. 4, pp. 633–642, 2012.
- [8] Q. Yang, "The relaxed CQ algorithm solving the split feasibility problem," *Inverse Problems*, vol. 20, no. 4, pp. 1261–1266, 2004.
- [9] Y. Yao, T. H. Kim, S. Chebbi, and H. K. Xu, "A modified extragradient method for the split feasibility and fixed point problems," *Journal of Nonlinear and Convex Analysis*, vol. 13, pp. 383–396, 2012.
- [10] J. Zhao and Q. Yang, "Several solution methods for the split feasibility problem," *Inverse Problems*, vol. 21, no. 5, pp. 1791–1799, 2005.
- [11] L.-C. Ceng, Q. H. Ansari, and J.-C. Yao, "Relaxed extragradient methods for finding minimum-norm solutions of the split feasibility problem," *Nonlinear Analysis. Theory, Methods & Applications A*, vol. 75, no. 4, pp. 2116–2125, 2012.
- [12] A. Moudafi, "A note on the split common fixed-point problem for quasi-nonexpansive operators," *Nonlinear Analysis. Theory, Methods & Applications A*, vol. 74, no. 12, pp. 4083–4087, 2011.
- [13] Z. H. He, "The split equilibrium problems and its convergence algorithms," *Journal of Inequality and Application*, vol. 2012, article 162, 2012.
- [14] A. Moudafi, "Split monotone variational inclusions," *Journal of Optimization Theory and Applications*, vol. 150, no. 2, pp. 275–283, 2011.
- [15] A. Moudafi, "The split common fixed-point problem for demi-contractive mappings," *Inverse Problems*, vol. 26, no. 5, pp. 587–600, 2010.
- [16] C. Byrne, Y. Censor, A. Gibali, and S. Reich, "The split common null point problem," *Journal of Nonlinear and Convex Analysis*, vol. 13, no. 4, pp. 759–775, 2012.
- [17] Y. Censor, A. Gibali, and S. Reich, "Algorithms for the split variational inequality problem," *Numerical Algorithms*, vol. 59, no. 2, pp. 301–323, 2012.
- [18] Z. H. He and W. S. Du, "On hybrid split problem and its nonlinear algorithms," *Fixed Point Theory and Applications*, vol. 2013, article 47, 2013.
- [19] S. S. Chang, L. Wang, Y. K. Tang, and L. Yang, "The split common fixed point problem for total asymptotically strictly pseudocontractive mappings," *Journal of Applied Mathematics*, vol. 2013, Article ID 385638, 13 pages, 2012.
- [20] P. L. Combettes and S. A. Hirstoaga, "Equilibrium programming in Hilbert spaces," *Journal of Nonlinear and Convex Analysis*, vol. 6, no. 1, pp. 117–136, 2005.
- [21] K. Geobel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, 1990.
- [22] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.