Research Article

Eigenvalues of a Class of Singular Boundary Value Problems of Impulsive Differential Equations in Banach Spaces

Daliang Zhao and Yansheng Liu

School of Mathematical Sciences, Shandong Normal University, Ji'nan, Shandong 250014, China

Correspondence should be addressed to Yansheng Liu; yanshliu@gmail.com

Received 26 May 2013; Accepted 28 October 2013; Published 23 February 2014

Academic Editor: Gestur Ólafsson

Copyright © 2014 D. Zhao and Y. Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is devoted to investigating the eigenvalue problems of a class of nonlinear impulsive singular boundary value problem in Banach spaces: $\mu x'' + f(t, x) = 0, t \in (0, 1), t \neq t_i; \Delta x|_{t=t_i} = \alpha_i x(t_i - 0), i = 1, 2, \dots, k; ax(0) - bx'(0) = \theta; cx(1) + dx'(1) = \theta$, where θ denotes the zero element of Banach space, $\Delta x|_{t=t_i} = x(t_i + 0) - x(t_i - 0), \alpha_i > -1, a, b, c, d \in R^+, \gamma = ac + ad + bc > 0, \mu$ is a parameter, and $f(t, x)$ may be singular at $t = 0, 1$ and $x = \theta$. The arguments are mainly based upon the theory of fixed point index, measure of noncompactness, and the special cone, which is constructed to overcome the singularity.

1. Introduction

Consider the following eigenvalue problems of singular boundary value problem (SBVP) with impulse in Banach space E :

$$\begin{aligned} \mu x'' + f(t, x) &= 0, \quad t \in (0, 1), \quad t \neq t_i; \\ \Delta x|_{t=t_i} &= \alpha_i x(t_i - 0), \quad i = 1, 2, \dots, k; \\ ax(0) - bx'(0) &= \theta; \\ cx(1) + dx'(1) &= \theta, \end{aligned} \quad (1)$$

where θ denotes the zero element of Banach space $E, \Delta x|_{t=t_i} = x(t_i + 0) - x(t_i - 0), 0 = t_0 < t_1 < t_2 < \dots < t_k < t_{k+1} = 1, \alpha_i > -1, a, b, c, d \in R^+, \gamma = ac + ad + bc > 0, \mu$ is a parameter, $f(t, x)$ may be singular at $t = 0, 1$ and $x = \theta$.

During the last few decades, the existence of positive solution of nonlinear singular boundary value problems has gained considerable popularity (see [1–12] and references therein). In recent years, there were also a lot of papers which dealt with eigenvalue problems (see [13–27]). Some of them considered singular case (see, for instance, [13–16, 18, 19], etc.).

On the other hand, as we know, the theory of impulsive differential equations has found its extensive applications in realistic mathematical modeling of a wide variety of practical situations such as physics, chemical technology, population dynamics, biotechnology, and economics. It has emerged as an important area of investigation in recent years (see [28–36] and references therein).

To the best of our knowledge, there is no paper studying the eigenvalue problems of the impulsive singular boundary value problem in Banach spaces. The main purpose of this paper is to fill this gap. By using the theory of fixed point index, measure of noncompactness, and the special cone which is constructed to overcome the singularity, we investigate the existence of eigenvalues of (1).

The main features of the present paper are as follows. By virtue of a special transformation, we first convert (1) into another solvable form such that the associated operator can be used to overcome the influence of impulse and parameter μ . Then a special cone is constructed to deal with the singularity of (1).

This paper is organized as follows. In Section 2, we provide some basic definitions, preliminaries facts, and lemmas. Meanwhile, some transformations are introduced to convert (1) into another solvable form. In Section 3, the main results

are presented and proved. Finally, an example is worked out to demonstrate the application of the main result.

2. Preliminaries and Conversion of (1)

Let P be a normal solid cone of real Banach space E . Without loss of generality, suppose the normal constant is 1. Let P^* denote the dual cone of P , $J = [0, 1]$, and $P_r = \{x \in P : \|x\| < r\}$, $\bar{P}_r = \{x \in P : \|x\| \leq r\}$ ($r > 0$). Denoted by $C[J, E]$ the Banach space of all continuous functions $x : J \rightarrow E$ with norm $\|x\|_C = \max_{t \in J} \|x(t)\|$.

We define $PC[J, E] = \{x : x \text{ is a map from } J \text{ into } E \text{ such that } x(t) \text{ is continuous at } t \neq t_i \text{ and left continuous at } t = t_i \text{ and its right limit at } t = t_i \text{ (denoted by } x(t_i^+)) \text{ exists for } i = 1, 2, \dots, m\}$, and $PC^1[J, E] = \{x \in PC[J, E] : x'(t) \text{ is continuous at } t \neq t_i \text{ and left continuous at } t = t_i \text{ and the right limit at } t = t_i \text{ (denoted by } x'(t_i^+)) \text{ exists for } i = 1, 2, \dots, m\}$.

Let $Y = \{y \in C^1[J, E] : ay(0) - by'(0) = \theta, cy(1) + dy'(1) = \theta\}$, and $X = \{x \in PC^1[J, E] : \Delta x|_{t=t_i} = \alpha_i x(t_i - 0), \Delta x'|_{t=t_i} = \alpha_i x'(t_i - 0), ax(0) - bx'(0) = \theta, cx(1) + dx'(1) = \theta\}$. It is well known that Y is a Banach space with the norm $\|y\|_1 = \max\{\max_{t \in J} \|y(t)\|, \max_{t \in J} \|y'(t)\|\}$.

Evidently, $PC[J, E]$ and $PC^1[J, E]$ are Banach spaces with norm $\|x\|_{PC} = \sup_{t \in J} \|x(t)\|$ and $\|x\|_{PC^1} = \max\{\sup_{t \in J} \|x(t)\|, \sup_{t \in J} \|x'(t)\|\}$, respectively. Furthermore, X is a closed subspace of $PC^1[J, E]$.

For each $y \in Y$, let $x(t) = \prod_{0 < t_i < t} (1 + \alpha_i) y(t)$; it is easy to see $x \in X$. Conversely, for each $x \in X$, let $y(t) = x(t) / \prod_{0 < t_i < t} (1 + \alpha_i)$; then $y \in Y$.

We convert (1) into another solvable form first.

Lemma 1. *If $x \in X$ is a solution of impulsive SBVP (1), then $y(t) = x(t) / \prod_{0 < t_i < t} (1 + \alpha_i)$ satisfies SBVP:*

$$\mu y'' + \frac{1}{\prod_{0 < t_i < t} (1 + \alpha_i)} f\left(t, \prod_{0 < t_i < t} (1 + \alpha_i) y(t)\right) = 0, \quad t \in (0, 1); \quad (2)$$

$$ay(0) - by'(0) = \theta;$$

$$cy(1) + dy'(1) = \theta.$$

Conversely, if $y \in Y$ is a solution of SBVP (2), then $x(t) = \prod_{0 < t_i < t} (1 + \alpha_i) y(t)$ is a solution of impulsive SBVP (1).

Lemma 2. *$y \in Y$ is a solution of SBVP (2) if and only if y is a solution of the integral equation:*

$$\begin{aligned} \mu y(t) = & \int_J \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s) \\ & \times f\left(t, \prod_{0 < t_i < s} (1 + \alpha_i) y(s)\right) ds, \end{aligned} \quad (3)$$

where

$$G(t, s) = \begin{cases} \gamma^{-1}(at + b)(c(1 - s) + d), & 0 \leq t \leq s \leq 1, \\ \gamma^{-1}(as + b)(c(1 - t) + d), & 0 \leq s \leq t \leq 1. \end{cases} \quad (4)$$

By Lemmas 1 and 2, we can obtain the following.

Lemma 3. *$x(t) \in PC[J, E] \cap X$ is a solution of impulsive SBVP (1) if and only if $x \in PC[J, E] \cap X$ is a solution of the integral equation:*

$$\mu x(t) = \int_J G^*(t, s) f(s, x(s)) ds, \quad (5)$$

where

$$G^*(t, s) = \frac{\prod_{0 < t_i < t} (1 + \alpha_i)}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s). \quad (6)$$

Let $\alpha(\cdot)$ and $\alpha_{pc}(\cdot)$ denoted the Kuratowski noncompactness measure of bounded sets in E and $PC[J, E]$, respectively. If nontrivial function $x \in PC[J, E]$ satisfies problem (5) for some $\mu \neq 0$, then μ is called an eigenvalue and x is called an eigenfunction of problem (1) corresponding to the eigenvalue μ .

It is well known that the following conclusions hold.

Lemma 4 (see [17]). *Let P be a cone in Banach space E , $P_r = \{x \in P : \|x\| < r\}$ ($r > 0$), and $\bar{P}_r = \{x \in P : \|x\| \leq r\}$. Let operator $A : \bar{P}_r \rightarrow P$ be a strict set contraction. If $\|Ax\| \geq \|x\|$ and $Ax \neq x$ for $x \in \partial P_r$, then $i(A, P_r, P) = 0$.*

Lemma 5 (see [17]). *$H \subset C[J, E]$ is relatively compact if and only if H is equicontinuous and for any $t \in J$, $H(t)$ is a relatively compact set in E .*

3. Main Results

For convenience, let

$$\sigma_* =: \min_{1 \leq j \leq k} \prod_{1 \leq i \leq j} (1 + \alpha_i), \quad \sigma^* =: \max_{1 \leq j \leq k} \prod_{1 \leq i \leq j} (1 + \alpha_i),$$

$$\Omega(t) = \frac{\varrho(t) \prod_{0 < t_i < t} (1 + \alpha_i)}{\sigma^*}, \quad (7)$$

$$\varrho(t) = \min \left\{ \frac{b + at}{b + a}, \frac{d + c(1 - t)}{d + c} \right\}, \quad \forall t \in (0, 1).$$

For the forthcoming analysis, we list the following assumptions:

$$(H_1) \quad f \in C[(0, 1) \times P \setminus \{\theta\}, P],$$

$$\|f(t, x)\| \leq g(t) \|l(x)\|, \quad t \in (0, 1), \quad x \in P \setminus \{\theta\}, \quad (8)$$

where $g : (0, 1) \rightarrow (0, +\infty)$, $l \in C[P \setminus \{\theta\}, P]$ satisfy

$$\int_0^1 G(s, s) g(s) l[\Omega(s)r, R] ds < +\infty, \quad \forall R > r > 0, \quad (9)$$

where $l[r, R] = \sup_{x \in \bar{P}_R \setminus P_r} \|l(x)\| < +\infty$.

(H₂) There exists a constant $L \geq 0$ such that

$$\begin{aligned} \alpha(f(t, D)) &\leq L\alpha(D), \quad \forall t \in (0, 1), \\ D &\subset \bar{P}_R \setminus P_r, \quad \forall R > r > 0, \\ L &< \frac{\sigma_*}{2M_0\sigma^*}, \quad M_0 = \max_{s \in J} \gamma^{-1}(as + b)(c(1-s) + d) > 0. \end{aligned} \quad (10)$$

(H₃) There exist $u_0 \in P \setminus \{\theta\}$, $[a_0, b_0] \subset (0, 1)$ such that

$$\begin{aligned} f(t, x) &\geq h(\|x\|)u_0, \quad t \in J_0 = [a_0, b_0], \\ \frac{2\sigma^*}{\|u_0\| \sigma_* \max_{t \in J} \int_{a_0}^{b_0} G(t, s) \Omega(s) ds} & & (11) \\ &< \liminf_{t \rightarrow 0^+} \frac{h(t)}{t} \leq +\infty, \end{aligned}$$

where $h \in C[R^+, R^+]$.

(H₄) There exist $\phi \in P^*$ and $[a_1, b_1] \subset (0, 1)$ such that $\phi(x) > 0$ for $x > \theta$ and

$$\liminf_{\|x\| \rightarrow \infty, x \in P} \frac{\phi(f(t, x))}{\phi(x)} = \xi(t) \quad (12)$$

uniformly for $t \in J_1 = [a_1, b_1]$, and $\int_{J_1} G(t, s)\xi(s)ds > \sigma^*/\sigma_*$.

(H₅) There exist $\phi \in P^*$ and $[a_1, b_1] \subset (0, 1)$ such that $\phi(x) > 0$ for $x > \theta$ and

$$\liminf_{\|x\| \rightarrow 0, x \in P} \frac{\phi(f(t, x))}{\phi(x)} = \xi(t) \quad (13)$$

uniformly for $t \in J_1 = [a_1, b_1]$, and $\int_{J_1} G(t, s)\xi(s)ds > \sigma^*/\sigma_*$.

Define

$$Q = \{x \in PC[J, P] : x(t) \geq \Omega(t)x(s), \forall t, s \in J\}. \quad (14)$$

It is easy to check that Q is a cone in space $PC[J, E]$ and $Q \subset PC[J, P]$. Let $Q_r = \{x \in Q : \|x\| < r\}$. As in [2], we can prove that

$$G(t, \tau) \geq \varrho(t)G(s, \tau), \quad \forall t, s, \tau \in [0, 1], \quad (15)$$

where $G(t, s)$ is defined in Lemma 2. So for all $t, s, \tau \in [0, 1]$, we have

$$\begin{aligned} G^*(t, s) &\geq \frac{\varrho(t) \prod_{0 < t_i < t} (1 + \alpha_i)}{\prod_{0 < t_i < \tau} (1 + \alpha_i)} \cdot \frac{\prod_{0 < t_i < \tau} (1 + \alpha_i)}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(\tau, s) \quad (16) \\ &\geq \Omega(t)G^*(\tau, s). \end{aligned}$$

To solve eigenvalues of impulsive SBVP (1), we first consider operator A associated with (5) and defined by

$$(Ax)(t) = \int_J G^*(t, s)f(s, x(s))ds, \quad \forall x \in Q \setminus \{\theta\}. \quad (17)$$

For $\forall x \in Q \setminus \{\theta\}$, from the definition of Q , we have

$$\Omega(t)\|x\|_{pc} \leq \|x(t)\| \leq \|x\|_{pc}, \quad t \in J. \quad (18)$$

By (H₁), we know

$$\begin{aligned} \|(Ax)(t)\| &\leq \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(s, s) \|f(s, x(s))\| ds \\ &\leq \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ &\quad \times G(s, s) g(s) \|l(x(s))\| ds \\ &\leq \frac{\sigma^*}{\sigma_*} \int_0^1 G(s, s) g(s) l[\Omega(s)\|x\|_{pc}, \|x\|_{pc}] ds < +\infty. \end{aligned} \quad (19)$$

So the operator A is well defined in $Q \setminus \{\theta\}$.

For the sake of overcoming the singularity, choose $e \in \text{int } P$ with $\|e\| = 1$ and consider the approximate problem of (17):

$$(A_m x)(t) = \int_0^1 G^*(t, s)f\left(s, x(s) + \frac{e}{m}\right)ds. \quad (20)$$

For any $x \in Q, t, \tau \in J$,

$$\begin{aligned} (A_m x)(t) &= \int_0^1 G^*(t, s)f\left(s, x(s) + \frac{e}{m}\right)ds \\ &\geq \Omega(t) \int_0^1 G^*(\tau, s)f\left(s, x(s) + \frac{e}{m}\right)ds \quad (21) \\ &= \Omega(t)(A_m x)(\tau). \end{aligned}$$

Hence $A_m Q \subset Q$.

Lemma 6. Let conditions (H₁) and (H₂) be satisfied; then for any $r > 0$, the operator A_m is a strict set contraction from Q into Q .

Proof. Obviously $A_m Q \subset Q$. Now we prove that A_m is continuous. Let $\|x_n - x\|_{pc} \rightarrow 0$ as $n \rightarrow \infty$ ($x_n, x \in \bar{Q}_r$); we have

$$\begin{aligned}
 & \| (A_m x_n)(t) - (A_m x)(t) \| \\
 & \leq \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s) \\
 & \quad \times \left\| f\left(s, x_n(s) + \frac{e}{m}\right) - f\left(s, x(s) + \frac{e}{m}\right) \right\| ds \\
 & \leq \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(s, s) g(s) \\
 & \quad \times \left(\left\| l\left(x_n(s) + \frac{e}{m}\right) \right\| + \left\| l\left(x(s) + \frac{e}{m}\right) \right\| \right) ds \\
 & \leq 2 \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(s, s) g(s) \\
 & \quad \times l[\Omega(s)r, r+1] ds \\
 & \leq 2 \frac{\sigma^*}{\sigma_*} \int_0^1 G(s, s) g(s) l[\Omega(s)r, r+1] ds.
 \end{aligned} \tag{22}$$

So (H_1) and the dominated convergence theorem imply that

$$\lim_{n \rightarrow \infty} (A_m x_n)(t) = (A_m x)(t). \tag{23}$$

We now show that

$$\lim_{n \rightarrow \infty} \|A_m x_n - A_m x\|_{pc} = 0. \tag{24}$$

In fact, if (24) is not true, then there exist a positive number ε_0 and a sequence $\{x_{n_i}\} \subset \{x_n\}$ such that

$$\|A_m x_{n_i} - A_m x\|_{pc} \geq \varepsilon_0 \quad (i = 1, 2, \dots). \tag{25}$$

Since $\{A_m x_n\}$ is relatively compact, there is a subsequence of $\{A_m x_{n_i}\}$ which converges to some $x_0 \in Q$. Without loss of generality, we may assume that $\{A_m x_{n_i}\}$ itself converges to x_0 ; that is,

$$\lim_{i \rightarrow \infty} \|A_m x_{n_i} - x_0\|_{pc} = 0. \tag{26}$$

By virtue of (23) and (26), we have $x_0 = A_m x$, and so (26) contradicts with (25). Hence (24) holds, and the continuity of A_m is proved.

By (H_1) , it is easy to see that A_m is bounded from Q into Q .

Now we will prove that the operator A_m is a strict set contraction from Q into Q . Let

$$(A_m^{(n)} x)(t) = \int_{1/n}^{1-(1/n)} G^*(t, s) f\left(s, x(s) + \frac{e}{m}\right) ds. \tag{27}$$

By (H_1) , as $n \rightarrow \infty$, we have

$$\begin{aligned}
 & \| (A_m x)(t) - (A_m^{(n)} x)(t) \| \\
 & \leq \int_0^{1/n} G^*(t, s) \left\| f\left(s, x(s) + \frac{e}{m}\right) \right\| ds \\
 & \quad + \int_{1-(1/n)}^1 G^*(t, s) \left\| f\left(s, x(s) + \frac{e}{m}\right) \right\| ds \rightarrow 0.
 \end{aligned} \tag{28}$$

So for any bounded set $U \subset Q$, we have

$$\lim_{n \rightarrow \infty} (A_m^{(n)} x)(t) = (A_m x)(t), \quad \forall x \in U, t \in J. \tag{29}$$

Hence

$$d_h((A_m^{(n)} U)(t), (A_m U)(t)) \rightarrow 0, \quad n \rightarrow \infty, \tag{30}$$

where $d_h(\cdot, \cdot)$ denotes the Hausdorff metric, which implies

$$\alpha((A_m^{(n)} U)(t)) \rightarrow \alpha((A_m U)(t)), \quad n \rightarrow \infty, t \in J. \tag{31}$$

For any $t \in J$, by (H_2) , we have

$$\begin{aligned}
 & \alpha((A_m^{(n)} U)(t)) \\
 & = \alpha\left(\left\{ \prod_{0 < t_i < t} (1 + \alpha_i) \right. \right. \\
 & \quad \times \int_{1/n}^{1-(1/n)} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s) \\
 & \quad \times \left. \left. f\left(s, x(s) + \frac{e}{m}\right) ds, x \in U \right\}\right) \\
 & \leq \left(1 - \frac{1}{2n}\right) \prod_{0 < t_i < t} (1 + \alpha_i) \\
 & \quad \times \alpha\left(\overline{co}\left\{ \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s) \right. \right. \\
 & \quad \times \left. \left. f\left(s, x(s) + \frac{e}{m}\right) : s \in \left[\frac{1}{n}, 1 - \frac{1}{n}\right], x \in U \right\}\right)
 \end{aligned}$$

$$\begin{aligned}
&\leq \prod_{0 < t_i < t} (1 + \alpha_i) \\
&\quad \times \alpha \left(\left\{ \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(s, s) \right. \right. \\
&\quad \left. \left. \times f \left(s, x(s) + \frac{e}{m} \right) : s \in \left[\frac{1}{n}, 1 - \frac{1}{n} \right], x \in U \right\} \right) \\
&\leq \frac{\sigma^*}{\sigma_*} M_0 \alpha(f(s, U(I_n))), \quad I_n = \left[\frac{1}{n}, 1 - \frac{1}{n} \right], s \in I_n, \\
&\leq LM_0 \frac{\sigma^*}{\sigma_*} \alpha(U(I_n)) \\
&\leq 2LM_0 \frac{\sigma^*}{\sigma_*} \alpha_{pc}(U) \\
&< \alpha_{pc}(U).
\end{aligned} \tag{32}$$

It is easy to show that $\alpha_{pc}(A_m(U)) = \sup_{t \in J} \alpha((A_m(U))(t))$. Therefore, we can have $\alpha_{pc}(A_m U) \leq 2LM_0(\sigma^*/\sigma_*)\alpha_{pc}(U) < \alpha_{pc}(U)$. Consequently, the operator A_m is a strict set contraction from Q into Q . The proof is thus completed. $\square$

Theorem 7. Let conditions (H_1) , (H_2) , and (H_3) be satisfied; then there exists a positive number δ such that for any given $r \in (0, \delta)$, the impulsive SBVP (1) has a nontrivial eigenfunction $x_r \in PC[J, P]$ with $\|x_r\|_{pc} = r$ corresponding to eigenvalue $\mu_r \geq 1$.

Proof. By Lemma 6, the operator A_m is a strict set contraction from $\overline{Q_r}$ into Q . Observing (H_3) , if $\liminf_{t \rightarrow 0^+} (h(t)/t) \neq +\infty$, we can choose a v_0 with

$$\frac{1}{2} \liminf_{t \rightarrow 0^+} \frac{h(t)}{t} \leq v_0 < \liminf_{t \rightarrow 0^+} \frac{h(t)}{t}, \tag{33}$$

and there is a $\delta > 0$ such that

$$h(t) \geq v_0 t, \quad \forall 0 < t < \delta. \tag{34}$$

Choose $r \in (0, \delta)$ and m sufficiently large such that $(1/m) < \delta - r$; we have

$$\begin{aligned}
&(A_m x)(t) \\
&= \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
&\quad \times G(t, s) f \left(s, x(s) + \frac{e}{m} \right) ds \\
&\geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_{a_0}^{b_0} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s)
\end{aligned}$$

$$\begin{aligned}
&\quad \times h \left(\left\| x(s) + \frac{e}{m} \right\| \right) u_0 ds \\
&\geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_{a_0}^{b_0} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
&\quad \times G(t, s) v_0 \|x(s)\| u_0 ds \\
&\geq \frac{\sigma_*}{\sigma^*} \int_{a_0}^{b_0} G(t, s) v_0 \Omega(s) \|x\|_{PC} u_0 ds.
\end{aligned} \tag{35}$$

So

$$\begin{aligned}
&\|A_m x\|_{PC} \\
&\geq \frac{\sigma_*}{\sigma^*} v_0 \|u_0\| \max_{t \in J} \int_{a_0}^{b_0} G(t, s) \Omega(s) ds \|x\|_{PC} > \|x\|_{PC}.
\end{aligned} \tag{36}$$

By virtue of Lemma 4, we have $i(A_m, Q_r, Q) = 0$. Since $i(\theta, Q_r, Q) = 1$, it follows from the homotopy invariance of fixed point index for strict set contraction that there exist $x_m \in \partial \overline{Q_r}$ and $\lambda_m \in (0, 1)$ such that

$$x_m = \lambda_m A_m x_m, \quad \text{i.e. } A_m x_m = \lambda_m^{-1} x_m, \tag{37}$$

where $m \geq m_0 = [1/(\delta - r)] + 1$.

Let $B = \{x_m : m \geq m_0\}$. Obviously, B is uniformly bounded. We shall show that B is equicontinuous. By virtue of boundary condition, we need to consider only the following eight cases.

Case 1. $b = d = 0$, the boundary value condition is $x(0) = x(1) = 0$. So we need only to show that $\{x_m(t)\}$ uniformly converge to θ with respect to $m \geq m_0$ as $t \rightarrow 0^+$, $t \rightarrow 1 - 0$, and B is equicontinuous on any closed subinterval of $(0, 1)$.

By (H_1) , let

$$\eta =: \int_0^1 G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds < +\infty. \tag{38}$$

Using the absolute continuity of integration, for any $\varepsilon > 0$, there exists a $\delta_1 > 0$, such that

$$\int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds < \varepsilon \tag{39}$$

as $|t_1 - t_2| < \delta_1, t_1, t_2 \in J$. Since

$$\begin{aligned}
x_m(t) &= \lambda_m \int_0^1 G^*(t, s) f \left(s, x_m(s) + \frac{e}{m} \right) ds \\
&\leq \lambda_m \frac{\sigma^*}{\sigma_*} \left[(1-t) \int_0^t s f \left(s, x_m(s) + \frac{e}{m} \right) ds \right. \\
&\quad \left. + t \int_t^1 (1-s) f \left(s, x_m(s) + \frac{e}{m} \right) ds \right],
\end{aligned} \tag{40}$$

we have

$$\begin{aligned}
& \|x_m(t)\| \\
& \leq \frac{\sigma^*}{\sigma_*} \left\{ (1-t) \int_0^t sg(s) \left\| l \left(x_m(s) + \frac{e}{m} \right) \right\| ds \right. \\
& \quad \left. + t \int_t^1 (1-s) g(s) \left\| l \left(x_m(s) + \frac{e}{m} \right) \right\| ds \right\} \quad (41) \\
& \leq \frac{\sigma^*}{\sigma_*} \left\{ (1-t) \int_0^t sg(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \right. \\
& \quad \left. + t \int_t^1 (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \right\}.
\end{aligned}$$

To prove that $\{x_m(t)\}$ uniformly converge to θ with respect to $m \geq m_0$ as $t \rightarrow 0^+$, we need only to show

$$\lim_{t \rightarrow 0^+} (1-t) \int_0^t sg(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds = 0, \quad (42)$$

$$\lim_{t \rightarrow 0^+} t \int_t^1 (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds = 0. \quad (43)$$

Notice that

$$\begin{aligned}
& (1-t) \int_0^t sg(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& \leq \int_0^t (1-s) sg(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \quad (44) \\
& = \int_0^t G(s, s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds.
\end{aligned}$$

This together with (39) implies that (42) holds. For ε, δ_1 in (39), choose

$$\delta_2 = \min \left\{ \delta_1, \frac{\varepsilon \delta_1}{\eta} \right\}. \quad (45)$$

Then for $t \in (0, \delta_2)$, (39) implies that

$$\begin{aligned}
& t \int_t^1 (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& \leq t \int_t^{\delta_1} (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds
\end{aligned}$$

$$\begin{aligned}
& + \int_{\delta_1}^1 \frac{t}{s} (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& \leq \int_t^{\delta_1} s (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& \quad + \frac{t}{\delta_1} \int_{\delta_1}^1 s (1-s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& = \int_t^{\delta_1} G(s, s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \quad (46) \\
& \quad + \frac{t}{\delta_1} \int_{\delta_1}^1 G(s, s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& \leq \varepsilon + \frac{t}{\delta_1} \int_0^1 G(s, s) g(s) l \left[\Omega(s)r, r + \frac{1}{m_0} \right] ds \\
& \leq 2\varepsilon.
\end{aligned}$$

Hence (43) holds. Very similarly, we can obtain that $\{x_m(t)\}$ uniformly converge to θ with respect to $m \geq m_0$ as $t \rightarrow 1-0$. Now, we show that B is equicontinuous on $[\zeta, 1-\zeta]$ for any $[\zeta, 1-\zeta] \subset (0, 1)$, $\zeta \in (0, 1/2)$. Notice that

$$\begin{aligned}
& x_m(t_2) - x_m(t_1) \\
& = \lambda_m \prod_{0 < t_i < t} (1 + \alpha_i) \\
& \quad \times \left[(1-t_2) \int_0^{t_2} s \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} f \left(s, x_m(s) + \frac{e}{m} \right) ds \right. \\
& \quad + t_2 \int_{t_2}^1 (1-s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} f \left(s, x_m(s) + \frac{e}{m} \right) ds \\
& \quad - (1-t_1) \int_0^{t_1} s \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} f \left(s, x_m(s) + \frac{e}{m} \right) ds \\
& \quad \left. - t_1 \int_{t_1}^1 (1-s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} f \left(s, x_m(s) + \frac{e}{m} \right) ds \right] \\
& = \lambda_m \prod_{0 < t_i < t} (1 + \alpha_i) \\
& \quad \times \left[(t_2 - t_1) \right. \\
& \quad \times \left(- \int_0^{t_2} s \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} f \left(s, x_m(s) + \frac{e}{m} \right) ds \right. \\
& \quad \left. + \int_{t_2}^1 (1-s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \right. \\
& \quad \left. \left. \times f \left(s, x_m(s) + \frac{e}{m} \right) ds \right) \right]
\end{aligned}$$

$$\begin{aligned}
& + (1 - t_1) \int_{t_1}^{t_2} s \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times f\left(s, x_m(s) + \frac{e}{m}\right) ds \\
& - t_1 \int_{t_1}^{t_2} (1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times f\left(s, x_m(s) + \frac{e}{m}\right) ds \Big].
\end{aligned} \tag{47}$$

So for any $t_1, t_2 \in [\zeta, 1 - \zeta]$, $t_2 > t_1$, $m \geq m_0$, we have

$$\begin{aligned}
& \|x_m(t_2) - x_m(t_1)\| \\
& \leq \prod_{0 < t_i < t} (1 + \alpha_i) \\
& \quad \times \left\{ (t_2 - t_1) \left[\int_0^{t_2} s \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \right. \right. \\
& \quad \times g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \\
& \quad + \int_{t_2}^1 (1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \Big] \\
& + (1 - t_1) \int_{t_1}^{t_2} s \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \\
& + t_1 \int_{t_1}^{t_2} (1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \Big\} \\
& \leq \prod_{0 < t_i < t} (1 + \alpha_i) \\
& \quad \times \left\{ (t_2 - t_1) \left(\frac{1}{1 - t_2} \int_0^{t_2} (1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \right. \right. \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \\
& + \frac{1}{t_2} \int_{t_2}^1 s(1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \Big)
\end{aligned}$$

$$\begin{aligned}
& + \frac{1 - t_1}{1 - t_2} \int_{t_1}^{t_2} s(1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \\
& + \int_{t_1}^{t_2} s(1 - s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \Big\} \\
& \leq \prod_{0 < t_i < t} (1 + \alpha_i) \\
& \quad \times \left\{ (t_2 - t_1) \left(\frac{1}{1 - t_2} \int_0^{t_2} G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \right. \right. \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \\
& + \frac{1}{t_2} \int_{t_2}^1 G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \Big) \\
& + \frac{1 - t_1}{1 - t_2} \int_{t_1}^{t_2} G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \\
& + \int_{t_1}^{t_2} G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \Big\} \\
& \leq \prod_{0 < t_i < t} (1 + \alpha_i) \\
& \quad \times \left\{ (t_2 - t_1) \left(\frac{1}{\zeta} \int_0^{t_2} G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \right. \right. \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \\
& + \frac{1}{\zeta} \int_{t_2}^1 G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \Big) \\
& + \left(\frac{1 - \zeta}{\zeta} + 1 \right) \int_{t_1}^{t_2} G(s, s) \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\
& \quad \times g(s) l\left[\Omega(s)r, r + \frac{1}{m_0}\right] ds \Big\}
\end{aligned}$$

$$\leq \frac{\sigma^*}{\sigma_*} \left\{ (t_2 - t_1) \frac{2\eta}{\zeta} + \frac{1}{\zeta} \int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \right\}. \quad (48)$$

This together with (39) implies that B is equicontinuous on J .

Case 2. $b = 0, d \neq 0, c \neq 0$, the boundary value condition is $x(0) = 0, cx(1) + dx'(1) = 0$. We need to show that $\{x_m(t)\}$ uniformly converge to θ with respect to $m \geq m_0$ as $t \rightarrow 0^+$. This can be obtained by the similar way in Case 1, so we omit it. Next, we show B is equicontinuous on $[t, 1]$ for any $t \in (0, 1)$. In fact, for any $t_1, t_2 \in [t, 1], t_2 > t_1, m \geq m_0$, we have

$$\begin{aligned} & \|x_m(t_2) - x_m(t_1)\| \\ & \leq \int_0^{t_1} |G^*(t_2, s) - G^*(t_1, s)| \left\| f\left(s, x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \quad + \int_{t_1}^{t_2} |G^*(t_2, s) - G^*(t_1, s)| \left\| f\left(s, x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \quad + \int_{t_2}^1 |G^*(t_2, s) - G^*(t_1, s)| \left\| f\left(s, x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \leq (t_2 - t_1) \frac{c}{c(1-s) + d} \frac{\sigma^*}{\sigma_*} \\ & \quad \times \int_0^{t_1} \frac{s(c(1-s) + d)}{c + d} g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \quad + 2 \frac{\sigma^*}{\sigma_*} \int_{t_1}^{t_2} G(s, s) g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \quad + (t_2 - t_1) \frac{1}{s} \frac{\sigma^*}{\sigma_*} \\ & \quad \times \int_{t_2}^1 \frac{s(c(1-s) + d)}{c + d} g(s) \left\| l\left(x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \leq (t_2 - t_1) \frac{c}{d} \frac{\sigma^*}{\sigma_*} \int_0^{t_1} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \quad + 2 \frac{\sigma^*}{\sigma_*} \int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \quad + (t_2 - t_1) \frac{1}{t} \frac{\sigma^*}{\sigma_*} \int_{t_2}^1 G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \leq (t_2 - t_1) \eta \frac{\sigma^*}{\sigma_*} \left(\frac{c}{d} + \frac{1}{t} \right) \\ & \quad + 2 \frac{\sigma^*}{\sigma_*} \int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds. \end{aligned} \quad (49)$$

This together with (39) implies that B is equicontinuous on J .

Case 3. $b = 0, d \neq 0, c = 0$, the boundary value condition is $x(0) = 0, x'(1) = 0$. The equicontinuity of B on J of this case can be obtained by the same way of Case 2, so we omit it.

Case 4. $d = 0, b \neq 0, a \neq 0$, the boundary value condition is $ax(0) - bx'(0) = 0, x(1) = 0$. Similarly, by Case 1, we can get that $\{x_m(t)\}$ uniformly converge to θ with respect to $m \geq m_0$ as $t \rightarrow 1^-$. For any $\varsigma \in (0, 1)$, we need to show that B is equicontinuous on $[0, \varsigma]$. In fact, for any $t_1, t_2 \in [0, \varsigma], t_2 > t_1, m \geq m_0$, we have

$$\begin{aligned} & \|x_m(t_2) - x_m(t_1)\| \\ & \leq \int_0^{t_1} |G^*(t_2, s) - G^*(t_1, s)| \left\| f\left(s, x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \quad + \int_{t_1}^{t_2} |G^*(t_2, s) - G^*(t_1, s)| \left\| f\left(s, x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \quad + \int_{t_2}^1 |G^*(t_2, s) - G^*(t_1, s)| \left\| f\left(s, x_m(s) + \frac{e}{m}\right) \right\| ds \\ & \leq (t_2 - t_1) \frac{1}{1-s} \frac{\sigma^*}{\sigma_*} \int_0^{t_1} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \quad + 2 \frac{\sigma^*}{\sigma_*} \int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \quad + (t_2 - t_1) \frac{a}{as + b} \frac{\sigma^*}{\sigma_*} \int_{t_2}^1 G(s, s) g(s) \\ & \quad \quad \times l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \leq (t_2 - t_1) \frac{1}{1-\varsigma} \frac{\sigma^*}{\sigma_*} \int_0^{t_1} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \quad + 2 \frac{\sigma^*}{\sigma_*} \int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \quad + (t_2 - t_1) \frac{b}{a} \frac{\sigma^*}{\sigma_*} \int_{t_2}^1 G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds \\ & \leq (t_2 - t_1) \eta \frac{\sigma^*}{\sigma_*} \left(\frac{1}{1-\varsigma} + \frac{b}{a} \right) \\ & \quad + 2 \frac{\sigma^*}{\sigma_*} \int_{t_1}^{t_2} G(s, s) g(s) l \left[\Omega(s) r, r + \frac{1}{m_0} \right] ds. \end{aligned} \quad (50)$$

This together with (39) implies that B is equicontinuous on J .

Case 5. $d = 0, b \neq 0, a = 0$, the boundary value condition is $x'(0) = 0, x(1) = 0$.

Case 6. $a = 0, bd \neq 0, c \neq 0$, the boundary value condition is $x'(0) = 0, cx(1) + dx'(1) = 0$.

Case 7. $a \neq 0, bd \neq 0, c = 0$, the boundary value condition is $ax(0) - bx'(0) = 0, x'(1) = 0$.

Case 8. $bd \neq 0, ac \neq 0$, the boundary value condition is $ax(0) - bx'(0) = 0, cx(1) + dx'(1) = 0$.

By the very similar methods in Cases 2 and 4, we can prove that B is equicontinuous on J in Cases 5–8. Hence, the proofs are omitted. Next we show that B is relatively compact. It is easy to see that

$$\begin{aligned} & \alpha(B(t)) \\ & \leq \alpha \left(\left\{ \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s) \right. \right. \\ & \quad \left. \left. \times f \left(s, x_m(s) + \frac{e}{m} \right) ds : m \geq m_0 \right\} \right) \\ & \leq 2 \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} G(t, s) \\ & \quad \times \alpha \left(\left\{ f \left(s, x_m(s) + \frac{e}{m} \right) : m \geq m_0 \right\} \right) ds \\ & \leq 2 \frac{\sigma_*^*}{\sigma_*} L \int_0^1 G(t, s) \alpha(B(s)) ds. \end{aligned} \quad (51)$$

Hence, we have

$$\begin{aligned} \alpha_{pc}(B) & \leq 2 \frac{\sigma_*^*}{\sigma_*} L \int_0^1 G(t, s) ds \alpha_{pc}(B) \\ & \leq 2 \frac{\sigma_*^*}{\sigma_*} L \int_0^1 G(s, s) ds \alpha_{pc}(B) \\ & \leq 2LM_0 \frac{\sigma_*^*}{\sigma_*} \alpha_{pc}(B). \end{aligned} \quad (52)$$

Observing $L < \sigma_*/2M_0\sigma^*$, so $\alpha_{pc}(B) = 0$, which means B is relatively compact.

It follows from Lemma 5 that there is a convergent subsequence of $\{x_m\}$, and without loss of generality, we may assume that $\{x_m\}$ itself converges to some $x_r \in \partial\bar{Q}_r$, and $\lim_{m \rightarrow \infty} \lambda_m = \lambda_r \in [0, 1]$. Hence (40) and the dominated convergence theorem imply that

$$x_r(t) = \lambda_r \int_0^1 G^*(t, s) f(s, x_r(s)) ds = \lambda_r Ax_r(t), \quad t \in J. \quad (53)$$

Obviously, $\lambda_r > 0$. Consequently, $Ax_r = \lambda_r^{-1} x_r = \mu_r x_r$, where $\mu_r = \lambda_r^{-1} \geq 1$.

The case $\liminf_{t \rightarrow 0^+} (h(t)/t) = +\infty$ can be proved similarly, so it is omitted. Then the theorem is proved. $\square$

Theorem 8. Let conditions (H_1) , (H_2) , and (H_4) be satisfied; then there exists a positive number M such that for any given $r > M/\min_{t \in J} \Omega(t)$, the impulsive SBVP (1) has a nontrivial eigenfunction $x_r \in PC[J, P]$ with $\|x_r\|_{PC} = r$ corresponding to eigenvalue $\mu_r \geq 1$.

Proof. By (H_4) , there exist $\varepsilon_0 > 0$ and $M > 0$ such that

$$\int_{a_1}^{b_1} G(t, s) (\xi(s) - \varepsilon_0) ds > \frac{\sigma_*^*}{\sigma_*}, \quad (54)$$

and $\phi(f(t, x)) \geq (\xi(t) - \varepsilon_0)\phi(x)$ for $t \in J_1 = [a_1, b_1]$, $x \in P$ with $\|x\| > M$.

Choose $r > M/\min_{t \in J} \Omega(t)$. If there exists $x_0 \in \partial\bar{Q}$ such that $x_0 \geq A_m x_0$, then

$$\|x_0(s)\| \geq \Omega(s) \|x_0\|_{pc} = \Omega(s) r > M, \quad s \in J. \quad (55)$$

Hence, for $t \in J_1$, we have

$$\begin{aligned} & \phi(x_0(t)) \\ & \geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ & \quad \times G(t, s) \phi \left(f \left(s, x_0(s) + \frac{e}{m} \right) \right) ds \\ & \geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_{J_1} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ & \quad \times G(t, s) (\xi(s) - \varepsilon_0) \phi \left(x_0(s) + \frac{e}{m} \right) ds \\ & \geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_{J_1} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ & \quad \times G(t, s) (\xi(s) - \varepsilon_0) \phi(x_0(s)) ds \\ & \geq \frac{\sigma_*^*}{\sigma_*} \int_{J_1} G(t, s) (\xi(s) - \varepsilon_0) \phi(x_0(s)) ds. \end{aligned} \quad (56)$$

Obviously,

$$\inf_{t \in J_1} \|x_0(t)\| \geq \Omega(t) \|x_0\|_{PC} = \Omega(t) r > 0. \quad (57)$$

The continuity of $x_0(t)$ implies that $\phi(x_0(t))$ is continuous on J_1 . Hence, this together with (57) and (H_4) implies that $\min_{t \in J_1} \phi(x_0(t)) > 0$. Observing (54) and (56), we can obtain a contradiction. Applying Lemma 4, we know that the fixed point index $i(A_m, Q_r, Q) = 0$. Since $i(\theta, Q_r, Q) = 1$, it follows from the homotopy invariance of fixed point index for strict set contraction that there exist $x_m \in \partial\bar{Q}_r$ and $0 < \lambda_m < 1$ such that

$$x_m = \lambda_m A_m x_m, \quad \text{i.e. } A_m x_m = \lambda_m^{-1} x_m = \mu_m x_m, \quad (58)$$

where $\mu_m = \lambda_m^{-1} > 1$. The rest of the proof is completely similar to that of Theorem 7. So it is omitted. $\square$

Theorem 9. Let conditions (H_1) , (H_2) , and (H_5) be satisfied; then there exists a positive number δ such that for any given $r \in (0, \delta)$, the impulsive SBVP (1) has a nontrivial eigenfunction $x_r \in PC[J, P]$ with $\|x_r\|_{PC} = r$ corresponding to eigenvalue $\mu_r \geq 1$.

Proof. By (H_5) , there exist $\varepsilon_1 > 0$ and $\delta > 0$ such that

$$\int_{J_1} G(t, s) (\xi(s) - \varepsilon_1) ds > 1, \quad (59)$$

and $\phi(f(t, x)) \geq (\xi(t) - \varepsilon_1)\phi(x)$ for $t \in J_1 = [a_1, b_1]$, $x \in P$ with $\|x\| < \delta$. Choose $r \in (0, \delta)$ and m such that $(1/m) < \delta - r$. If there exists $x_0 \in \partial Q_r$ such that $x_0 \geq A_m x_0$, then for $t \in J_1$, we have

$$\begin{aligned} & \phi(x_0(t)) \\ & \geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_0^1 \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ & \quad \times G(t, s) \phi\left(f\left(s, x_0(s) + \frac{e}{m}\right)\right) ds \\ & \geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_{J_1} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ & \quad \times G(t, s) (\xi(s) - \varepsilon_1) \phi\left(x_0(s) + \frac{e}{m}\right) ds \\ & \geq \prod_{0 < t_i < t} (1 + \alpha_i) \int_{J_1} \frac{1}{\prod_{0 < t_i < s} (1 + \alpha_i)} \\ & \quad \times G(t, s) (\xi(s) - \varepsilon_1) \phi(x_0(s)) ds \\ & \geq \frac{\sigma^*}{\sigma^*} \int_{J_1} G(t, s) (\xi(s) - \varepsilon_1) \phi(x_0) ds. \end{aligned} \quad (60)$$

Obviously,

$$\inf_{t \in J_1} \|x_0(t)\| \geq \Omega(t) \|x_0\|_{PC} = \Omega(t) r > 0. \quad (61)$$

The continuity of $x_0(t)$ implies that $\phi(x_0(t))$ is continuous on J_1 . Hence, this together with (61) and (H_5) implies that $\min_{t \in J_1} \phi(x_0(t)) > 0$. Observing (59) and (60), then we can obtain a contradiction. Applying Lemma 4, we know that the fixed point index $i(A_m, Q_r, Q) = 0$. Since $i(\theta, Q_r, Q) = 1$, it follows from the homotopy invariance of fixed point index for strict set contraction that there exist $x_m \in \partial Q_r$ and $0 < \lambda_m < 1$ such that

$$x_m = \lambda_m A_m x_m, \quad \text{i.e. } A_m x_m = \lambda_m^{-1} x_m = \mu_m x_m, \quad (62)$$

where $\mu_m = \lambda_m^{-1} > 1$. The rest of the proof is completely similar to that of Theorem 7. So it is omitted. $\square$

Example 10. Consider the following impulsive SBVP:

$$\begin{aligned} \mu x_n'' + \frac{\pi}{\sqrt{t(1-t)}} \left(\frac{1}{3n} (\sqrt{t} x_n + e^{2+x_{n+1}}) + \frac{\arctan t}{\sqrt{n} \|x\|} \right) &= 0, \\ t \in (0, 1), \quad t \neq \frac{1}{5}; \\ \Delta x_n|_{t=1/5} &= \frac{2}{5} x_n, \\ 2x_n(0) - x_n'(0) &= \theta; \\ x_n(1) &= \theta, \quad (n = 1, 2, 3, \dots). \end{aligned} \quad (63)$$

Conclusion. There exists a positive number δ such that for any given $r \in (0, \delta)$, impulsive SBVP (63) has a nontrivial, nonnegative eigenfunction x_{rn} ($n = 1, 2, 3, \dots$) satisfying $\sup_n x_{rn}(t) < \infty$ for $0 \leq t \leq 1$ and $\max_{t \in [0, 1]} \sup_n x_{rn}(t) = r$ corresponding to $\mu_r \geq 1$.

Proof. Let $E = l^\infty = \{x = (x_1, \dots, x_n, \dots) : \sup_n |x_n| < \infty\}$ with norm $\|x\| = \sup_n |x_n|$ and $P = \{x = (x_1, \dots, x_n, \dots) \in l^\infty : x_n \geq 0, n = 1, 2, 3, \dots\}$. Then, P is a normal solid cone of E with normal constant 1, and system (63) can be regarded as an impulsive SBVP in $E = l^\infty$ of the form (1), where $x(t) = (x_1(t), x_2(t), x_3(t), \dots)$, $f(t, x) = (f_1, f_2, f_3, \dots)$, and

$$f_n(t, x) = \frac{\pi}{\sqrt{t(1-t)}} \left(\frac{1}{3n} (\sqrt{t} x_n + e^{2+x_{n+1}}) + \frac{\arctan t}{\sqrt{n} \|x\|} \right). \quad (64)$$

It is easy to see that $f(t, x)$ is singular at $t = 0, 1$ and $x = \theta$.

Let $g(t) = \pi/\sqrt{t(1-t)}$ and $l(x) = (l_1(x), l_2(x), l_3(x), \dots)$, where

$$l_n(x) = \frac{1}{3n} (x_n + e^{2+x_{n+1}}) + \frac{\pi}{4\sqrt{n} \|x\|}. \quad (65)$$

So we have

$$\|f(t, x)\| \leq g(t) \|l(x)\|, \quad t \in (0, 1), \quad x \in P \setminus \{\theta\},$$

$$l[r, R] = \sup_{x \in \bar{P}_R \setminus P_r} \|l(x)\| < R + e^{2+R} + \frac{\pi}{4r} < +\infty, \quad (66)$$

$$\forall R > r > 0.$$

By virtue of $\int_0^1 (\pi/\sqrt{t(1-t)}) dt = \pi^2$, $G(s, s) = (1/3)(1 + 2s)(1 - s)$, and

$$M_0 = \max_{s \in J} G(s, s) = \frac{3}{8}, \quad (67)$$

$$\Omega(s) = \min \left\{ \frac{1+2s}{3}, 1-s \right\} \geq \frac{1}{3} (1+2s)(1-s),$$

we can obtain

$$\begin{aligned} & \int_0^1 G(s, s) g(s) l[\Omega(s) r, R] ds \\ & \leq \frac{3\pi^2}{8} (R + e^{2+R}) + \frac{\pi^3}{4r} < +\infty. \end{aligned} \quad (68)$$

So condition (H_1) is satisfied. On the other hand, by the diagonal method of choosing subsequence, we can see that $L = 0$ in condition (H_2) ; that is, condition (H_2) holds.

We now show that condition (H_3) is satisfied for

$$J_0 = \left[\frac{1}{2}, \frac{9}{10} \right], \quad h(t) = \frac{1}{t},$$

$$u_0 = \left(2\pi \arctan \frac{1}{2}, \frac{2}{\sqrt{2}}\pi \arctan \frac{1}{2}, \frac{2}{\sqrt{3}}\pi \arctan \frac{1}{2}, \dots \right). \quad (69)$$

In fact,

$$f_n(t, x) \geq \frac{\pi}{\sqrt{t(1-t)}} \frac{\arctan t}{\sqrt{n}\|x\|} \geq \frac{2}{\sqrt{n}}\pi \arctan \frac{1}{2} \cdot \frac{1}{\|x\|}, \quad (70)$$

$$\forall t \in \left[\frac{1}{2}, \frac{9}{10} \right],$$

which means that condition (H_3) is satisfied for

$$u_0 = \left(2\pi \arctan \frac{1}{2}, \frac{2}{\sqrt{2}}\pi \arctan \frac{1}{2}, \frac{2}{\sqrt{3}}\pi \arctan \frac{1}{2}, \dots \right),$$

$$J_0 = \left[\frac{1}{2}, \frac{9}{10} \right], \quad h(t) = \frac{1}{t}; \quad (71)$$

that is, condition (H_3) is satisfied.

Hence, our conclusion follows from Theorem 7. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors sincerely thank the reviewers for their useful comments and valuable suggestions on the original paper. This research is supported by NNSF of China (11171192) and Natural Science Foundation of Shandong Province (ZR2013AM005).

References

- [1] Z. Wei, "Positive solution of singular Dirichlet boundary value problems for second order differential equation system," *Journal of Mathematical Analysis and Applications*, vol. 328, no. 2, pp. 1255–1267, 2007.
- [2] Y. Liu and B. Yan, "Multiple solutions of singular boundary value problems for differential systems," *Journal of Mathematical Analysis and Applications*, vol. 287, no. 2, pp. 540–556, 2003.
- [3] X. Xian, D. O'Regan, and C. Yanfang, "Structure of positive solution sets of semi-positone singular boundary value problems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 7-8, pp. 3535–3550, 2010.
- [4] X. Zhang and M. Feng, "Positive solutions for a class of 2nth-order singular boundary value problems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 4, pp. 1287–1298, 2008.
- [5] R. P. Agarwal and D. O'Regan, "Nonlinear superlinear singular and nonsingular second order boundary value problems," *Journal of Differential Equations*, vol. 143, no. 1, pp. 60–95, 1998.
- [6] D. Jiang, "Upper and lower solutions method and a superlinear singular boundary value problem," *Computers & Mathematics with Applications*, vol. 44, no. 3-4, pp. 323–337, 2002.
- [7] Z. Wei and C. Pang, "Positive solutions of nonresonant singular boundary value problem of second order differential equations," *Nagoya Mathematical Journal*, vol. 162, pp. 127–148, 2001.
- [8] Y. Cui and Y. Zou, "Positive solutions of nonlinear singular boundary value problems in abstract spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 1, pp. 287–294, 2008.
- [9] X. Xu and J. Ma, "A note on singular nonlinear boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 293, no. 1, pp. 108–124, 2004.
- [10] S. Staněk, "Positive solutions of singular Dirichlet and periodic boundary value problems," *Computers & Mathematics with Applications*, vol. 43, no. 6-7, pp. 681–692, 2002.
- [11] X. Hao, L. Liu, Y. Wu, and N. Xu, "Multiple positive solutions for singular n th-order nonlocal boundary value problems in Banach spaces," *Computers & Mathematics with Applications*, vol. 61, no. 7, pp. 1880–1890, 2011.
- [12] H. Su, L. Liu, and Y. Wu, "Positive solutions for Sturm-Liouville boundary value problems in a Banach space," *Abstract and Applied Analysis*, vol. 2012, Article ID 572172, 11 pages, 2012.
- [13] X. Liu, W. Feng, and H. Liu, "Non-resonance and eigenvalues of nonlinear singular boundary value problems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 50, no. 7, pp. 995–1012, 2002.
- [14] J. Henderson and W. Yin, "Positive solutions and nonlinear eigenvalue problems for functional differential equations," *Applied Mathematics Letters*, vol. 12, no. 2, pp. 63–68, 1999.
- [15] X. Zhang and L. Liu, "Eigenvalue of fourth-order m -point boundary value problem with derivatives," *Computers & Mathematics with Applications*, vol. 56, no. 1, pp. 172–185, 2008.
- [16] D. Zhao and Y. Liu, "Eigenvalues of nonlinear singular boundary value problems," *International Journal of Mathematics and Computation*, vol. 22, no. 1, pp. 34–46, 2014.
- [17] D. Guo, V. Lakshmikantham, and X. Liu, *Nonlinear Integral Equations in Abstract Spaces*, vol. 373, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1996.
- [18] B. Chanane, "Computing the eigenvalues of singular Sturm-Liouville problems using the regularized sampling method," *Applied Mathematics and Computation*, vol. 184, no. 2, pp. 972–978, 2007.
- [19] K. Q. Lan, "Multiple eigenvalues for singular Hammerstein integral equations with applications to boundary value problems," *Journal of Computational and Applied Mathematics*, vol. 189, no. 1-2, pp. 109–119, 2006.
- [20] J. Ji and B. Yang, "Eigenvalue comparisons for boundary value problems for second order difference equations," *Journal of Mathematical Analysis and Applications*, vol. 320, no. 2, pp. 964–972, 2006.
- [21] J. Henderson and W. Yin, "Positive solutions and nonlinear eigenvalue problems for functional differential equations," *Applied Mathematics Letters: An International Journal of Rapid Publication*, vol. 12, no. 2, pp. 63–68, 1999.
- [22] J. Suo and W. Wang, "Eigenvalues of a class of regular fourth-order Sturm-Liouville problems," *Applied Mathematics and Computation*, vol. 218, no. 19, pp. 9716–9729, 2012.

- [23] H. Coşkun and N. Bayram, "Asymptotics of eigenvalues for regular Sturm-Liouville problems with eigenvalue parameter in the boundary condition," *Journal of Mathematical Analysis and Applications*, vol. 306, no. 2, pp. 548–566, 2005.
- [24] L. Aceto, P. Ghelardoni, and C. Magherini, "Boundary value methods as an extension of Numerov's method for Sturm-Liouville eigenvalue estimates," *Applied Numerical Mathematics*, vol. 59, no. 7, pp. 1644–1656, 2009.
- [25] P. Ghelardoni, "Approximations of Sturm-Liouville eigenvalues using boundary value methods," *Applied Numerical Mathematics*, vol. 23, no. 3, pp. 311–325, 1997.
- [26] S. Somali and V. Oger, "Improvement of eigenvalues of Sturm-Liouville problem with t -periodic boundary conditions," *Journal of Computational and Applied Mathematics*, vol. 180, no. 2, pp. 433–441, 2005.
- [27] Q. Yao, "Positive solutions for eigenvalue problems of fourth-order elastic beam equations," *Applied Mathematics Letters*, vol. 17, no. 2, pp. 237–243, 2004.
- [28] Z. Liu, J. Han, and L. Fang, "Integral boundary value problems for first order integro-differential equations with impulsive integral conditions," *Computers & Mathematics with Applications*, vol. 61, no. 10, pp. 3035–3043, 2011.
- [29] J. Tariboon, "Boundary value problems for first order functional differential equations with impulsive integral conditions," *Journal of Computational and Applied Mathematics*, vol. 234, no. 8, pp. 2411–2419, 2010.
- [30] R. Liang and J. Shen, "Periodic boundary value problem for second-order impulsive functional differential equations," *Applied Mathematics and Computation*, vol. 193, no. 2, pp. 560–571, 2007.
- [31] X. Liu and D. Guo, "Method of upper and lower solutions for second-order impulsive integro-differential equations in a Banach space," *Computers & Mathematics with Applications*, vol. 38, no. 3–4, pp. 213–223, 1999.
- [32] K. Teng and C. Zhang, "Existence of solution to boundary value problem for impulsive differential equations," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 5, pp. 4431–4441, 2010.
- [33] L. Chen and J. Sun, "Boundary value problem of second order impulsive functional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 1, pp. 708–720, 2006.
- [34] Y. Xu and H. Zhang, "Positive solutions of an infinite boundary value problem for n th-order nonlinear impulsive singular integro-differential equations in Banach spaces," *Applied Mathematics and Computation*, vol. 218, pp. 5806–5818, 2012.
- [35] W. Li and G. Song, "Nonlinear boundary value problem for second order impulsive integro-differential equations of mixed type in Banach space," *Computers & Mathematics with Applications*, vol. 56, no. 5, pp. 1372–1381, 2008.
- [36] D. Guo, "Positive solutions of an infinite boundary value problem for n th-order nonlinear impulsive singular integro-differential equations in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 5, pp. 2078–2090, 2009.