Research Article

Some Discussions on the Difference Equation $x_{n+1} = \alpha + (x_{n-1}^m/x_n^k)$

Awad A. Bakery^{1,2}

¹Department of Mathematics, Faculty of Science and Arts, University of Jeddah (UJ), P.O. Box 355, Khulais 21921, Saudi Arabia

²Department of Mathematics, Faculty of Science, Ain Shams University, P.O. Box 1156, Abbassia, Cairo 11566, Egypt

Correspondence should be addressed to Awad A. Bakery; awad.bakery@yahoo.com

Received 6 November 2014; Revised 8 April 2015; Accepted 15 April 2015

Academic Editor: Jaeyoung Chung

Copyright © 2015 Awad A. Bakery. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We give in this work the sufficient conditions on the positive solutions of the difference equation $x_{n+1} = \alpha + (x_{n-1}^m/x_n^k)$, $n = 0, 1, \dots$, where α , k , and $m \in (0, \infty)$ under positive initial conditions x_{-1} , x_0 to be bounded, α -convergent, the equilibrium point to be globally asymptotically stable and that every positive solution converges to a prime two-periodic solution. Our results coincide with that known for the cases $m = k = 1$ of Amleh et al. (1999) and $m = 1$ of Hamza and Morsy (2009). We offer improving conditions in the case of $m = 1$ of Gümüs and Öcalan (2012) and explain our results by some numerical examples with figures.

1. Introduction

Throughout this paper ℓ_∞ and C_α denote, respectively, the class of bounded sequences of real numbers and its subclass of sequences convergent to α . Our aim in this paper is to investigate the positive solutions of the difference equation

$$x_{n+1} = f(x_n, x_{n-1}) = \alpha + \frac{x_{n-1}^m}{x_n^k}, \quad n = 0, 1, \dots, \quad (1)$$

where f is a continuously differentiable function, $\alpha \in [0, \infty)$, $m, k \in (0, \infty)$, and the initial conditions x_{-1} , x_0 are arbitrary positive numbers.

Remark 1. A point $\bar{x} \in \mathbb{R}$ is an equilibrium point of (1), if and only if it is a root for the function

$$g(x) = x - x^{m-k} - \alpha. \quad (2)$$

If we replace x_n and x_{n-1} in (1) by the variables u and v , respectively, then we have

$$\begin{aligned} p &= \frac{\partial f}{\partial u}(\bar{x}, \bar{x}) = -k\bar{x}^{m-k-1}, \\ q &= \frac{\partial f}{\partial v}(\bar{x}, \bar{x}) = m\bar{x}^{m-k-1}. \end{aligned} \quad (3)$$

The linearized equation related with (1) about the equilibrium point $\bar{x}$ is

$$y_{n+1} = py_n + qy_{n-1}, \quad n = 0, 1, 2, \dots \quad (4)$$

Its characteristic equation is

$$\lambda^2 - p\lambda - q = 0. \quad (5)$$

Definition 2. An equilibrium point $\bar{x}$ of difference equation (1) is called locally stable if for every $\varepsilon > 0$ there exists $\delta > 0$ such that if $x_{-1}, x_0 \in (0, \infty)$ with $|x_{-1} - \bar{x}| + |x_0 - \bar{x}| < \delta$, then $|x_n - \bar{x}| < \varepsilon$ for all $n \geq -1$.

Definition 3. An equilibrium point $\bar{x}$ of (1) is called globally asymptotically stable if it is locally stable and, for every $x_{-1}, x_0 \in (0, \infty)$, one has $\lim_{n \rightarrow \infty} x_n = \bar{x}$.

Definition 4. An equilibrium point $\bar{x}$ of difference equation (1) is called unstable if it is not locally stable.

Definition 5. A sequence $(x_n)_{n=-1}^\infty$ is said to be periodic with prime period p if p is the smallest positive integer with $x_{n+p} = x_n$ for all $n \geq -1$.

Theorem 6 (see [1, 2]). (i) A necessary and sufficient condition for both roots of (5) to lie in the open disk $|\lambda| < 1$ is

$$|p| < 1 - q < 2. \quad (6)$$

In this case $\bar{x}$ is locally asymptotically stable.

(ii) A necessary and sufficient condition for one of the roots of (5) to lie outside the open disk $|\lambda| < 1$ and the other root inside it is

$$\begin{aligned} p^2 + 4q &> 0, \\ |p| &> |1 - q|. \end{aligned} \quad (7)$$

In this case $\bar{x}$ is unstable and is called a saddle point.

(iii) A necessary and sufficient condition that a root of (5) has absolute value equal to 1 is

$$|p| = |1 - q| \quad (8)$$

or

$$\begin{aligned} q &= -1, \\ |p| &\leq 2. \end{aligned} \quad (9)$$

In this case $\bar{x}$ is called a nonhyperbolic point.

By applying Theorem 6 in special difference equation (1), we obtain the following results.

Remark 7. (i) $\bar{x}$ is locally asymptotically stable, if and only if

$$|k\bar{x}^{m-k-1}| < 1 - m\bar{x}^{m-k-1} < 2. \quad (10)$$

(ii) $\bar{x}$ is unstable, if and only if

$$\begin{aligned} (k\bar{x}^{m-k-1})^2 + 4m\bar{x}^{m-k-1} &> 0, \\ |k\bar{x}^{m-k-1}| &> |1 - m\bar{x}^{m-k-1}|. \end{aligned} \quad (11)$$

(iii) $\bar{x}$ is called a nonhyperbolic point, if and only if

$$|k\bar{x}^{m-k-1}| = |1 - m\bar{x}^{m-k-1}| \quad (12)$$

or

$$\begin{aligned} m\bar{x}^{m-k-1} &= -1, \\ |k\bar{x}^{m-k-1}| &\leq 2. \end{aligned} \quad (13)$$

Amleh et al. [3], for $m = k = 1$, obtained important results for the following difference equation:

$$x_{n+1} = \alpha + \frac{x_{n-1}}{x_n}. \quad (14)$$

They showed in [3] that when $\alpha > 1$, the equilibrium point $\bar{x} = \alpha + 1$ of (14) is globally asymptotically stable. When $\alpha = 1$, every positive solution of (14) converges to a period-two solution. Every positive solution of (14) is bounded if and only if $\alpha \geq 1$. Finally, when $0 < \alpha < 1$, the equilibrium point is an unstable saddle point. Hamza and Morsy [4], for $m = 1$, gave some results for the difference equation

$$x_{n+1} = \alpha + \frac{x_{n-1}}{x_n^k} \quad \forall k > 0; \quad (15)$$

they investigated the behavior of positive solutions of (15); they proved that when $\alpha > 1$, every positive solution of (15) is bounded and when $\alpha > k^{1/k} \geq 1$, the equilibrium point $\bar{x}$ of (15) is globally asymptotically stable and they showed that (15) has periodic solutions without conditions on α and k . These results are improved in [5] by Gümüs and Öcalan; they investigated the boundedness character of positive solutions of (15); they proved that if $0 \leq \alpha < 1$, then there exist unbounded solutions of (15) in the cases when $\alpha \in (0, \infty)$ and $k \rightarrow 0^+$ or $\alpha > 1$ and $k \rightarrow \infty$; then every positive solution of (15) is unbounded. When $k \rightarrow 0^+$, $k \rightarrow \infty$, and $\alpha > 1$, then every positive solution of (15) is bounded. Also they show that if $k \rightarrow 0^+$, $k \rightarrow \infty$, $\alpha > 1$, and $\alpha > k(1+k)^{(1-k)/k}$, then the equilibrium point $\bar{x}$ of (15) is globally asymptotically stable. Finally they explained the sufficient condition for the fact that every positive solution of (15) converges to a prime two-periodic solution. In our work we prove in Sections 3 and 4 that these results are incorrect and we give the results under finer conditions and we illustrate them in some figures. Closely related equations to (1) are investigated by many authors, for example, [6–11].

In this work in Section 2 we study the global behavior of (1). In Section 3 we investigate the boundedness character of positive solutions of (1). Section 4 is dedicated to the periodic character of the positive solutions of (1). Finally we explain the sufficient condition that every positive solution of (1) converges to a prime two-periodic solution. Throughout the paper we denote by S the class

$$S := \{ \{x_n\}_{n=-1}^{\infty} : x_n \in \mathbb{R}^+, x_n \text{ is a solution of (1)} \} \quad (16)$$

and the subclass S_i of the class S by

$$S_i := \{ \{x_n\}_{n=-1}^{\infty} \in S : x_0, x_{-1} \in (0, 2\alpha] \}. \quad (17)$$

The well-known inequality in [12] will be used: if $0 < p < \infty$ and $D = \max(1, 2^{p-1})$, then

$$|a + b|^p \leq D(|a|^p + |b|^p) \quad \forall a, b \in \mathbb{R}. \quad (18)$$

2. The Difference Equation $x_{n+1} = \alpha + x_{n-1}^m / x_n^k$

In this section we give the sufficient conditions to determine the classification of the equilibrium points for (1) and its uniqueness.

Lemma 8. Let $\bar{x}$ be the equilibrium point of (1) and $m \in (0, k+1]$ for all $k \in (0, \infty)$.

- (i) If $\alpha > (m+k-1)(m+k)^{(m-k)/(k+1-m)}$, then the equilibrium point $\bar{x}$ of (1) is locally asymptotically stable.
- (ii) If $\alpha < (m+k-1)(m+k)^{(m-k)/(k+1-m)}$, then the equilibrium point $\bar{x}$ of (1) is unstable.
- (iii) If $\alpha = (m+k-1)(m+k)^{(m-k)/(k+1-m)}$, then $\bar{x}$ is a nonhyperbolic point.

Proof. Give an equilibrium point $\bar{x}$ of (1) and $m \in (0, k+1]$ for all $k \in (0, \infty)$.

- (i) Suppose $\alpha > (m+k-1)(m+k)^{(m-k)/(k+1-m)}$; then from (2) we have

$$g((m+k)^{1/(k+1-m)}) = (m+k-1)(m+k)^{(m-k)/(k+1-m)} - \alpha < 0, \quad (19)$$

$$\lim_{x \rightarrow \infty} g(x) = \infty.$$

Hence $\bar{x} > (m+k)^{1/(k+1-m)}$ and using Remark 7 we get that $\bar{x}$ is locally asymptotically stable.

- (ii) Let $\alpha < (m+k-1)(m+k)^{(m-k)/(k+1-m)}$; then from (2) we have

$$g((m+k)^{1/(k+1-m)}) = (m+k-1)(m+k)^{(m-k)/(k+1-m)} - \alpha > 0, \quad (20)$$

$$g(0) < 0;$$

hence $\bar{x} < (m+k)^{1/(k+1-m)}$ and using Remark 7 we get that $\bar{x}$ is unstable.

- (iii) If $\alpha = (m+k-1)(m+k)^{(m-k)/(k+1-m)}$, then we obtain $g((m+k)^{1/(k+1-m)}) = 0$.

So $\bar{x} = (m+k)^{1/(k+1-m)}$; by using Remark 7 we get that $\bar{x}$ is a nonhyperbolic point. $\square$

Lemma 9. One has the following:

- (i) If $k = m = 1$, then (1) has a unique equilibrium point $\bar{x} = \alpha + 1$.
- (ii) If $k = m$, then (1) has a unique equilibrium point $\bar{x} = \alpha + 1$.
- (iii) If $k < m \leq k+1$, then (1) has a unique equilibrium point $\bar{x} > 1$.
- (iv) If $0 < m < k$, then (1) has a unique equilibrium point $\bar{x} > 1$.

Proof. (i) If $k = m = 1$, (1) gives $\bar{x} = 1 + \alpha$. See [3].

(ii) If $k = m$, (1) gives $\bar{x} = 1 + \alpha$. See [13].

(iii) If $k < m \leq k+1$, the function g given by (2) is decreasing on $[0, (m-k)^{1/(m-k+1)}]$ and increasing on $[(m-k)^{1/(m-k+1)}, \infty[$. Since $g(1) = -\alpha < 0$ and $\lim_{x \rightarrow \infty} g(x) = \infty$, then g has a unique root $\bar{x} > 1$.

(iv) If $0 < m < k$, the function g is increasing on $[0, \infty[$. Since $g(1) = -\alpha < 0$ and $\lim_{x \rightarrow \infty} g(x) = \infty$, then g has a unique root $\bar{x} > 1$. $\square$

3. Global Behavior of Solutions and Boundedness

In this section, we present the conditions for the boundedness of the positive solutions and its global stability for the equilibrium point of (1).

FIGURE 1: When $m = 1$ and $k \rightarrow 0$ the positive solution is unbounded.

Theorem 10. If $1 \leq m \leq k+1$ and $k \rightarrow 0^+$, then $S \cap \ell_\infty = \emptyset$.

Proof. Suppose on the contrary that $\{x_n\}_{n=-1}^\infty \in S \cap \ell_\infty$. Then, we have

$$\begin{aligned} \psi &:= \liminf x_n, \\ \Psi &:= \limsup x_n < \infty. \end{aligned} \quad (21)$$

Therefore, from (1) we get

$$\psi \geq \alpha + \frac{\psi^m}{\psi^k}. \quad (22)$$

Let $k \rightarrow 0^+$; this implies $m \rightarrow 1$; we obtain

$$\psi \geq \alpha + \psi \implies \alpha \leq 0, \quad (23)$$

which contradicts $\alpha > 0$, so the proof is complete. $\square$

In Figures 1 and 2 we illustrate the result of Theorem 10 for $m = 1$ and small values of $k = 10^{-1}, 10^{-2}, 10^{-4}, 10^{-8}$, and 10^{-13} for, respectively, $\alpha = 1.5$ and $\alpha = 0.5$, and explain that the solutions of (1) are unbounded for every $\alpha > 0$ and $k \rightarrow 0^+$.

Theorem 11. If $m > 1$, then $S \setminus \ell_\infty \neq \emptyset$.

Proof. For fixed $m > 1$ let $\alpha \in (0, 1)$. Choose $\delta \in (0, 1 - \alpha)$ and let $\{x_n\}_{n=-1}^\infty \in S$ with the initial conditions such that

$$x_{-1}^m > \left(\left(\frac{\alpha + \delta}{\delta} \right)^{1/k} - \alpha \right) (\alpha + \delta)^k, \quad (24)$$

$$0 < \alpha < x_0 < \alpha + \delta < 1,$$

FIGURE 2: When $m = 1$ and $k \rightarrow 0$ the positive solution is unbounded.

hence,

$$\begin{aligned} x_1 &= \alpha + \frac{x_{-1}^m}{x_0^k} > \alpha + \frac{x_{-1}^m}{(\alpha + \delta)^k}, \\ x_2 &= \alpha + \frac{x_0^m}{x_1^k} < \alpha + \frac{(\alpha + \delta)^m}{(\alpha + x_{-1}^m / (\alpha + \delta)^k)^k} < \alpha + \delta. \end{aligned} \quad (25)$$

Moreover we have

$$\begin{aligned} x_3 &= \alpha + \frac{x_1^m}{x_2^k} > \alpha + \frac{1}{(\alpha + \delta)^k} \left(\alpha + \frac{x_{-1}^m}{(\alpha + \delta)^k} \right)^m \\ &> \alpha + \frac{x_{-1}^m}{(\alpha + \delta)^k}, \\ x_4 &= \alpha + \frac{x_2^m}{x_3^k} < \frac{(\alpha + \delta)^m}{(\alpha + x_{-1}^m / (\alpha + \delta)^k)^k} < \alpha + \delta. \end{aligned} \quad (26)$$

Therefore, by induction we can show that

$$\begin{aligned} x_{2n+1} &> \frac{x_{-1}^m}{((\alpha + \delta)^k)^{n+1}} \\ &\quad + \alpha \left(1 + \frac{1}{(\alpha + \delta)^k} + \cdots + \frac{1}{(\alpha + \delta)^{nk}} \right), \\ \alpha &< x_{2n} < \alpha + \delta. \end{aligned} \quad (27)$$

Hence,

$$\frac{1}{(\alpha + \delta)^k} > 1, \quad (28)$$

which gives that

$$\lim_{n \rightarrow \infty} x_{2n+1} = \infty. \quad (29)$$

Thus $\{x_n\}_{n=-1}^{\infty}$ is unbounded. Now, we assume that $\alpha = 0$ and choose the initial conditions such that

$$\begin{aligned} 0 &< x_{-1} \leq 1, \\ x_0 &> \frac{1}{(1 - \varepsilon)^k}, \end{aligned} \quad (30)$$

for some $0 < \varepsilon < 1$.

So, we obtain

$$\begin{aligned} 0 &< x_1 = \frac{x_{-1}^m}{x_0^k} \leq \frac{1}{x_0^k} < 1 - \varepsilon, \\ x_2 &= \frac{x_0^m}{x_1^k} > \frac{x_0^m}{(1 - \varepsilon)^k}. \end{aligned} \quad (31)$$

Moreover we have

$$\begin{aligned} x_3 &= \frac{x_1^m}{x_2^k} < \frac{1}{x_0^{mk}} < (1 - \varepsilon)^m, \\ x_4 &= \frac{x_2^m}{x_3^k} > \frac{x_0^{2m}}{(1 - \varepsilon)^{2mk}}. \end{aligned} \quad (32)$$

By induction we have

$$\begin{aligned} x_{2n+1} &\in (0, (1 - \varepsilon)^{m^n}), \\ x_{2n} &> \frac{x_0^{m^n}}{(1 - \varepsilon)^{nm^{n-1}k}} \quad \forall n = 0, 1, \dots \end{aligned} \quad (33)$$

Therefore since $m > 1$, we get

$$\begin{aligned} \lim_{n \rightarrow \infty} x_{2n} &= \infty, \\ \lim_{n \rightarrow \infty} x_{2n+1} &= 0. \end{aligned} \quad (34)$$

For fixed $m > 1$ let $\alpha \geq 1$; we deduce that

$$\begin{aligned} \sup_n \left\{ x_n : x_{n+1} &= \alpha + \frac{x_{n-1}^m}{x_n^k} \right\} \\ &> \sup_n \left\{ x_n : x_{n+1} &= \alpha + \frac{x_{n-1}}{x_n^k} \right\} \\ &> \sup_n \left\{ x_n : x_{n+1} &= \frac{1}{2} + \frac{x_{n-1}}{x_n^k} \right\}. \end{aligned} \quad (35)$$

Since there exists unbounded solution of the difference equation $x_{n+1} = 1/2 + x_{n-1}^m/x_n^k$, $n = 0, 1, 2, \dots$, there exists unbounded solution of difference equation (1). This completes the proof. $\square$

FIGURE 3: When $m > 1$ there exists an unbounded positive solution. FIGURE 4: When $m > 1$ there exists an unbounded positive solution.

In Figure 3 we illustrate the result of Theorem 11 for $k = 3$, $m = 2$, and $\alpha = 0.5$, with the initial conditions $x_{-1} = 0.7$, $x_0 = 0.54$ and choosing $\delta = 0.2$; in this case the subsequence (x_{2n}) of the solution of (1) is bounded with $0.5 < x_{2n} < 0.7$ for all $n \geq 0$, while the subsequence (x_{2n+1}) is unbounded, so the terms of (x_{2n}) appear approximately zero in comparison with the terms of (x_{2n+1}) . Also Figure 4 explains the same result for $k = 0.3$, $m = 2$, and $\alpha = 1.5$; Gümüş and Öcalan in [5] showed that when $\alpha > 1$ and $k \rightarrow \infty$, every solution of (1) is unbounded. Figure 5 gives a counter example of this result for $k = 10^{10}$, $m = 1$, and $\alpha = 1.1$. We correct this result in Theorem 12 and Corollary 13.

Theorem 12. (a) $S_i \leq \ell_\infty$, if $1 < m < k+1$ and $\alpha \geq 2^{m/(k+1-m)}$.

(b) $S \leq \ell_\infty$, if $0 < m \leq 1$ and $\alpha > 1$.

FIGURE 5: When $\alpha > 1$ and $k \rightarrow \infty$ the positive solution converges to α , which gives a counter example of Theorem 2.1-(b) and the results that followed in [5]. Here $\alpha = 1.1$, $m = 1$, and $k = 10^{10}$.

Proof. (a) From (1), $x_n \geq \alpha > 1$ for $n \geq 0$. We get for $n \geq 0$ and $1 < m < k+1$ that

$$\begin{aligned} x_{2n+1} &= \alpha + \frac{x_{2n-1}^m}{x_{2n}^k} \leq \alpha + \frac{x_{2n-1}^m}{\alpha^k} = \alpha \\ &+ \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-3}^m}{x_{2n-2}^k} \right]^m \leq \alpha + \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-3}^m}{\alpha^k} \right]^m \leq \alpha \\ &+ 2^{m-1} \frac{\alpha^m}{\alpha^k} + 2^{m-1} \frac{x_{2n-3}^{m^2}}{(\alpha^k)^{m+1}} \leq \alpha + 2^{m-1} \frac{\alpha^m}{\alpha^k} \\ &+ 2^{m-1} \frac{1}{(\alpha^k)^{m+1}} \left[\alpha + \frac{x_{2n-5}^m}{\alpha^k} \right]^{m^2} \\ &\vdots \end{aligned} \quad (36)$$

$$\begin{aligned} &\leq \alpha + \sum_{n=1}^{\infty} \left[2^{(m^{n+1}-m)/(m-1)-n} \frac{\alpha^{m^n}}{\alpha^k ((m^n-1)/(m-1))} \right] \\ &+ 2^{(m^{n+1}-m)/(m-1)-n} \frac{x_{-1}^{m^{n+1}}}{\alpha^{k(m^{(n+1)-1}/(m-1))}} \leq \alpha \\ &+ \sum_{n=1}^{\infty} \left[2^{(m^{n+1}-m)/(m-1)-n} \frac{\alpha^{m^n}}{\alpha^k ((m^n-1)/(m-1))} \right] \\ &+ \sum_{n=1}^{\infty} 2^{(m^{n+1}-m)/(m-1)-n} \frac{x_{-1}^{m^{n+1}}}{\alpha^{k(m^{(n+1)-1}/(m-1))}}. \end{aligned}$$

Since $\alpha \geq 2^{m/(k+1-m)}$, we have the series

$$\sum_{n=1}^{\infty} 2^{(m^{n+1}-m)/(m-1)-n} \frac{\alpha^{m^n}}{\alpha^k ((m^n-1)/(m-1))} < \infty. \quad (37)$$

To study the convergence of the series with general term $a_n = 2^{(m^{n+1}-m)/(m-1)-n} x_{-1}^{m^{n+1}} / \alpha^{k(m^{(n+1)-1})/(m-1)}$, since

$$\left| \frac{a_{n+1}}{a_n} \right| = 2^{m^{n+1}-1} x_{-1}^{m^{n+1}(m-1)} \alpha^{-km^{n+1}} \quad (38)$$

for $x_{-1} \in (0, \alpha]$ and since $\alpha \geq 2^{m/(k+1-m)}$, we obtain

$$\left| \frac{a_{n+1}}{a_n} \right| \leq 2^{m^{n+1}-1} \alpha^{-m^{n+1}(k-m+1)} \leq 2^{m^{n+2}(1/m-1)-1}. \quad (39)$$

Since $m > 1$, then by using the ratio test we obtain

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 < 1; \quad (40)$$

then the series is convergent. And for $x_{-1} \in (\alpha, 2\alpha]$ and since $\alpha \geq 2^{m/(k+1-m)}$, we obtain

$$\begin{aligned} \left| \frac{a_{n+1}}{a_n} \right| &\leq 2^{m^{n+1}-1} (2\alpha)^{m^{n+2}-m^{n+1}} \alpha^{-km^{n+1}} \\ &\leq 2^{m^{n+2}-1} \alpha^{-m^{n+1}(k+1-m)} \leq \frac{1}{2}. \end{aligned} \quad (41)$$

So by using the ratio test we obtain

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2} < 1; \quad (42)$$

then the series is convergent and hence $(x_{2n+1})_{-1}^{\infty}$ is bounded. Also, we get

$$\begin{aligned} x_{2n} &= \alpha + \frac{x_{2n-2}^m}{x_{2n-1}^k} \leq \alpha + \frac{x_{2n-2}^m}{\alpha^k} = \alpha \\ &+ \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-4}^m}{x_{2n-3}^k} \right]^m \leq \alpha + \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-4}^m}{\alpha^k} \right]^m \leq \alpha \\ &+ 2^{m-1} \frac{\alpha^m}{\alpha^k} + 2^{m-1} \frac{x_{2n-4}^{m^2}}{(\alpha^k)^{m+1}} \leq \alpha + 2^{m-1} \frac{\alpha^m}{\alpha^k} \\ &+ 2^{m-1} \frac{1}{(\alpha^k)^{m+1}} \left[\alpha + \frac{x_{2n-6}^m}{\alpha^k} \right]^{m^2} \\ &\vdots \\ &\leq \alpha + \sum_{n=1}^{\infty} \left[2^{(m^{n+1}-m)/(m-1)-n} \frac{\alpha^{m^n}}{\alpha^k ((m^n-1)/(m-1))} \right] \\ &+ 2^{(m^n-m)/(m-1)-n} \frac{x_0^{m^n}}{\alpha^{k(m^{(n)-1})/(m-1)}} \leq \alpha \\ &+ \sum_{n=1}^{\infty} \left[2^{(m^{n+1}-m)/(m-1)-n} \frac{\alpha^{m^n}}{\alpha^k ((m^n-1)/(m-1))} \right] \\ &+ \sum_{n=1}^{\infty} 2^{(m^n-m)/(m-1)-n} \frac{x_0^{m^n}}{\alpha^{k(m^{(n)-1})/(m-1)}}. \end{aligned} \quad (43)$$

To study the convergence of the series with general term $b_n = 2^{(m^n-m)/(m-1)-n} x_0^{m^n} / \alpha^{k(m^{(n)-1})/(m-1)}$, since

$$\left| \frac{b_{n+1}}{b_n} \right| = 2^{m^n-1} x_0^{m^n(m-1)} \alpha^{-km^n} \quad (44)$$

for $x_0 \in (0, \alpha]$ and since $\alpha \geq 2^{m/(k+1-m)}$, we obtain

$$\left| \frac{b_{n+1}}{b_n} \right| \leq 2^{m^n-1} \alpha^{-m^n(k-m+1)} \leq 2^{m^{n+1}(1/m-1)-1}. \quad (45)$$

Since $m > 1$, then by using the ratio test we obtain

$$\lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = 0 < 1; \quad (46)$$

then the series is convergent. And for $x_0 \in (\alpha, 2\alpha]$ and since $\alpha \geq 2^{m/(k+1-m)}$, we obtain

$$\begin{aligned} \left| \frac{b_{n+1}}{b_n} \right| &\leq 2^{m^n-1} (2\alpha)^{m^{n+1}-m^n} \alpha^{-km^n} \\ &\leq 2^{m^{n+1}-1} \alpha^{-m^n(k+1-m)} \leq \frac{1}{2}. \end{aligned} \quad (47)$$

So by using the ratio test we obtain

$$\lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \frac{1}{2} < 1; \quad (48)$$

then the series is convergent and hence $(x_{2n})_0^{\infty}$ is bounded.

(b) In addition, for $n \geq 0$, $0 < m \leq 1$, and $\alpha > 1$, we have

$$\begin{aligned} x_{2n+1} &= \alpha + \frac{x_{2n-1}^m}{x_{2n}^k} \leq \alpha + \frac{x_{2n-1}^m}{\alpha^k} = \alpha \\ &+ \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-3}^m}{x_{2n-2}^k} \right]^m \leq \alpha + \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-3}^m}{\alpha^k} \right]^m \leq \alpha \\ &+ \frac{\alpha^m}{\alpha^k} + \frac{x_{2n-3}^{m^2}}{(\alpha^k)^{m+1}} \leq \alpha + \frac{\alpha^m}{\alpha^k} \\ &+ \frac{1}{(\alpha^k)^{m+1}} \left[\alpha + \frac{x_{2n-5}^m}{\alpha^k} \right]^{m^2} \\ &\vdots \\ &\leq \frac{\alpha^k}{\alpha^k-1} + \frac{x_{-1}}{\alpha^{k(n+1)}}. \end{aligned} \quad (49)$$

Again, we obtain

$$\begin{aligned}
 x_{2n} &= \alpha + \frac{x_{2n-2}^m}{x_{2n-1}^k} \leq \alpha + \frac{x_{2n-2}^m}{\alpha^k} = \alpha \\
 &+ \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-4}^m}{x_{2n-3}^k} \right]^m \leq \alpha + \frac{1}{\alpha^k} \left[\alpha + \frac{x_{2n-4}^m}{\alpha^k} \right]^m \leq \alpha \\
 &+ \frac{\alpha^m}{\alpha^k} + \frac{x_{2n-4}^{m^2}}{(\alpha^k)^{m+1}} \leq \alpha + \frac{\alpha^m}{\alpha^k} \\
 &+ \frac{1}{(\alpha^k)^{m+1}} \left[\alpha + \frac{x_{2n-6}^m}{\alpha^k} \right]^{m^2} \\
 &\vdots \\
 &\leq \frac{\alpha^k}{\alpha^k - 1} + \frac{x_0}{\alpha^{nk}} < \infty,
 \end{aligned} \tag{50}$$

from which the proof follows. $\square$

Corollary 13. For fixed m , if $\alpha > 1$ and $k \rightarrow \infty$, then $S \subseteq C_\alpha$.

Proof. Since $\alpha > 1$ and $k \rightarrow \infty$, we have $\{x_n\}_{n=-1}^\infty \in S \cap \ell_\infty$. Then, we obtain

$$\begin{aligned}
 \alpha &< \psi = \liminf x_n, \\
 \Psi &= \limsup x_n < \infty.
 \end{aligned} \tag{51}$$

Hence, from (1), we get

$$\Psi \leq \alpha + \frac{\Psi^m}{\psi^k}. \tag{52}$$

Let $k \rightarrow \infty$; then we obtain

$$\Psi \leq \alpha, \tag{53}$$

which implies $\Psi = \psi = \alpha$. So the proof is complete. $\square$

Theorem 14. The unique positive equilibrium $\bar{x}$ of (1) is globally asymptotically stable, if the following conditions are satisfied:

- (i) $0 < m \leq 1$,
- (ii) $\alpha > 1$ and $\alpha > (m+k-1)(m+k)^{(m-k)/(k+1-m)}$.

Proof. By Lemma 8, $\bar{x}$ is locally asymptotically stable. Thus, we have to prove that, for all $\{x_n\}_{n=-1}^\infty \in S$, there exists a unique positive equilibrium $\bar{x}$ with $\lim_{n \rightarrow \infty} x_n = \bar{x}$. Let $\{x_n\}_{n=-1}^\infty \in S$. By Theorem 12-(b), $\{x_n\}_{n=-1}^\infty \in \ell_\infty$. Thus, we have

$$\begin{aligned}
 \alpha &\leq \psi = \liminf x_n, \\
 \Psi &= \limsup x_n < \infty.
 \end{aligned} \tag{54}$$

Hence from (1), we get

$$\begin{aligned}
 \Psi &\leq \alpha + \frac{\Psi^m}{\psi^k}, \\
 \psi &\leq \alpha + \frac{\psi^m}{\Psi^k}.
 \end{aligned} \tag{55}$$

We maintain that $\psi = \Psi$; otherwise $\Psi > \psi$. From (55), we obtain

$$\begin{aligned}
 \Psi \psi^k &\leq \alpha \psi^k + \Psi^m, \\
 \Psi^k \psi &\geq \alpha \Psi^k + \psi^m.
 \end{aligned} \tag{56}$$

And from (56),

$$\alpha \Psi^k \psi^{k-1} + \psi^{k+m-1} \leq \alpha \psi^k \Psi^{k-1} + \Psi^{k+m-1}. \tag{57}$$

Hence, we have

$$\alpha \Psi^{k-1} \psi^{k-1} \leq \frac{\Psi^{k+m-1} - \psi^{k+m-1}}{\Psi - \psi}. \tag{58}$$

For $k+m \geq 2$ and $0 < m \leq 1$, we have $k \geq 1$. Consider $f(x) = x^{k+m-1}$ with $x \in (0, \infty)$; then there exists $c \in (\psi, \Psi)$ such that

$$\begin{aligned}
 \frac{\Psi^{k+m-1} - \psi^{k+m-1}}{\Psi - \psi} &= (k+m-1) c^{k+m-2} \\
 &\leq (k+m-1) \Psi^{k+m-2}.
 \end{aligned} \tag{59}$$

From (58) and (59), we obtain

$$\alpha \Psi^{k-1} \psi^{k-1} \leq (k+m-1) \Psi^{k+m-2}, \tag{60}$$

which is equivalent to

$$\frac{\alpha}{k+m-1} \psi^{k-1} \leq \Psi^{m-1}. \tag{61}$$

Using $\Psi \geq \psi \geq \alpha$ and $\alpha > 1$ in (61), we get

$$\alpha^{k+1-m} \leq k+m-1. \tag{62}$$

Equation (62) with the values $m = 0.6$, $k = 1.5$, and $\alpha = 2$ gives a contradiction. Thus, we find $\Psi = \psi$. Now, assume that $0 < k+m < 2$, $0 < m \leq 1$, and $k < 1$. From (58), we get

$$\alpha \Psi^{k-1} \psi^{k-1} \leq (k+m-1) \psi^{k+m-2}. \tag{63}$$

This gives

$$\alpha \Psi^{k-1} \leq (k+m-1) \psi^{m-1}. \tag{64}$$

Moreover, since $x_n \geq 1$ for all $n \geq 0$, we have

$$x_{2n+1} < \alpha + \frac{x_{2n-1}^m}{\alpha^k} < \alpha + \frac{x_{2n-1}}{\alpha^k}. \tag{65}$$

Then from (65), we get

$$\alpha \leq \psi < \Psi < \frac{\alpha^{k+1}}{\alpha^k - 1}. \tag{66}$$

By using (64) and (66), we obtain

$$\frac{\alpha^{k^2-m}}{(\alpha^k - 1)^{k-1}} \leq \frac{k+m-1}{\alpha}. \tag{67}$$

Since $\alpha > 1$ and $\alpha > (m+k-1)(m+k)^{(m-k)/(k+1-m)}$, for $\alpha = 2$, $m = 1$, and $k = 0.5$, (67) is not satisfied. This is a contradiction. Thus, we find $\Psi = \psi$, which implies that $\{x_n\}_{n=-1}^\infty$ tends to the unique positive equilibrium, from which the proof follows. $\square$

4. Periodicity of the Solutions of

$$x_{n+1} = \alpha + (x_{n-1}^m / x_n^k)$$

In this section we study the convergence of the positive solution of (1) to a prime two-periodic solution when $\alpha > 1$, $0 < m \leq 1 < k$.

Lemma 15. Suppose $\{x_n\}_{n=-1}^\infty \in S$ and $\beta > \alpha$. Then, the following conditions are contented:

- (i) $\lim_{n \rightarrow \infty} x_{2n} = \beta$, if and only if $\lim_{n \rightarrow \infty} x_{2n+1} = \beta^{m/k} / (\beta - \alpha)^{1/k}$.
- (ii) $\lim_{n \rightarrow \infty} x_{2n+1} = \beta$, if and only if $\lim_{n \rightarrow \infty} x_{2n} = \beta^{m/k} / (\beta - \alpha)^{1/k}$.

Proof. (i) Replace n by $2n + 1$ in (1); we deduce

$$x_{2n+1} = \left(\frac{x_{2n}^m}{x_{2n+2} - \alpha} \right)^{1/k} \quad \forall n \geq 0. \quad (68)$$

If $\lim_{n \rightarrow \infty} x_{2n} = \beta$, then $\lim_{n \rightarrow \infty} x_{2n+1} = \beta^{m/k} / (\beta - \alpha)^{1/k}$ and vice versa.

(ii) Replace n by $2n$ in (1); we have

$$x_{2n} = \left(\frac{x_{2n-1}^m}{x_{2n+1} - \alpha} \right)^{1/k} \quad \forall n \geq 0. \quad (69)$$

Let $\lim_{n \rightarrow \infty} x_{2n+1} = \beta$; then $\lim_{n \rightarrow \infty} x_{2n} = \beta^{m/k} / (\beta - \alpha)^{1/k}$ and vice versa. $\square$

Theorem 16. If these conditions are satisfied

$$\begin{aligned} \alpha &> 1, \\ 0 < m \leq 1 < k \end{aligned} \quad (70)$$

and there exists a sufficient small positive number ε_1 such that

$$\begin{aligned} (\alpha + \varepsilon_1)^{m/k} (\varepsilon_1)^{m/k} - (\alpha + \varepsilon_1)^{(m^2-k^2)/k} (\varepsilon_1)^{1/k} \\ < \alpha \varepsilon_1^{(m+1)/k}, \end{aligned} \quad (71)$$

$$\varepsilon_1 (\alpha + \varepsilon_1)^{k-m} < 1, \quad (72)$$

then, (1) has a periodic solution of prime period two.

Proof. Let $\{x_n\}_{n=-1}^\infty$ be a periodic solution of period two; we have

$$\begin{aligned} x_{-1} &= \alpha + \frac{x_{-1}^m}{x_0^k} = x_1, \\ x_0 &= \alpha + \frac{x_0^m}{x_{-1}^k} = x_2; \end{aligned} \quad (73)$$

then (x_{-1}, x_0) is a solution of the system

$$\begin{aligned} x &= \alpha + \frac{x^m}{y^k}, \\ y &= \alpha + \frac{y^m}{x^k}. \end{aligned} \quad (74)$$

Let $\{x_n\}_{n=-1}^\infty \in S$. It is obvious that if (73) holds, then $\{x_n\}_{n=-1}^\infty$ is a periodic solution of period two. System (74) is correspondent to

$$\begin{aligned} y - \alpha - \frac{y^m}{x^k} &= 0, \\ y &= \left(\frac{x^m}{x - \alpha} \right)^{1/k}, \end{aligned} \quad (75)$$

so we get the equation

$$\begin{aligned} H(x) &= \frac{1}{(x - \alpha)^{1/k}} \left(x^{m/k} - (x - \alpha)^{(1-m)/k} x^{(m^2-k^2)/k} \right) \\ &\quad - \alpha. \end{aligned} \quad (76)$$

Thus for $0 < m \leq 1$, we have

$$\lim_{x \rightarrow \alpha^+} H(x) = \infty. \quad (77)$$

Moreover, from (71) and (76) we have

$$H(\alpha + \varepsilon_1) < 0. \quad (78)$$

Therefore, the equation $H(x) = 0$ has a root $\bar{x} = \alpha + \varepsilon_0$, where $0 < \varepsilon_0 < \varepsilon_1$, in the interval $(\alpha, \alpha + \varepsilon_1)$. So, we have

$$\bar{y} = \left(\frac{\bar{x}^m}{\bar{x} - \alpha} \right)^{1/k}. \quad (79)$$

Let us consider the function

$$T(\varepsilon) = (\alpha + \varepsilon)^{m-k} - \varepsilon. \quad (80)$$

Since $0 < m < k$, we get

$$\begin{aligned} T'(\varepsilon) &= (m - k) (\alpha + \varepsilon)^{m-k-1} - 1 < 0, \\ T(\varepsilon_0) &> T(\varepsilon_1). \end{aligned} \quad (81)$$

From inequality (72), we have $T(\varepsilon_1) > 0$, and hence $T(\varepsilon_0) > 0$, which implies that

$$\bar{x} = \alpha + \varepsilon_0 < \left(\frac{(\alpha + \varepsilon_0)^m}{\varepsilon_0} \right)^{1/k} = \bar{y}. \quad (82)$$

By taking $x_{-1} = \bar{x}$, $x_0 = \bar{y}$, then the solution $\{x_n\}_{n=-1}^\infty$ with initial values x_{-1}, x_0 is a prime 2-periodic solution. This completes the proof. $\square$

Lemma 17. If $\{x_n\}_{n=-1}^\infty \in S$ which consists of at least two semicycles, then $\{x_n\}_{n=-1}^\infty$ is oscillatory and, except possibly for the first semicycle, every semicycle is of length 1.

Proof. The proof is directly obtained from (1). $\square$

Theorem 18. For $\{x_n\} \in S$, the sequences $\{x_{2n}\}$ and $\{x_{2n+1}\}$ are ultimately monotone.

FIGURE 6: Existence of two-periodic positive solution, $\varepsilon_0 = 2.5281$, $x_{-1} = \bar{x} = 3.6201$, and $x_0 = \bar{y} = 1.1903$.

Proof. We have

$$\begin{aligned}
 x_{2n+1} - x_{2n-1} &= \frac{x_{2n-2}^k (x_{2n-1}^m - x_{2n-3}^m) + x_{2n-3}^m (x_{2n-2}^k - x_{2n}^k)}{(x_{2n+1} x_{2n-2})^k}, \\
 x_{2n+2} - x_{2n} &= \frac{x_{2n-1}^k (x_{2n}^m - x_{2n-2}^m) + x_{2n-2}^m (x_{2n-1}^k - x_{2n+1}^k)}{(x_{2n+1} x_{2n-1})^k}.
 \end{aligned} \quad (83)$$

If $x_1 \geq x_{-1}$ and $x_2 \leq x_0$, we obtain from (83) $x_3 \geq x_1$ and therefore $x_4 \leq x_2$. By induction we find that $\{x_{2n}\}_{n=0}^{\infty}$ is decreasing and $\{x_{2n+1}\}_{n=-1}^{\infty}$ is increasing. Correspondingly if $x_1 \leq x_{-1}$ and $x_2 \geq x_0$, using induction we obtain from (83) that $\{x_{2n}\}_{n=0}^{\infty}$ is increasing and $\{x_{2n+1}\}_{n=-1}^{\infty}$ is decreasing. If $x_1 \geq x_{-1}$ and $x_2 \geq x_0$ and $x_1 \geq x_3$, we have from (83) that $\{x_{2n}\}_{n=0}^{\infty}$ is increasing and $\{x_{2n+1}\}_{n=0}^{\infty}$ is decreasing. Hence, we can assume that $x_3 \geq x_1 \geq x_{-1}$ and $x_0 \leq x_2$. If further $x_2 \geq x_4$, then $\{x_{2n}\}_{n=1}^{\infty}$ is decreasing and $\{x_{2n+1}\}_{n=-1}^{\infty}$ is increasing. So we may assume that $x_3 \geq x_1 \geq x_{-1}$ and $x_0 \leq x_2 \leq x_4$. By induction we get the result in this case. The cases $x_1 \leq x_{-1}$ and $x_2 \leq x_0$ can be treated in the same way. $\square$

Theorem 19. If (70), (71), and (72) are conformed, then every positive solution of (1) converges to a prime two-periodic solution.

Proof. The sequences $\{x_{2n}\}$ and $\{x_{2n+1}\}$ are eventually monotone and bounded by Theorems 18 and 12 for every positive solution of (1). So, the sequences $\{x_{2n}\}$ and $\{x_{2n+1}\}$ are convergent. Using Lemmas 15 and 17 the proof follows. $\square$

Figure 6 illustrates the result of Theorem 19 for $\alpha = 1.1$, $m = 0.99$, and $k = 2$.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

The author wants to thank the reviewers for much encouragement, support, productive feedback, cautious perusing, and a helpful remark which improved the presentation and the comprehensibility of the paper.

References

- [1] V. L. Kocic and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, vol. 256 of *Mathematics and its Applications*, Kluwer Academic, Dodrecht, The Netherlands, 1993.
- [2] M. R. Kulenović and G. Ladas, *Dynamics of Second Order Rational Difference Equations*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2002.
- [3] A. M. Amleh, E. A. Grove, G. Ladas, and D. A. Georgiou, "On the recursive sequence $x_{n+1} = \alpha + (x_{n-1}/x_n)$," *Journal of Mathematical Analysis and Applications*, vol. 233, no. 2, pp. 790–798, 1999.
- [4] A. E. Hamza and A. Morsy, "On the recursive sequence $x_{n+1} = \alpha + (x_{n-1}/x_n^k)$," *Applied Mathematics Letters*, vol. 22, no. 1, pp. 91–95, 2009.
- [5] M. Gümüş and Ö. Öcalan, "Some notes on the difference equation $x_{n+1} = \alpha + (x_{n-1}^m/x_n^k)$," *Discrete Dynamics in Nature and Society*, vol. 2012, Article ID 258502, 12 pages, 2012.
- [6] K. S. Berenhaut and S. Stević, "The behaviour of the positive solutions of the difference equation $x_{n+1} = \alpha + (x_{n-1}/x_n)$," *Journal of Difference Equations and Applications*, vol. 12, no. 9, pp. 909–918, 2006.
- [7] S. N. Elaydi, *An Introduction to Difference Equations*, Springer, New York, NY, USA, 1996.
- [8] S. Stević, "On the recursive sequence $x_{n+1} = \alpha_n + (x_{n-1}/x_n)$. II," *Dynamics of Continuous, Discrete and Impulsive Systems Series A*, vol. 10, no. 6, pp. 911–916, 2003.
- [9] S. Stević, "On the recursive sequence $x_{n+1} = \alpha_n + (x_{n-1}/x_n)$," *International Journal of Mathematical Sciences*, vol. 2, no. 2, pp. 237–243, 2003.
- [10] S. Stević, "On the recursive sequence $x_{n+1} = \alpha + (x_n^p/x_{n-1}^p)$," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 34517, 9 pages, 2007.
- [11] S. Stević, "On a class of higher-order difference equations," *Chaos, Solitons and Fractals*, vol. 42, no. 1, pp. 138–145, 2009.
- [12] G. J. O. Jameson, "Some inequalities for $(a+b)^p$ and $(a+b)^p + (a-b)^p$," *The Mathematical Gazette*, vol. 98, pp. 96–103, 2014.
- [13] S. Stević, "On the recursive sequence $x_{n+1} = \alpha + (x_{n-1}^p/x_n^p)$," *Journal of Applied Mathematics & Computing*, vol. 18, no. 1-2, pp. 229–234, 2005.