Research Article

Approximative Compactness and Radon-Nikodym Property in w^* Nearly Dentable Banach Spaces and Applications

Shaoqiang Shang¹ and Yunan Cui²

¹Department of Mathematics, Northeast Forestry University, Harbin 150040, China

²Department of Mathematics, Harbin University of Science and Technology, Harbin 150080, China

Correspondence should be addressed to Shaoqiang Shang; sqshang@163.com

Received 3 March 2015; Revised 8 April 2015; Accepted 18 April 2015

Academic Editor: Ismat Beg

Copyright © 2015 S. Shang and Y. Cui. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Authors define w^* nearly dentable Banach space. Authors study Radon-Nikodym property, approximative compactness and continuity metric projector operator in w^* nearly dentable space. Moreover, authors obtain some examples of w^* nearly dentable space in Orlicz function spaces. Finally, by the method of geometry of Banach spaces, authors give important applications of w^* nearly dentability in generalized inverse theory of Banach space.

1. Introduction

Let $(X, \|\cdot\|)$ be a real Banach space. $S(X)$ and $B(X)$ denote the unit sphere and the unit ball, respectively. By X^* we denote the dual space of X . Let N, R , and R^+ denote the set of natural numbers, reals, and nonnegative reals, respectively. $A(x) = \{x^* : x^* \in S(X^*), x^*(x) = 1 = \|x\|\}$ and $B_r(x) = \{y \in X : \|y - x\| < r\}$. By $x_n \xrightarrow{w} x$ denote that $\{x_n\}_{n=1}^\infty$ is weakly convergent to x . $x_n \xrightarrow{w^*} x^*$ denote that $\{x_n^*\}_{n=1}^\infty$ is weakly* convergent to x^* . The weak topology of X is denoted by $\sigma(X^*, X)$ or (X, w) . The weak* topology of X^* is denoted by $\sigma(X, X^*)$ or (X^*, w^*) . $C(\overline{C^w}, \overline{C^{w^*}})$ denote closed hull of C (weak closed hull, closed hull weak* closed hull). $\text{dist}(x, C)$ denote distance of x and C .

The study of geometric and topological properties of unit balls of Banach spaces plays a central rule in the geometry of Banach spaces. Almost all properties of Banach spaces, such as convexity, smoothness, nonsquareness, and the Radon-Nikodym property, can be viewed as properties of the unit ball. We should mention here that there are many topics studying behavior of ball collections. For example, the Mazur intersection property, the packing sphere problem of unit balls, the measure of noncompactness with respect to topological degree, and the ball topology have also brought

great attention of many mathematicians. In 2006, Cheng defined ball-covering property of Banach space (see [1]). In [2], Chen and Cheng proved analytical characterizations of Mazur intersection property.

Starting with a different viewpoint, a notion of nearly dentable space is introduced by Shang et al. in [3].

Definition 1. Banach space X is called nearly dentable if, for any $f \in S(X^*)$ and any open set $U_{A_f} \supset A_f$, we have $A_f \neq \emptyset$ and $A_f \cap \overline{C\sigma}(B(X) \setminus U_{A_f}) = \emptyset$, where $A_f = \{x : f(x) = \|x\| = \|f\| = 1\}$.

By James theorem, it is easy to see that nearly dentable space is reflexive.

Definition 2. A nonempty subset C of X is said to be approximatively compact if for any $\{y_n\}_{n=1}^\infty \subset C$ and any $x \in X$ satisfying $\|x - y_n\| \rightarrow \inf_{y \in C} \|x - y\|$ ($n \rightarrow \infty$), then $\{y_n\}_{n=1}^\infty$ has a subsequence converging to an element in C . X is called approximatively compact if every nonempty closed convex subset of X is approximatively compact.

Let us present the history of the approximative compactness and related notions. This notion has been introduced by Jefimow and Stechkin in [4] as a property of Banach spaces,

which guarantees the existence of the best approximation element in a nonempty closed convex set C for any $x \in X$. In [3] it was established (without proof) that the approximative compactness of C in a rotund Banach space X gives also the continuity of the projector operator P_C . Fully rotund Banach spaces were defined by Fan and Glicksberg in [5]. In [6] it has been proved that fully rotund Banach spaces are approximately compact. In [7] it has been proved that approximative compactness of a Banach space X means that X is reflexive and it has the Kadec-Klee property (called also the property H). In 2007, Chen et al. (see [8]) proved that a nonempty closed convex C of a midpoint locally uniformly rotund space is approximately compact if and only if C is a Chebyshev set and the metric projection operator P_C is continuous. In 1972, Oshman (see [9]) proved that the metric projection operator P_C is upper semicontinuous if C is approximatively compact subset of a Banach space X .

Let D be a nonempty open convex subset of X and f a real-valued continuous convex function on D . f is said to be Gateaux differentiable at the point x in D if the limit

$$f'(x)(y) = \lim_{t \rightarrow 0} \frac{f(x+ty) - f(x)}{t} \quad (*)$$

exists for all $y \in X$. When this is the case, the limit is a continuous linear function of y , denoted by $f'(x)$. If the difference quotient in $(*)$ converges to $f'(x)(y)$ uniformly for y in the unit ball, then f is said to be Fréchet differentiable at x . X is called a weak Asplund space (Asplund space) or said to have the weak Asplund property if, for every f and D as above, f is "generically" Gateaux (Frechet) differentiable; that is, there exists a dense G_δ subset G of D such that f is Gateaux (Frechet) differentiable at each point of G .

The paper is organized as follows. In Section 1, authors define w^* nearly dentable Banach space and some necessary definitions and notations are collected. In Section 2, authors prove that if X^* is a w^* nearly dentable space, then X is an Asplund space if and only if $A_E(x) = \{x^* \in S(E^*) : x^*(x) = 1 = \|x\|\}$ is a separable subset of E^* , where E is a separable closed subspace of X . In Section 3, authors prove that the following statements are equivalent: (1) every weak* hyperplane of X^* is approximatively compact; (2) if C is a weak* open convex set, then $\overline{C^{w^*}}$ is approximatively compact; (3) X^* is w^* nearly dentable space and $A(x)$ is compact for any $x \in S(X)$. Moreover, authors also prove that if X^* is w^* nearly dentable space, then (1) for any weak* open convex set C metric projector operator $P_{\overline{C^{w^*}}}$ is upper semicontinuous; (2) For any weak* hyperplane D^* metric projector operator P_{D^*} is upper semicontinuous. In Section 4, authors prove that there exists a Banach space X such that X^* is not w^* nearly dentable Banach space. Moreover, authors also prove that there exists a Banach space X such that X^* is w^* nearly dentable Banach space and not nearly dentable Banach space. The main results in this paper have important applications to the ill-posed equation theory and the generalized inverse theory. In Section 5, authors give an important application of w^* nearly dentability in generalized inverse theory of Banach space.

Definition 3. X^* is called w^* nearly dentable space, if, for any $x \in S(X)$ and open set $U_{A(x)} \supset A(x)$, we have $A(x) \cap \overline{co}^{w^*}(B(X^*) \setminus U_{A(x)}) = \emptyset$.

Let us recall some definitions and some lemmas which will be used in the further part of the paper.

Definition 4 (see [8]). A subset C is said to be proximal if $P_C(x) \neq \emptyset$ for all $x \in X$. C is said to be semi-Chebyshev if $P_C(x)$ is at most a singleton for all $x \in X$. C is said to be Chebyshev if it is proximal and semi-Chebyshev.

Definition 5. A Banach space X is said to have the Radon-Nikodym property whenever if (T, Σ, μ) is a nonatomic measure space and ν is a vector measure on Σ with values in X which is absolutely continuous with respect to μ and has bounded variation; then there exists $f \in L_1(X)$ such that, for any $A \in \Sigma$,

$$\nu(A) = \int_A f(t) dt. \quad (1)$$

It is well known that X is reflexive if and only if each closed convex subset of X is proximal. It is well known that X is Asplund space if and only if X^* has the Radon-Nikodym property.

Definition 6 (see [10]). Set-valued mapping $F : X \rightarrow Y$ is called upper semicontinuous at x_0 , if, for each norm open set W with $F(x_0) \subset W$, there exists a norm neighborhood U of x_0 such that $F(x) \subset W$ for all x in U . F is called lower continuous at x_0 , if, for any $y \in F(x_0)$ and any $\{x_n\}_{n=1}^\infty$ in X with $x_n \rightarrow x_0$, there exists $y_n \in F(x_n)$ such that $y_n \rightarrow y$ as $n \rightarrow \infty$. F is called continuous at x_0 , if F is upper semicontinuous and is lower continuous at x_0 .

Definition 7. $H^* \subset X^*$ is called weak* hyperplane, if there exist $x \in S(X)$ and $\lambda \in R$ such that $H^* = \{x^* \in X^* : x^*(x) = \lambda\}$.

Lemma 8 (see [11]). Suppose that X is a Banach space. Then the following statements are equivalent:

- (1) X is Asplund space.
- (2) If E is a separable closed subspace of X , then E^* is a separable space.
- (3) X^* has the Radon-Nikodym property.

Lemma 9 (see [12]). Let $B \subset S(X^*)$ and $B \cap A(x)$ be not an empty set for all $x \in S(X)$. Assume that $B \subset \overline{co}(F + \alpha B(X^*))$ for some countable set $F \subset X^*$ and some $0 < \alpha < 1$. Then X^* is the closed linear span of F .

2. w^* Nearly Dentability and Radon-Nikodym Property in Banach Space

Theorem 10. Suppose that X^* is a w^* nearly dentable space. Then the following statements are equivalent:

- (1) X is a Asplund space.

- (2) X^* has the Radon-Nikodym property.
- (3) If E is a separable closed subspace of X , then E^* is a separable space.
- (4) If E is a separable closed subspace of X , then $A_E(x) = \{x^* \in S(E^*) : x^*(x) = 1 = \|x\|\}$ is a separable subset of E^* .

Proof. (4) $\Rightarrow$ (1) Let E be a separable closed subspace of X and $\{x_n\}_{n=1}^\infty$ be a dense subset of $S(E)$. Then, for any $x \in S(E)$, we have

$$\inf \{\text{dist}(A_E(x_n), A_E(x)) : n \in N\} = 0. \quad (2)$$

In fact, suppose that $\inf \{\text{dist}(A_E(x_n), A_E(x)) : n \in N\} > 0$. Then, for any $\varepsilon > 0$, there exist $x^* \in A(x)$ and $x_i \in \{x_n\}_{n=1}^\infty$ such that $\inf \{\text{dist}(A(x_n), A(x)) : n \in N\} + \varepsilon > \|x_i^* - x^*\|$, where $x_i^* \in A(x_i)$. This implies that

$$\begin{aligned} \inf \{\text{dist}(A(x_n), A(x)) : n \in N\} + \varepsilon \\ > \|x_i^* - x^*\| \geq \|(x_i^* - x^*)|_E\| \geq \|x_i^*|_E - x^*|_E\| \\ \geq \inf \{\text{dist}(A_E(x_n), A_E(x)) : n \in N\}. \end{aligned} \quad (3)$$

Since ε is arbitrary, we obtain that

$$\begin{aligned} \inf \{\text{dist}(A(x_n), A(x)) : n \in N\} \\ \geq \inf \{\text{dist}(A_E(x_n), A_E(x)) : n \in N\} > 0. \end{aligned} \quad (4)$$

This implies that there exists $r > 0$ such that $\inf \{\text{dist}(A(x_n), A(x)) : n \in N\} \geq 2r$. Hence we define an open set

$$U_{A(x)} = \bigcup_{x^* \in A(x)} \{y^* \in X^* : \|y^* - x^*\| < r\}. \quad (5)$$

Since X^* is a w^* nearly dentable space, we have $A(x) \cap \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)}) = \emptyset$. Since $\overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)})$ is a weak* compact set, by the separation theorem of locally convex space, there exists $x \in S(X)$ such that

$$\begin{aligned} \inf \{x(x^*) : x^* \in A(x)\} \\ > \sup \{x(x^*) : x^* \in \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)})\}. \end{aligned} \quad (6)$$

Since the sequence $\{x_n\}_{n=1}^\infty$ is a dense subset of $S(E)$, we may assume without loss of generality that $x_n \rightarrow x$ as $n \rightarrow \infty$. Pick $x_n^* \in A(x_n)$. Then

$$\begin{aligned} |x_n^*(x) - 1| &= |x_n^*(x) - x_n^*(x_n)| \leq \|x_n^*\| \cdot \|x_n - x\| \\ &\rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (7)$$

Moreover, we may assume without loss of generality that if $m \neq n$, then $x_n^* \neq x_m^*$. Since $B(X^*)$ is weak* compact, there exists $x_0^* \in B(X^*)$ such that x_0^* is a weak* accumulation point of $\{x_n^*\}_{n=1}^\infty$. This implies that $x_0^*(x) = 1$. Then $x_0^* \in A(x)$. Therefore, by formula (6), we obtain that

$$x(x_0^*) > \sup \{x(x^*) : x^* \in \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)})\}. \quad (8)$$

Since $A(x_n) \subset \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)})$ and $x_0^* \in A(x)$, by formula (6), we have $x(x_0^*) > \sup \{x(x^*) : x^* \in \{x_n^*\}\}$,

which contradicts $x_\alpha^* \xrightarrow{w^*} x_0^*$. Hence formula (2) is true. Since $A_E(x)$ is a separable subset of E^* , we obtain that there exists a countable dense subset D_n of $A_E(x_n)$ for each $n \in N$. Then $D = \bigcup_{n=1}^\infty D_n$ is a countable set. Pick $\alpha \in (0, 1)$. Therefore, by formula (2), we obtain that there exist $\alpha(x) \in A_E(x)$ and $n(x) \in N$ such that $\text{dist}(A_E(x_{n(x)}), \alpha(x)) < \alpha$ for any $x \in S(E)$. Hence, for every $x \in S(E)$, we obtain that

$$B = \{\alpha(x) : x \in S(E)\} \subset S(E^*), \quad (9)$$

$$B \cap A_E(x) \neq \emptyset$$

for any $x \in S(E)$. Since $\text{dist}(A_E(x_{n(x)}), \alpha(x)) < \alpha$ and $D_{n(x)}$ is a countable dense subset of $A_E(x_{n(x)})$, we obtain that

$$\begin{aligned} \text{dist}(\alpha(x), D) &\leq \text{dist}(\alpha(x), D_{n(x)}) \\ &= \text{dist}(\alpha(x), A_E(x_{n(x)})) < \alpha \end{aligned} \quad (10)$$

for any $x \in S(E)$. This implies that

$$\alpha(x) \in D + \alpha B(E^*) \subset \overline{\text{co}}(D + \alpha B(E^*)) \quad (11)$$

for any $x \in S(E)$. Noticing that $B = \{\alpha(x) : x \in S(E)\}$, we have

$$B \subset D + \alpha B(E^*) \subset \overline{\text{co}}(D + \alpha B(E^*)). \quad (12)$$

By Lemma 9, we obtain that $E^* = \overline{\text{span}}(D)$, and so E^* is separable.

This means that if E is a separable closed subspace of X , then E^* is a separable space. By Lemma 15, we obtain that X is Asplund space and X^* has the Radon-Nikodym property.

(1) $\Rightarrow$ (4). Since X^* has the Radon-Nikodym property, we obtain that if E is a separable closed subspace of X , then E^* is a separable space. Hence $A_E(x) = \{x^* \in S(E^*) : x^*(x) = 1 = \|x\|\}$ is a separable subset of E^* .

Therefore, by Lemma 8, we obtain that Theorem 10 is true, which completes the proof. $\square$

3. w^* Nearly Dentability and Approximative Compactness and Continuity of Metric Projector Operator in Banach Spaces

Theorem 11. Suppose that X is a Banach space. Then the following statements are equivalent:

- (1) Every weak* hyperplane of X^* is approximatively compact.
- (2) If $C \subset X^*$ is a weak* open convex set, then $\overline{C}^{w^*}$ is approximatively compact.
- (3) X^* is w^* nearly dentable space and $A(x)$ is compact for any $x \in S(X)$.

Proof. (3) $\Rightarrow$ (2). (a) Let $x_n^*(x_0) \rightarrow 1$ as $n \rightarrow \infty$, where $x_n^* \in S(X^*)$ and $x_0 \in S(X)$. Since $B(X^*)$ is weak* compact, there exists $x_0^* \in B(X^*)$ such that x_0^* is a weak* accumulation

point of $\{x_n^*\}_{n=1}^\infty$. Hence there exists a net $\{x_\alpha^*\} \subset \{x_n^*\}_{n=1}^\infty$ such that $x_\alpha^* \xrightarrow{w^*} x_0^*$ and $x_0^* \in A(x_0)$. We next will prove that the sequence $\{x_n^*\}_{n=1}^\infty$ is relatively compact. The proof requires considering a few cases separately.

Case I ($\{x_n^*\}_{n=1}^\infty \cap A(x_0) = \emptyset$). Suppose that $\{x_n^*\}_{n=1}^\infty$ is not relatively compact. Then there exists a subsequence $\{x_n^*\}_{n=1}^\infty$ denoted again by $\{x_n^*\}_{n=1}^\infty$ such that $\{x_n^*\}_{n=1}^\infty$ does not contain convergent subsequence. This implies that every point of $A(x_0)$ is not accumulation point of $\{x_n^*\}_{n=1}^\infty$. It is easy to see that for any $x^* \in A(x_0)$, there exists $\varepsilon_{x^*} > 0$ such that $\{x_n^*\}_{n=1}^\infty \cap \{y^* : \|y^* - x^*\| < \varepsilon_{x^*}\} = \emptyset$. Put

$$U_{A(x_0)} = \bigcup_{x^* \in A(x_0)} \{y^* : \|y^* - x^*\| < \varepsilon_{x^*}\}. \quad (13)$$

Then, it is easy to see that $U_{A(x_0)} \cap \{x_n^*\}_{n=1}^\infty = \emptyset$. Moreover, since $B(X^*)$ is a weak* compact convex set, we obtain that $\overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x_0)})$ is a weak* compact convex set. Since X^* is a w^* nearly dentable space, we obtain that $A(x_0) \cap \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x_0)}) = \emptyset$. By separation theorem, there exist $y \in X$ and $r > 0$ such that

$$\begin{aligned} & \inf \{x^*(y) : x^* \in A(x_0)\} - r \\ & \geq \sup \{x^*(y) : x^* \in \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x_0)})\}. \end{aligned} \quad (14)$$

From the previous proof, there exists $\{x_\alpha^*\} \subset \{x_n^*\}_{n=1}^\infty$ such that $x_\alpha^* \xrightarrow{w^*} x_0^* \in A(x_0)$. By $\{x_n^*\}_{n=1}^\infty \cap U_{A(x_0)} = \emptyset$, we have

$$\begin{aligned} \{x_\alpha^*\} & \subset \{x_n^*\}_{n=1}^\infty \subset B(X^*) \setminus U_{A(x_0)} \\ & \subset \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x_0)}). \end{aligned} \quad (15)$$

Therefore, by formulas (14) and (15), we obtain that $x_0^*(y) - r \geq \sup \{x^*(y) : x^* \in \{x_\alpha^*\}\}$. This implies that $x_\alpha^* \xrightarrow{w^*} x_0^* \in A(x_0)$ is impossible, a contradiction. Hence $\{x_n^*\}_{n=1}^\infty$ is relatively compact.

Case II ($\{x_n^*\}_{n=1}^\infty \cap A(x_0) \neq \emptyset$). From the previous proof, $\{x_n^*\}_{n=1}^\infty \setminus A(x_0)$ is relatively compact. Since $A(x_0)$ is compact, we obtain that $\{x_n^*\}_{n=1}^\infty \cap A(x_0)$ is relatively compact. Noticing that $\{x_n^*\}_{n=1}^\infty = (\{x_n^*\}_{n=1}^\infty \cap A(x_0)) \cup (\{x_n^*\}_{n=1}^\infty \setminus A(x_0))$, we obtain that $\{x_n^*\}_{n=1}^\infty$ is relatively compact.

Let C be a weak* open convex set and $H^* = \overline{C}^{w^*}$. Then, for any $x^* \notin H^*$, we may assume without loss of generality that $x^* = 0$. We next will prove that if $\|0 - y_n^*\| \rightarrow \text{dist}(x^*, H^*)$ as $n \rightarrow \infty$, then the sequence $\{y_n^*\}_{n=1}^\infty$ has a subsequence converging to an element in H^* , where $\{y_n^*\}_{n=1}^\infty \subset H^*$. Since H^* is a weak* closed set, by proof of theorem 2.1 of [13], we obtain that H^* is a proximal set and $\text{dist}(0, H^*) = r > 0$. Pick $y_0^* \in P_{H^*}(0)$. Then $\overline{B_r(0)} \cap C = \emptyset$. In fact, suppose that $\overline{B_r(0)} \cap C \neq \emptyset$. Then there exist $z^* \in \overline{B_r(0)} \cap C$ and $\delta > 0$ such that $B_{2\delta}(z^*) \subset C$ and $\|z^*\| > 4\delta$. Since

$$\left\| \frac{\|z^*\| - \delta}{\|z^*\|} z^* - z^* \right\| = \left(1 - \frac{\|z^*\| - \delta}{\|z^*\|} \right) \|z^*\| = \delta, \quad (16)$$

we obtain that

$$\frac{\|z^*\| - \delta}{\|z^*\|} z^* \in C \subset H^*, \quad (17)$$

$$\text{dist}(0, H^*) \leq \left\| 0 - \frac{\|z^*\| - \delta}{\|z^*\|} z^* \right\| < \|z^*\| \leq r,$$

a contradiction. Hence

$$r = \text{dist}(0, P_{H^*}(0)) = \|y_0^*\|,$$

$$\overline{B_r(0)} \cap C = \emptyset, \quad (18)$$

$$\overline{B_r(0)} \cap H^* = P_{H^*}(0).$$

Noticing that $(X^*, w^*)^* = X$ and separation theorem of locally convex space, there exists $x \in S(X)$ such that

$$\begin{aligned} & \sup \{x(y^*) : y^* \in C\} \\ & \leq \inf \{x(y^*) : y^* \in \overline{B_r(0)} = \overline{B_r(0)}^{w^*}\} \\ & = -\|x\| \cdot \|y_0^*\|. \end{aligned} \quad (19)$$

This implies that

$$\begin{aligned} & \sup \{x(y^*) : y^* \in H^*\} \\ & \leq \inf \{x(y^*) : y^* \in B_r(0)\} = -\|x\| \cdot \|y_0^*\|. \end{aligned} \quad (20)$$

Since $y_0^* \in P_{H^*}(0) \subset H^*$, we obtain that

$$\begin{aligned} -\|x\| \cdot \|y_0^*\| & \leq x(y_0^*) \leq \sup \{x(y^*) : y^* \in H^*\} \\ & \leq \inf \{x(y^*) : y^* \in B_r(0)\} \\ & = -\|x\| \cdot \|y_0^*\|, \\ -\|x\| \cdot \|y_0^*\| & = x(y_0^*) = \sup \{x(y^*) : y^* \in H^*\}. \end{aligned} \quad (21)$$

This means that the inequality $x(y_0^*) \geq x(y_n^*)$ holds. Therefore

$$\begin{aligned} \|0 - y_0^*\| & = x(0 - y_0^*) \leq x(0 - y_n^*) \leq \|0 - y_n^*\| \\ & \rightarrow \text{dist}(0, H^*) = \|0 - y_0^*\| = \|y_0^*\|. \end{aligned} \quad (22)$$

Hence $\|y_n^*\| \rightarrow \|y_0^*\|$ and $x(0 - y_n^*) \rightarrow \|0 - y_0^*\|$ as $n \rightarrow \infty$. Furthermore, we have

$$\begin{aligned} x\left(-\frac{y_n^*}{\|y_n^*\|} + \frac{y_n^*}{\|y_0^*\|}\right) & = \left(\frac{1}{\|y_0^*\|} - \frac{1}{\|y_n^*\|}\right) \cdot x(y_n^*) \\ & \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned} \quad (23)$$

which shows that $x(-y_n^*/\|y_n^*\|) \rightarrow 1$ as $n \rightarrow \infty$. From the previous proof, we obtain that the sequence $\{-y_n^*/\|y_n^*\|\}_{n=1}^\infty$ is relatively compact. Noticing that $\|y_n^*\| \rightarrow \|y_0^*\|$ as $n \rightarrow \infty$, we obtain that the sequence $\{y_n^*\}_{n=1}^\infty$ is relatively compact. This implies that the sequence $\{y_n^*\}_{n=1}^\infty$ has a subsequence converging to an element in H^* . Hence we obtain that H^* is approximatively compact.

(2) $\Rightarrow$ (1). Let $D^* = \{x^* \in X^* : x^*(x_0) = 1\}$ be a weak* hyperplane. Then it is easy to see that $D_1^* = \{x^* \in X^* : x^*(x_0) \geq 1\}$ and $D_2^* = \{x^* \in X^* : x^*(x_0) \leq 1\}$ are weak* closed convex sets. Moreover, it is easy to see that if $y^*(x_0) > 1$, then $P_{D^*}(y^*) = P_{D_1^*}(y^*)$ and if $y^*(x_0) < 1$, then $P_{D^*}(y^*) = P_{D_2^*}(y^*)$. Since D_1^* and D_2^* are approximatively compact, we obtain that D^* is approximatively compact.

(1) $\Rightarrow$ (3). For any $x \in S(X)$, we define a weak* hyperplane $H^* = \{x^* : x^*(x) = \|x^*\| = 1\}$. Let $\{y_n^*\}_{n=1}^\infty \subset A(x)$. Then $\|0 - y_n^*\| = \text{dist}(0, H^*) = 1$. Hence $\{y_n^*\}_{n=1}^\infty$ has a subsequence converging to $y_0^* \in H^*$. Moreover, it is easy to see that $A(x)$ is a closed set. Hence $A(x)$ is compact. We claim that

$$\begin{aligned} r &= r(x, U_{A(x)}) \\ &= \inf \{\text{dist}(x^*, A(x)) : x^* \in B(X^*) \setminus U_{A(x)}\} > 0 \end{aligned} \quad (24)$$

for any open set $U_{A(x)} \supset A(x)$. In fact, suppose that $r = 0$. Then there exists $\{x_n^*\}_{n=1}^\infty \subset B(X^*) \setminus U_{A(x)}$ such that $\text{dist}(x_n^*, A(x)) \rightarrow 0$ as $n \rightarrow \infty$. Since $A(x)$ is compact, there exists $\{y_n^*\}_{n=1}^\infty \subset A(x)$ such that $\|x_n^* - y_n^*\| = \text{dist}(x_n^*, A(x)) \rightarrow 0$. Since $A(x)$ is compact, without loss of generality, we may assume that $\{y_n^*\}_{n=1}^\infty$ is a Cauchy sequence. Let $y_n^* \rightarrow y_0^*$ as $n \rightarrow \infty$. Then $y_0^* \in A(x)$. Hence there exists $c > 0$ such that $B_c(y_0^*) \subset U_{A(x)}$. Moreover, we have

$$0 \leq \|x_n^* - y_0^*\| \leq \|x_n^* - y_n^*\| + \|y_n^* - y_0^*\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (25)$$

Therefore, $x_n^* \in B_c(y_0^*) \subset U_{A(x)}$, when n is large enough, which contradicts $\{x_n^*\}_{n=1}^\infty \subset B(X^*) \setminus U_{A(x)}$. Hence, for any open set $U_{A(x)} \supset A(x)$, we have $r = r(x, U_{A(x)}) > 0$. We next will prove that there exists $m > 0$ such that $1 > y^*(x) + m$ whenever $y^* \in B(X^*) \setminus U_{A(x)}$. Otherwise, there exists $\{h_n^*\}_{n=1}^\infty \subset B(X^*) \setminus U_{A(x)}$ such that $h_n^*(x) \rightarrow 1$. It is easy to see that $\|h_n^*\| \rightarrow 1$. Let $z_n^* = h_n^*/\|h_n^*\|$. Then it is easy to see that $z_n^*(x) \rightarrow 1$. Hence we define a weak* hyperplane $H^* = \{x^* \in X^* : x^*(x) = 1\}$. Then it is easy to see that $\text{dist}(0, H^*) = 1$. Hence, for any $e^* \in \overline{H^*}^{w^*}$, there exists a net $\{e_\alpha^*\}_{\alpha \in \Delta} \subset H^*$ such that $e_\alpha^* \xrightarrow{w^*} e^*$. Moreover, it is easy to see that $1 = e_\alpha^*(x) = e^*(x)$. This implies that $e^* \in H^*$. Hence we obtain that H^* is a weak* closed set. By the proof of Theorem 2.1 of [13], we know that if H^* is a weak* closed set, then H^* is a proximal set. Hence there exists $y_n^* \in H^*$ such that $\|z_n^* - y_n^*\| = \text{dist}(z_n^*, H^*)$. Pick $x_0^* \in H^*$. Then $H^* - x_0^* = \{x^* \in X^* : x^*(x) = 0\}$. Since

$$\begin{aligned} \|z_n^* - y_n^*\| &= \text{dist}(z_n^*, H^*) = \text{dist}(z_n^* - x_0^*, H^* - x_0^*) \\ &= |(z_n^* - x_0^*)(x)| \rightarrow 0, \end{aligned} \quad (26)$$

we obtain that

$$\|y_n^*\| \leq \|z_n^*\| + \|z_n^* - y_n^*\| \rightarrow 1 = \text{dist}(0, H^*). \quad (27)$$

Therefore, by $\|y_n^*\| \geq 1$ and formula (27), we obtain that

$$\lim_{n \rightarrow \infty} \|0 - y_n^*\| = \lim_{n \rightarrow \infty} \|y_n^*\| = 1 = \text{dist}(0, H^*). \quad (28)$$

Since every weak* hyperplane of X^* is approximatively compact, we obtain that $\{y_n^*\}_{n=1}^\infty$ has a subsequence converging to an element in H^* . By formula (26), we obtain that $\{z_n^*\}_{n=1}^\infty$ has a subsequence converging to a element in H^* . Hence we may assume without loss of generality that $z_n^* \rightarrow z_0^*$. Then $\|z_0^*\| \leq 1$. By $z_n^*(x) \rightarrow z_0^*(x)$ and $z_n^*(x) \rightarrow 1$, we have $z_0^*(x) = 1$. This implies that $z_0^* \in A(x)$. Then $\text{dist}(z_n^*, A(x)) \rightarrow 0$ as $n \rightarrow \infty$. By $z_n^* = h_n^*/\|h_n^*\|$ and $\|h_n^*\| \rightarrow 1$, we obtain that $\text{dist}(h_n^*, A(x)) \rightarrow 0$ as $n \rightarrow \infty$. Moreover, we have proved that $r = r(x, U_{A(x)}) > 0$ for any open set $U_{A(x)} \supset A(x)$, a contradiction. This implies that there exists $m > 0$ such that $1 > y^*(x) + m$ whenever $y^* \in B(X^*) \setminus U_{A(x)}$. Hence, for any $x^* \in A(x)$, we have

$$\begin{aligned} 1 - m &\geq \sup \{y^*(x) : y^* \in B(X^*) \setminus U_{A(x)}\} \\ &= \sup \{y^*(x) : y^* \in \text{co}(B(X^*) \setminus U_{A(x)})\} \\ &= \sup \{y^*(x) : y^* \in \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)})\}. \end{aligned} \quad (29)$$

This implies that $x^* \notin \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)})$. Therefore, by the arbitrariness of x^* , we have $A(x) \cap \overline{\text{co}}^{w^*}(B(X^*) \setminus U_{A(x)}) = \emptyset$. Hence X^* is w^* nearly dentable space. This completes the proof. $\square$

Theorem 12. Suppose that X^* is w^* nearly dentable space. Then,

- (1) for any weak* open convex set C , metric projector operator $P_{\overline{C}^{w^*}}$ is upper semicontinuous;
- (2) for any weak* hyperplane D^* , metric projector operator P_{D^*} is upper semicontinuous.

Proof. (1) Let C be a weak* open convex set and $H^* = \overline{C}^{w^*}$. Suppose that metric projector operator P_{H^*} is not upper semicontinuous. Then we without loss of generality may assume that metric projector operator P_{H^*} is not upper semicontinuous in origin. Hence there exist a sequence $\{x_n^*\}_{n=1}^\infty$ and an open set $U_{P_{H^*}(0)} \supset P_{H^*}(0)$ such that $P_{H^*}(x_n^*) \notin U_{P_{H^*}(0)}$, and $x_n^* \rightarrow 0$ as $n \rightarrow \infty$. This implies that there exists $y_n^* \in P_{H^*}(x_n^*)$ such that $y_n^* \notin U_{P_{H^*}(0)}$. We will derive a contradiction for each of the following two cases.

Case I. Let $0 \in H^*$. Then, by $x_n^* \rightarrow 0$, we have $\text{dist}(x_n^*, H^*) \rightarrow 0$ as $n \rightarrow \infty$. Moreover, by $y_n^* \in P_{H^*}(x_n^*)$, we have $\|x_n^* - y_n^*\| = \text{dist}(x_n^*, H^*) \rightarrow 0$ as $n \rightarrow \infty$. Therefore,

$$0 \leq \|y_n^*\| \leq \|x_n^*\| + \|x_n^* - y_n^*\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (30)$$

This implies that $y_n^* \rightarrow 0$ as $n \rightarrow \infty$. Hence there exists $n \in N$ such that $y_n^* \in U_{P_{H^*}(0)}$, a contradiction.

Case II. Let $0 \notin H^*$. Then $d = \text{dist}(0, H^*) > 0$, $\overline{B_d(0)} \cap C = \emptyset$, and $\overline{B_d(0)} \cap H^* = P_{H^*}(0)$. Therefore, by $(X^*, w^*)^* = X$ and the separation theorem of locally convex space, there exists $x \in S(X)$ such that $\sup\{x(y^*) : y^* \in \overline{B_d(0)}\} \leq \inf\{x(y^*) : y^* \in C\}$. This implies that

$$\begin{aligned} &\sup \{x(y^*) : y^* \in \overline{B_d(0)}\} \\ &\leq \inf \{x(y^*) : y^* \in H^*\} = d. \end{aligned} \quad (31)$$

Therefore, by $P_{H^*}(0) \subset H^*$, we obtain that $x^*(x) = d$ for any $x^* \in P_{H^*}(0)$. Since $P_{H^*}(0) \subset \overline{B_d(0)}$, we obtain that $\{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\} \supset P_{H^*}(0)$. Since the distance function is continuous, we obtain that

$$\|x_n^* - y_n^*\| = \text{dist}(x_n^*, H^*) \longrightarrow \text{dist}(0, H^*) = d \quad (32)$$

as $n \longrightarrow \infty$.

Moreover, we have

$$0 \leq \|y_n^*\| - \|x_n^* - y_n^*\| \leq \|x_n^*\| \longrightarrow 0 \quad \text{as } n \longrightarrow \infty. \quad (33)$$

Therefore, by formulas (32) and (33), we have $\|y_n^*\| \rightarrow d$. By $y_n^* \notin U_{P_{H^*}(0)}$, we get that every subsequence $\{y_n^*\}_{n=1}^\infty$ does not converge to point of $P_{H^*}(0)$. Moreover, by $y_n^* \notin U_{P_{H^*}(0)}$, we have $\|y_n^*\| > d$. Hence, for any $y^* \in P_{H^*}(0)$, there exists $\varepsilon_{y^*} > 0$ such that $\{y_n^*\}_{n=1}^\infty \cap \{x^* : \|x^* - y^*\| < \varepsilon_{y^*}\} = \emptyset$. Since H^* is closed convex set and $(\{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\} \setminus P_{H^*}(0)) \cap H^* = \emptyset$, we obtain that for any $y^* \in \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\} \setminus P_{H^*}(0)$, there exists $\varepsilon_{y^*} > 0$ such that $\text{dist}(y^*, H^*) = 2\varepsilon_{y^*}$. This implies that $\{y_n^*\}_{n=1}^\infty \cap \{x^* : \|x^* - y^*\| < \varepsilon_{y^*}\} = \emptyset$. Noticing that

$$\begin{aligned} & (\{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\} \setminus P_{H^*}(0)) \cup P_{H^*} \\ &= \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}, \end{aligned} \quad (34)$$

we obtain that, for any $y^* \in \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}$, there exists $\varepsilon_{y^*} > 0$ such that

$$\{y_n^*\}_{n=1}^\infty \cap \{x^* \in X^* : \|x^* - y^*\| < \varepsilon_{y^*}\} = \emptyset. \quad (35)$$

Therefore, by $\|y_n^*\| \rightarrow d$ and $\|y_n^*\| > d$, we have

$$\left\| \frac{d}{\|y_n^*\|} \cdot y_n^* - y_n^* \right\| = \left| \frac{d}{\|y_n^*\|} - 1 \right| \cdot \|y_n^*\| \longrightarrow 0 \quad (36)$$

as $n \longrightarrow \infty$.

Let us define sequence $\{z_n^*\}_{n=1}^\infty$, where

$$z_n^* = \frac{n}{n+1} \cdot \frac{d}{\|y_n^*\|} \cdot y_n^* \quad n \in N. \quad (37)$$

Then

$$\begin{aligned} \|z_n^*\| &= \left\| \frac{n}{n+1} \cdot \frac{d}{\|y_n^*\|} \cdot y_n^* \right\| = \left(\frac{n}{n+1} \cdot \frac{d}{\|y_n^*\|} \right) \cdot \|y_n^*\| \\ &= \frac{nd}{n+1} < d. \end{aligned} \quad (38)$$

Noticing that $x \in S(X)$, we obtain that $z_n^*(x) < d$. This implies that $z_n^* \notin \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}$. Since $\|y_n^*\| \rightarrow d$, we have

$$\begin{aligned} \|z_n^* - y_n^*\| &= \left\| \frac{n}{n+1} \cdot \frac{d}{\|y_n^*\|} \cdot y_n^* - y_n^* \right\| \\ &= \left| \frac{n}{n+1} \cdot \frac{d}{\|y_n^*\|} - 1 \right| \cdot \|y_n^*\| \longrightarrow 0. \end{aligned} \quad (39)$$

Hence there exists $n_{y^*} \in N$ such that $\|z_n^* - y_n^*\| < (1/4)\varepsilon_{y^*}$ whenever $n > n_{y^*}$. Therefore, by formula (35), we have $\|y_n^* - y^*\| \geq \varepsilon_{y^*}$. Hence if $n > n_{y^*}$, then

$$\begin{aligned} \|y^* - z_n^*\| &\geq \|y^* - y_n^*\| - \|y_n^* - z_n^*\| \geq \varepsilon_{y^*} - \frac{1}{4}\varepsilon_{y^*} \\ &= \frac{3}{4}\varepsilon_{y^*}. \end{aligned} \quad (40)$$

Since $z_n^* \notin \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}$ and $y^* \in \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}$, we have $\min\{\|y^* - z_1^*\|, \|y^* - z_2^*\|, \dots, \|y^* - z_{n_{y^*}}^*\|\} > 0$. Put

$$\begin{aligned} \eta_{y^*} &= \frac{1}{2} \min \left\{ \frac{3}{4}\varepsilon_{y^*}, \right. \\ &\quad \left. \min \left\{ \|y^* - z_1^*\|, \|y^* - z_2^*\|, \dots, \|y^* - z_{n_{y^*}}^*\| \right\} \right\}. \end{aligned} \quad (41)$$

Then, by formulas (40) and (41), we have $\{z_n^*\}_{n=1}^\infty \cap \{x^* : \|x^* - y^*\| < \varepsilon_{y^*}\} = \emptyset$. Moreover, by the arbitrariness of y^* , we have

$$\begin{aligned} & \{z_n^*\}_{n=1}^\infty \\ & \cap \bigcup_{y^* \in \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}} \{x^* : \|x^* - y^*\| < \varepsilon_{y^*}\} \\ &= \emptyset. \end{aligned} \quad (42)$$

Put

$$\begin{aligned} \overline{P_{H^*}(0)} &= \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}, \\ V_{\overline{P_{H^*}(0)}} &= \bigcup_{y^* \in \overline{P_{H^*}(0)}} \{x^* : \|x^* - y^*\| < \varepsilon_{y^*}\}. \end{aligned} \quad (43)$$

Then, by formulas (42) and (43), we have $\{z_n^*\}_{n=1}^\infty \cap V_{\overline{P_{H^*}(0)}} = \emptyset$.

Since the distance function is continuous, we have $\|x_n^* - y_n^*\| = \text{dist}(x_n^*, H^*) \rightarrow \text{dist}(0, H^*)$ as $n \rightarrow \infty$. Moreover, we have $x_n^* \rightarrow 0$ as $n \rightarrow \infty$. This implies that $\|0 - y_n^*\| \rightarrow \text{dist}(0, H^*)$ as $n \rightarrow \infty$. Therefore, by $\|z_n^* - y_n^*\| \rightarrow 0$ as $n \rightarrow \infty$, we obtain that

$$\begin{aligned} \|0 - z_n^*\| &\leq \|0 - y_n^*\| + \|y_n^* - z_n^*\| \longrightarrow \text{dist}(0, H^*) \\ &\text{as } n \longrightarrow \infty. \end{aligned} \quad (44)$$

Since $\|0 - z_n^*\| \geq \text{dist}(0, H^*)$, we obtain that $\|0 - z_n^*\| \rightarrow \text{dist}(0, H^*)$ as $n \rightarrow \infty$. By

$$\begin{aligned} \frac{1}{d} y_n^*(x) &= 1, \\ z_n^* &= \frac{n}{n+1} \cdot \frac{d}{\|y_n^*\|} \cdot y_n^*, \end{aligned} \quad (45)$$

we have $z_n^*(x)/d \rightarrow 1$ as $n \rightarrow \infty$. Since $B(X^*)$ is weak* compact, there exists $z_0^*/d \in B(X^*)$ such that z_0^*/d is a weak* accumulation point of $\{z_n^*/d\}_{n=1}^\infty$. Hence there exists a net

$\{z_n^*\}_{n=1}^\infty \subset \{z_n^*\}_{n=1}^\infty$ such that $z_n^* \xrightarrow{w^*} z_0^*$ and $z_0^*/d \in A(x)$. Since $V_{P_{H^*}(0)} \supset \{x^* : x^*(x) = d, x^* \in \overline{B_d(0)}\}$, we obtain that

$$\frac{1}{d} V_{P_{H^*}(0)} \supset \frac{1}{d} \{x^* \in \overline{B_d(0)} : x^*(x) = d\} \quad (46)$$

$$= \{x^* \in B(X^*) : x^*(x) = 1\} = A(x).$$

Let $U_{A(x)} = (1/d)V_{P_{H^*}(0)}$. Then, by $\{z_n^*\}_{n=1}^\infty \cap V_{P_{H^*}(0)} = \emptyset$ and $\{z_n^*\}_{n=1}^\infty \subset \overline{B_d(0)}$, we have $z_n^* \in \overline{B_d(0)} \setminus V_{P_{H^*}(0)}$. This implies that $z_n^* \in \overline{B_d(0)}$ and $z_n^* \notin V_{P_{H^*}(0)}$. Therefore,

$$\frac{1}{d} z_n^* \in \frac{1}{d} \overline{B_d(0)} = B(X^*), \quad (47)$$

$$\frac{1}{d} z_n^* \notin \frac{1}{d} V_{P_{H^*}(0)} = U_{A(x)}.$$

This implies that $z_n^*/d \in B(X^*) \setminus U_{A(x)}$. Since X^* is a w^* nearly dentable space, we obtain that $A(x) \cap \overline{co}^{w^*}(B(X^*) \setminus U_{A(x)}) = \emptyset$. Since $B(X^*)$ is w^* compact, it is easy to see that $\overline{co}^{w^*}(B(X^*) \setminus U_{A(x)})$ is w^* compact. Moreover, since $A(x)$ is w^* closed convex set, by the separation theorem of locally convex space, there exist $y \in X$ and $r > 0$ such that

$$\inf \{x^*(y) : x^* \in A(x)\} - r$$

$$\geq \sup \{x^*(y) : x^* \in \overline{co}^{w^*}(B(X^*) \setminus U_{A(x)})\}. \quad (48)$$

Therefore, by $z_n^* \xrightarrow{w^*} z_0^*$, we obtain that $z_n^*/d \xrightarrow{w^*} z_0^*/d$. Noticing that $z_n^*/d \in B(X^*) \setminus U_{A(x)}$, we obtain that

$$\left\{ \frac{1}{d} z_n^* \right\}_{n=1}^\infty \subset \left\{ \frac{1}{d} z_n^* \right\}_{n=1}^\infty \subset B(X^*) \setminus U_{A(x)} \quad (49)$$

$$\subset \overline{co}^{w^*}(B(X^*) \setminus U_{A(x)}).$$

Therefore, by formulas (48)-(49) and $z_0^*/d \in A(x)$, we have $(z_0^*/d)(y) - r \geq \sup \{x^*(y) : x^* \in \{z_n^*/d\}_{n=1}^\infty\}$, which contradicts $z_n^*/d \xrightarrow{w^*} z_0^*/d$. Hence the metric projector operator P_{H^*} is upper semicontinuous.

By the proof (3) $\Rightarrow$ (1) of Theorem 11, it is easy to see that (2) is true. This completes the proof. $\square$

4. Some Examples of w^* Nearly Dentable Space in Orlicz Function Space

In the section, authors prove that there exists a Banach space X^* such that X^* is not a w^* nearly dentable Banach space. Moreover, authors also prove that there exists a Banach space X^* such that X^* is a w^* nearly dentable Banach space and not a nearly dentable Banach space. First, we give the definition of Orlicz function space.

Definition 13. $M : R \rightarrow R$ is called an N -function if it has the following properties:

- (1) M is even, convex, and $M(0) = 0$;
- (2) $M(u) > 0$ for all $u \neq 0$;
- (3) $\lim_{u \rightarrow 0} M(u)/u = 0$ and $\lim_{u \rightarrow \infty} M(u)/u = \infty$.

Let $p(u)$ denote the right derivative of $M(u)$ at $u \in R^+$ and let $q(v)$ be the generalized inverse function of $p(u)$ defined on R^+ by $q(v) = \sup_{u \geq 0} \{u \geq 0 : p(u) \leq v\}$. Then we call $N(v) = \int_0^{|v|} q(s)ds$ the complementary function of M . It is well known that there holds the Young inequality $uv \leq M(u) + N(v)$ and $uv = M(u) + N(v) \Leftrightarrow u = |q(v)| \operatorname{sign} v$ or $v = |p(u)| \operatorname{sign} u$. Moreover, it is well known that M and N are complementary to each other. A continuous function $M : R \rightarrow R$ is called strictly convex if $2M((u+v)/2) < M(u) + M(v)$ for all $u \neq v$.

Let (G, Σ, μ) be a finite nonatomic and complete measure space. Denote by p and q the right derivative of M and N , respectively. We define

$$\rho_M(x) = \int_G M(x(t)) dt,$$

$$L_M = \{x(t) : \rho_M(\lambda x) < \infty \text{ for some } \lambda > 0\}, \quad (50)$$

$$E_M = \{x(t) : \rho_M(\lambda x) < \infty \forall \lambda > 0\}.$$

It is well known that the Orlicz function space L_M is a Banach space when it is equipped with the Luxemburg norm

$$\|x\| = \inf \left\{ \lambda > 0 : \rho_M\left(\frac{x}{\lambda}\right) \leq 1 \right\} \quad (51)$$

or equipped with the Amemiya-Orlicz norm

$$\|x\|^0 = \inf_{k>0} \frac{1}{k} [1 + \rho_M(kx)]. \quad (52)$$

L_M, E_M denote Orlicz function spaces equipped with the Luxemburg norm. L_M^0, E_M^0 denote Orlicz function spaces equipped with the Orlicz norm. It is well known that $(E_N)^* = L_M^0, (E_N^0)^* = L_M$.

Definition 14 (see [13]). We say that an N -function $M \in \Delta_2(N \in \nabla_2)$ if there exist $K > 2$ and $u_0 \geq 0$ such that $M(2u) \leq KM(u)$ whenever $u \geq u_0$.

Lemma 15 (see [13, 14]). Let L_M be Orlicz function space. Then the following statements are equivalent:

- (1) Every weakly* hyperplane of L_M is approximatively compact;
- (2) E_N^0 is smooth space and $M \in \Delta_2$.

Lemma 16 (see [15]). $L_M^0(L_M)$ has the Radon-Nikodym property if and only if $M \in \Delta_2$.

Lemma 17 (see [15]). L_M^0 is a strictly convex space if and only if M is strictly convex.

Lemma 18 (see [15]). $L_M(L_M^0)$ is reflexive if and only if $M \in \Delta_2 \cap \nabla_2$.

Theorem 19. *If every weakly* hyperplane of L_M^0 is approximatively compact, then $M \in \Delta_2$.*

Proof. Since every weak* hyperplane of L_M^0 is approximatively compact, we obtain that L_M^0 has the Radon-Nikodym property (see [14]). Therefore, by Lemma 16, we have $M \in \Delta_2$. This completes the proof. $\square$

Example 20. If M is strictly convex and $M \notin \Delta_2$, then L_M^0 is not w^* nearly dentable. In fact, suppose that L_M^0 is w^* nearly dentable. Since M is strictly convex, we obtain that L_M^0 is a strictly convex space by Lemma 17. Since $(E_N)^* = L_M^0$, we obtain that E_N is a smooth space. This implies that if E is a separable closed subspace of E_N , then $A_E(x) = \{x^* \in S(E^*) : x^*(x) = 1 = \|x\|\}$ is a singleton. Therefore, by Theorem 10, we obtain that L_M^0 has the Radon-Nikodym property. Then, by Lemma 16, we obtain that $M \in \Delta_2$, a contradiction.

Example 21. If M is strictly convex, $M \in \Delta_2$ and $N \notin \Delta_2$, then L_M is w^* nearly dentable not nearly dentable. In fact, by Theorem 19, we obtain that every weakly* hyperplane of L_M^0 is approximatively compact. Therefore, by Theorem 11, we obtain that L_M is w^* nearly dentable. Since $N \notin \Delta_2$, we obtain that L_M is not reflexive. This implies that L_M is not nearly dentable.

5. Applications to Theory of Generalized Inverses of Banach Space

Let T be a linear bounded operator from X into Y . Let $D(T)$, $R(T)$ and $N(T)$ denote the domain, range, and null space of T , respectively. If $N(T) \neq \{0\}$ or $R(T) \neq Y$, the operator equation $Tx = y$ is generally ill-posed. In applications, one usually looks for the best approximative solution (b.a.s.) to the equation (see [16]).

A point $x_0 \in D(T)$ is called the best approximative solution to the operator equation $Tx = y$, if

$$\begin{aligned} \|Tx_0 - y\| &= \inf \{\|Tx - y\| : x \in D(T)\}, \\ \|x_0\| &= \min \left\{ \|v\| : v \in D(T), \|Tv - y\| \right. \\ &= \left. \inf_{x \in D(T)} \|Tx - y\| \right\}, \end{aligned} \quad (53)$$

where $y \in Y$ (see [16]). Nashed and Votruba [16] introduced the concept of the (set-valued) metric generalized inverse T .

Definition 22. Let X, Y be Banach spaces and let T be a linear operator from X to Y . The set-valued mapping $T^\partial : Y \rightarrow X$ defined by

$$\begin{aligned} T^\partial(y) &= \{x_0 \in D(T) : \\ &x_0 \text{ is a best approximation solution to } T(x) = y\} \end{aligned} \quad (54)$$

for any $y \in D(T^\partial)$ is called the (set-valued) metric generalized inverse of T , where

$$\begin{aligned} D(T^\partial) &= \{y \in Y : T(x) \\ &= y \text{ has a best approximation solution in } X\}. \end{aligned} \quad (55)$$

Definition 23. $H(A, B)$ is said to be Hausdorff distance of A and B if

$$\begin{aligned} H(A, B) &= \max \left\{ \sup_{a \in A} \left\{ \inf_{b \in B} \|a - b\| \right\}, \sup_{b \in B} \left\{ \inf_{a \in A} \|a - b\| \right\} \right\}. \end{aligned} \quad (56)$$

During the last three decades, the linear generalized inverses of linear operators in Banach spaces and their applications have been investigated by many authors. In 2008, Chen et al. [8], under some assumption on the geometric structure of Banach space, gave the criteria for the existence of continuous metric generalized inverse. In 2005, Wei et al. [17] presented componentwise condition numbers for the problems of Moore-Penrose generalized matrix inverses and linear least squares. For the selection of the metric generalized inverse of linear operator in Banach space, in 2008, Hudzik et al. [18] obtained a bounded homogeneous selection.

Theorem 24. *Let X^* be a w^* nearly dentable space, Y^* a nearly dentable space, $N(T)$ a one-dimensional subspace of X , and $R(T^*)$ a closed semi-Chebychev set. Then $D((T^*)^\partial) = X^*$. Moreover, if for any $y_0^* \in (T^*)^\partial(x_0^*)$ there exists $f \in S(X^{**})$ such that A_f is compact and $y_0^*/\|y_0^*\| \in A_f$, then*

- (1) $(T^*)^\partial(x_0^*)$ is compact and $(T^*)^\partial$ is upper semicontinuous at x_0^* ;
- (2) $(T^*)^\partial$ is lower semicontinuous at x_0^* if and only if the function $g(x^*) = H((T^*)^\partial(x^*), (T^*)^\partial(x_0^*))$ is lower semicontinuous at x_0^* .

Proof. Since $R(T^*)$ is closed, we obtain that $R(T^*) = N(T)^\perp = \{x^* \in X^* : x^*(x) = 0, x \in N(T)\}$ by the closed range theorem. Since $\dim N(T) = 1$, there exists $x \in S(X) \cap N(T)$ such that $R(T^*) = \{x^* \in X^* : x^*(x) = 0\}$. This implies that $R(T^*)$ is a weakly* hyperplane of X^* . By proof of Theorem 2.1 of [13], we obtain that $R(T^*)$ is a proximal set. Since $R(T^*)$ is a semi-Chebychev set, we obtain that $R(T^*)$ is a Chebychev set. Let $x^* \in X^*$. Then there exists a unique $z^* \in R(T^*)$ such that $z^* \in P_{R(T^*)}(x^*)$. Hence there exists $y^* \in Y^*$ such that $(T^*)^{-1}(z^*) = y^* + N(T^*)$. Since Y^* is a nearly dentable space, we obtain that Y^* is reflexive. This implies that $N(T^*)$ is a proximal set. Hence we obtain that $D((T^*)^\partial) = X^*$.

(1) We will prove that the set $(T^*)^\partial(x_0^*)$ is compact. Let $x_n^* \rightarrow x_0^*$. Since X^* is a w^* nearly dentable space and $R(T^*)$ is a weakly* hyperplane of X^* , by Theorem 12, we obtain that metric projector operator $P_{R(T^*)}$ is upper semicontinuous. Moreover, since $R(T^*)$ is a Chebychev set, we obtain that metric projector operator $P_{R(T^*)}$ is continuous; that is, $P_{R(T^*)}(x_n^*) \rightarrow P_{R(T^*)}(x_0^*)$ as $n \rightarrow \infty$. Since T is linear

bounded operator, we obtain that $N(T^*)$ is a closed subspace. Put

$$\overline{T^*} : \left[\frac{Y^*}{N(T^*)} \right] \longrightarrow R(T^*), \quad \overline{T^*}[y^*] = T^*(y^*). \quad (57)$$

Then, by the inverse operator theorem, we obtain that

$$\begin{aligned} [y_n^*] &= \overline{T^*}^{-1}(P_{R(T^*)}(x_n^*)) \longrightarrow \overline{T^*}^{-1}(P_{R(T^*)}(x_0^*)) \\ &= [y_0^*] \quad \text{as } n \longrightarrow \infty. \end{aligned} \quad (58)$$

This implies that $\|[y_n^*] - [y_0^*]\| \rightarrow 0$ as $n \rightarrow \infty$. Since, for any $y_0^* \in (T^*)^\partial(x_0^*)$, there exists $f \in S(X^{**})$ such that $y_0^*/\|y_0^*\| \in A_f$, we have $0 \notin (T^*)^\partial(x_0^*)$. Moreover, we may assume without loss of generality that $\|y_n^* - y_0^*\| \rightarrow 0$ as $n \rightarrow \infty$. Therefore, by the definition of set-valued metric generalized inverse, we obtain that

$$\begin{aligned} (T^*)^\partial(x_n^*) &= y_n^* - P_{N(T^*)}(y_n^*), \\ (T^*)^\partial(x_0^*) &= y_0^* - P_{N(T^*)}(y_0^*). \end{aligned} \quad (59)$$

Suppose that $P_{N(T^*)}(y_0^*)$ is not compact. Then there exists a sequence $\{z_n^*\}_{n=1}^\infty \subset P_{N(T^*)}(y_0^*)$ such that $\{z_n^*\}_{n=1}^\infty$ has no Cauchy subsequence. Since Y^* is a nearly dentable space, we obtain that Y^* is reflexive. Hence we may assume without loss of generality that $z_n^* \xrightarrow{w} z_0^*$. Noticing that $P_{N(T^*)}(y_0^*)$ is a closed convex set, we obtain that $P_{N(T^*)}(y_0^*)$ is a weakly closed convex set. Then $z_0^* \in P_{N(T^*)}(y_0^*)$. Put

$$\begin{aligned} h_n^* &= \frac{y_0^* - z_n^*}{\|y_0^* - z_n^*\|}, \\ h_0^* &= \frac{y_0^* - z_0^*}{\|y_0^* - z_0^*\|}. \end{aligned} \quad (60)$$

Noticing that $z_n^* \in P_{N(T^*)}(y_0^*)$, $z_0^* \in P_{N(T^*)}(y_0^*)$, and $0 \notin (T^*)^\partial(x_0^*)$, we obtain that $\|y_0^* - z_n^*\| = \|y_0^* - z_0^*\| \neq 0$. By $z_n^* \xrightarrow{w} z_0^*$, we obtain that $h_n^* \xrightarrow{w} h_0^*$. Moreover, it is easy to see that $\|y_0^* - z_0^*\|h_0^* \in (T^*)^\partial(x_0^*)$. Then there exists $f \in S(X^{**})$ such that A_f is compact and $h_0^* \in A_f$. We will derive a contradiction for each of the following two cases.

Case I. Let $\{h_n^*\}_{n=1}^\infty \cap A_f$ be a finite set. Then we may assume without loss of generality that $\{h_n^*\}_{n=1}^\infty \cap A_f = \emptyset$. Hence, for any $y^* \in A_f$, there exists $\eta_{y^*} > 0$ such that $\{h_n^*\}_{n=1}^\infty \cap B(y^*, \eta_{y^*}) = \emptyset$, where $f(h_0^*) = 1 = \|f\|$. Hence there exists an open set $U_{A_f} = \cup_{y^* \in A_f} B(y^*, \eta_{y^*})$ such that $U_{A_f} \cap \{h_n^*\}_{n=1}^\infty = \emptyset$. Since Y^* is a nearly dentable space, we obtain that $A_f \cap \overline{\text{co}}(B(X) \setminus U_{A_f}) = \emptyset$. By the separation theorem, there exists $g \in Y^{**}$ such that

$$\begin{aligned} \inf \{g(y^*) : y^* \in A_f\} \\ - \sup \{g(y^*) : y^* \in \overline{\text{co}}(B(X) \setminus U_{A_f})\} > 0. \end{aligned} \quad (61)$$

Noticing that $\{h_n^*\}_{n=1}^\infty \subset \overline{\text{co}}(B(X) \setminus U_{A_f})$ and $h_0^* \in A_f$, we obtain that $g(h_0^*) - \sup\{g(y^*) : y^* \in \{h_n^*\}_{n=1}^\infty\} > 0$, which contradict $h_n^* \xrightarrow{w} h_0^*$ as $n \rightarrow \infty$. This implies that the sequence $\{h_n^*\}_{n=1}^\infty$ has a Cauchy subsequence.

Case II. Let $\{h_n^*\}_{n=1}^\infty \cap A_f$ be an infinite set. Then there exists a subsequence $\{h_{n_l}^*\}_{l=1}^\infty$ of $\{h_n^*\}_{n=1}^\infty$ such that $\{h_{n_l}^*\}_{l=1}^\infty \subset A_f$. Since A_f is a compact set, we obtain that $\{h_{n_l}^*\}_{l=1}^\infty \subset A_f$ is relatively compact. This implies that $\{h_n^*\}_{n=1}^\infty$ has a Cauchy subsequence. Hence $P_{N(T^*)}(y_0^*)$ is compact. By $(T^*)^\partial(x_0^*) = y_0^* - P_{N(T^*)}(y_0^*)$, we obtain that $(T^*)^\partial(x_0^*)$ is compact.

We next will prove that the set-valued mapping $(T^*)^\partial$ is upper semicontinuous at x_0^* . Let W be an open set and $P_{N(T^*)}(y_0^*) \subset W$. We claim that there exists $\delta > 0$ such that

$$\bigcup_{z^* \in P_{N(T^*)}(y_0^*)} \{y^* \in Y^* : \|y^* - z^*\| < 2\delta\} \subset W. \quad (62)$$

Otherwise, there exists a sequence $\{h_n^*\}_{n=1}^\infty \subset P_{N(T^*)}(y_0^*)$ such that $B(h_n^*, 1/n) \not\subset W$. Let $\{h_k^*\}_{k=1}^\infty$ be a Cauchy sequence.

Noticing that $h_n^* \xrightarrow{w} h_0^*$, we have $h_{n_k}^* \rightarrow h_0^*$ as $k \rightarrow \infty$. Then

$$\lim_{k \rightarrow \infty} z_{n_k}^* = y_0^* - \|y_0^* - z_0^*\|h_0^* \in P_{N(T^*)}(y_0^*). \quad (63)$$

Hence there exists $r > 0$ such that $B(y_0^* - \|y_0^* - z_0^*\|h_0^*, r) \subset W$. Since $z_{n_k}^* \rightarrow y_0^* - \|y_0^* - z_0^*\|h_0^*$, there exists $k_0 \in \mathbb{N}$ such that

$$\begin{aligned} \|(y_0^* - \|y_0^* - z_0^*\|h_0^*) - z_{n_{k_0}}^*\| &< \frac{1}{4}r, \\ \frac{1}{n_{k_0}} &< \frac{1}{4}r. \end{aligned} \quad (64)$$

Hence, if $\|y^* - z_{n_{k_0}}^*\| \leq 1/n_{k_0}$, then

$$\begin{aligned} &\|y^* - (y_0^* - \|y_0^* - z_0^*\|h_0^*)\| \\ &\leq \|y^* - z_{n_{k_0}}^*\| + \|z_{n_{k_0}}^* - (y_0^* - \|y_0^* - z_0^*\|h_0^*)\| \\ &\leq \frac{1}{n_{k_0}} + \frac{1}{4}r < r. \end{aligned} \quad (65)$$

This means that if $\|y^* - z_{n_{k_0}}^*\| \leq 1/n_{k_0}$, then $y^* \in B(z_{n_{k_0}}^*, 1/n_{k_0}) \subset B(y_0^* - \|y_0^* - z_0^*\|h_0^*, r) \subset W$, a contradiction. Hence there exists $\delta > 0$ such that

$$\bigcup_{z^* \in P_{N(T^*)}(y_0^*)} \{y^* \in Y^* : \|y^* - z^*\| < 2\delta\} \subset W. \quad (66)$$

Let

$$\begin{aligned} W_1 &= \bigcup_{z^* \in P_{N(T^*)}(y_0^*)} \{y^* \in Y^* : \|y^* - z^*\| < \delta\} \\ &\supset P_{N(T^*)}(y_0^*). \end{aligned} \quad (67)$$

Since Y^* is a w^* nearly dentable space, we obtain that the metric projector operator $P_{N(T^*)}$ is upper semicontinuous. Since

metric projector operator $P_{N(T^*)}$ is upper semicontinuous and $\|y_n^* - y_0^*\| \rightarrow 0$, there exists $n_1 \in \mathbb{N}$ such that $P_{N(T^*)}(y_n^*) \subset W_1$ whenever $n > n_1$. This means that $y_0^* - P_{N(T^*)}(y_n^*) \subset y_0^* - W_1$ whenever $n > n_1$. Moreover, by $\|y_n^* - y_0^*\| \rightarrow 0$, there exists $n_2 \in \mathbb{N}$ such that $\|y_n^* - y_0^*\| < \delta/4$ whenever $n > n_1$. Let $N_1 = \max\{n_1, n_2\}$. We claim that $y_n^* - P_{N(T^*)}(y_n^*) \subset y_0^* - W$ whenever $n > N_1$. In fact, if $z^* \in y_n^* - P_{N(T^*)}(y_n^*)$, then there exists $h_n^* \in P_{N(T^*)}(y_n^*)$ such that $z^* = y_n^* - h_n^*$. Let $l_n^* = y_n^* - y_0^*$. Then $z^* = y_0^* - (h_n^* - l_n^*)$ and $\|l_n^*\| < \delta/4$. Since $h_n^* \in P_{N(T^*)}(y_n^*) \subset W_1$, there exists $v_n^* \in P_{N(T^*)}(y_0^*)$ such that $\|h_n^* - v_n^*\| < \delta$. Then

$$\|h_n^* - l_n^* - v_n^*\| \leq \|h_n^* - v_n^*\| + \|l_n^*\| \leq \delta + \frac{1}{4}\delta < 2\delta. \quad (68)$$

This implies that $h_n^* - l_n^* \in W$. Then $z^* = y_0^* - (h_n^* - l_n^*) \in y_0^* - W$. This implies that $y_n^* - P_{N(T^*)}(y_n^*) \subset y_0^* - W$. Therefore, by formula (59), we have $(T^*)^\partial(x_n^*) \subset y_0^* - W$.

Let V be an open set and $(T^*)^\partial(x_0^*) \subset V$. Then, by formula (59), we have $y_0^* - V \supset P_{N(T^*)}(y_0^*)$. Hence there exists natural number N_2 such that $(T^*)^\partial(x_n^*) \subset y_0^* - (y_0^* - V)$ whenever $n > N_2$. This implies that $(T^*)^\partial(x_n^*) \subset V$ whenever $n > N_2$. Hence we obtain that the set-valued mapping T^∂ is upper semicontinuous at x_0^* .

(2) Let $(T^*)^\partial$ be lower semicontinuous at x_0^* . It is easy to see that $g(x_0^*) = 0$. Suppose that $g(x^*)$ is not lower semicontinuous at x_0^* . Then there exist a sequence $\{x_n^*\}_{n=1}^\infty$ and $\varepsilon_0 > 0$ such that $x_n^* \rightarrow x_0^*$ and

$$\liminf_{n \rightarrow \infty} g(x_n^*) = \liminf_{n \rightarrow \infty} H\left((T^*)^\partial(x_n^*), (T^*)^\partial(x_0^*)\right) > \varepsilon_0. \quad (69)$$

Since $(T^*)^\partial$ is lower semicontinuous at x_0^* , we obtain that, for any $y^* \in (T^*)^\partial(x_0^*)$, there exists $y_n^* \in (T^*)^\partial(x_n^*)$ such that $y_n^* \rightarrow y^*$ as $n \rightarrow \infty$. This implies that for any $y^* \in (T^*)^\partial(x_0^*)$, we have

$$\lim_{n \rightarrow \infty} \inf_{z^* \in (T^*)^\partial(x_n^*)} \|z^* - y^*\| = 0. \quad (70)$$

We claim that

$$\lim_{n \rightarrow \infty} \sup_{y^* \in (T^*)^\partial(x_0^*)} \left\{ \inf_{z^* \in (T^*)^\partial(x_n^*)} \|z^* - y^*\| \right\} = 0. \quad (71)$$

In fact, suppose that formula (71) is not true. Since $(T^*)^\partial(x_0^*)$ is a compact set, there exist $y_n^* \in (T^*)^\partial(x_0^*)$, $z_n^* \in (T^*)^\partial(x_n^*)$, and $r > 0$ such that

$$\begin{aligned} \inf_{z^* \in (T^*)^\partial(x_n^*)} \|z^* - y_n^*\| &= \|z_n^* - y_n^*\|, \\ \limsup_{n \rightarrow \infty} \|z_n^* - y_n^*\| &> 2r. \end{aligned} \quad (72)$$

Since $(T^*)^\partial(x_0^*)$ is compact, we may assume that $y_n^* \rightarrow y_0^* \in (T^*)^\partial(x_0^*)$ as $n \rightarrow \infty$. Therefore, by formula (72), we have

$\limsup_{n \rightarrow \infty} \|z_n^* - y_0^*\| > r$. Moreover, by formula (71), we obtain that

$$\begin{aligned} 0 &= \limsup_{n \rightarrow \infty} \inf_{z^* \in (T^*)^\partial(x_n^*)} \|z^* - y_0^*\| = \limsup_{n \rightarrow \infty} \|z_n^* - y_0^*\| \\ &> r, \end{aligned} \quad (73)$$

a contradiction. Hence formula (71) is true.

Since $(T^*)^\partial$ is upper semicontinuous at x_0^* , we obtain that for any $\varepsilon > 0$, there exists N_1 such that

$$(T^*)^\partial(x_n^*) \subset \bigcup_{y^* \in (T^*)^\partial(x_0^*)} \{z^* \in Y^* : \|z^* - y^*\| < \varepsilon\} \quad (74)$$

whenever $n > N_1$. This implies that for any $y_n^* \in (T^*)^\partial(x_n^*)$, we obtain that $\text{dist}(y_n^*, (T^*)^\partial(x_0^*)) \rightarrow 0$ as $n \rightarrow \infty$. Hence, for any $y_n^* \in (T^*)^\partial(x_n^*)$, we obtain that

$$\begin{aligned} \lim_{n \rightarrow \infty} \inf_{y_n^* \in (T^*)^\partial(x_n^*)} \|y_n^* - z^*\| \\ = \lim_{n \rightarrow \infty} \text{dist}(y_n^*, (T^*)^\partial(x_0^*)) = 0. \end{aligned} \quad (75)$$

We claim that

$$\lim_{n \rightarrow \infty} \sup_{y^* \in (T^*)^\partial(x_0^*)} \left\{ \inf_{z^* \in (T^*)^\partial(x_n^*)} \|y^* - z^*\| \right\} = 0. \quad (76)$$

Otherwise, there exist $y_n^* \in (T^*)^\partial(x_n^*)$, $z_n^* \in (T^*)^\partial(x_0^*)$, and $d > 0$ such that

$$\inf_{z^* \in (T^*)^\partial(x_0^*)} \|y_n^* - z^*\| = \|y_n^* - z_n^*\|, \quad (77)$$

$$\limsup_{n \rightarrow \infty} \|y_n^* - z_n^*\| > 2d.$$

Moreover, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|y_n^* - z_n^*\| &= \lim_{n \rightarrow \infty} \inf_{z^* \in (T^*)^\partial(x_0^*)} \|y_n^* - z^*\| \\ &= \lim_{n \rightarrow \infty} \text{dist}(y_n^*, (T^*)^\partial(x_0^*)) = 0, \end{aligned} \quad (78)$$

a contradiction. This implies that formula (76) is true. Therefore, by formulas (71) and (76), we have

$$\limsup_{n \rightarrow \infty} H\left((T^*)^\partial(x_n^*), (T^*)^\partial(x_0^*)\right) = 0, \quad (79)$$

which contradict formula (69). Hence

$$\begin{aligned} \liminf_{x^* \rightarrow x_0^*} g(x^*) &= \liminf_{x^* \rightarrow x_0^*} H\left((T^*)^\partial(x^*), (T^*)^\partial(x_0^*)\right) \\ &= 0. \end{aligned} \quad (80)$$

This implies that the function $g(x^*) = H((T^*)^\partial(x^*), (T^*)^\partial(x_0^*))$ is lower semicontinuous at x_0^* .

On the other hand, if $g(x^*)$ is lower semicontinuous at x_0^* , then $\liminf_{x^* \rightarrow x_0^*} g(x^*) = \liminf_{x^* \rightarrow x_0^*} H((T^*)^\partial(x^*), (T^*)^\partial(x_0^*)) = 0$. Hence, if $x_n^* \rightarrow x_0^*$, then

$$\lim_{n \rightarrow \infty} \sup_{y^* \in (T^*)^\partial(x_0^*)} \left\{ \inf_{z^* \in (T^*)^\partial(x_n^*)} \|z^* - y^*\| \right\} = 0. \quad (81)$$

Hence, for any $y^* \in (T^*)^\partial(x_0^*)$, there exists $z_n^* \in (T^*)^\partial(x_n^*)$ such that

$$\lim_{n \rightarrow \infty} \|z_n^* - y^*\| = \lim_{n \rightarrow \infty} \inf_{z^* \in (T^*)^\partial(x_n^*)} \|z^* - y^*\| = 0. \quad (82)$$

This implies that $(T^*)^\partial$ is lower semicontinuous at x_0^* . This completes the proof. $\square$

Theorem 25. Let X^* be a w^* nearly dentable space, Y^* approximatively compact, $N(T)$ a one-dimensional subspace, and $R(T^*)$ a closed semi-Chebyshev set. Then

- (1) $D((T^*)^\partial) = X^*$ and $(T^*)^\partial$ is upper semicontinuous;
- (2) $(T^*)^\partial$ is lower semicontinuous at x_0^* if and only if the function $g(x^*) = H((T^*)^\partial(x^*), (T^*)^\partial(x_0^*))$ is lower semicontinuous at x_0^* .

Proof. Let $x_0^* \in X^*$. Since Y^* is approximatively compact, we obtain that, for any $y_0^* \in (T^*)^\partial(x_0^*)$, there exists $f \in S(X^{**})$ such that A_f is compact and $y_0^*/\|y_0^*\| \in A_f$. Moreover, since Y^* is approximatively compact, we obtain that Y^* is a nearly dentable space. By Theorem 24, Theorem 25 is true, which completes the proof. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This work is supported by “China Natural Science Fund under Grant 11401084 and Scientific Research Foundation for the Returned Overseas Chinese Scholars of Heilongjiang Province (Grant no. LC201402)”.

References

- [1] L. Cheng, “Ball-covering property of Banach spaces,” *Israel Journal of Mathematics*, vol. 156, pp. 111–123, 2006.
- [2] L. Chen and L. Cheng, “Analytic characterizations of Mazur’s intersection property via convex functions,” *Journal of Functional Analysis*, vol. 262, no. 11, pp. 4731–4745, 2012.
- [3] S. Shang, Y. Cui, and Y. Fu, “Nearly dentability and approximative compactness and continuity of metric projector in Banach spaces,” *Scientia Sinica Mathematica*, vol. 41, no. 9, pp. 815–825, 2011 (Chinese).
- [4] N. W. Jefimow and S. B. Stechkin, “Approximative compactness and Cheby-shev sets,” *Soviet Mathematics*, vol. 2, pp. 1226–1228, 1961.
- [5] K. Fan and I. Glicksberg, “Fully convex normed linear spaces,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 41, pp. 947–953, 1955.
- [6] H. Hudzik and B. X. Wang, “Approximative compactness in Orlicz spaces,” *Journal of Approximation Theory*, vol. 95, no. 1, pp. 82–89, 1998.
- [7] H. Hudzik, W. Kowalewski, and G. Lewicki, “Approximate compactness and full rotundity in Musielak-Orlicz spaces and Lorentz-Orlicz spaces,” *Zeitschrift für Analysis und ihre Anwendungen*, vol. 25, no. 2, pp. 163–192, 2006.
- [8] S. Chen, H. Hudzik, W. Kowalewski, Y. Wang, and M. Wisła, “Approximative compactness and continuity of metric projector in Banach spaces and applications,” *Science in China, Series A: Mathematics*, vol. 51, no. 2, pp. 293–303, 2008.
- [9] E. V. Oshman, “Characterization of subspaces with continuous metric projection into a normed linear space,” *Soviet Mathematics*, vol. 13, pp. 1521–1524, 1972.
- [10] J. Aubin and H. Frankowska, *Set-Valued Analysis*, vol. 2, Birkhäuser, Boston, Mass, USA, 1990.
- [11] R. R. Phelps, *Convex Functions, Monotone Operators and Differentiability*, vol. 1364 of *Lecture Notes in Mathematics*, Springer, New York, NY, USA, 1989.
- [12] M. D. Contreras and R. Paya, “On upper semicontinuity of duality mappings,” *Proceedings of the American Mathematical Society*, vol. 121, no. 2, pp. 451–459, 1994.
- [13] S. Shang, Y. Cui, and Y. Fu, “Smoothness and approximative compactness in Orlicz function spaces,” *Banach Journal of Mathematical Analysis*, vol. 8, no. 1, pp. 26–38, 2014.
- [14] S. Shang and Y. Cui, “Approximative compactness and Asplund property in Banach function spaces and in Orlicz-BOchner spaces in particular with application,” *Journal of Mathematical Analysis and Applications*, vol. 421, no. 2, pp. 1377–1395, 2015.
- [15] S. T. Chen, *Geometry of Orlicz Spaces*, Dissertationes Mathematicae, Istitute of Mathematics, Warszawa, Poland, 1996.
- [16] M. Z. Nashed and G. F. Votruba, “A unified approach to generalized inverses of linear operators: II Extremal and proximal properties,” *Bulletin of the American Mathematical Society*, vol. 80, pp. 831–835, 1974.
- [17] Y. M. Wei, W. Xu, S. Z. Qiao et al., “Componentwise condition numbers for the problems of Moore-Penrose generalized matrix inversion and linear least squares,” *Numerical Mathematics: A Journal of Chinese Universities (English Series)*, vol. 14, no. 3, pp. 277–286, 2005.
- [18] H. Hudzik, Y. W. Wang, and W. Zheng, “Criteria for the metric generalized inverse and its selections in Banach spaces,” *Set-Valued Analysis*, vol. 16, no. 1, pp. 51–65, 2008.