Research Article

Legume Lectin FRIL Preserves Neural Progenitor Cells in Suspension Culture In Vitro

Hailei Yao, Xiaoyan Xie, Yanhua Li, Dongmei Wang, Shu Han, Shuangshuang Shi, Xue Nan, Cixian Bai, Yunfang Wang, and Xuetao Pei

Stem Cells and Regenerative Medicine Lab, Beijing Institute of Transfusion Medicine, Beijing 100850, China

Correspondence should be addressed to Xuetao Pei, peixt@nic.bmi.ac.cn

Received 25 February 2008; Accepted 3 May 2008

Recommended by Yang Liu

In vitro maintenance of stem cells is crucial for many clinical applications. Stem cell preservation factor FRIL (Flt3 receptor-interacting lectin) is a plant lectin extracted from *Dolichos Lablab* and has been found preserve hematopoietic stem cells in vitro for a month in our previous studies. To investigate whether FRIL can preserve neural progenitor cells (NPCs), it was supplemented into serum-free suspension culture media. FRIL made NPC grow slowly, induced cell adhesion, and delayed neurospheres formation. However, FRIL did not initiate NPC differentiation according to immunofluorescence and semiquantitative RT-PCR results. In conclusion, FRIL could also preserve neural progenitor cells in vitro by inhibiting both cell proliferation and differentiation.

Copyright © 2008 Hailei Yao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Stem cells are unique cells that retain the ability to divide and proliferate throughout postnatal life to provide progenitor cells that can progressively become committed to produce specialized cells. Many cytokines have been reported as inducers of stem cell proliferation and differentiation, one of them is FL (Flt3 ligand) which can prolong maintenance of primitive human cord blood cells in stromal-free suspension cultures [1–4]. However, FL-containing cultures still cause the loss of repopulating capacity of primitive cells [5–10].

We have extracted and identified a new legume lectin from Hyacinth bean [11]. It is a ligand of receptor tyrosine kinase Flt3 which is strictly expressed in hematopoietic and neural cell lines [12–14]. Gabriella has reported a new flt3 ligand which has the ability to preserve hematopoietic progenitor cells in vitro for 4 weeks [15]. The gene and protein sequences of our protein are quite similar to Gabriella's protein [11], besides our protein is also capable of long time in vitro preservation of hematopoietic progenitor cells. According to previous reports [15–19], we named our protein FRIL too and found that FRIL could maintain hematopoietic stem cell in G_0/G_1 phase for at least 4 weeks.

Low activation of FRIL's receptor Flt3 is considered the key mechanism through which FRIL preserves cells. Further experiments showed that FRIL inhibited hematopoietic stem/progenitor cell (HSPC) apoptosis by upregulating p53, reduced activation of MAPK and phosphorylation of STAT5, disturbed formation of AP-1 complex, and in the end suppressed the expression/activation of cell cycle proteins [20–22].

Over the last decade, neural stem cell research provided penetrating insights into plasticity and regenerative medicine. Stem cells have been isolated from both embryonic and adult nervous system [23, 24]. Although with the help of extrinsic signaling factors progress has been made in understanding fundamental stem cell properties, processes that determine cell fate or that maintain a cell's primitive state are still unknown. As a following experiment to find out whether FRIL can preserve neural progenitor cells as it does on HSPCs, we first identified the isolated neural progenitor cells (NPCs) and their flt3 expressing levels, then we supplemented FRIL into different NPC suspension culture media, observed cell morphological changes, analyzed cell growth and cell surface marker expression, finally we used semiquantitative RT-PCR to verify the changes in gene expression.

2. MATERIALS AND METHODS

2.1. NPCs isolation and cell culture

E13.5 SD rats (experimental animals department, Peking University, Beijing, China) were sacrificed, then fetal embryonic telencephalon and mesencephalon were dissected under the dissecting microscope, after digestion in the sterilized tube, cells were seeded in 25 cm² flask for general culture and passage. Meanwhile statistically supplemented EGF(20 ng/mL), bFGF(20 ng/mL), and FRIL(300 µg/mL, 1:1000) into basic culture media(98% DMEM/F12 (1:1) and 2% B27/N2) to form eight groups of culture media (control, EGF, bFGF, FRIL, EGF&bFGF, EGF&FRIL, bFGF&FRIL, EGF&bFGF&FRIL). After second passage, equal numbers of NPCs were seeded with these different culture media. Furthermore, some NPCs were seeded in differentiation media which contains 5% fetal bovine serum. Culture media comprised a low percent of glucose and did not contain mannose which may exerted negative effect on FRIL's function [15].

2.2. Identification of NPCs and expression of *flt3* on NPCs

P2 neurospheres were transferred into poly-L-lysine (pll) precoated 24-well plate, incubated for 8 hours at 37°C, and then stained by nestin for immunofluorescence for the purpose of NPCs identification. Meanwhile, some other neurospheres were treated with HRP conjugated FRIL in order to find out the proportion of *flt3* expressed on NPCs.

2.3. Cell counting assay

Cells were harvested at day 7 following seeding, resuspended with PBS after digestion and centrifuge, and then seeded on 96-well plate together with 100 µL solution and 10 µL CCK-8 (Cat. no. CK04, Dojindo Laboratories Co, Japan.) per well. Plate was incubated at 37°C, for 4 hours, then the absorbance at 450 nm was measured using microplate reader with a reference wavelength at 650 nm (Bio-Rad Model 550). Data was analyzed by SAS 8.2 software.

2.4. Morphological observation

P3 NPCs were harvested and digested into single cell, cells were counted and seeded into 25 cm² flasks with a density of 5×10^4 per mL. Each flask was supplemented with only one media described above, and NPC morphology was observed everyday.

2.5. Immunofluorescence assay

NPCs were seeded into pll precoated 24-well plate and incubated at 37°C for 8 hours, fixed with 4% paraformaldehyde for 20 minutes, washed with PBS 5 minutes three times, 0.3% Triton X-100 20 minutes, PBS 5 minutes three times, then were incubated with primary antibody for 2 hours at 37°C, washed, and incubated with secondary antibody for 30 minutes at 37°C. Primary antibodies that were used

include nestin (Mouse Anti-Nestin Monoclonal Antibody, Cat. no. MAB353, Chemicon, Inc, USA.), GFAP (Cat. no. ZA-0117, Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd.), NSE (RABBIT ANTI-NSE, Cat. no. BA0535, Boster Biotechnology Co. Ltd.), Tubulin (Monoclonal Anti-β-Tubulin III antibody produced in mouse Clone SDL3D10, ascites fluid, Cat. no. T 8660, Sigma-Aldrich Co, USA.), O4 (Mouse Anti-Oligodendrocyte Marker O4 Monoclonal Antibody, Cat. no. MAB345, Chemicon, Inc. USA.). All primary antibodies were diluted with PBS and 10% normal goat serum. Secondary antibodies included FITC (Fluorescein-Conjugated) AffiniPure Goat Anti-Mouse IgG (H+L) (Cat. no. ZF-0312, Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd.), Rhodamine (TRITC)-Conjugated AffiniPure Goat Anti-Mouse IgG (H+L) (Cat. no. ZF-0313, Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd.), Rhodamine (TRITC)-conjugated Affinipure Goat Anti-Rabbit IgG (H+L) (Cat. no. ZF-0316, Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd.). When cells were stained for nestin together with GFAP, Tubulin or NSE, primary and secondary antibodies for each were added simultaneously. 4',6'-Diamidino-2-phenylindole dihydrochloride (DAPI; 1:10000) was used in all fluorescence facilitating cell counts. Images for cell counts were captured from six equally spaced, predetermined areas of each well under fluorescence microscope. NPCs, neurons, glials, and other cells on the images were counted and analyzed.

2.6. Semiquantitative RT-PCR assay

mRNA was isolated using trizol kit according to manufacture protocols, then mRNA was reverse transcribed in a final volume of 100 µL, 42°C 5 minutes, 42°C 1 hour, 94°C 5 minutes, -20°C. Then, *ACTIN* F 5'TACCACTGGCATCG RGATGGACT 3' R 5'TCCTTCTGCATCCTGTCCGCAAT 3' 94°C 30 seconds, 59°C 30 seconds, 72°C 30 seconds, 25 cycles, 72°C 5 minutes, 4°C. *GFAP* F 5'GAAGAAAA CCGCATCACCAT3' R 5'GCACACCTCACATCACATCC3' 94°C 30 seconds, 57°C 30 seconds, 72°C 30 seconds, 25 cycles, 72°C 5 minutes, 4°C. *Nestin* F 5'GGAGCCATTGTG GTCTACTGA3' R 5'TCCCACCGCTGTTGATTT3' 94°C 30sec, 56°C 30 seconds, 72°C 30 seconds, 28 cycles, 72°C 5 minutes, 4°C. *NeuroD1* F 5'ACCTGCTGCCAGAGTTTTA 3' R 5'CAGAGGCTACCGAGGACTTG3' 94°C 30 seconds, 59°C 30 seconds, 72°C 30 seconds, 35 cycles, 72°C 5 minutes, 4°C. *NeuroD2* F 5'GGGACTCGCCTTCTCTCTCT3' R 5'CTATCCCCGAACTCAGCAG3' 94°C 30 seconds, 59°C 30 seconds, 72°C 30 seconds, 40 cycles, 72°C 5 minutes, 4°C. *NeuroD3* F 5'CAGTAGTCCCTCGGCTTCAG3' R 5'TAGACTGGGGCAGGAAAGAA3' 94°C 30 seconds, 59°C 30 seconds, 72°C 30 seconds, 40 cycles, 72°C 5 minutes, 4°C. *Ngn3* F 5'AGGGGACACACGATTAGCAG3' R 5'GGTCTCTTGGGACACTTGA3' 94°C 30 seconds, 59°C 30 seconds, 72°C 30 seconds, 38 cycles, 72°C 5 minutes, 4°C. *Notch* F 5'GGTGGACATTGACGAGTGTG3' R 5'CCCTTGAGGCATAAGCAGAG3' 94°C 30 seconds, 59°C 30 seconds, 72°C 30 seconds, 27 cycles, 72°C 5 minutes, 4°C. 20 µL of each samples of PCRs were run on 10% TAE PAGE stained with ethidium bromide. Gel image were saved as

FIGURE 1: Neural progenitor cell identification and Flt3 expression. (a) Neurospheres under bright field. (b) Flt3-HRP staining on neural spheres. (c) nestin immunofluorescence on neurospheres, RED nestin, BLUE DAPI. (d) Nestin immunofluorescence on single NPC, GREEN nestin, BLUE DAPI.

FIGURE 2: Cell counting assay. Cells were cultured in eight media with different growth factors combination represented in figure by control, FRIL, bFGF, bFGF&FRIL, EGF, EGF&FRIL, bFGF&EGF, bFGF&EGF&FRIL. Cell numbers were transferred into absorption value obtained from microplate reader Bio-Rad Model 550.

TIFF files using AlphaImager3300 (Serial no. 981878, Alpha Inc, China.). Gel band quantitation analysis was performed via analysis tools of AlphaImager3300.

3. RESULTS

3.1. NPCs identification, flt3 expression, and neural progenitor cell growth assay

Primer (P0) cells began to form regular spheres on day 2 after dissection (Figure 1(a)), although some cell death was evident by the presence of cell fragments. More than 98% of the P2 cells were nestin-positive, indicating they were NPCs (Figures 1(c), 1(d)). While more than 95% of cells were positive for FRIL-HRP stain, indicating that most NPCs expressed flt3 protein (Figure 1(b)).

Then it came to the NPCs in vitro growth assay. FRIL can promote NPCs growth ($P < .0001$). However, FRIL is

less effective than bFGF ($P < .0001$) and EGF ($P < .0001$) (Figure 2).

3.2. FRIL delayed formation of neurospheres and induced cell adhesion

When seeded, NPCs formed spheres later in media with FRIL than those without FRIL. Although there were neurospheres in FRIL containing media, these spheres were smaller than those formed in media without FRIL (Figures 3(b), 3(d), 3(f), 3(h)). Especially in FRIL medium, there were many single/two cells and some small neurospheres which mostly had 3 or 4 cells (Figure 3(b)).

After P3 NPCs were seeded, there was no difference on cell growth in media with/without FRIL from day 1 to day 4 after seeding. However, after day 4, cells in FRIL containing media adhered to the flask surface (Figure 3). And this phenomenon took place again when NPCs were cultured after passage.

NPCs cultured in differentiation media showed difference from day 3 to day 7 after seeding. Cells in serum media seemed to begin to grow on day 3 and covered the bottom of well/flask on day 6/day 7, while cells in FRIL and serum containing media still spotted on the bottom or just formed some colonies. However, from day 7 on cells in FRIL and serum-containing media reached confluency. Most cells showed glial-like morphology. 1~3 neuron-like cells per well existed in serum-containing media, while 13~21 cells were present in FRIL and serum-containing media (data not shown).

3.3. FRIL reduced GFAP expression while maintained NSE expression

We immunostained cells, nestin for NPCs, GFAP for astrocytes, NSE/tubulin for neurons, and O4 for oligodendrocytes. Fewer FRIL treated NPCs differentiated into astrocytes than those not treated by FRIL, despite the fact that they had adhered to the plates (Figure 4). Oligodendrocytes were not found. NSE positive cells were not found in culture media containing EGF and EGF&FRIL too. The number of NSE positive cells in FRIL containing media was slightly higher than that without FRIL, although NSE positive rates were

FIGURE 3: morphological observation of NPCs cultured in different media. (a) NPCs in control medium. (b) NPCs in FRIL medium. (c) NPCs in bFGF medium. (d) NPCs in bFGF&FRIL medium. (e) NPCs in EGF medium. (f) NPCs in EGF&FRIL medium. (g) NPCs in bFGF&EGF medium. (h) NPCs in bFGF&EGF&FRIL medium.

FIGURE 4: GFAP and nestin immunofluorescence on NPCs cultured in different media. (a) NPCs in control medium. (b) NPCs in FRIL medium. (c) NPCs in bFGF medium. (d) NPCs in bFGF&FRIL medium. (e) NPCs in EGF medium. (f) NPCs in EGF&FRIL medium. (g) NPCs in bFGF&EGF medium. (h) NPCs in bFGF&EGF&FRIL medium. GREEN nestin, RED GFAP, BLUE DAPI.

FIGURE 5: semiquantitative RT-PCR of different genes in NPCs cultured in different media. Lane: (C) NPCs in control medium. (F) NPCs in FRIL medium. (b) NPCs in bFGF medium. (E) NPCs in EGF medium. (bF) NPCs in bFGF&FRIL medium. (bE) NPCs in bFGF&EGF medium. (EF) NPCs in EGF&FRIL medium. (bEF) NPCs in bFGF&EGF&FRIL medium. (S) NPCs in serum medium. (FS) NPCs in serum&FRIL medium. Row: *ngn3*, neurogenin3. NOTCH, notch. NeuroD3, NeuroD3. NeuroD2, NeuroD2. NeuroD1, NeuroD1. Nestin, nestin. GFAP, GFAP. ACTIN, actin.

small. In addition, tubulin positive cells were not found in any media.

3.4. Semiquantitative RT-PCR analysis affirmed FRIL inhibition in glial differentiation

On the basis of previous findings [25–27], we chose 5 genes, namely, *GFAP* and *Notch* (an indicator of GFAP), *NeuroD1*, *NeuroD2*, *NeuroD3*, and *Ngn3* (which serves as upstream regulator of *NeuroD*) to examine gene expression of cells in eight different cell culture media. *GFAP* and *nestin* were expressed in each group, *nestin* expression was similar, while *GFAP* expression in FRIL containing groups was lower than those without FRIL (Figure 5). *Notch* was not found in control and serum-treated groups, and its expression was quite opposite to that of *GFAP*. Neuronal gene expression was weak, there was no *NeuroD3* expression, *NeuroD2* was only expressed in 3 media, although expression in *NeuroD3* and *Ngn3* was much higher in FRIL and serum media than in serum media. Although FRIL might have no function on neuronal differentiation, these RT-PCR results suggested that FRIL inhibited neural progenitor cells from autodifferentiating into glials.

4. DISCUSSION

98.6% nestin positive cells showed that to some extent all cells at P2 were NPCs. 95.7% *flt3* expression confirmed the similarities between NPCs and HSCs, although this contradicted to Brazel's findings in which *flt3* was expressed only in NPCs in dorsal root ganglions [12]. The similarities

between NPCs and HSCs [28, 29], especially *flt3* expression, gave us hints on FRIL-NPCs study.

Previous studies showed that the plant lectin FRIL was successfully extracted from *Dolichos lablab* using a mannose affinity column [11, 15]. Although mannose or α -methyl α -D-mannoside were reported exerting a negative effects, FRIL had the ability of preserving HSCs in vitro for a long time [21, 22]. In this study, we reported that FRIL could also preserve neural progenitor cells as it does on hematopoietic stem/progenitor cells, it slowed NPC aggregation and growth, on the other hand it inhibited NPC glial autodifferentiation, although it might not affect NPC neuronal autodifferentiation. At present, the common widely studied neural differentiation pathways are Notch, Shh, and so forth [29, 30]. Based on these pathways, both neuronal and glial RT-PCR results are in coincident with the results of immunofluorescence. No matter what kinds of development in which FRIL is involved, the mechanism is still unclear and further researches in signal pathways should be done. Depending on our previous studies that FRIL can affect activities of MAPK, STAT5, and p53, FRIL neural mechanical researches that may be more complicated could be performed.

There have been reports that stem cells can be induced differentiating into neural cells, however cytokines that effectively control these cells are still inadequate. FRIL's ability to functionally preserve hematopoietic and neural cells may significantly extend the range and time for manipulating these cells. Experiments are underway to find out whether FRIL can improve applications in neural stem cell transplantation, treatment of neural degenerative diseases, and cell therapy, and FRIL's properties of functionally select, preserve, and synchronize hematopoietic cell population could also give some hints.

ACKNOWLEDGMENTS

The authors would like to thank Professor Janet Benson (Lovelace Respiratory Research Institute, NM, USA) for her advice on this manuscript revision. X. Pei was supported by National High Technology Research and Development Program of China (no. 2006AA02A107), the Major State Basic Research Program of China (no. 2005CB522702).

REFERENCES

- [1] C. Hannum, J. Culpepper, D. Campbell, et al., "Ligand for FLT3/FLK2 receptor tyrosine kinase regulates growth of haematopoietic stem cells and is encoded by variant RNAs," *Nature*, vol. 368, no. 6472, pp. 643–648, 1994.
- [2] L. S. Rusten, S. D. Lyman, O. P. Veiby, and S. E. W. Jacobsen, "The FLT3 ligand is a direct and potent stimulator of the growth of primitive and committed human CD34⁺ bone marrow progenitor cells in vitro," *Blood*, vol. 87, no. 4, pp. 1317–1325, 1996.
- [3] S. Hudak, B. Hunte, J. Culpepper, et al., "FLT3/FLK2 ligand promotes the growth of murine stem cells and the expansion of colony-forming cells and spleen colony-forming units," *Blood*, vol. 85, no. 10, pp. 2747–2755, 1995.

- [4] H. J. McKenna, K. L. Stocking, R. E. Miller, et al., "Mice lacking flt3 ligand have deficient hematopoiesis affecting hematopoietic progenitor cells, dendritic cells, and natural killer cells," *Blood*, vol. 95, no. 11, pp. 3489–3497, 2000.
- [5] H. G. Drexler, "Expression of FLT3 receptor and response to FLT3 ligand by leukemic cells," *Leukemia*, vol. 10, no. 4, pp. 588–599, 1996.
- [6] J. E. Rasko, D. Metcalf, M. T. Rossner, C. G. Begley, and N. A. Nicola, "The flt3/flk-2 ligand: receptor distribution and action on murine haemopoietic cell survival and proliferation," *Leukemia*, vol. 9, no. 12, pp. 2058–2065, 1995.
- [7] S. D. Lyman, L. James, T. Vanden Bos, et al., "Molecular cloning of a ligand for the flt3/flk-2 tyrosine kinase receptor: a proliferative factor for primitive hematopoietic cells," *Cell*, vol. 75, no. 6, pp. 1157–1167, 1993.
- [8] E. Maraskovsky, E. Daro, E. Roux, et al., "In vivo generation of human dendritic cell subsets by Flt3 ligand," *Blood*, vol. 96, no. 3, pp. 878–884, 2000.
- [9] H. J. McKenna, "Role of hematopoietic growth factors/flt3 ligand in expansion and regulation of dendritic cells," *Current Opinion in Hematology*, vol. 8, no. 3, pp. 149–154, 2001.
- [10] A. Wodnar-Filipowicz, "Flt3 ligand: role in control of hematopoietic and immune functions of the bone marrow," *News in Physiological Sciences*, vol. 18, no. 6, pp. 247–251, 2003.
- [11] X. Xie, C. Xie, J. Li, C.-X. Bai, and X. Pei, "Extraction, purification and function analysis of a new lectin which can maintain the multipotential of hematopoietic stem/progenitor cells in vitro," *Sheng Wu Hua Xue Yu Sheng Wu Wu Li Jin Zhan*, vol. 31, no. 2, pp. 134–140, 2004.
- [12] C. Y. Brazel, M. H. Ducceschi, B. Pytowski, and S. W. Levison, "The FLT3 tyrosine kinase receptor inhibits neural stem/progenitor cell proliferation and collaborates with NGF to promote neuronal survival," *Molecular and Cellular Neuroscience*, vol. 18, no. 4, pp. 381–393, 2001.
- [13] A. Ito, S. Hirota, Y. Kitamura, and S. Nomura, "Developmental expression of flt3 mRNA in the mouse brain," *Journal of Molecular Neuroscience*, vol. 4, no. 4, pp. 235–243, 1993.
- [14] O. deLapeyrière, P. Naquet, J. Planche, et al., "Expression of Flt3 tyrosine kinase receptor gene in mouse hematopoietic and nervous tissues," *Differentiation*, vol. 58, no. 5, pp. 351–359, 1995.
- [15] G. Colucci, J. G. Moore, M. Feldman, and M. J. Chrispeels, "cDNA cloning of FRIL, a lectin from *Dolichos lablab*, that preserves hematopoietic progenitors in suspension culture," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 2, pp. 646–650, 1999.
- [16] O. Kollet, J. G. Moore, R. Aviram, et al., "The plant lectin FRIL supports prolonged in vitro maintenance of quiescent human cord blood CD34⁺CD38^{-low}/SCID repopulating stem cells," *Experimental Hematology*, vol. 28, no. 6, pp. 726–736, 2000.
- [17] L. K. Kari, B. Scott, and C. Yung, "Characterization of FRIL, a mannose binding lectin of *Dolichos lablab* (Hyacinth Bean)." *Journal of Biological Chemistry*, vol. 269, no. 29, pp. 18789–18793, 1994.
- [18] L. R. Gowda, H. S. Savithri, and D. R. Rao, "The complete primary structure of a unique mannose/glucose-specific lectin from field bean (*Dolichos lablab*)," *Journal of Biological Chemistry*, vol. 269, no. 29, pp. 18789–18793, 1994.
- [19] T. W. Hamelryck, J. G. Moore, M. J. Chrispeels, R. Loris, and L. Wyns, "The role of weak protein-protein interactions in multivalent lectin-carbohydrate binding: crystal structure of cross-linked FRIL," *Journal of Molecular Biology*, vol. 299, no. 4, pp. 875–883, 2000.
- [20] C. Xie and X. Pei, "The mechanisms of a new plant lectin FRIL from *Dolichos lablab* that preserves quiescent hematopoietic stem cells in suspension culture," *Sheng li ke xue jin zhan [Progress in physiology]*, vol. 34, no. 2, pp. 127–131, 2003.
- [21] X. Xie, C. Xie, W. Shi, et al., "The maintenance of cord blood CD34⁺ progenitor cells with plant lectin FRIL in vitro and the expression of related cell cycle modulator HTm4 and HTm4S," *Sheng Li Xue Bao*, vol. 56, no. 3, pp. 306–312, 2004.
- [22] X. Xie, C. Xie, J. Li, et al., "Studies on the mechanism that how could lectin FRIL preserve hematopoietic stem/progenitor cells in vitro," *Chinese Journal of Hematology*, vol. 25, pp. 501–504, 2004.
- [23] Y.-C. Hsu, D.-C. Lee, and I.-M. Chiu, "Neural stem cells, neural progenitors, and neurotrophic factors," *Cell Transplantation*, vol. 16, no. 2, pp. 133–150, 2007.
- [24] L. Dang and V. Tropepe, "Neural induction and neural stem cell development," *Regenerative Medicine*, vol. 1, no. 5, pp. 635–652, 2006.
- [25] Y. Sun, M. Nadal-Vicens, S. Misono, et al., "Neurogenin promotes neurogenesis and inhibits glial differentiation by independent mechanisms," *Cell*, vol. 104, no. 3, pp. 365–376, 2001.
- [26] S. J. Morrison, "Neuronal differentiation: proneural genes inhibit gliogenesis," *Current Biology*, vol. 11, no. 9, pp. R349–R351, 2001.
- [27] M. Nieto, C. Schuurmans, O. Britz, and F. Guillemot, "Neural bHLH genes control the neuronal versus glial fate decision in cortical progenitors," *Neuron*, vol. 29, no. 2, pp. 401–403, 2001.
- [28] E. Cau and S. W. Wilson, "Ash1a and Neurogenin1 function downstream of Floating head to regulate epiphyseal neurogenesis," *Development*, vol. 130, no. 11, pp. 2455–2466, 2003.
- [29] V. A. Terskikh, C. M. Easterday, L. Li, et al., "From hematopoiesis to neuropoiesis: evidence of overlapping genetic programs," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 14, pp. 7934–7939, 2001.
- [30] C. R. R. Bjornson, R. L. Rietze, B. A. Reynolds, M. C. Magli, and A. L. Vescovi, "Turning brain into blood: a hematopoietic fate adopted by adult neural stem cells in vivo," *Science*, vol. 283, no. 5401, pp. 534–537, 1999.