Review Article

Blood-Brain Barrier Disruption Induced by Chronic Sleep Loss: Low-Grade Inflammation May Be the Link

**G. Hurtado-Alvarado,¹ E. Domínguez-Salazar,¹ L. Pavon,²
J. Velázquez-Moctezuma,¹ and B. Gómez-González¹**

¹*Area of Neurosciences, Department of Biology of Reproduction, CBS, Universidad Autónoma Metropolitana-Iztapalapa, Mexico City, Mexico*

²*Department of Psychoimmunology, National Institute of Psychiatry “Ramón de la Fuente”, Mexico City, Mexico*

Correspondence should be addressed to B. Gómez-González; bgomezglez@gmail.com

Received 26 March 2016; Revised 14 August 2016; Accepted 24 August 2016

Academic Editor: Douglas C. Hooper

Copyright © 2016 G. Hurtado-Alvarado et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sleep is a vital phenomenon related to immunomodulation at the central and peripheral level. Sleep deficient in duration and/or quality is a common problem in the modern society and is considered a risk factor to develop neurodegenerative diseases. Sleep loss in rodents induces blood-brain barrier disruption and the underlying mechanism is still unknown. Several reports indicate that sleep loss induces a systemic low-grade inflammation characterized by the release of several molecules, such as cytokines, chemokines, and acute-phase proteins; all of them may promote changes in cellular components of the blood-brain barrier, particularly on brain endothelial cells. In the present review we discuss the role of inflammatory mediators that increase during sleep loss and their association with general disturbances in peripheral endothelium and epithelium and how those inflammatory mediators may alter the blood-brain barrier. Finally, this manuscript proposes a hypothetical mechanism by which sleep loss may induce blood-brain barrier disruption, emphasizing the regulatory effect of inflammatory molecules on tight junction proteins.

1. Introduction

Almost all of our knowledge about the effect of inflammatory events on blood-brain barrier is related to chronic diseases or acute events, in which exacerbated responses to pathogens are present. The role of low-grade inflammation in the generation or exacerbation of neuropathologies is recently explored because several conditions such as obesity and diabetes concur with this inflammatory status during long-term periods and, perhaps, it may be related to systemic and central comorbidities. Most, if not all, pathologies are associated with sleep disturbances. Sleep loss *per se*, including sleep deprivation, sleep restriction, or sleep fragmentation (see Table 1 for a full differentiation between the concepts), generates a pathogen-independent low-grade inflammatory status. Here, we will review (1) the inflammatory mediators that increase during periods of sleep loss and their association with general disturbances in peripheral endothelium and epithelium and (2) how those inflammatory mediators might

alter the blood-brain barrier during sleep loss. With the evidence presented in this review, we propose a hypothetical mechanism by which sleep restriction could induce blood-brain barrier disruption, emphasizing the effect of inflammatory molecules on tight junction maintenance.

2. Sleep Loss as an Inflammatory Event

Sleep is one of the most widely observed phenomena in mammals and is recognized to play a vital regulatory role in a number of physiological and psychological systems [1, 2]. The paramount role of sleep in the physiology of animal models and humans is evident by the effects of sleep loss. Serious physiological consequences of sleep loss include decreased neurogenesis, cognitive dysfunction (deficits in learning, memory, and decision-making), metabolic alterations, cardiovascular diseases, immune disturbances, and blood-brain barrier disruption [1–8]. Both chronic and acute sleep loss associate with energy balance disturbances [9] and changes

TABLE 1: Sleep loss procedures.

	Human procedures	Duration	Animal models	Duration
Sleep deprivation (SD)	(i) Shift working [117] (ii) Voluntary SD [117]	(i) Several days (ii) 12–90 h	(i) Modified multiple platform method (REM SD) [118] (ii) Gentle SD (total SD) [119] (iii) Disk-over-water method (total and selective SD) [120]	(i) 3–96 h (ii) 3–96 h (iii) 3–96 h
Sleep restriction (SR)	(i) Voluntarily SR [117]	(i) 3–5 h	(i) Modified multiple platform method [7] (ii) Rotating bar at the bottom of the house-cage [8]	(i) 20 h of SD plus 4 h of daily sleep recovery (ii) 18 h of SD plus 6 h of daily sleep recovery
Sleep fragmentation (SF)	(i) Obstructive apnoea patients [117] (ii) The elderly [117]	(i) Several days (ii) Several days	(i) Gentle manipulation coupled to EEG recording [119] (ii) Disk-over-water method [120]	(i) 1 to several days (ii) 1 to several days

Sleep deprivation consists of sleep loss without sleep opportunity along a short period; sleep restriction consists of a reduction in total sleep time with short periods of sleep opportunity; and sleep fragmentation consists of multiple awakenings during sleep time.

in cellular and humoral immunity [10, 11]; however, the direct mechanism by which sleep induces a low-grade inflammatory status is unclear. Experimental research has demonstrated that acute and chronic sleep loss result in impairments in the immune response, characterized by deficits in the cellular component (both in number and in function) and increased levels of proinflammatory mediators, such as tumor necrosis factor- α (TNF- α), interleukin- 1β (IL- 1β), IL-6, IL-17A, and C-reactive protein (CRP) (for details of the cytokine levels related to varying periods of sleep loss see [12]). In addition to immune-derived inflammatory mediators, sleep loss also increases the levels of other inflammatory molecules such as cyclooxygenase-2 (COX-2) [8], nitric oxide synthase (NOS), endothelin-1 (ET-1), vascular endothelial growth factor (VEGF), and insulin-like growth factor-1 (IGF-1) [8, 13].

The major aim of this review is to discuss the role of low-grade inflammation in the blood-brain barrier disruption induced by sleep loss; nevertheless, because endothelial cells form the blood-brain barrier we considered it relevant also to discuss the effect of sleep loss on peripheral endothelial and epithelial cells as early markers of inflammation.

3. Peripheral Endothelial and Epithelial Disturbances Induced by Sleep Loss

Endothelial and epithelial cells form protecting barriers in the central nervous system but also in the periphery. Several pathological states are known to target peripheral epithelial and/or endothelial barriers; therefore the knowledge of regulatory mechanisms in those peripheral barriers may contribute to improving the understanding of central barriers. Among the pathologies affecting body barriers, those involving infections and also diabetes, cardiovascular diseases, psoriasis, and cancer are associated with sleep disturbances [14–16]. Here, we present evidence regarding the disrupting effect of sleep loss on peripheral epithelial and endothelial cells.

3.1. Humans. When fluid compartmentalization goes awry, homeostasis is altered and the possibility exists of induction of inflammation by microorganism invasion and even of tumor microenvironment induction [14]. In humans sleep restriction increases sympathetic activity and, concomitantly, causes endothelial dysfunction at the venous level [17]; the effect may be mediated via endothelin-1 (ET-1) because ET-1-mediated vasoconstriction is greater in adults with short sleep duration (less than 7 h per night) than in those with normal sleep duration (7–9 h per night) [18]. ET-1 is the most potent vasoconstrictor peptide released by the endothelium. The link between sleep restriction and increased ET-1 activity is not clear, but the role of the inflammatory status induced by sleep loss may partially explain this association. In this way, inflammatory cytokines, insulin, and epinephrine altered during sleep loss have each been shown to increase ET-1 in hypertensive subjects [19]. The cytokines that may increase in sleep-deprived humans (e.g., TNF- α , IL- 1β , and IL-6) raise arterial vascular tone via endothelin receptors [20]. Several

reports indicate that sleep loss induces vascular alterations related to inflammatory markers (for a review see [21]). Some studies have tried to clarify the underlying mechanism; for instance, sleep deprivation in humans induced magnesium deficiency [22], which produces arterial constriction, and is a possible cause of myocardial damage [22]. Other barriers are not yet studied in sleep-deprived or sleep-restricted humans, but some studies indicate that sleep deficiency alters skin conductance [23].

3.2. Animal Models. Animal models currently used in sleep research include those that model shift work by totally sleep depriving rodents; human sleep deficiency by sleep restricting; and sleep loss-associated with pathologies, such as obstructive apnoea, by promoting sleep fragmentation. Contrary to the human studies, in the case of animal models, several studies have identified negative effects of sleep loss on peripheral endothelia and epithelia. For instance, sleep fragmentation in mice (20 weeks) induces vascular endothelial dysfunction and mild blood pressure increases. Those physiological effects are accompanied by morphological vessel changes characterized by elastic fiber disruption and disorganization, increased recruitment of inflammatory cells to the vessel wall, and increased plasma levels of IL-6 [24]. In rats, total sleep deprivation reduces endothelial-dependent cutaneous vasodilation. This endothelial dysfunction is independent of blood pressure and sympathetic activity but is associated with changes in NOS and COX pathways [25].

The effect of sleep loss on physical barriers such as the intestinal barrier or blood-testis barrier is not reported; however, gut bacteria are present in blood after sleep deprivation [26] and both sleep-deprived and sleep-restricted rats exhibit lower sperm viabilities associated with an increase in endothelial NOS expression [27]. Those data suggest that sleep loss also might alter the physiology of the above-mentioned barriers with the ensuing tissue damage.

4. Blood-Brain Barrier Impairment Induced by Sleep Loss

We reported for the first time that sleep restriction induces blood-brain barrier hyperpermeability in rats [7]. We used a procedure consisting of 20-hour sleep deprivation plus 4 hours of sleep opportunity during 10 consecutive days; because a reduction in total sleep time is observed, it is named sleep restriction. In our conditions, rapid eye movement (REM) sleep is fully suppressed and non-REM sleep is 30% reduced since the first day of sleep restriction. In those conditions we showed a widespread breakdown of the blood-brain barrier [7]. We described that brief periods of sleep opportunity (40 to 120 minutes) induced a progressive recovery of blood-brain barrier permeability to Evans blue (>60 000 Da) in the majority of brain regions studied, with exception of the hippocampus and cerebellum [7]. We also observed that in the hippocampus the number of pinocytic vesicles increased threefold. In a subsequent study, mice were subjected to sleep restriction for 6 days in a rotatory bar for 12 hours per day. Sleep restriction by this method induced REM

sleep loss in the first 3 days with partial REM sleep recovery afterwards; at the end of the 6th day of sleep restriction, there was 13.3% increase of wakefulness, 10.2% reduction of non-REM sleep, and 2.1% reduction of REM sleep [8]. Under these conditions, increased blood-brain barrier permeability to sodium fluorescein, a low molecular-weight tracer, was observed; sleep recovery by 24 hours fully reverted the effect. In the same way, sleep restriction decreased the mRNA levels of the tight junction proteins claudin-5, zonula occludens-2 (ZO-2), and occludin [8]. In the first study [7] a yoked control was included to avoid any potential confounding effects of stress on blood-brain barrier permeability; rats were placed on large platforms during the same period of time as sleep-restricted subjects and despite being in the same stressful conditions as the sleep-restricted subjects they have a fully functional blood-brain barrier [7]. The second study did not include a yoked control, a newly developed sleep deprivation method was used that involves a rotating bar at the bottom of the house-cage with random changes of direction; this method may certainly be stressful to the rodents due to the presence of forced exercise; however, our recent results replicate their findings (Hurtado-Alvarado et al. personal communication). Therefore, the evidence of changes in the blood-brain barrier integrity induced by sleep loss is substantial and inflammatory molecules appear to play a key role in the mechanism subjacent to this phenomenon.

5. Role of Inflammatory Mediators Released during Sleep Loss in Blood-Brain Barrier Physiology

The increase in the levels of inflammatory mediators during chronic sleep loss may be related to blood-brain barrier disruption because several previous reports show that *per se* those inflammatory molecules affect the integrity of the blood-brain barrier (see Table 2 for a summary).

5.1. Proinflammatory Cytokines Involved in Sleep and Blood-Brain Barrier Modulation

5.1.1. Tumor Necrosis Factor- α . Tumor necrosis factor- α (TNF- α) is a protein synthesized mainly by monocytes and macrophages that plays an essential role in the initial activation of the immune system. In the central nervous system TNF- α is a multipotent cytokine produced by neurons, glia, and microvascular endothelial cells that is implicated in several physiological events, such as memory consolidation and sleep regulation. TNF- α is also a potent regulator of blood-brain barrier permeability. The role of TNF- α as an inducer of blood-brain barrier disruption includes its overexpression in microglia, astrocytes, and microvascular endothelial cells [28].

Several reports indicate that sleep loss increases the plasma and brain levels of TNF- α [29–33], the mRNA expression of TNF- α in the brain [33, 34], the spontaneous production of TNF- α in lymphocytes [35], and the mRNA expression of TNF- α in peritoneal and epididymal adipose tissue [36, 37]. Despite the fact that the changes in TNF- α

induced by sleep loss are 2 to 5 times higher compared to rats sleeping *ad libitum*, the levels are below those reported in the case of infectious diseases; however, the chronic exposure to this inflammatory mediator may underlie the sleep-induced blood-brain barrier dysfunction.

The effect of TNF- α in endothelial cells is well studied. *In vivo* and *in vitro* studies report an increase in the permeability of microvascular endothelial cells after the administration of TNF- α in both animal models and human cell lines [38–41]. Nonetheless, the TNF- α levels used in those studies are 100,000 times higher compared to concentrations reported under sleep loss conditions. The lower dose of TNF- α used in *in vitro* studies (1 ng/mL) results in a transendothelial electric resistance (TEER) reduction at 60 minutes after treatment with TEER recovery at 210 minutes after administration, which is similar to the results observed using higher doses of TNF- α (50, 100 ng/mL), suggesting that the effect mediated by TNF- α receptors is saturable [42].

While we can infer that peripheral changes mediate the main effect of TNF- α on blood-brain barrier, we must not ignore the fact that TNF- α levels also increase in the brain. In this way, it is known that after the administration of TNF- α (250 ng) in the lateral ventricle an increase in the transport from cerebrospinal fluid (CSF) to blood of 125 I-human serum albumin is observed in rats, which demonstrates that TNF- α promotes the clearance of macromolecules from the CSF to the venous blood [43]. Taking into consideration that the restorative function of non-REM sleep may be a consequence of the enhanced removal of waste products accumulated in the awaking brain via the glymphatic system [44], the TNF- α increase during sleep loss may contribute to the clearance of toxins by efflux of potentially neurotoxic waste products via the blood-brain barrier. Interestingly, in the brain, sleep restriction increases the mRNA expression of TNF- α in a region-dependent manner in the mouse [45], suggesting that if TNF- α regulates the microvascular brain endothelial cells from inside the brain, it may do it in specific areas, such as the somatosensory and frontal cortices, which indicates that blood-brain barrier regulation by inflammatory molecules is heterogeneous (a finding reported by us in the case of blood-brain barrier changes induced by sleep loss and recovery; see [7]).

Another example of TNF- α role in blood-brain barrier regulation during peripheral inflammation occurs after the induction of acute pancreatitis in rats, where an increase in TNF- α levels is observed as early as 6 hours after pancreatitis induction and at the same time increases the blood-brain barrier permeability to sodium fluorescein (365 Da) in the hippocampus and cerebellum as well as to Evans blue in the hippocampus, basal nuclei, and cerebellum. In the case of the low molecular-weight tracer the normal blood-brain barrier permeability reestablishes at 24 hours after induction, while, for Evans blue, reestablishment occurs 48 hours after induction [46]. We also observed region-dependent effects of sleep loss and recovery on blood-brain barrier integrity; for instance, in the cerebellum the hyperpermeability remained even after sleep opportunity periods of 40–120 minutes; meanwhile the cortex recovered the normal blood-brain barrier permeability at the same time points [7]. Therefore,

TABLE 2: Inflammatory mediators released during sleep loss that may potentially regulate blood-brain barrier integrity.

Inflammatory mediator	General changes during sleep loss	General effects on blood-brain barrier
TNF- α	↑ circulating levels in human and rodents [29–32]	↑ blood-brain barrier permeability in <i>in vivo</i> and <i>in vitro</i> models (rodent and human brain endothelial cells) [38–40]
	↑ mRNA expression in mice brain [33]	↑ efflux of albumin from brain to blood [43]
		↓ ZO-1 expression [103]
IL-1 β		↑ MMP-9 protein expression [106]
	↑ circulating levels in human and rodents [3, 4, 29, 49]	↑ blood-brain barrier permeability in <i>in vivo</i> and <i>in vitro</i> models (rodent and human brain endothelial cells) [42, 54]
	↑ mRNA expression in mice brain [45]	↓ TEER of primary cultures of brain endothelial cells and human brain endothelial cells [42, 54]
IL-6		↑ production of PGE and COX [57]
	↑ circulating levels in human after chronic sleep loss [64, 65]	↓ ZO-1 expression [103]
	↓ circulating levels in humans [62]	↓ TEER in cerebrovascular endothelial cells from rats at higher doses but not at lower doses [42]
IL-17A	↑ circulating levels during sleep recovery in humans [69]	↓ blood-brain barrier permeability in ischemic brain in rodents [71]
	↑ mRNA expression in human PBMC [50, 67]	
	↑ circulating levels in rodents [29]	
CRP	↑ mRNA expression in human PBMC [50]	↑ blood-brain barrier permeability in <i>in vivo</i> and <i>in vitro</i> models (rodent and human brain endothelial cells) [78, 79]
		↑ blood-brain barrier permeability in <i>in vivo</i> and <i>in vitro</i> models (rodent and human brain endothelial cells) [85]
	↑ circulating levels in humans and rodents [4, 8, 50, 67, 81–83]	↑ ROS production in brain endothelial cells [86]

TNF: tumor necrosis factor; IL: interleukin; CRP: C-reactive protein; ZO: zonula occludens; MMP-9: matrix metalloproteinase-9; PBMC: peripheral blood mononuclear cells; ROS: reactive oxygen species; COX: cyclooxygenase; and TEER: transendothelial electric resistance.

the cerebellum could be considered as a highly susceptible region to inflammatory mediators such as TNF- α [47] in comparison with other brain regions (e.g., the hippocampus and cortex). The differential distribution of TNF- α receptors in the brain may explain why TNF- α regulates blood-brain barrier function in a region-dependent manner; however, it is also possible that other molecules may have synergic effects with TNF- α to regulate blood-brain barrier physiology.

5.1.2. Interleukin-1 Beta. IL-1 β is the prototypical signal molecule for neuroimmune communication. Classically, phagocytic cells in response to inflammatory stimuli release IL-1 β ; in the brain IL-1 β activates the regions involved in the generation of hyperthermia [48]. Similar to the effect of TNF- α , IL-1 β administration promotes sleep in mammals [1] and sleep deprivation has been shown to increase serum IL-1 β levels both in humans and in animal models [3, 4, 29, 49]. In addition, sleep loss induces IL-1 β gene expression in the brain [34, 45], cardiac muscle, and adipose tissue [36] and on phytohaemagglutinin (PHA) activated peripheral blood mononuclear cells (PBMC) [50]. In the case of the brain, several reports indicate that the expression of the IL-1 receptor-1 (IL-1R1) in endothelial cells is high in the pre-optic area, subfornical organ, and supraoptic hypothalamus, while a lesser expression is found in the paraventricular hypothalamus, cerebral cortex, nucleus of the solitary tract, ventrolateral medulla, trigeminal and hypoglossal motor nuclei, and the area postrema [51–53].

In *in vitro* models of blood-brain barrier, IL-1 β (in doses of 5, 100, and 1000 ng/mL) decreases the TEER similar to the levels observed after TNF- α administration [42, 54]. IL-1 β also promotes the release of IL-6 and prostaglandin E (PGE₂) in rat brain endothelial cells [55]. Likely, *in vivo* studies have shown that IL-1 induces sickness behaviour mediated by endothelial IL-1R1 activation in rats [56]; the probable mechanism may be the induction of COX-2 in brain endothelial cells after IL-1R1 activation with the concomitant increase in the synthesis of PGE₂ [57].

IL-1 β may have a key role in blood-brain barrier dysfunction during sleep loss because it has been reported that sleep loss increases IL-1 β gene expression in the cerebral cortex, hippocampus, and basal forebrain [45]. In addition, IL-1 β released from activated microglia increases blood-brain barrier permeability; this effect may depend on the suppression of astrocyte-derived signals that maintain blood-brain barrier integrity (e.g., sonic hedgehog, SHH) [58]. IL-1 β action on blood-brain barrier may induce the expression of other inflammatory mediators produced by microglia and astroglia. For instance, the lack of IL-1R1 specifically in endothelial cells precluded the brain increase of IL-1 β , TNF- α , and IL-6 in stressed rats despite the presence of reactive microglia [59, 60], which places IL-1 β and its receptor on endothelial cells as central mediators of brain inflammatory responses. Hence, the role of IL-1 β in blood-brain barrier could be mainly related to endothelial-glial interactions [61].

5.1.3. Interleukin-6. Sleep onset is associated with an increase in circulating levels of IL-6 [62]; nevertheless, the potential

role of IL-6 in sleep regulation is controversial, and it may take a secondary role as compared to its primary role in the acute-phase response [63]. Some studies indicate an increase of IL-6 circulating levels in sleep-deprived subjects [64–66] and also in gene expression in immune cells [35, 50, 67], whereas others report a delay in the sleep-related peak of plasma IL-6 in sleep-restricted subjects [62]. Even some authors report that plasma levels of IL-6 are maintained without change despite sleep loss [30, 68]. Some studies also show that sleep recovery after total sleep deprivation increases plasma levels of IL-6 [69]; however, others found that in immune cells IL-6 levels remain unchanged during sleep recovery [50]. IL-6 is a pleiotropic cytokine key for immune regulation and if secreted during sleep loss and recovery may have neuroprotective effects; indeed, it has been reported that IL-6 appears to be neuroprotective and is involved in endothelial survival after shear stress [70]. However, given the high variability of IL-6 after sleep loss and recovery, the role of IL-6 as a possible modulator of blood-brain barrier during sleep is unclear. It is necessary to elucidate the precise changes in IL-6 levels both centrally and peripherally to clarify the role of IL-6 in blood-brain barrier modulation during sleep.

IL-6 has pyrogenic effects when endogenously released during systemic inflammation; it achieves this function by its binding to IL-6 receptor α (IL-6R α) on brain endothelial cells and the subsequent induction of PGE synthesis. However, those effects require high levels of IL-6 (>1 ng/mL). In humans, IL-6 serum levels were less than 100 pg/mL and the normal levels for IL-6 in CSF are around 10 pg/mL, significantly lesser than those measured in several *in vitro* and *in vivo* experiments [70]. For instance, treatment with 50 or 500 ng of IL-6 reduced the infarct volumes and symptoms of neurological deficit in a rat model of cerebral ischemia [71]. In addition, the administration of IL-6 decreased the blood-brain barrier permeability to Evans blue by suppressing the expression of matrix metalloproteinase-9 (MMP-9) [71]. The role of IL-6 as well as TNF- α and IL-1 β may depend on the brain region, for example, the stimulation with lipopolysaccharide (LPS) induces in the brain the expression of the IL-6 receptor (IL-6R) in the cortex and hippocampus but not in the cerebellum [72]. Therefore, considering IL-6 a proinflammatory cytokine it is possible to suggest that its role in blood-brain barrier physiology during sleep loss may be related to the modulation of the expression of other proinflammatory cytokines.

5.1.4. Interleukin-17A. Th17 cells have been identified as a subset of T helper lymphocytes characterized by the production of a number of cytokines including IL-17A, IL-17F, and IL-22. Th17 cells have emerged as a key factor in the pathogenesis of autoimmune disorders. For instance, high expression of IL-17A is associated with autoimmune inflammatory diseases including multiple sclerosis [73], rheumatoid arthritis [74], inflammatory bowel disease [75], and systemic lupus erythematosus [76]. During sleep loss a subtle increase of IL-17A is reported (from 0.5 to 3 ng/mL in rat) [29]. IL-17A high levels were found in plasma even after 24 hours of sleep recovery in sleep-restricted rats [29]. Sleep loss also

increases the mRNA and protein expression of IL-17A on PHA activated PBMC in humans [50].

Particularly, the receptor for IL-17A is expressed in epithelial and endothelial cells and promotes the expression of inflammatory mediators such as IL-6 and chemokines [77]. IL-17A induces epithelial and endothelial dysfunction; it decreases the TEER and concomitantly increases tracer permeability; the mechanism is mediated through tight junction disruption [77]. Finally, from *in vitro* experiments it is known that IL-17A increases endothelial cell permeability at 10 or 100 ng/mL doses [78, 79]. These data suggest that IL-17A might be involved in blood-brain barrier disruption during sleep loss.

5.2. Other Inflammatory Molecules Altered during Sleep Loss and Their Role in Blood-Brain Barrier Regulation

5.2.1. C-Reactive Protein. C-reactive protein (CRP) is the major acute-phase protein involved in the resistance to microbes and autoimmune diseases and is an important risk marker of cardiovascular and cerebrovascular disorders. The plasma levels of CRP increase faster and at higher magnitude than other acute-phase proteins [80]. Sleep loss increases the circulating levels of CRP (0.5 µg/mL), which is associated with increased risk of cardiovascular disease and stroke [4, 8, 50, 67, 81, 81–83].

The synthesis of CRP in the liver is controlled by proinflammatory cytokines, including TNF- α , IL-1 β , IL-6, and IL-17A [82, 84]. CRP (10–20 µg/mL) induces blood-brain barrier disruption [85] because brain endothelial cells express high levels of CRP receptors (CD16 and CD32) and also because brain endothelial cells express high levels of the p22phox subunit of the NAD(P)H-oxidase. The high expression of both exacerbates the generation of reactive oxygen species (ROS) with the resultant oxidation of tight junction proteins [86].

5.2.2. Intercellular Adhesion Molecule-1 (ICAM-1). The expression of ICAM-1 in endothelial cells is pivotal in supporting lymphocyte migration across the vascular endothelium [87]. ICAM-1 associates with an endothelial cytoskeleton fraction, suggesting that ICAM-1 redistribution is an early event in the signalling cascade during inflammatory events, particularly in lymphocyte transmigration [87]. The expression of endothelial cell adhesion molecules increases in the central nervous system during inflammation secondary to pathogen intracerebral administration (e.g., *Corynebacterium parvum*). Brain vessels located in the centre of the cellular infiltrate began to express markers of fenestrate endothelium such as the endothelial-specific expression of MECA32 suggesting an altered functional status of the endothelial cell [88]. Abundant ICAM-1 expression has been observed after IL-1 or TNF- α stimulation of cultured heart endothelial cells [89].

Elevated levels of ICAM-1 may contribute to cardiovascular disease and are associated with obstructive sleep apnoea (OSA) and obesity, in which sleep deficiency is present [90]. In the same way, it has been shown that patients with diabetes mellitus type 2 and poor sleep present higher morbidity of cardiovascular diseases than diabetes mellitus patients

sleeping normally; those patients also present higher plasma levels of ICAM-1 [91]. ICAM-1 higher serum levels were also found during the sleep recovery period after 40 hours of total sleep deprivation in healthy men [69]. Therefore it seems that the mediator between poor sleep (with bad quality and poor sleep recovery) and higher risk for cardiovascular diseases is ICAM-1.

5.2.3. Vascular Endothelial Growth Factor. Inflammation is characterized by upregulation of vascular endothelial growth factor (VEGF). In *in vivo* experiments, increases in VEGF during neuroinflammation (e.g., in experimental autoimmune encephalomyelitis (EAE)) are accompanied with increased blood-brain barrier permeability and decreased expression of tight junction proteins (e.g., claudin-5 and occludin). Likely, VEGF administration to human brain endothelial cells increases permeability of the monolayer and downregulates claudin-5 and occludin, but not junctional adhesion molecule-1 (JAM-1), cingulin, peripheral plasma membrane protein (CASK), or ZO-1 [92].

Given the role of VEGF in regulating blood-brain barrier during neuroinflammation, it may participate in generating the vascular changes associated with sleep loss. Indeed, it has been shown that VEGF is overexpressed in OSA patients and it is generally considered that VEGF increases are associated with hypoxia events [93]. However, OSA patients also have severe sleep fragmentation; therefore, in addition to chronic intermittent hypoxia, VEGF changes may be related to sleep loss [94]. In fact, in a study with major depressive disorder patients, sleep deprivation increased VEGF plasma levels [95].

5.2.4. Insulin-Like Growth Factor-1. Sleep deprivation decreases IGF-1 levels in rats and humans and one night of sleep recovery is sufficient to restore its basal levels [96]. The neuroprotective effects of IGF-1 are unclear but it is known that IGF-1 receptors are present in brain endothelial cells, microglia, and astroglia and even in neurons [97]. Indeed, it has been suggested that IGF-1 may promote neuroprotection by acting on the blood-brain barrier; in an experimental model of ischemic stroke IGF-1 reduced the inflammatory infiltrate in the brain [97]. In an *in vitro* experiment with brain endothelial cells IGF-1 reverted the hyperpermeability to bovine serum albumin induced by oxygen-glucose deprivation (an *in vitro* model of ischemic stroke) [97].

Changes on inflammatory molecules during sleep loss are well described but we do not know what the source of those alterations is. In this way the role of microbiota could appear a good candidate to induce the low-grade proinflammatory status during sleep loss.

6. A Brief View of the Microbiota and Barriers Dysfunction as a Possible Source of Inflammatory Mediators in Sleep-Deficient Subjects

The source of inflammatory mediators during sleep loss remains unclear; however, microbiota may play a key role

in this event. In other conditions that exhibit low-grade systemic inflammation, such as chronic depression, obesity, and diabetes, evidence from murine models initially suggested a role for the gut microbiota in the generation of low-grade inflammation, with the consequent increased risk of endothelial and epithelial dysfunction [98, 99]. For instance, changes in gut microbiota composition increase intestinal permeability [100]. In the same way, during sleep deprivation gut microbiota has been detected in blood, suggesting the induction of systemic inflammation and deficits in gut epithelial permeability [26]. In addition, preclinical evidence from germ-free mice suggests that the microbiota can also modulate the blood-brain barrier; exposure of germ-free adult mice to the faecal microbiota from pathogen-free donors decreased the blood-brain barrier permeability and increased the expression of tight junction proteins in brain endothelial cells [101], therefore strengthening the hypothesis that the blood-brain barrier may also be sensible to changes in the gut microbiota composition [100]. The candidate pathways to induce barriers dysfunction under altered gut microbiota composition include serotonin, cytokines, toll-like receptor activation, and short chain fatty acids [100]. Moreover, the inflammatory response subsequent to microbiota-induced barriers disruption may underlie the sleep loss-related cognitive deficits and the exacerbation of neurological disorders such as depression [100].

These data might support the theory of a coevolution between sleep and blood-brain barrier proposed by Korth in 1995 [102]. Because the brain and blood-brain barrier react sensitively to the exposure to bacterial cell wall constituents and sleep is regulated by gut microbiota products, Korth proposed that low amounts of bacterial cell wall constituents that induce sleep under sleep loss conditions, by themselves or by cytokine production, increase the blood-brain barrier permeability ensuing their passage into the brain [102].

7. Molecular Mechanisms by Which Inflammatory Mediators Might Induce Blood-Brain Barrier Disruption during Sleep Loss

Cytokines and other inflammatory mediators induce blood-brain barrier disruption through mechanisms involving signalling pathways that converge in the disorganization of tight junctions (Figure 1). For instance, it has been reported that proinflammatory cytokines, including TNF- α and IL-1 β , decreased ZO-1 expression and ZO-1-occludin coassociation, concomitant to increased ZO-1 phosphorylation in tyrosine and threonine residues [103]. Those effects are presumably mediated by ROS [103]. ZO-1 phosphorylation in tyrosine residues is also observed after VEGF administration [104]. In this way, VEGF-A also promotes disruption of blood-brain barrier by downregulating the expression of claudin-5 and occludin [92]. Low cytokine concentrations (>1 ng/mL) led to activation of effector caspases via c-Jun N-terminal kinases (JNK) and protein kinase C (PKC) signalling pathways, increased paracellular flux, and redistribution of ZO-1 and VE-cadherin but failed to induce apoptosis [105]. In addition

to caspase-3, TNF- α activates the production of MMP-9 [106], which is also associated with high levels of IL-1 β in brain parenchyma [107].

TNF- α activates the NF κ B signalling pathway, leading to increased PGE levels via COX-2 [108]. COX-2 plays a crucial role in the inflammatory response of the blood-brain barrier (for review see [109]); particularly COX-2 derived PGE₂ increases blood-brain barrier permeability [110]. Other cytokines, such as IL-1, use other signalling pathways that finally converge in COX-2 induction; particularly, the IL-1 receptor-1 (IL-1R1) signals via the p38 mitogen-activated protein kinase (MAPK) and the c-Jun pathway to induce COX-2 synthesis, whereas activation of the IL-6 receptor leads to COX-2 expression through activation of signal transducer and activator of transcription-3 (STAT-3) [111]. The activation of NF κ B by TNF- α and IL-1 β is also correlated with COX-2 expression in microvascular endothelial cells. Indeed, both I κ B α and COX-2 are expressed within the same endothelial cells, suggesting a potential interaction between the transcription factor and COX-2 expression in the cerebral endothelium of animals with systemic inflammation [112].

TNF- α and IL-1 β promote the release of CRP. The putative mechanism by which CRP increases blood-brain barrier permeability is by its action on CD16/CD32 receptors present in the cell membrane of brain endothelial cells [85]. This association activates the Myosin Light Chain (MLC) phosphorylation by MLC-kinase (MLCK) and the activation of p38-MAPK, with the subsequent formation of actin stress fibers [85]. Brain endothelial cells express the p22phox subunit located in the cell membrane; this enzyme uses NADH or NADPH as the electron donor for the single electron reduction of oxygen to produce ROS during CRP stimulation [86]. The assembly of active NADPH oxidase requires translocation of cytosolic subunits, p47phox, p67phox, and Rac1 (a cytosolic GTPase), to the plasma membrane, where they interact with gp91phox and p22phox and associate with other membrane cofactors to form a functional enzyme complex [113]. In addition, CRP stimulation also disorganizes ZO-1 via MLCK and ROS production [85]. In this way, IL-17A also induces NADPH oxidase- or xanthine oxidase-dependent ROS production and downregulates the expression of occludin by activation of MLCK [79].

The signalling of inflammatory mediators and particularly NADPH oxidase may promote the upregulation of adhesion molecules such as ICAM-1 via JAK/epidermal growth factor receptor (EGFR) signalling [113] contributing to a possible leukocyte infiltration. Therefore, these changes may be deemed as the mechanisms involved in brain endothelial cell dysfunction during sleep loss.

8. Conclusion and Future Directions

We propose that inflammatory mediators increased during chronic sleep loss might promote blood-brain barrier disruption (Figures 1 and 2). For aims of clarity the hypothesis does not explicitly distinguish between REM and non-REM sleep and we know that other molecules altered during sleep loss also should be studied because they may have a potent role in the blood-brain barrier disruption such as adenosine

FIGURE 1: Potential inflammatory mediators participating in the regulation of blood-brain barrier permeability during sleep loss. The figure shows the platform method to induce sleep loss in the rat. Chronic sleep restriction increases blood-brain barrier permeability to circulating molecules (e.g., Evans blue) and sleep recovery promotes restoration of normal blood-brain barrier permeability. Inflammatory mediators with barrier regulation properties, such as tumor necrosis factor- α (TNF- α), vascular endothelial growth factor (VEGF), interleukin-1 β (IL-1 β), and IL-17A, are released during sleep loss conditions and some of them return to basal levels during sleep recovery; others, like IL-17A and TNF- α , are maintained at high levels despite sleep recovery. The barrier changes induced by inflammatory mediators may lead to neuroinflammation and potentially may underlie the cognitive impairments induced by sleep loss.

[114] and hormones [115]. In interpreting these data, a number of factors need to be considered. For instance, the cellular components of the blood-brain barrier that promote inflammation in the brain, such as microglia and astroglia, in addition to regulating blood-brain barrier may also be affecting several brain functions during sleep and sleep loss. On the other hand, pericytes have a unique synergistic relationship with brain endothelial cells in the regulation of capillary permeability through secretion of inflammatory mediators including cytokines, chemokines, nitric oxide, and matrix metalloproteinases. Those inflammatory mediators released during sleep restriction may directly induce pericyte detachment from the vessel wall ensuing blood-brain barrier disruption (for review see Hurtado-Alvarado, 2014 [116]).

Summarizing, chronic sleep loss induces systemic low-grade inflammation that may be related to epithelial and

endothelial disturbances both at the systemic and at the central level. Particularly, the role of inflammatory mediators in the blood-brain barrier disruption induced by sleep loss might explain the cognitive impairment associated with sleep loss. The systemic and local effect of inflammatory molecules accumulated during chronic sleep loss should be taken into account for the study of general consequences of sleep deficiency including the risk of developing neurologic and neurodegenerative diseases.

Abbreviations

CSF:	Cerebrospinal fluid
COX:	Cyclooxygenase
CRP:	C-reactive protein
CXCL-1:	Chemokine (C-X-C motif) ligand 1

FIGURE 2: Hypothetical molecular mechanisms mediating sleep loss effect on blood-brain barrier permeability. During sleep loss the increase of soluble inflammatory mediators such as TNF- α , IL-1 β , IL-17A, CRP, and VEGF activates several membrane receptors that converge in cellular pathways hallmark of inflammation, for example, the NF κ B pathway. The final outcome involves the phosphorylation of tight junction proteins and the generation of actin fiber stress. But also other pathways are potentially activated, such as the NADPH oxidase pathway, leading to ROS generation and the subsequent lipoxidation and protein oxidation. The activation of transcription factors (eg., NF κ B) and their translocation to the nuclei may promote the transcription of inflammatory-related genes (eg., ICAM-1, prostaglandins, and matrix metalloproteinases (MMP)) as well as death-related genes (eg., caspase 3) and the repression of genes involved in the maintenance of the barrier properties (eg., claudin-5). Conjointly, all those pathways could lead to increased blood-brain barrier permeability during chronic sleep loss. Cldn5: claudin-5, COX: cyclooxygenase, CRP: C-reactive protein, Fc γ receptor: fragment crystallizable region, ICAM-1: intracellular adhesion molecule-1, IL: interleukin, NADPHox: nicotinamide adenine dinucleotide phosphate oxidase, NF κ B: nuclear factor kappa-light-chain-enhancer, MMP: matrix metalloproteinase, MLC: myosin light chain, MLCK: myosin light chain kinase, PGE: prostaglandin, PKC: protein kinase C, sICAM: soluble ICAM, sTNF α : soluble TNF- α receptor, VEGF: vascular endothelial growth factor, TNF: tumor necrosis factor, and ZO: zonula occludens.

EGFR: Epidermal growth factor receptor
 ET-1: Endothelin-1
 ICAM-1: Intracellular adhesion molecule-1
 IGF-1: Insulin-like growth factor-1
 I κ B α : Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha
 IL: Interleukin
 JNK: c-Jun N-terminal kinase

NADPH: Nicotinamide adenine dinucleotide phosphate
 NF κ B: Nuclear factor kappa-light-chain-enhancer
 NOS: Nitric oxide synthase
 MMP: Matrix metalloproteinase
 MLC: Myosin light chain
 MLCK: Myosin light chain kinase
 PBMC: Peripheral blood mononuclear cells
 PGE: Prostaglandin

PHA: Phytohaemagglutinin
 PKC: Protein kinase C
 REM: Rapid eye movement
 STAT: Signal transducer and activator of transcription
 TEER: Transendothelial resistance
 TNF: Tumor necrosis factor
 VEGF: Vascular endothelial growth factor
 ZO: Zonula occludens.

Competing Interests

The authors declare that there are no competing interests.

Authors' Contributions

G. Hurtado-Alvarado took part in the conception and design of the review, critically revised the manuscript, and took primary responsibility of writing the manuscript. B. Gómez-González took part in the conception of the review and drafted and critically revised the manuscript. L. Pavon, E. Domínguez-Salazar, and J. Velázquez-Moctezuma drafted and critically revised the paper. All authors read and approved the final manuscript.

Acknowledgments

B. Gómez-González received funding from CONACyT (Grant CB-2012-180792). All the authors form part of the “network for the study of the neuro-immune-endocrine alterations induced by sleep loss” supported by PRODEP-SEP.

References

- [1] J. M. Krueger, F. Obál Jr., and J. Fang, “Why we sleep: a theoretical view of sleep function,” *Sleep Medicine Reviews*, vol. 3, no. 2, pp. 119–129, 1999.
- [2] J. M. Siegel, “Do all animals sleep?” *Trends in Neurosciences*, vol. 31, no. 4, pp. 208–213, 2008.
- [3] D. F. Dinges, S. D. Douglas, S. Hamarman, L. Zaugg, and S. Kapoor, “Sleep deprivation and human immune function,” *Advances in Neuroimmunology*, vol. 5, no. 2, pp. 97–110, 1995.
- [4] D. J. Frey, M. Fleshner, and K. P. Wright Jr., “The effects of 40 hours of total sleep deprivation on inflammatory markers in healthy young adults,” *Brain, Behavior, and Immunity*, vol. 21, no. 8, pp. 1050–1057, 2007.
- [5] P. Meerlo, R. E. Mistlberger, B. L. Jacobs, H. C. Heller, and D. McGinty, “New neurons in the adult brain: the role of sleep and consequences of sleep loss,” *Sleep Medicine Reviews*, vol. 13, no. 3, pp. 187–194, 2009.
- [6] J. G. McCoy and R. E. Strecker, “The cognitive cost of sleep lost,” *Neurobiology of Learning and Memory*, vol. 96, no. 4, pp. 564–582, 2011.
- [7] B. Gómez-González, G. Hurtado-Alvarado, E. Esqueda-León, R. Santana-Miranda, J. Á. Rojas-Zamorano, and J. Velázquez-Moctezuma, “REM sleep loss and recovery regulates blood-brain barrier function,” *Current Neurovascular Research*, vol. 10, no. 3, pp. 197–207, 2013.
- [8] J. He, H. Hsueh, Y. He, A. J. Kastin, Y. Wang, and W. Pan, “Sleep restriction impairs blood-brain barrier function,” *The Journal of Neuroscience*, vol. 34, no. 44, pp. 14697–14706, 2014.
- [9] T. Porkka-Heiskanen, K.-M. Zitting, and H.-K. Wigren, “Sleep, its regulation and possible mechanisms of sleep disturbances,” *Acta Physiologica*, vol. 208, no. 4, pp. 311–328, 2013.
- [10] A. Zager, M. L. Andersen, F. S. Ruiz, I. B. Antunes, and S. Tufik, “Effects of acute and chronic sleep loss on immune modulation of rats,” *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology*, vol. 293, no. 1, pp. R504–R509, 2007.
- [11] F. S. Ruiz, M. L. Andersen, R. C. S. Martins, A. Zager, J. D. Lopes, and S. Tufik, “Immune alterations after selective rapid eye movement or total sleep deprivation in healthy male volunteers,” *Innate Immunity*, vol. 18, no. 1, pp. 44–54, 2012.
- [12] G. Hurtado-Alvarado, L. Pavón, S. A. Castillo-García et al., “Sleep loss as a factor to induce cellular and molecular inflammatory variations,” *Clinical & Developmental Immunology*, vol. 2013, Article ID 801341, 14 pages, 2013.
- [13] L. Ibrahim, W. Duncan, D. A. Luckenbaugh, P. Yuan, R. Machado-Vieira, and C. A. Zarate Jr., “Rapid antidepressant changes with sleep deprivation in major depressive disorder are associated with changes in vascular endothelial growth factor (VEGF): a pilot study,” *Brain Research Bulletin*, vol. 86, no. 1–2, pp. 129–133, 2011.
- [14] J. M. Mullin, N. Agostino, E. Rendon-Huerta, and J. J. Thornton, “Keynote review: epithelial and endothelial barriers in human disease,” *Drug Discovery Today*, vol. 10, no. 6, pp. 395–408, 2005.
- [15] T. Anothaisintawee, S. Reutrakul, E. Van Cauter, and A. Thakkinstian, “Sleep disturbances compared to traditional risk factors for diabetes development: systematic review and meta-analysis,” *Sleep Medicine Reviews*, vol. 30, pp. 11–24, 2016.
- [16] M. A. Gupta, F. C. Simpson, and A. K. Gupta, “Psoriasis and sleep disorders: a systematic review,” *Sleep Medicine Reviews*, vol. 29, pp. 63–75, 2016.
- [17] J. L. Dettoni, F. M. Consolim-Colombo, L. F. Drager et al., “Cardiovascular effects of partial sleep deprivation in healthy volunteers,” *Journal of Applied Physiology*, vol. 113, no. 2, pp. 232–236, 2012.
- [18] B. R. Weil, M. L. Mestek, C. M. Westby et al., “Short sleep duration is associated with enhanced endothelin-1 vasoconstrictor tone,” *Canadian Journal of Physiology and Pharmacology*, vol. 88, no. 8, pp. 777–781, 2010.
- [19] E. L. Schiffrin, “Vascular endothelin in hypertension,” *Vascular Pharmacology*, vol. 43, no. 1, pp. 19–29, 2005.
- [20] M. Woods, J. A. Mitchell, E. G. Wood et al., “Endothelin-1 is induced by cytokines in human vascular smooth muscle cells: evidence for intracellular endothelin-converting enzyme,” *Molecular Pharmacology*, vol. 55, no. 5, pp. 902–909, 1999.
- [21] M. Kohansieh and A. N. Makaryus, “Sleep deficiency and deprivation leading to cardiovascular disease,” *International Journal of Hypertension*, vol. 2015, Article ID 615681, 5 pages, 2015.
- [22] B. Takase, T. Akima, A. Uehata, F. Ohsuzu, and A. Kurita, “Effect of chronic stress and sleep deprivation on both flow-mediated dilation in the brachial artery and the intracellular magnesium level in humans,” *Clinical Cardiology*, vol. 27, no. 4, pp. 223–227, 2004.
- [23] J. C. J. Liu, S. Verhulst, S. A. A. Massar, and M. W. L. Chee, “Sleep deprived and sweating it out: the effects of total sleep deprivation on skin conductance reactivity to psychosocial stress,” *Sleep*, vol. 38, no. 1, pp. 155–159, 2015.

- [24] A. Carreras, S. X. Zhang, E. Peris et al., "Chronic sleep fragmentation induces endothelial dysfunction and structural vascular changes in mice," *Sleep*, vol. 37, no. 11, pp. 1817–1824, 2014.
- [25] F. Sauvet, G. Florence, P. Van Beers et al., "Total sleep deprivation alters endothelial function in rats: a nonsympathetic mechanism," *SLEEP*, vol. 37, no. 3, pp. 465–473, 2014.
- [26] C. A. Everson and L. A. Toth, "Systemic bacterial invasion induced by sleep deprivation," *American Journal of Physiology—Regulatory Integrative and Comparative Physiology*, vol. 278, no. 4, pp. R905–R916, 2000.
- [27] T. A. Alvarenga, C. Hirotsu, R. Mazaro-Costa, S. Tufik, and M. L. Andersen, "Impairment of male reproductive function after sleep deprivation," *Fertility and Sterility*, vol. 103, no. 5, pp. 1355–1362.e1, 2015.
- [28] T. Deboer, A. Fontana, and I. Tobler, "Tumor necrosis factor (TNF) ligand and TNF receptor deficiency affects sleep and the sleep EEG," *Journal of Neurophysiology*, vol. 88, no. 2, pp. 839–846, 2002.
- [29] S. Yehuda, B. Sredni, R. L. Carasso, and D. Kenigsbuch-Sredni, "REM sleep deprivation in rats results in inflammation and interleukin-17 elevation," *Journal of Interferon and Cytokine Research*, vol. 29, no. 7, pp. 393–398, 2009.
- [30] M. Chennaoui, F. Sauvet, C. Drogou et al., "Effect of one night of sleep loss on changes in tumor necrosis factor alpha (TNF- α) levels in healthy men," *Cytokine*, vol. 56, no. 2, pp. 318–324, 2011.
- [31] D. P. Venancio and D. Suchecki, "Prolonged REM sleep restriction induces metabolic syndrome-related changes: mediation by pro-inflammatory cytokines," *Brain, Behavior, and Immunity*, vol. 47, pp. 109–117, 2015.
- [32] A. Wolkow, B. Aisbett, J. Reynolds, S. A. Ferguson, and L. C. Main, "Acute psychophysiological relationships between mood, inflammatory and cortisol changes in response to simulated physical firefighting work and sleep restriction," *Applied Psychophysiology and Biofeedback*, vol. 41, no. 2, pp. 165–180, 2016.
- [33] V. Ramesh, D. Nair, S. X. L. Zhang et al., "Disrupted sleep without sleep curtailment induces sleepiness and cognitive dysfunction via the tumor necrosis factor- α pathway," *Journal of Neuroinflammation*, vol. 9, article 91, 2012.
- [34] Q. Zhao, X. Xie, Y. Fan et al., "Phenotypic dysregulation of microglial activation in young offspring rats with maternal sleep deprivation-induced cognitive impairment," *Scientific Reports*, vol. 5, article 9513, 2015.
- [35] M. R. Irwin, T. Witaran, M. Caudill, R. Olmstead, and E. C. Breen, "Sleep loss activates cellular inflammation and signal transducer and activator of transcription (STAT) family proteins in humans," *Brain, Behavior, and Immunity*, vol. 47, pp. 86–92, 2015.
- [36] J. E. Dumaine and N. T. Ashley, "Acute sleep fragmentation induces tissue-specific changes in cytokine gene expression and increases serum corticosterone concentration," *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology*, vol. 308, no. 12, pp. R1062–R1069, 2015.
- [37] E. M. de Oliveira, B. Visniauskas, S. Sandri et al., "Late effects of sleep restriction: potentiating weight gain and insulin resistance arising from a high-fat diet in mice," *Obesity*, vol. 23, no. 2, pp. 391–398, 2015.
- [38] G. A. Rosenberg, E. Y. Estrada, J. E. Dencoff, and W. G. Stetler-Stevenson, "Tumor necrosis factor- α -induced gelatinase B causes delayed opening of the blood-brain barrier: an expanded therapeutic window," *Brain Research*, vol. 703, no. 1-2, pp. 151–155, 1995.
- [39] K. S. Mark and D. W. Miller, "Increased permeability of primary cultured brain microvessel endothelial cell monolayers following TNF- α exposure," *Life Sciences*, vol. 64, no. 21, pp. 1941–1953, 1999.
- [40] J. Peng, F. He, C. Zhang, X. Deng, and F. Yin, "Protein kinase C- α signals PI15RhoGEF phosphorylation and RhoA activation in TNF- α -induced mouse brain microvascular endothelial cell barrier dysfunction," *Journal of Neuroinflammation*, vol. 8, article 28, 2011.
- [41] J. Matsumoto, F. Takata, T. Machida et al., "Tumor necrosis factor- α -stimulated brain pericytes possess a unique cytokine and chemokine release profile and enhance microglial activation," *Neuroscience Letters*, vol. 578, pp. 133–138, 2014.
- [42] H. E. de Vries, M. C. M. Blom-Roosemalen, M. van Oosten et al., "The influence of cytokines on the integrity of the blood-brain barrier in vitro," *Journal of Neuroimmunology*, vol. 64, no. 1, pp. 37–43, 1996.
- [43] J. B. Dickstein, H. Moldofsky, and J. B. Hay, "Brain-blood permeability: TNF-alpha promotes escape of protein tracer from CSF to blood," *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology*, vol. 279, no. 1, pp. R148–R151, 2000.
- [44] L. Xie, H. Kang, Q. Xu et al., "Sleep drives metabolite clearance from the adult brain," *Science*, vol. 342, no. 6156, pp. 373–377, 2013.
- [45] M. R. Zielinski, Y. Kim, S. A. Karpova, R. W. McCarley, R. E. Strecker, and D. Gerashchenko, "Chronic sleep restriction elevates brain interleukin-1 beta and tumor necrosis factor-alpha and attenuates brain-derived neurotrophic factor expression," *Neuroscience Letters*, vol. 580, pp. 27–31, 2014.
- [46] G. Farkas, J. Márton, Z. Nagy et al., "Experimental acute pancreatitis results in increased blood-brain barrier permeability in the rat: a potential role for tumor necrosis factor and interleukin 6," *Neuroscience Letters*, vol. 242, no. 3, pp. 147–150, 1998.
- [47] C. Silwedel and C. Förster, "Differential susceptibility of cerebral and cerebellar murine brain microvascular endothelial cells to loss of barrier properties in response to inflammatory stimuli," *Journal of Neuroimmunology*, vol. 179, no. 1-2, pp. 37–45, 2006.
- [48] D. L. Bornstein, "Leukocytic pyrogen: a major mediator of the acute phase reaction," *Annals of the New York Academy of Sciences*, vol. 389, no. 1, pp. 323–337, 1982.
- [49] J. Hu, Z. Chen, C. P. Gorczynski et al., "Sleep-deprived mice show altered cytokine production manifest by perturbations in serum IL-1ra, TNFa, and IL-6 levels," *Brain, Behavior, and Immunity*, vol. 17, no. 6, pp. 498–504, 2003.
- [50] W. M. A. van Leeuwen, M. Lehto, P. Karisola et al., "Sleep restriction increases the risk of developing cardiovascular diseases by augmenting proinflammatory responses through IL-17 and CRP," *PLoS ONE*, vol. 4, no. 2, Article ID e4589, 2009.
- [51] A. Ericsson, C. Liu, R. P. Hart, and P. E. Sawchenko, "Type 1 interleukin-1 receptor in the rat brain: distribution, regulation, and relationship to sites of IL-1-induced cellular activation," *Journal of Comparative Neurology*, vol. 361, no. 4, pp. 681–698, 1995.
- [52] J. P. Konsman and A. Blomqvist, "Brain blood vessel distribution of interleukin-1 type 1 receptor corresponds to patterns of jun phosphorylation and cyclooxygenase-2 induction by inflammatory stimuli," *Society for Neuroscience*, vol. 27, pp. 843–814, 2001.
- [53] J. P. Konsman, S. Vignes, L. Mackerlova, A. Bristow, and A. Blomqvist, "Rat brain vascular distribution of interleukin-1

- type-1 receptor immunoreactivity: relationship to patterns of inducible cyclooxygenase expression by peripheral inflammatory stimuli," *Journal of Comparative Neurology*, vol. 472, no. 1, pp. 113–129, 2004.
- [54] R. A. Skinner, R. M. Gibson, N. J. Rothwell, E. Pinteaux, and J. I. Penny, "Transport of interleukin-1 across cerebrovascular endothelial cells," *British Journal of Pharmacology*, vol. 156, no. 7, pp. 1115–1123, 2009.
- [55] A.-M. Van Dam, H. E. De Vries, J. Kuiper et al., "Interleukin-1 receptors on rat brain endothelial cells: a role in neuroimmune interaction?" *The FASEB Journal*, vol. 10, no. 2, pp. 351–356, 1996.
- [56] S. Ching, H. Zhang, N. Belevych et al., "Endothelial-specific knockdown of interleukin-1 (IL-1) type 1 receptor differentially alters CNS responses to IL-1 depending on its route of administration," *The Journal of Neuroscience*, vol. 27, no. 39, pp. 10476–10486, 2007.
- [57] N. Quan, L. He, and W. Lai, "Endothelial activation is an intermediate step for peripheral lipopolysaccharide induced activation of paraventricular nucleus," *Brain Research Bulletin*, vol. 59, no. 6, pp. 447–452, 2003.
- [58] Y. Wang, S. Jin, Y. Sonobe et al., "Interleukin-1 β induces blood-brain barrier disruption by downregulating sonic hedgehog in astrocytes," *PLoS ONE*, vol. 9, no. 10, Article ID e110024, 2014.
- [59] E. S. Wohleb, M. L. Hanke, A. W. Corona et al., " β -Adrenergic receptor antagonism prevents anxiety-like behavior and microglial reactivity induced by repeated social defeat," *The Journal of Neuroscience*, vol. 31, no. 17, pp. 6277–6288, 2011.
- [60] E. S. Wohleb, J. M. Patterson, V. Sharma, N. Quan, J. P. Godbout, and J. F. Sheridan, "Knockdown of interleukin-1 receptor type-1 on endothelial cells attenuated stress-induced neuroinflammation and prevented anxiety-like behavior," *The Journal of Neuroscience*, vol. 34, no. 7, pp. 2583–2591, 2014.
- [61] M.-A. Lécuyer, H. Kebir, and A. Prat, "Glial influences on BBB functions and molecular players in immune cell trafficking," *Biochimica et Biophysica Acta (BBA)—Molecular Basis of Disease*, vol. 1862, no. 3, pp. 472–482, 2016.
- [62] L. Redwine, R. L. Hauger, J. C. Gillin, and M. Irwin, "Effects of sleep and sleep deprivation on interleukin-6, growth hormone, cortisol, and melatonin levels in humans," *The Journal of Clinical Endocrinology & Metabolism*, vol. 85, no. 10, pp. 3597–3603, 2000.
- [63] E. Späth-Schwalbe, K. Hansen, F. Schmidt et al., "Acute effects of recombinant human interleukin-6 on endocrine and central nervous sleep functions in healthy men," *Journal of Clinical Endocrinology and Metabolism*, vol. 83, no. 5, pp. 1573–1579, 1998.
- [64] A. N. Vgontzas, M. Zoumakis, D. A. Papanicolaou et al., "Chronic insomnia is associated with a shift of interleukin-6 and tumor necrosis factor secretion from nighttime to daytime," *Metabolism*, vol. 51, no. 7, pp. 887–892, 2002.
- [65] M. Haack, E. Sanchez, and J. M. Mullington, "Elevated inflammatory markers in response to prolonged sleep restriction are associated with increased pain experience in healthy volunteers," *Sleep*, vol. 30, no. 9, pp. 1145–1152, 2007.
- [66] S. Abdelmalek, N. Souissi, H. Chtourou et al., "Effects of partial sleep deprivation on proinflammatory cytokines, growth hormone, and steroid hormone concentrations during repeated brief sprint interval exercise," *Chronobiology International*, vol. 30, no. 4, pp. 502–509, 2013.
- [67] M. R. Irwin, M. Wang, C. O. Campomayor, A. Collado-Hidalgo, and S. Cole, "Sleep deprivation and activation of morning levels of cellular and genomic markers of inflammation," *Archives of Internal Medicine*, vol. 166, no. 16, pp. 1756–1762, 2006.
- [68] J. Guess, J. B. Burch, K. Ogoussan et al., "Circadian disruption, Per3, and human cytokine secretion," *Integrative Cancer Therapies*, vol. 8, no. 4, pp. 329–336, 2009.
- [69] F. Sauvet, G. Leftheriotis, D. Gomez-Merino et al., "Effect of acute sleep deprivation on vascular function in healthy subjects," *Journal of Applied Physiology*, vol. 108, no. 1, pp. 68–75, 2010.
- [70] L. Krizanac-Bengez, M. Kapural, F. Parkinson et al., "Effects of transient loss of shear stress on blood-brain barrier endothelium: role of nitric oxide and IL-6," *Brain Research*, vol. 977, no. 2, pp. 239–246, 2003.
- [71] Q. Feng, Y. I. Wang, and Y. Yang, "Neuroprotective effect of interleukin-6 in a rat model of cerebral ischemia," *Experimental and Therapeutic Medicine*, vol. 9, no. 5, pp. 1695–1701, 2015.
- [72] L. Vallières and S. Rivest, "Regulation of the genes encoding interleukin-6, its receptor, and gp130 in the rat brain in response to the immune activator lipopolysaccharide and the proinflammatory cytokine interleukin-1 β ," *Journal of Neurochemistry*, vol. 69, no. 4, pp. 1668–1683, 1997.
- [73] D. Matusevicius, P. Kivisäkk, B. He et al., "Interleukin-17 mRNA expression in blood and CSF mononuclear cells is augmented in multiple sclerosis," *Multiple Sclerosis*, vol. 5, no. 2, pp. 101–104, 1999.
- [74] E. Lubberts, L. A. B. Joosten, B. Oppers et al., "IL-1-independent role of IL-17 in synovial inflammation and joint destruction during collagen-induced arthritis," *The Journal of Immunology*, vol. 167, no. 2, pp. 1004–1013, 2001.
- [75] D. Yen, J. Cheung, H. Scheerens et al., "IL-23 is essential for T cell-mediated colitis and promotes inflammation via IL-17 and IL-6," *The Journal of Clinical Investigation*, vol. 116, no. 5, pp. 1310–1316, 2006.
- [76] C. K. Wong, C. Y. Ho, E. K. Li, and C. W. K. Lam, "Elevation of proinflammatory cytokine (IL-18, IL-17, IL-12) and Th2 cytokine (IL-4) concentrations in patients with systemic lupus erythematosus," *Lupus*, vol. 9, no. 8, pp. 589–593, 2000.
- [77] Y. Chen, P. Yang, F. Li, and A. Kijlstra, "The effects of Th17 cytokines on the inflammatory mediator production and barrier function of ARPE-19 cells," *PLoS ONE*, vol. 6, no. 3, Article ID e18139, 2011.
- [78] H. Kebir, K. Kreymborg, I. Ifergan et al., "Human TH17 lymphocytes promote blood-brain barrier disruption and central nervous system inflammation," *Nature Medicine*, vol. 13, no. 10, pp. 1173–1175, 2007.
- [79] J. Huppert, D. Closhen, A. Croxford et al., "Cellular mechanisms of IL-17-induced blood-brain barrier disruption," *The FASEB Journal*, vol. 24, no. 4, pp. 1023–1034, 2010.
- [80] A. J. Szalai and M. A. McCrory, "Varied biologic functions of c-reactive protein: lessons learned from transgenic mice," *Immunologic Research*, vol. 26, no. 1–3, pp. 279–287, 2002.
- [81] H. K. Meier-Ewert, P. M. Ridker, N. Rifai et al., "Effect of sleep loss on C-Reactive protein, an inflammatory marker of cardiovascular risk," *Journal of the American College of Cardiology*, vol. 43, no. 4, pp. 678–683, 2004.
- [82] S. R. Patel, X. Zhu, A. Storfer-Isser et al., "Sleep duration and biomarkers of inflammation," *Sleep*, vol. 32, no. 2, pp. 200–204, 2009.
- [83] M. L. Okun, M. Coussons-Read, and M. Hall, "Disturbed sleep is associated with increased C-reactive protein in young women," *Brain, Behavior, and Immunity*, vol. 23, no. 3, pp. 351–354, 2009.

- [84] J. V. Castell, M. J. Gómez-Lechón, M. David, R. Fabra, R. Trullenque, and P. C. Heinrich, "Acute-phase response of human hepatocytes: regulation of acute-phase protein synthesis by interleukin-6," *Hepatology*, vol. 12, no. 5, pp. 1179–1186, 1990.
- [85] C. R. W. Kuhlmann, L. Librizzi, D. Closhen et al., "Mechanisms of C-reactive protein-induced blood-brain barrier disruption," *Stroke*, vol. 40, no. 4, pp. 1458–1466, 2009.
- [86] D. Closhen, B. Bender, H. J. Luhmann, and C. R. W. Kuhlmann, "CRP-induced levels of oxidative stress are higher in brain than aortic endothelial cells," *Cytokine*, vol. 50, no. 2, pp. 117–120, 2010.
- [87] C. Amos, I. A. Romero, C. Schültze et al., "Cross-linking of brain endothelial intercellular adhesion molecule (ICAM)-1 induces association of ICAM-1 with detergent-insoluble cytoskeletal fraction," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 21, no. 5, pp. 810–816, 2001.
- [88] B. Engelhardt, F. K. Conley, and E. C. Butcher, "Cell adhesion molecules on vessels during inflammation in the mouse central nervous system," *Journal of Neuroimmunology*, vol. 51, no. 2, pp. 199–208, 1994.
- [89] B. Devaux, D. Scholz, A. Hirche, W. P. Klovekorn, and J. Schaper, "Upregulation of cell adhesion molecules and the presence of low grade inflammation in human chronic heart failure," *European Heart Journal*, vol. 18, no. 3, pp. 470–479, 1997.
- [90] V. M. Pak, B. T. Keenan, N. Jackson et al., "Adhesion molecule increases in sleep apnea: beneficial effect of positive airway pressure and moderation by obesity," *International Journal of Obesity*, vol. 39, no. 3, pp. 472–479, 2015.
- [91] L.-L. Meng, Y.-Z. Tang, C.-L. Ni et al., "Impact of inflammatory markers on the relationship between sleep quality and incident cardiovascular events in type 2 diabetes," *Journal of Diabetes and its Complications*, vol. 29, no. 7, pp. 882–886, 2015.
- [92] A. T. Argaw, B. T. Gurfein, Y. Zhang, A. Zameer, and G. R. John, "VEGF-mediated disruption of endothelial CLN-5 promotes blood-brain barrier breakdown," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 6, pp. 1977–1982, 2009.
- [93] P. Stiefel, M. A. Sánchez-Armengol, J. Villar, A. Vallejo-Vaz, R. Moreno-Luna, and F. Capote, "Obstructive sleep apnea syndrome, vascular pathology, endothelial function and endothelial cells and circulating microparticles," *Archives of Medical Research*, vol. 44, no. 6, pp. 409–414, 2013.
- [94] D. C. Lim and A. I. Pack, "Obstructive sleep apnea and cognitive impairment: addressing the blood-brain barrier," *Sleep Medicine Reviews*, vol. 18, no. 1, pp. 35–48, 2014.
- [95] L. Ibrahim, W. Duncan, D. A. Luckenbaugh, P. Yuan, R. Machado-Vieira, and C. A. Zarate, "Rapid antidepressant changes with sleep deprivation in major depressive disorder are associated with changes in vascular endothelial growth factor (VEGF): a pilot study," *Brain Research Bulletin*, vol. 86, no. 1–2, pp. 129–133, 2011.
- [96] H. Chen, Z. Zhao, J. Zheng et al., "The effect of IGF-1 on symptoms of sleep deprivation in a rat model of inflammatory heart disease and metabolic syndrome," *Biochemical and Biophysical Research Communications*, vol. 446, no. 4, pp. 843–849, 2014.
- [97] S. Bake, A. K. Okoreeh, R. C. Alaniz, and F. Sohrabji, "Insulin-Like Growth Factor (IGF)-I modulates endothelial blood-brain barrier function in ischemic middle-aged female rats," *Endocrinology*, vol. 157, no. 1, pp. 61–69, 2016.
- [98] A. J. Park, J. Collins, P. A. Blennerhassett et al., "Altered colonic function and microbiota profile in a mouse model of chronic depression," *Neurogastroenterology and Motility*, vol. 25, no. 9, pp. 733–e575, 2013.
- [99] E. Patterson, P. M. Ryan, J. F. Cryan et al., "Gut microbiota, obesity and diabetes," *Postgraduate Medical Journal*, vol. 92, no. 1087, pp. 286–300, 2016.
- [100] J. R. Kelly, P. J. Kennedy, J. F. Cryan, T. G. Dinan, G. Clarke, and N. P. Hyland, "Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related psychiatric disorders," *Frontiers in Cellular Neuroscience*, vol. 9, article 392, 2015.
- [101] V. Braniste, M. Al-Asmakh, C. Kowal et al., "The gut microbiota influences blood-brain barrier permeability in mice," *Science Translational Medicine*, vol. 19, no. 6, pp. 1–24, 2014.
- [102] C. Korth, "A co-evolutionary theory of sleep," *Medical Hypotheses*, vol. 45, no. 3, pp. 304–310, 1995.
- [103] K. D. Rochfort and P. M. Cummins, "Cytokine-mediated dysregulation of zonula occludens-1 properties in human brain microvascular endothelium," *Microvascular Research*, vol. 100, pp. 48–53, 2015.
- [104] D. A. Antonetti, A. J. Barber, L. A. Hollinger, E. B. Wolpert, and T. W. Gardner, "Vascular endothelial growth factor induces rapid phosphorylation of tight junction proteins occludin and zonula occluden 1. A potential mechanism for vascular permeability in diabetic retinopathy and tumors," *The Journal of Biological Chemistry*, vol. 274, no. 33, pp. 23463–23467, 1999.
- [105] M. A. Lopez-Ramirez, R. Fischer, C. C. Torres-Badillo et al., "Role of caspases in cytokine-induced barrier breakdown in human brain endothelial cells," *The Journal of Immunology*, vol. 189, no. 6, pp. 3130–3139, 2012.
- [106] K. Wiggins-Dohlvik, M. Merriman, C. A. Shaji et al., "Tumor necrosis factor- α disruption of brain endothelial cell barrier is mediated through matrix metalloproteinase-9," *The American Journal of Surgery*, vol. 208, no. 6, pp. 954–960, 2014.
- [107] W. Tian and T. R. Kyriakides, "Matrix metalloproteinase-9 deficiency leads to prolonged foreign body response in the brain associated with increased IL-1 β levels and leakage of the blood-brain barrier," *Matrix Biology*, vol. 28, no. 3, pp. 148–159, 2009.
- [108] W. J. Trickler, W. G. Mayhan, and D. W. Miller, "Brain microvessel endothelial cell responses to tumor necrosis factor- α involve a nuclear factor kappa B (NF- κ B) signal transduction pathway," *Brain Research*, vol. 1048, no. 1–2, pp. 24–31, 2005.
- [109] D. Engblom, M. Ek, S. Saha, A. Ericsson-Dahlstrand, P.-J. Jakobsson, and A. Blomqvist, "Prostaglandins as inflammatory messengers across the blood-brain barrier," *Journal of Molecular Medicine*, vol. 80, no. 1, pp. 5–15, 2002.
- [110] D. J. Jaworowicz Jr., P. J. Korytko, S. S. Lakhman, and K. M. K. Boje, "Nitric oxide and prostaglandin E₂ formation parallels blood-brain barrier disruption in an experimental rat model of bacterial meningitis," *Brain Research Bulletin*, vol. 46, no. 6, pp. 541–546, 1998.
- [111] A. Eskilsson, E. Mirrasekhian, S. Dufour, M. Schwaninger, D. Engblom, and A. Blomqvist, "Immune-induced fever is mediated by IL-6 receptors on brain endothelial cells coupled to stat3-dependent induction of brain endothelial prostaglandin synthesis," *The Journal of Neuroscience*, vol. 34, no. 48, pp. 15957–15961, 2014.
- [112] N. Laflamme, S. Lacroix, and S. Rivest, "An essential role of interleukin-1 β in mediating NF- κ B activity and COX-2 transcription in cells of the blood-brain barrier in response to a systemic and localized inflammation but not during endotoxemia," *The Journal of Neuroscience*, vol. 19, no. 24, pp. 10923–10930, 1999.

- [113] S. Y. Eum, I. Andras, B. Hennig, and M. Toborek, "NADPH oxidase and lipid raft-associated redox signaling are required for PCB153-induced upregulation of cell adhesion molecules in human brain endothelial cells," *Toxicology and Applied Pharmacology*, vol. 240, no. 2, pp. 299–305, 2009.
- [114] J. H. Mills, L. F. Thompson, C. Mueller et al., "CD73 is required for efficient entry of lymphocytes into the central nervous system during experimental autoimmune encephalomyelitis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 27, pp. 9325–9330, 2008.
- [115] W. Pan and A. J. Kastin, "The blood-brain barrier: regulatory roles in wakefulness and sleep," *The Neuroscientist*, 2016.
- [116] G. Hurtado-Alvarado, A. M. Cabañas-Morales, and B. Gómez-González, "Pericytes: brain-immune interface modulators," *Frontiers in Integrative Neuroscience*, vol. 7, article 80, 2014.
- [117] A. C. Reynolds and S. Banks, "Total sleep deprivation, chronic sleep restriction and sleep disruption," *Progress in Brain Research*, vol. 185, pp. 91–103, 2010.
- [118] D. Suchecki and S. Tufik, "Social stability attenuates the stress in the modified multiple platform method for paradoxical sleep deprivation in the rat," *Physiology & Behavior*, vol. 68, no. 3, pp. 309–316, 2000.
- [119] A. Rechtschaffen, B. M. Bergmann, C. A. Everson, C. A. Kushida, and M. A. Gilliland, "Sleep deprivation in the rat: I. Conceptual issues," *Sleep*, vol. 12, no. 1, pp. 1–4, 1989.
- [120] A. Rechtschaffen and B. M. Bergmann, "Sleep deprivation in the rat by the disk-over-water method," *Behavioural Brain Research*, vol. 69, no. 1-2, pp. 55–63, 1995.