

Research Article

Proteomic Identification of Heat Shock-Induced Danger Signals in a Melanoma Cell Lysate Used in Dendritic Cell-Based Cancer Immunotherapy

Fermín E. González ,^{1,2} Alexey Chernobrovkin,³ Cristián Pereda,^{2,4} Tamara García-Salum,^{5,6} Andrés Tittarelli,^{2,4} Mercedes N. López,^{2,4} Flavio Salazar-Onfray,^{2,4} and Roman A. Zubarev ³

¹Laboratory of Experimental Immunology & Cancer, Faculty of Dentistry, Universidad de Chile, 8380492 Santiago, Chile

²Millennium Institute on Immunology and Immunotherapy, Institute of Biomedical Sciences, Faculty of Medicine, Universidad de Chile, 8380453 Santiago, Chile

³Department of Medical Biochemistry and Biophysics, Karolinska Institute, 17177 Stockholm, Sweden

⁴Disciplinary Program of Immunology, Institute of Biomedical Sciences, Faculty of Medicine, Universidad de Chile, 8380453 Santiago, Chile

⁵Laboratory of Molecular Virology, Departamento de Enfermedades Infecciosas e Inmunología Pediátrica, School of Medicine, Pontificia Universidad Católica de Chile, 8331150 Santiago, Chile

⁶Millennium Institute on Immunology and Immunotherapy, Pontificia Universidad Católica de Chile, 8331150 Santiago, Chile

Correspondence should be addressed to Fermín E. González; fgonzalez@uchile.cl and Roman A. Zubarev; roman.zubarev@ki.se

Received 29 June 2017; Revised 28 November 2017; Accepted 11 December 2017; Published 18 March 2018

Academic Editor: Aurelia Rughetti

Copyright © 2018 Fermín E. González et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Autologous dendritic cells (DCs) loaded with cancer cell-derived lysates have become a promising tool in cancer immunotherapy. During the last decade, we demonstrated that vaccination of advanced melanoma patients with autologous tumor antigen presenting cells (TAPCells) loaded with an allogeneic heat shock- (HS-) conditioned melanoma cell-derived lysate (called TRIMEL) is able to induce an antitumor immune response associated with a prolonged patient survival. TRIMEL provides not only a broad spectrum of potential melanoma-associated antigens but also danger signals that are crucial in the induction of a committed mature DC phenotype. However, potential changes induced by heat conditioning on the proteome of TRIMEL are still unknown. The identification of newly or differentially expressed proteins under defined stress conditions is relevant for understanding the lysate immunogenicity. Here, we characterized the proteomic profile of TRIMEL in response to HS treatment. A quantitative label-free proteome analysis of over 2800 proteins was performed, with 91 proteins that were found to be regulated by HS treatment: 18 proteins were overexpressed and 73 underexpressed. Additionally, 32 proteins were only identified in the HS-treated TRIMEL and 26 in non HS-conditioned samples. One protein from the overexpressed group and two proteins from the HS-exclusive group were previously described as potential damage-associated molecular patterns (DAMPs). Some of the HS-induced proteins, such as haptoglobin, could be also considered as DAMPs and candidates for further immunological analysis in the establishment of new putative danger signals with immunostimulatory functions.

1. Introduction

Dendritic cells (DCs) are professional antigen presenting cells (APCs) that, upon encountering antigens (Ags) and proper sensing of danger signals, such as pathogen-associated molecular patterns (PAMPs) and/or damage-

associated molecular patterns (DAMPs) in the tissue microenvironment, efficiently trigger adaptive immunity against pathogens and tumors [1–6], thus establishing a link between the innate and adaptive immunity [7]. Over the past decade, autologous DC-based immunotherapy against cancer has become a safe and reliable therapeutic approach, especially

for solid tumors [8]. We have previously shown that immunotherapy using autologous *ex vivo*-generated tumor antigen presenting cells (TAPCells) from cytokine-activated monocytes (AM), and loaded with an original melanoma cell-derived lysate (referred to as TRIMEL), generated from three human melanoma cell lines, induces T cell-mediated immune responses and increased survival time of stage IV malignant melanoma (MM) patients [9–13]. In addition, more than 60% of treated patients showed a delayed type IV hypersensitivity (DTH) reaction against TRIMEL, indicating the development of an immunological memory. Importantly, positive DTH response correlated with prolonged survival of treated malignant melanoma patients [10–12]. Furthermore, we have observed that TAPCells vaccination induces differential response patterns of specific regulatory cell subpopulations in patients' peripheral blood leucocytes [10, 13]. These data strongly support an important role of TRIMEL in the *ex vivo* education of immunotherapeutic TAPCells and, in turn, in their capacity to trigger an *in vivo* antitumor immune response.

Despite these positive outcomes, around 40% of treated patients do not respond to the therapy (considering their DTH response) and have the same survival rate as nontreated ones [9–11]. This lack of response could be explained, at least in part, by carrying the 896 A>G *TLR4* gene polymorphism [12], an absence of sufficient immunogenic danger signals or a deficient timing in the input of danger signals to DCs [11], either during the *ex vivo* TAPCells generation or after their injection, which could induce deficiencies in migration, antigen processing, and/or presentation by inoculated cells.

In vitro, human DCs loaded with melanoma cells that were heat-treated at 42°C before being killed showed more efficient cross-priming to naive human CD8⁺ T cells than DCs loaded with unheated killed melanoma cells [14]. These heat-treated melanoma cells expressed enhanced amounts of the heat shock protein (HSP) 70, and the enhanced cross-priming could be reproduced by overexpression of Hsp70 in melanoma cells [14]. In this regard, we have previously shown that the TRIMEL lysate can induce a mature and committed DC phenotype from AM cells [11, 15]. Moreover, we have also demonstrated that the HS treatment of melanoma cells before their final lysis for TRIMEL generation increases calreticulin (CALR) plasma membrane translocation and induces the release of high mobility group box 1 (HMGB1) protein [11] and two well-described DAMPs [16, 17]. Importantly, *in vitro*-generated DCs from melanoma patients stimulated with TRIMEL induced a fivefold increase of IFN- γ release by a melanoma-specific cytotoxic T cell clone, compared to APCs stimulated with a non-HS-treated melanoma cell lysate [11], indicating the importance of the HS treatment in the capacity of TRIMEL to induce DCs with immunostimulatory properties. Both CALR and HMGB1 mobilizations were associated with enhanced DCs' maturation and with an efficient antigen cross-presentation capacity, respectively [11]. Additionally, HMGB1 from TRIMEL colocalizes with the receptor TLR4 on THP-1 cell surface, and the blockade of TLR4 in AM inhibits the expression of maturation-associated markers, proinflammatory cytokines,

and CCR7 chemokine receptor induced by TRIMEL [12]. Moreover, DCs' ability to migrate to draining lymph nodes, a relevant prerequisite for its clinical efficacy, is also increased upon TRIMEL stimulation [18]. Taken together, these data strongly support that TRIMEL would contain not only HMGB1 and CALR but also other proteins or factors with DAMP functions, which contribute to its capacity to induce the TAPCells phenotype and their therapeutic performance. In this context, identifying the proteome changes in the lysate TRIMEL in response to HS would help to better understand TRIMEL's capacity to induce the *in vitro/ex vivo* DC maturation.

2. Material and Methods

2.1. Patients and Healthy Donors. Peripheral blood mono-nuclear cells (PBMC) were obtained by a leukapheresis procedure from four advanced (stage IV) MM patients previously treated using a reported TAPCells vaccination protocol [19]. Additionally, PBMC from six healthy donors, from the Blood Bank Service, Clinical Hospital, Universidad de Chile, were obtained. The present study was performed in agreement with the Helsinki Declaration and approved by the Bioethical Committee for Human Research of the Clinical Hospital, Universidad de Chile. All patients and healthy donors signed an informed consent form.

2.2. Cell Lines, Melanoma Cell Lysate TRIMEL, and HS Conditioning. The allogeneic cell lysate TRIMEL was prepared as previously described [10, 11]. Briefly, three different melanoma cell lines (MEL-1, MEL-2, and MEL-3), established from three tumor-infiltrated lymph nodes from metastatic HLA-A2⁺ stage IV melanoma patients and those positive for several melanoma-associated antigens, were cultured in RPMI-1640 medium (Gibco, Austria) supplemented with 10% (*v/v*) fetal bovine serum (FBS, Gibco/BRL), 10 µg/mL streptomycin, and 100 mg/mL penicillin (Sigma, CA, USA), until 95% confluence. Cells were subcultured every 2-3 days. Before use, all the cell lines were tested by PCR techniques, to check the absence of potentially infecting virus or mycoplasma. The presence of contaminating bacteria was also ruled out by periodical culture testing in agar.

The cells were mixed in equal proportions (1×10^7 cells for each cell line), resuspended in the therapeutic AIM-V medium (Gibco, CA, USA) at a concentration of 4×10^6 cells/mL, HS-treated by incubating the cells one hour at 42°C, then two hours at 37°C, and finally lysed by performing three freeze-thaw cycles using liquid nitrogen. In order to perform the proteomic analysis, before the lysing step, part of the cell mixture was washed three times with PBS and frozen as pellets at -80°C until further proteomic analysis. Five independently produced batches for the complete lysate TRIMEL, with and without HS conditioning, were prepared (a total of 10 samples).

2.3. In Vitro Human DC Generation. PBMC of melanoma patients and healthy donors were cultured in serum-free therapeutic AIM-V medium at a concentration

of 13×10^6 cells/mL in six-well plates (BD Biosciences, Hershey, PA, USA) at 37°C and 5% CO₂ for 2 hours. Thereafter, nonadherent cells were removed and the adherents (monocytes) were maintained and incubated for 22 additional hours in the presence of 500 U/mL recombinant human IL-4 (rhIL-4) and 800 U/mL of GM-CSF (US Biological, Swampscott, MA, USA). The obtained cytokine-activated monocytes (AM), which showed an immature DC-like phenotype, were then stimulated for 24 additional hours with 100 µg/mL of TRIMEL or the lysate without HS conditioning.

2.4. Flow Cytometry Analysis. The cells were phenotypically characterized by flow cytometry using the following conjugated antibodies (Abs): mouse anti-human-HLA-ABC-FITC, HLA-DR-FITC, CD80-FITC, and CD11c-PE-Cy7 (eBioscience, San Diego, CA, USA). Briefly, cells were gently removed from the culture plates using cell scrapers. Then, the cells were centrifuged at 1000 rpm for 5 minutes at 4°C, washed with PBS, and incubated with Abs for 30 minutes. After being washed twice with PBS, samples were acquired on a FACSCalibur (BD Biosciences, Hershey, PA, USA) and analyzed using FlowJo software (Tree Star Inc., OR, USA). All the analyses were made in the CD11c⁺ cell population of each condition and sample.

2.5. Cell Lysis and Protein Extraction and Digestion

2.5.1. Cell Lysis and Protein Extraction. A cell pellet containing 4×10^6 cells was resuspended in 1 mL of lysis solution (0.2% ProteaseMax/10% acetonitrile (ACN)/50 mM ammonium bicarbonate (AmBic)). Cell lysis was performed over 10 minutes with the aid of rigorous vortexing. The lysate was kept at 95°C for 5 minutes and then subjected to 15 minutes sonication (30% amplitude, 3:3 pulse) with a Bransonic sonicator. Samples were centrifuged at 14,000 rpm over 7 minutes at room temperature and the precipitate was discarded. The total concentration of proteins was determined using a bicinchoninic acid assay (Pierce BCA assay kit, Thermo Fisher Scientific Inc.).

2.5.2. In-Solution Digestion. Proteins were reduced by adding DTT to a final concentration of 10 mM and incubation for 30 minutes at 50°C, then alkylated via incubation with iodoacetamide for 30 minutes at room temperature. Proteins (80 µg) were digested by adding 2 µg of trypsin (Sequencing Grade Modified Trypsin, Promega) and incubated at 37°C for 9 hours. The digest was rigorously vortexed over 5 minutes. Digestion was terminated by the addition of 5% acetic acid. Samples were cleaned and desalted using C18 StageTips (Thermo Fisher Scientific Inc.), dried using a SpeedVac and resuspended in water with 0.1% formic acid.

2.6. Mass Spectrometry (MS). Peptide mixture was injected into an Ultimate 3000 nanoflow LC system (Thermo Scientific, USA) in-line coupled to a Q Exactive mass spectrometer (Thermo Scientific). The chromatographic separation of the peptides was achieved using a 25 cm long in-house packed column (C18-AQ ReproSil-Pur®, Dr. Maisch GmbH, Germany) at 55°C with the following gradient: 4–30% ACN

in 89 minutes, 26–95% ACN for 5 minutes, and 95% ACN for 8 minutes all at a flow rate of 250 nL/minutes.

The MS acquisition method comprised one full scan survey ranging from m/z 300 to m/z 1650 acquired with a resolution of $R=140,000$ at m/z 200 and AGC target value of 5×10^6 , followed by data-dependent higher-energy collisional dissociation fragmentation scans from a maximum of 16 most intense precursor ions with a charge state ≥ 2 . For dependent scans, the following parameters were used: precursor isolation width 4 Da, AGC target value of 2×10^5 , and normalized collision energy of 26. Scans were acquired in profile mode with a resolution of $R=17,500$.

2.7. Protein Identification and Quantification. The MS raw data were analyzed with the MaxQuant software (version 1.5.3.30). A false discovery rate (FDR) of 0.01 for proteins and peptides and a minimum peptide length of six amino acids were required. Mass accuracy of the precursor ions was improved by the time-dependent recalibration algorithm of MaxQuant. The Andromeda search engine was used to search the MS/MS spectra against the Uniprot human database (containing 90,482 entries) combined with 262 common contaminants and concatenated with the reversed versions of all sequences. Enzyme specificity was set to trypsin. Further modifications were cysteine carbamidomethylation (fixed) as well as protein N-terminal acetylation, asparagine and glutamine deamidation, and methionine oxidation (variable). A maximum of two missed cleavages were allowed. Peptide identification was based on a search with an initial mass deviation of the precursor ion of up to 7 ppm. The fragment mass tolerance was set to 20 ppm on the m/z scale. Only proteins quantified with at least two peptides were considered for quantitation.

2.8. Bioinformatics and Statistical Analysis. Analysis of variance (ANOVA) and the Kruskal-Wallis test for nonparametric variables were used to compare significance of the differences in maturation marker expressions between studied groups. Differences were considered statistically significant at $p < 0.05$. The analyses were performed using GraphPad Prism 5 software (GraphPad Software Inc., USA).

Analysis of the data provided by MaxQuant was performed in the R scripting and statistical environment. Differences in relative protein abundances between heat-treated and control samples were assessed by moderated *t*-test using limma package [20]. Benjamini-Hochberg correction for multiple comparisons was used.

Gene set enrichment analysis and visualization of protein-protein interaction networks was performed using STRING software (<http://string-db.org/>) [21]. Each group of proteins—overexpressed, underexpressed, exclusively expressed in TRIMEL, and exclusively expressed in non-treated (no-HS) samples—was analyzed separately.

3. Results

3.1. The HS-Conditioning Contributes to the In Vitro Capacity of TRIMEL to Induce a Mature Phenotype on Human DCs. We have previously demonstrated that the




FIGURE 1: The HS conditioning of TRIMEL melanoma cells contributes to its *in vitro* DC maturation capacity. Representative density plots (a) and statistical quantification (b) of the DC-associated marker expression MHC-I, MHC-II, and CD80 in primary human cytokine-activated monocytes stimulated with TRIMEL (HS), or with the same lysate generated without heat shock conditioning (no-HS) (100 µg/mL) or without lysate (unstimulated (Unst)). (b) The quantification of the maturation marker expression considered the % positive cells, the geometric mean fluorescence intensity (gMFI) of the positive cells, and the integrated MFI (iMFI: % positive cells × gMFI of positive cells/100). The expression of surface markers was assessed by flow cytometry (CD11c⁺ cells were gated). Data represent three independent experiments with PBMC derived from three different stage IV MM patients. (c) Bars indicate the average fold induction and standard deviation (SD) of the iMFI of DC markers relative to monocytes stimulated with no-HS lysate. **p* < 0.05 and ***p* < 0.01.

addition of TRIMEL to primary human AM cells mediated up to fourfold induction of several surface markers associated with DC maturation such as MHC-I, MHC-II, CD80, CD83, and CD86 [11]. In addition, TRIMEL could also significantly induce a twofold increase in the expression of MHC-II, CD83, and CCR7 molecules in monocyte/macrophage THP-1 cells, generating a DC-like phenotype as compared with the unstimulated control cells [18].

In order to evaluate the contribution of the HS conditioning of melanoma cells that generate TRIMEL to its capability in inducing a mature DC phenotype, we stimulated primary human AM cells with TRIMEL and the same lysate without the HS conditioning during 24 hours. All the canonical DC maturation-associated markers evaluated—MHC-I, MHC-II, and CD80—showed a higher percentage of positive cells in TRIMEL-stimulated cells when compared with control cells stimulated with the lysate generated with nontreated (no-HS) melanoma cells (Figure 1(a)). In addition, CD80 expression was significantly higher in cells

stimulated with TRIMEL when compared with primary AM cells stimulated with nontreated melanoma cell-derived lysates (Figures 1(b) and 1(c)).

3.2. Proteomic Analysis of TRIMEL Showed Proteins Differentially Regulated by HS, Some of Them with Previously Described DAMP Function. Proteomic analysis of the melanoma-derived lysate TRIMEL and nontreated (no-HS) lysates identified a total of 2798 proteins, 2740 of which were identified in both groups of samples, with and without HS conditioning (Supp. Figure 1). A principal component analysis clearly separates the samples by its HS conditioning (Figure 2(a)). In order to visualize changes in the protein expression induced by HS, proteomic data were visualized on a “volcano plot” (Figure 2(b)). Considering the regulated proteins by HS conditioning, a hierarchical clustering of proteins with the largest expression fold changes and *p* value < 0.01 was performed (Figure 3). As showed in Figure 3(a), a clearly distinctive protein expression profile for both groups of samples (TRIMEL (HS) and nontreated




FIGURE 2: Quantitative proteomic comparison of TRIMEL lysates with (HS) and without (no-HS) heat shock conditioning. (a) Principal components analysis (PCA). (b) Volcano plot representation of moderated *t*-test analysis. Each point represents one protein plotted by log₂ fold change (FC, average of four samples) versus minus logarithm of the *q*-value (Bemjamini-Hochberg corrected *p* value). The horizontal bar represents a *q*-value cutoff of 0.01. Red dots indicate the proteins selected for further analysis.

(no-HS) lysates) was found. Considering a selection criteria of *p* value < 0.01 or abs(log₂ (FC)) > 1 as a cutoff, 18 proteins were selected as significantly more abundant in the melanoma-derived lysate TRIMEL (with HS conditioning) when compared with the nontreated (no-HS) samples (Figure 3(b) and Table 1), being haptoglobin (HP) one of the most overexpressed protein, since it fulfilled both selection criteria. Importantly, when analyzing this group, the protein U2 snRNP-associated SURP motif-containing protein (U2SURP) was found with previously described DAMP function [16, 17] (Table 2).

STRING analysis of protein interactions among the overexpressed proteins, showed protein-protein relationships just among the proteins U2SURP, CPFS3, HNRNPL, and HNRNPA3 (Figure 4(a)). Of note, HP is not involved in the cluster of protein interaction identified by our analysis. This analysis was done considering only 17 proteins because in one case, a group of proteins was identified (proteins RPS27A, UBB, UBC, UBA52, and UBBP4). It means that the set of peptides matches all these proteins so we could not distinguish among them.

On the other hand, 73 proteins were significantly less abundant in TRIMEL (HS-conditioned) samples compared with non-HS-conditioned ones (Supp. Table 1). Of note, among this group of proteins, heat shock protein family A (Hsp70) member 4 (HSPA4) and ribosomal protein S19 (RPS19) are proteins previously described as DAMPs (Supp. Table 1). Remarkably, HMGB1, a well-known protein with an extensively described DAMP function, did not change in its abundance by HS-conditioning. Of note, protein-protein interaction analysis by STRING showed direct interaction between RPS19 and proteins from the

translational machinery like ETF1, BTF3, EEF2, EIF1, EIF3J, and EIF4E proteins (Supp. Figure 2).

3.3. Expression Profile Analysis of TRIMEL Showed Proteins Exclusively Identified in TRIMEL and in No-HS-Conditioned Lysates. Our proteomic analysis also revealed that there was a group of 32 proteins only identified in the lysate TRIMEL (HS-conditioned) (Table 3). Among proteins only identified in TRIMEL samples, histone cluster 2 H2A family member c (HIST2H2AC) and histone cluster 2 H2A family member a3 (HIST2H2AA3) have been previously described to possess DAMP function (Table 2). STRING analysis showed the direct interaction between these two proteins with ANAPC1, RRP8, and POLR1B and indirectly with LTN1, NSUN5, and TRMT112 (Figure 4(b)). In addition, a group of 26 proteins were only identified in nontreated (no-HS) samples (Supp. Table 2). Notably, when we analyzed the group of proteins exclusively identified in nontreated samples (no-HS), we did not find proteins with reported DAMP function. STRING analysis of this group of proteins showed a main interaction group among proteins WDR82, PPP1R2, PPP3CB, PPP3CA, and EPS15 (Supp. Figure 3).

4. Discussion

During recent years, intact cancer cells and cancer cell-derived lysates have been extensively used in different cell-based immunotherapies against cancer. This is mainly because they constitute not only a broad source for tumor-associated antigens but also for several and biochemically diverse molecules with immunomodulatory activity. Indeed, *ex vivo* educated DCs using tumor cell-derived lysates have




FIGURE 3: Expression profiles of selected proteins from TRIMEL and regulated by HS treatment. (a) Heat map with hierarchical clustering of proteins differentially expressed between TRIMEL (HS) and non-HS samples using a cutoff at $p < 0.01$. Protein names are displayed on the right and below is depicted an augmented section of the 18 HS-overexpressed proteins. Red, overexpressed; blue, underexpressed; and white, no change. The color-coded scale is indicated at the top of the chart. (b) Log-transformed relative protein expression of the 18 proteins regulated by HS treatment. The text and table only refer to 17 proteins because in one case a protein group was identified (RPS27A, UBB, UBC, UBA52, and UBBP4). It means the set of peptides matches all these five proteins and we cannot distinguish them here.

become an important approach in cancer immunotherapy, especially in the treatment of solid tumors [8]. We have previously demonstrated the capacity of the allogeneic HS-conditioned lysate TRIMEL to induce a mature DC phenotype on *ex vivo* generated TAPCells. In turn, it is able to trigger an *in vivo* antitumor immunity in advanced MM patients [10–12]. In this context, characterization of the proteomic profile changes induced by HS would help to identify more proteins and protein-protein interactions involved in DC maturation process triggered by their stimulation with cancer cell lysates.

Here, we showed that HS conditioning of melanoma cancer cells belonging to TRIMEL is responsible, at least in part, for the TRIMEL maturation capacity on DC phenotype. In this regard, in a previous study, we have shown that HS conditioning is able to induce the secretion of the DAMP protein HMGB1 by melanoma cells as well as the mobilization of CALR to plasma membrane, a well-known “eat me” signal for phagocytic cells [11]. In the current study, CALR was found among proteins slightly overexpressed after HS

conditioning (p value = 0.0259; logFC = 0.25), suggesting that HS treatment not only mobilizes this protein towards the plasma membrane of melanoma cells but also induces its expression by these cells. However, the nuclear protein HMGB1 did not change its abundance upon HS (p value = 0.5610; logFC = −0.14), indicating that this stimulus is only able to induce its secretion but not its expression by melanoma cells.

Interestingly, melanoma cells upon HS treatment underexpressed more proteins than the ones they overexpressed. This observation could be explained, at least in part, by the fact that HS constitutes a stress factor and, therefore, cells under HS enter in a metabolic state that can alter cellular protein homeostasis. In this context, Hsp70 has been involved in the modulation of the protein synthetic machinery, switching from a degradation phase to the protein synthesis phase [22]. Here, we found proteins belonging to the HSP family differentially regulated by HS. Indeed, heat shock protein family A (Hsp70) member 4 (HSPA4) was significantly underexpressed upon HS treatment and, on the

TABLE 1: Currently known functions of selected gene-proteins upregulated by HS. The protein lists was previously filtered by $p < 0.01$ or $\text{abs}(\log_2(\text{FC})) > 1$.

Gene	ID (NCBI)	Full name (NCBI)	Function (gene ontology)	Reference
CELF1	10658	CUGBP Elav-like family member 1	(i) BRE; RNA; mRNA; pre-mRNA; protein and translation repressor activity, nucleic acid binding	[43–48]
CPSF3	51692	Cleavage and polyadenylation specific factor 3	(i) Protein binding	[49]
FAM195B	348262	MAPK regulated corepressor interacting protein 1	(i) Protein binding	[50, 51]
GPNMB	10457	Glycoprotein NMB	(i) Chemoattractant and receptor ligand activity (ii) Heparin; protein and syndecan binding	[52–54]
HNRNPA3	220988	Heterogeneous nuclear ribonucleoprotein A3	(i) RNA and protein binding	[44, 45, 55]
HNRNPL	3191	Heterogeneous nuclear ribonucleoprotein L	(i) RNA; pre-mRNA; protein and transcription regulatory region DNA binding	[44, 45, 56–58]
HP	3240	Haptoglobin	(i) Hemoglobin and protein binding (ii) ATPase; GTPase activator; microfilament motor and NOT protein homodimerization activity	[59, 60]
MYO9B	4650	Myosin IXB	(ii) ADP; ATP; Rho GTPase; Roundabout; actin; calmodulin and protein binding	[61–65]
PIR	8544	Pirin	(i) Quercetin 2,3-dioxygenase and transcription cofactor activity (ii) Metal ion and protein binding	[66–69]
PPAP2C	8612	Phospholipid phosphatase 2	(i) Phosphoprotein phosphatase activity (ii) Protein binding	[70, 71]
PRCP	5547	Prolylcarboxypeptidase	(i) Protein binding	[72]
PRKD3	23683	Protein kinase D3	(i) Kinase activity (ii) Protein binding	[73, 74]
PTPN12	5782	Protein tyrosine phosphatase, nonreceptor type 12	(i) Nonmembrane spanning protein tyrosine phosphatase; phosphoprotein phosphatase and protein tyrosine phosphatase activity (ii) SH3 domain and protein binding	[75–80]
TOM1L2	146691	Target of myb1 like 2 membrane trafficking protein	(i) Clathrin; protein and protein kinase binding	[81, 82]
U2SURP	23350	U2 snRNP associated SURP motif-containing protein	(i) RNA and protein binding	[45, 83]
UTRN	7402	Utrophin	(i) Actin; integrin; protein; protein kinase and vinculin binding	[84, 85]
ZNF638	27332	Zinc finger protein 638	(i) RNA and double-stranded DNA binding	[45, 86]

ID, identification number; NCBI, National Center for Biotechnology Information.

TABLE 2: Protein with related/putative DAMP functions that were overexpressed and exclusively expressed in TRIMEL treated with HS.

Gene	ID (NCBI)	Full name (NCBI)	Protein subgroup	Reference
FLNC	2318	Filamin C	Exclusive	[87]
HIST2H2AA3/ HIST2H2AC	8337/ 8338	Histone cluster 2 H2A family member a3/ Histone cluster 2 H2A family member c	Exclusive	[17]
HP	3240	Haptoglobin	Overexpressed	[42]
RRP8	23378	Ribosomal RNA processing 8, methyltransferase, homolog (yeast)	Exclusive	[88]
U2SURP	23350	U2 snRNP associated SURP domain containing	Overexpressed	[17]

ID, identification number; NCBI, National Center for Biotechnology Information.




FIGURE 4: Overexpressed proteins by HS in the lysate TRIMEL. (a) Protein-protein interaction network of HS-overexpressed proteins from the lysate TRIMEL. Considering a cutoff $p < 0.01$ or $\text{abs}(\log_2(\text{FC})) > 1$, 18 proteins were significantly more abundant in TRIMEL lysate compared to no-HS-treated lysates; however, only 17 were considered for STRING analysis. Red arrows indicate the top one overexpressed protein HP and U2SURP a previously described DAMP protein. (b) Protein-protein interaction network of the 32 proteins exclusively identified in TRIMEL samples. Red arrows indicate the exclusively identified DAMPs (HIST2H2AC, HIST2H2AA3, RRP8, and FLNC) in TRIMEL samples. Line color indicates the type of interaction evidence. Known interactions: cyan, from curated databases and pink, experimentally determined. Predicted interactions: green, gene neighborhood. Others: yellow, textmining; black, coexpression. Interaction confidence score, 0.4 (medium).

contrary, heat shock protein family A (Hsp70) member 9 (HSPA9), heat shock protein family D (Hsp60) member 1 (HSPD1), and heat shock protein family E (Hsp10) member 1 (HSPE1) were slightly overexpressed. Related with this, one of the significantly underexpressed proteins was PSME1 (proteasome activator complex subunit 1), which is a regulator of proteasome activity [23], suggesting that HS treatment inhibits protein degradation in melanoma cells and, in turn, can contribute to modify protein homeostasis. In addition, several proteins involved in translational machinery, like EIF1, EIF3J, and EIF4E, and different 40S ribosomal proteins are among the underexpressed group of proteins. A less abundance of these proteins could contribute to the inhibition of the translation of different downstream proteins. Interestingly, some of these translation factors, like EIF4E, have been described to be downregulated under heat stress response during exercise [24]. On the other hand, the group of proteins overexpressed/exclusive in no-HS samples could be also relevant to be analyzed. Indeed, transcription factor binding to IGHM enhancer 3 (TFE3) has been also associated with stress response by promoting cell adaptation to nutrient deprivation by upregulating transcription of numerous autophagic and lysosomal genes [25].

The main protein-protein interaction among overexpressed proteins involves U2SURP, CPSF3, HNRNPL, and HNRNPA3 proteins. HNRNPL (heterogeneous nuclear ribonucleoprotein L) and HNRNPA3 (heterogeneous nuclear ribonucleoprotein A3) are members of the HNRNP family that regulate different pre-mRNA and mature mRNA transcription [26]. Importantly, HNRNPL has been recently

associated with aggressiveness and poor prognosis in different malignancies such as colorectal cancer, hepatocellular carcinoma, and bladder cancer [27–29]. On the contrary, and without interactions with other overexpressed proteins, PTPN12 (tyrosine-protein phosphatase nonreceptor type 12) is a tumor suppressor protein and has been associated with overall survival in esophageal squamous cell carcinoma patients and non-small-cell lung cancer [30, 31].

Currently, and despite the high research activity in this field, there is no consensus about DAMPs' immunomodulatory effects (i.e., promoting either antitumor immunity or cancer progression), as well as whether they can be divided based on the timing of their functions on APCs: early-stage effect-related DAMPs, that is, DAMPs inducing chemotaxis, phagocytosis, and proinflammatory cytokine production; or late-stage effect-related DAMPs, that is, DAMPs inducing migration, costimulatory molecules expression, and tumor-associated antigen cross-presentation. In this context, TRIMEL could be considered a source for initial danger signals (or early-stage DAMPs) to be sensed by immature DC which, in turn, are able to sense further signals *in vivo* after its injection into MM patients. Additionally, and in line with the concept recently coined by Yatim and colleagues [32], the DAMPs carried by TRIMEL could be considered as both inducible DAMPs (iDAMPs) and constitutive DAMPs (cDAMPs). Indeed, the six proteins from TRIMEL with described/putative DAMP function (two from the overexpressed group of proteins and four from proteins exclusively identified in TRIMEL) as well as CALR can be considered as an example of iDAMPs, and HMGB1,

TABLE 3: Currently known functions of gene-proteins exclusively identified in HS-conditioned samples (TRIMEL).

Gene	ID (NCBI)	Full name (NCBI)	Function (gene ontology)	Reference
AHSG	197	Alpha 2-HS glycoprotein	(i) Kinase inhibitor activity	[89]
ANAPC1	64682	Anaphase promoting complex subunit 1		
ARID2	196528	AT-rich interaction domain 2	(i) Protein binding	[90]
ATAD3B	83858	ATPase family, AAA domain containing 3B		
CLTB	1212	Clathrin light chain B	(i) Protein binding	[91]
CNIH	10175	Cornichon family AMPA receptor auxiliary protein 1		
DOCK3	1795	Dedicator of cytokinesis 3	(i) Protein binding	[92]
EEF1E1	9521	Eukaryotic translation elongation factor 1 epsilon 1	(i) Protein binding	[93]
FLNC	2318	Filamin C	(i) Ankyrin; cytoskeletal protein and protein binding	[94–96]
FUNDC2	65991	FUN14 domain containing 2		
GK2	2712	Glycerol kinase 2	(i) Glycerol kinase activity	[97]
HEXIM1	10614	Hexamethylene bisacetamide inducible 1	(i) Cyclin-dependent protein serine/threonine kinase inhibitor activity (ii) 7SK snRNA; protein and snRNA binding	[98–101]
HIST2H2AA3/ HIST2H2AC	8337/ 8338	Histone cluster 2 H2A family member a3/ Histone cluster 2 H2A family member c	(i) 3-Hydroxyacyl-CoA dehydrogenase; 3-oxoacyl-[acyl-carrier-protein] reductase (NADH) and estradiol 17-beta-dehydrogenase activity (ii) NADH and protein binding	
HSD17B8	7923	Hydroxysteroid 17-beta dehydrogenase 8		[102–104]
INPPL1	3636	Inositol polyphosphate phosphatase like 1	(i) SH2 domain and protein binding	[105, 106]
ISOC2	79763	Isochorismatase domain containing 2	(i) Protein binding	[107]
KIFC1	3833	Kinesin family member C1	(i) Microtubule motor activity (ii) ATP binding	[108]
LTN1	26046	Listerin E3 ubiquitin protein ligase 1	(i) Protein binding	[109]
NSUN5	55695	NOP2/Sun RNA methyltransferase family member 5	(i) RNA binding	[44, 45]
OAS1	4938	2'-5'-Oligoadenylate synthetase 1	(i) 2'-5'-Oligoadenylate synthetase activity (ii) ATP; double-stranded RNA and protein binding	[93, 110–112]
POLR1B	84172	RNA polymerase I subunit B	(i) Protein binding	[113]
PPP2R4	5524	Protein phosphatase 2 phosphatase activator	(i) Contributes to ATPase; protein heterodimerization; protein homodimerization; protein phosphatase regulator and protein tyrosine phosphatase activator activity (ii) ATP; protein, protein phosphatase 2A and receptor binding	[93, 114–116]
RRP8	23378	Ribosomal RNA processing 8, methyltransferase, homolog (yeast)	(i) S-Adenosylmethionine-dependent methyltransferase activity (ii) RNA; methylated histone and protein binding	[44, 45, 117]
SIGMAR1	10280	Sigma nonopioid intracellular receptor 1	(i) Drug binding	[118]
SPATA5	166378	Spermatogenesis associated 5		
SREK1	140890	Splicing regulatory glutamic acid and lysine rich protein 1	(i) RNA and protein binding	[45, 93]

TABLE 3: Continued.

Gene	ID (NCBI)	Full name (NCBI)	Function (gene ontology)	Reference
SYNGR1	9145	Synaptogyrin 1	(i) Protein binding	[93]
TMEM87A	25963	Transmembrane protein 87A		
TRMT112	51504	tRNA methyltransferase 11-2 homolog (<i>S. cerevisiae</i>)	(i) Protein methyltransferase activity (ii) Protein binding	[119–121]
UAP1	6675	UDP-N-acetylglucosamine pyrophosphorylase 1	(i) Identical protein binding	[93]
XPO6	23214	Exportin 6	(i) Protein transporter activity (ii) Protein binding	[122]

ID, identification number; NCBI, National Center for Biotechnology Information.

previously described as being also relevant for TRIMEL properties [11], could be considered as a cDAMP. These proteins contribute, probably in a synergic way, to the ability of the lysate TRIMEL to *ex vivo* induce a mature phenotype in therapeutic DCs (TAPCells) and could be responsible, at least in part, for the clinical effect of these cells in treated MM patients.

One of the main overexpressed proteins by HS conditioning of the melanoma cells belonging the lysate TRIMEL was HP, a plasmatic glycoprotein with a molecular weight of 38 kDa. The main function of HP is binding haemoglobin (Hb), forming a stable complex HP-Hb, which is cleared via CD163-mediated endocytosis and thus preventing the oxidative tissue damage induced by free haemoglobin [33, 34]. In fact, this protein-protein interaction described between HP and Hb was also confirmed by our STRING analysis, where the only interaction of HP was with HBD and HBB proteins. Moreover, it has been described that HP has a protective role in T cell-mediated inflammatory skin diseases [35]. In addition, it has been previously suggested as a biomarker for early diagnosis in ovarian cancer [36, 37], and its fucosylated form is considered a diagnosis and postsurgical prognosis biomarker in pancreatic and colorectal cancer, respectively [38, 39]. Importantly, during the last years, the capacity of HP to activate DCs was shown in a murine skin transplantation model [40], and recently, the same group showed an amplifying role of HP in inflammation after cardiac transplantation in a murine model, demonstrating a relevant interaction between this protein and the immune system [41]. Interestingly, HP also binds to HMGB1 forming a HP-HMGB1 complex, which elicits the secretion of anti-inflammatory enzymes (e.g., heme oxygenase-1) and cytokines (e.g., IL-10) in WT but not in CD163-deficient macrophages [42], indicating a regulatory function of HP. In this context, in order to confirm HP as a DAMP molecule, further experiments should be focused on the interaction of HP with immune receptors, such as pattern recognition receptors (PRRs), with APCs and other immune cell types.

This study constitutes a conceptual approach in order to identify DAMPs that are induced by HS, which is a fundamental step in TRIMEL generation and in its capacity to induce *ex vivo/in vitro* DC maturation. We have shown that the clinically used lysate TRIMEL carries at least six proteins

with previously described or putative DAMP function. These proteins, induced by HS conditioning of the melanoma cells before their lysis for TRIMEL generation, could be considered as iDAMPs and, therefore, involved in the capacity of TRIMEL to induce the *ex vivo* maturation of TAPCells and their *in vivo* clinical performance in vaccinated patients [10]. Importantly, there are several other proteins in the lysate that have been over- or exclusively expressed upon HS treatment and, therefore, are potential candidates to be confirmed as DAMPs such as HP. DC maturation is a very complex process, which strongly depends on the amount and quality of different signals that are sensed by DCs from either physiologic and pathologic microenvironments [2, 8]. Biochemically, some of these signals are proteins, nucleic acids, metabolites, and extracellular matrix-derived molecules, among others, constituting an even more complex scenario. Related to this, the lysate TRIMEL must contain several nonprotein factors that also contribute to its capacity to induce DC maturation. However, the specific contribution of these factors on TRIMEL capacity to induce the *ex vivo/in vitro* DC maturation is still unknown. Further analysis focused on the determination of the amount and relative contribution of different DAMPs in inducing a mature phenotype in human DC by clinically used cancer cell-derived lysates would help to design new strategies for efficiently activating *ex vivo*-generated DCs and, in turn, developing more effective DC-based immunotherapies against cancer.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this article.

Acknowledgments

This work was supported by a research grant from the Chilean National Fund for Scientific and Technological Development (FONDECYT 11130607, to Fermín E. González; FONDEF ID16I10148, to Mercedes N. López; and FONDECYT 1171213 to Flavio Salazar-Onfray). The authors would like to acknowledge the help from the Proteomics Karolinska Core Facility (PK/KI). The authors also thank Mr. Juan Fernandez, from the language and translation services of the Faculty of Dentistry, for

kindly proofreading and checking the English language of this paper.

Supplementary Materials

Supplementary Figure 1: Venn-diagram illustrating the number of identified proteins. Supplementary Figure 2: protein-protein interaction network from downregulated proteins by HS in the lysate TRIMEL. Supplementary Figure 3: protein-protein interaction network of proteins exclusively expressed in no-HS-conditioned melanoma-derived lysates samples. Supplementary Table 1: currently known functions of selected gene-proteins downregulated by HS. Supplementary Table 2: currently known functions of gene-proteins exclusively expressed by non-HS samples. Supplementary list of references. (*Supplementary Materials*)

References


- [1] G. Schuler, B. Schuler-Thurner, and R. M. Steinman, "The use of dendritic cells in cancer immunotherapy," *Current Opinion in Immunology*, vol. 15, no. 2, pp. 138–147, 2003.
- [2] J. Banchereau, S. Paczesny, P. Blanco et al., "Dendritic cells: controllers of the immune system and a new promise for immunotherapy," *Annals of the New York Academy of Sciences*, vol. 987, no. 1, pp. 180–187, 2003.
- [3] K. Palucka and J. Banchereau, "Cancer immunotherapy via dendritic cells," *Nature Reviews Cancer*, vol. 12, no. 4, pp. 265–277, 2012.
- [4] N. Schmitt, R. Morita, L. Bourdery et al., "Human dendritic cells induce the differentiation of interleukin-21-producing T follicular helper-like cells through interleukin-12," *Immunity*, vol. 31, no. 1, pp. 158–169, 2009.
- [5] J. I. Mayordomo, T. Zorina, W. J. Storkus et al., "Bone marrow-derived dendritic cells pulsed with synthetic tumour peptides elicit protective and therapeutic antitumour immunity," *Nature Medicine*, vol. 1, no. 12, pp. 1297–1302, 1995.
- [6] M. Ferrantini, I. Capone, and F. Belardelli, "Dendritic cells and cytokines in immune rejection of cancer," *Cytokine & Growth Factor Reviews*, vol. 19, no. 1, pp. 93–107, 2008.
- [7] K. Hoebe, E. Janssen, and B. Beutler, "The interface between innate and adaptive immunity," *Nature Immunology*, vol. 5, no. 10, pp. 971–974, 2004.
- [8] F. E. González, A. Gleisner, F. Falcon-Beas, F. Osorio, M. N. López, and F. Salazar-Onfray, "Tumor cell lysates as immunogenic sources for cancer vaccine design," *Human Vaccines & Immunotherapeutics*, vol. 10, no. 11, pp. 3261–3269, 2014.
- [9] A. Escobar, M. Lopez, A. Serrano et al., "Dendritic cell immunizations alone or combined with low doses of interleukin-2 induce specific immune responses in melanoma patients," *Clinical & Experimental Immunology*, vol. 142, no. 3, pp. 555–568, 2005.
- [10] M. N. Lopez, C. Pereda, G. Segal et al., "Prolonged survival of dendritic cell-vaccinated melanoma patients correlates with tumor-specific delayed type IV hypersensitivity response and reduction of tumor growth factor β -expressing T cells," *Journal of Clinical Oncology*, vol. 27, no. 6, pp. 945–952, 2009.
- [11] R. Aguilera, C. Saffie, A. Tittarelli et al., "Heat-shock induction of tumor-derived danger signals mediates rapid monocyte differentiation into clinically effective dendritic cells," *Clinical Cancer Research*, vol. 17, no. 8, pp. 2474–2483, 2011.
- [12] A. Tittarelli, F. E. González, C. Pereda et al., "Toll-like receptor 4 gene polymorphism influences dendritic cell in vitro function and clinical outcomes in vaccinated melanoma patients," *Cancer Immunology, Immunotherapy*, vol. 61, no. 11, pp. 2067–2077, 2012.
- [13] C. Duran-Aniotz, G. Segal, L. Salazar et al., "The immunological response and post-treatment survival of DC-vaccinated melanoma patients are associated with increased Th1/Th17 and reduced Th3 cytokine responses," *Cancer Immunology, Immunotherapy*, vol. 62, no. 4, pp. 761–772, 2013.
- [14] H. Shi, T. Cao, J. E. Connolly et al., "Hyperthermia enhances CTL cross-priming," *Journal of Immunology*, vol. 176, no. 4, pp. 2134–2141, 2006.
- [15] A. Mendoza-Naranjo, P. J. Saez, C. C. Johansson et al., "Functional gap junctions facilitate melanoma antigen transfer and cross-presentation between human dendritic cells," *Journal of Immunology*, vol. 178, no. 11, pp. 6949–6957, 2007.
- [16] G. Y. Chen and G. Nunez, "Sterile inflammation: sensing and reacting to damage," *Nature Reviews Immunology*, vol. 10, no. 12, pp. 826–837, 2010.
- [17] A. D. Garg, L. Galluzzi, L. Apetoh et al., "Molecular and translational classifications of DAMPs in immunogenic cell death," *Frontiers in Immunology*, vol. 6, p. 588, 2015.
- [18] F. E. González, C. Ortiz, M. Reyes et al., "Melanoma cell lysate induces CCR7 expression and in vivo migration to draining lymph nodes of therapeutic human dendritic cells," *Immunology*, vol. 142, no. 3, pp. 396–405, 2014.
- [19] P. J. Tacken, I. J. de Vries, R. Torensma, and C. G. Figg, "Dendritic-cell immunotherapy: from ex vivo loading to in vivo targeting," *Nature Reviews Immunology*, vol. 7, no. 10, pp. 790–802, 2007.
- [20] M. E. Ritchie, B. Phipson, D. Wu et al., "Limma powers differential expression analyses for RNA-sequencing and microarray studies," *Nucleic Acids Research*, vol. 43, no. 7, article e47, 2015.
- [21] D. Szklarczyk, A. Franceschini, S. Wyder et al., "STRING v10: protein-protein interaction networks, integrated over the tree of life," *Nucleic Acids Research*, vol. 43, no. D1, pp. D447–D452, 2015.
- [22] K. Dokladny, O. B. Myers, and P. L. Moseley, "Heat shock response and autophagy-cooperation and control," *Autophagy*, vol. 11, no. 2, pp. 200–213, 2015.
- [23] N. Vigneron and B. J. Van den Eynde, "Proteasome subtypes and regulators in the processing of antigenic peptides presented by class I molecules of the major histocompatibility complex," *Biomolecules*, vol. 4, no. 4, pp. 994–1025, 2014.
- [24] C. Jamart, M. Francaux, G. Y. Millet, L. Deldicque, D. Frère, and L. Féasson, "Modulation of autophagy and ubiquitin-proteasome pathways during ultra-endurance running," *Journal of Applied Physiology*, vol. 112, no. 9, pp. 1529–1537, 2012.
- [25] J. A. Martina, H. I. Diab, L. Lishu et al., "The nutrient-responsive transcription factor TFE3 promotes autophagy, lysosomal biogenesis, and clearance of cellular debris," *Science Signaling*, vol. 7, no. 309, article ra9, 2014.

- [26] T. Geuens, D. Bouhy, and V. Timmerman, "The hnRNP family: insights into their role in health and disease," *Human Genetics*, vol. 135, no. 8, pp. 851–867, 2016.
- [27] N. R. Hope and G. I. Murray, "The expression profile of RNA-binding proteins in primary and metastatic colorectal cancer: relationship of heterogeneous nuclear ribonucleoproteins with prognosis," *Human Pathology*, vol. 42, no. 3, pp. 393–402, 2011.
- [28] W. Y. Yau, H. C. Shih, M. H. Tsai, J. C. Sheu, C. H. Chen, and L. P. Chow, "Autoantibody recognition of an N-terminal epitope of hnRNP L marks the risk for developing HBV-related hepatocellular carcinoma," *Journal of Proteomics*, vol. 94, pp. 346–358, 2013.
- [29] D. Lv, H. Wu, R. Xing et al., "HnRNP-L mediates bladder cancer progression by inhibiting apoptotic signaling and enhancing MAPK signaling pathways," *Oncotarget*, vol. 8, no. 8, pp. 13586–13599, 2017.
- [30] X. Cao, Y. Li, R. Z. Luo et al., "Tyrosine-protein phosphatase nonreceptor type 12 is a novel prognostic biomarker for esophageal squamous cell carcinoma," *The Annals of Thoracic Surgery*, vol. 93, no. 5, pp. 1674–1680, 2012.
- [31] X. Cao, Y. Z. Chen, R. Z. Luo et al., "Tyrosine-protein phosphatase non-receptor type 12 expression is a good prognostic factor in resectable non-small cell lung cancer," *Oncotarget*, vol. 6, no. 13, pp. 11704–11713, 2015.
- [32] N. Yatim, S. Cullen, and M. L. Albert, "Dying cells actively regulate adaptive immune responses," *Nature Reviews Immunology*, vol. 17, no. 4, pp. 262–275, 2017.
- [33] R. Asleh, S. Marsh, M. Shilkret et al., "Genetically determined heterogeneity in hemoglobin scavenging and susceptibility to diabetic cardiovascular disease," *Circulation Research*, vol. 92, no. 11, pp. 1193–1200, 2003.
- [34] M. J. Nielsen, H. J. Moller, and S. K. Moestrup, "Hemoglobin and heme scavenger receptors," *Antioxidants & Redox Signaling*, vol. 12, no. 2, pp. 261–273, 2010.
- [35] Y. Xie, Y. Li, Q. Zhang, M. J. Stiller, C. L. A. Wang, and J. W. Strelein, "Haptoglobin is a natural regulator of Langerhans cell function in the skin," *Journal of Dermatological Science*, vol. 24, no. 1, pp. 25–37, 2000.
- [36] B. Ye, D. W. Cramer, S. J. Skates et al., "Haptoglobin- α subunit as potential serum biomarker in ovarian cancer: identification and characterization using proteomic profiling and mass spectrometry," *Clinical Cancer Research*, vol. 9, no. 8, pp. 2904–2911, 2003.
- [37] N. Ahmed, G. Barker, K. T. Oliva et al., "Proteomic-based identification of haptoglobin-1 precursor as a novel circulating biomarker of ovarian cancer," *British Journal of Cancer*, vol. 91, no. 1, pp. 129–140, 2004.
- [38] N. Okuyama, Y. Ide, M. Nakano et al., "Fucosylated haptoglobin is a novel marker for pancreatic cancer: a detailed analysis of the oligosaccharide structure and a possible mechanism for fucosylation," *International Journal of Cancer*, vol. 118, no. 11, pp. 2803–2808, 2006.
- [39] Y. Takeda, S. Shinzaki, K. Okudo, K. Moriwaki, K. Murata, and E. Miyoshi, "Fucosylated haptoglobin is a novel type of cancer biomarker linked to the prognosis after an operation in colorectal cancer," *Cancer*, vol. 118, no. 12, pp. 3036–3043, 2012.
- [40] H. Shen, Y. Song, C. M. Colangelo et al., "Haptoglobin activates innate immunity to enhance acute transplant rejection in mice," *The Journal of Clinical Investigation*, vol. 122, no. 1, pp. 383–387, 2012.
- [41] H. Shen, E. Heuzey, D. N. Mori et al., "Haptoglobin enhances cardiac transplant rejection," *Circulation Research*, vol. 116, no. 10, pp. 1670–1679, 2015.
- [42] H. Yang, H. Wang, Y. A. Levine et al., "Identification of CD163 as an antiinflammatory receptor for HMGB1-haptoglobin complexes," *JCI Insight*, vol. 1, no. 7, 2016.
- [43] P. J. Good, Q. Chen, S. J. Warner, and D. C. Herring, "A family of human RNA-binding proteins related to the Drosophila Bruno translational regulator," *The Journal of Biological Chemistry*, vol. 275, no. 37, pp. 28583–28592, 2000.
- [44] A. Castello, B. Fischer, K. Eichelbaum et al., "Insights into RNA biology from an atlas of mammalian mRNA-binding proteins," *Cell*, vol. 149, no. 6, pp. 1393–1406, 2012.
- [45] A. G. Baltz, M. Munschauer, B. Schwanhausser et al., "The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts," *Molecular Cell*, vol. 46, no. 5, pp. 674–690, 2012.
- [46] P. Iakova, G. L. Wang, L. Timchenko et al., "Competition of CUGBP1 and calreticulin for the regulation of p21 translation determines cell fate," *The EMBO Journal*, vol. 23, no. 2, pp. 406–417, 2004.
- [47] A. N. Ladd, N. Charlet-B, and T. A. Cooper, "The CELF family of RNA binding proteins is implicated in cell-specific and developmentally regulated alternative splicing," *Molecular and Cellular Biology*, vol. 21, no. 4, pp. 1285–1296, 2001.
- [48] S. Paul, W. Dansithong, D. Kim et al., "Interaction of muscle-blind, CUG-BP1 and hnRNP H proteins in DM1-associated aberrant IR splicing," *The EMBO Journal*, vol. 25, no. 18, pp. 4271–4283, 2006.
- [49] R. S. Laishram and R. A. Anderson, "The poly A polymerase star-PAP controls 3'-end cleavage by promoting CPSF interaction and specificity toward the pre-mRNA," *The EMBO Journal*, vol. 29, no. 24, pp. 4132–4145, 2010.
- [50] K. Ichikawa, Y. Kubota, T. Nakamura et al., "MCRIP1, an ERK substrate, mediates ERK-induced gene silencing during epithelial-mesenchymal transition by regulating the co-repressor CtBP," *Molecular Cell*, vol. 58, no. 1, pp. 35–46, 2015.
- [51] R. Bish, N. Cuevas-Polo, Z. Cheng et al., "Comprehensive protein interactome analysis of a key RNA helicase: detection of novel stress granule proteins," *Biomolecules*, vol. 5, no. 3, pp. 1441–1466, 2015.
- [52] A. A. Rose, M. G. Annis, Z. Dong et al., "ADAM10 releases a soluble form of the GPNMB/osteoactivin extracellular domain with angiogenic properties," *PLoS One*, vol. 5, no. 8, article e12093, 2010.
- [53] J. S. Chung, M. Bonkobara, M. Tomihari, P. D. Cruz Jr, and K. Ariizumi, "The DC-HIL/syndecan-4 pathway inhibits human allogeneic T-cell responses," *European Journal of Immunology*, vol. 39, no. 4, pp. 965–974, 2009.
- [54] A. Lin, C. Li, Z. Xing et al., "The LINK-A lncRNA activates normoxic HIF1 α signalling in triple-negative breast cancer," *Nature Cell Biology*, vol. 18, no. 2, pp. 213–224, 2016.
- [55] M. A. Todd and D. J. Picketts, "PHF6 interacts with the nucleosome remodeling and deacetylation (NuRD) complex," *Journal of Proteome Research*, vol. 11, no. 8, pp. 4326–4337, 2012.
- [56] T. J. Loh, S. Cho, H. Moon et al., "hnRNP L inhibits CD44 V10 exon splicing through interacting with its upstream intron," *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, vol. 1849, no. 6, pp. 743–750, 2015.


- [57] P. Yao, A. A. Potdar, P. S. Ray et al., "The HILDA complex coordinates a conditional switch in the 3'-untranslated region of the VEGFA mRNA," *PLoS Biology*, vol. 11, no. 8, article e1001635, 2013.
- [58] D. T. Kuninger, T. Izumi, J. Papaconstantinou, and S. Mitra, "Human AP-endonuclease 1 and hnRNP-L interact with a nCaRE-like repressor element in the AP-endonuclease 1 promoter," *Nucleic Acids Research*, vol. 30, no. 3, pp. 823–829, 2002.
- [59] A. Kapralov, I. I. Vlasova, W. Feng et al., "Peroxidase activity of hemoglobin-haptoglobin complexes: covalent aggregation and oxidative stress in plasma and macrophages," *The Journal of Biological Chemistry*, vol. 284, no. 44, pp. 30395–30407, 2009.
- [60] L. Ciglianò, C. R. Pugliese, M. S. Spagnuolo, R. Palumbo, and P. Abrescia, "Haptoglobin binds the antiatherogenic protein apolipoprotein E - impairment of apolipoprotein E stimulation of both lecithin:cholesterol acyltransferase activity and cholesterol uptake by hepatocytes," *The FEBS Journal*, vol. 276, no. 21, pp. 6158–6171, 2009.
- [61] T. Kambara and M. Ikebe, "A unique ATP hydrolysis mechanism of single-headed processive myosin, myosin IX," *The Journal of Biological Chemistry*, vol. 281, no. 8, pp. 4949–4957, 2006.
- [62] R. Kong, F. Yi, P. Wen et al., "Myo9b is a key player in SLIT/ROBO-mediated lung tumor suppression," *The Journal of Clinical Investigation*, vol. 125, no. 12, pp. 4407–4420, 2015.
- [63] M. Nishikawa, S. Nishikawa, A. Inoue, A. H. Iwane, T. Yanagida, and M. Ikebe, "A unique mechanism for the processive movement of single-headed myosin-IX," *Biochemical and Biophysical Research Communications*, vol. 343, no. 4, pp. 1159–1164, 2006.
- [64] A. Inoue, J. Saito, R. Ikebe, and M. Ikebe, "Myosin IXb is a single-headed minus-end-directed processive motor," *Nature Cell Biology*, vol. 4, no. 4, pp. 302–306, 2002.
- [65] O. D. Weiner, M. C. Rentel, A. Ott et al., "Hem-1 complexes are essential for Rac activation, actin polymerization, and myosin regulation during neutrophil chemotaxis," *PLoS Biology*, vol. 4, no. 2, article e38, 2006.
- [66] M. Adams and Z. Jia, "Structural and biochemical analysis reveal pirins to possess quercetinase activity," *The Journal of Biological Chemistry*, vol. 280, no. 31, pp. 28675–28682, 2005.
- [67] F. Liu, I. Rehmani, S. Esaki et al., "Pirin is an iron-dependent redox regulator of NF- κ B," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 110, no. 24, pp. 9722–9727, 2013.
- [68] I. Miyazaki, S. Simizu, H. Okumura, S. Takagi, and H. Osada, "A small-molecule inhibitor shows that pirin regulates migration of melanoma cells," *Nature Chemical Biology*, vol. 6, no. 9, pp. 667–673, 2010.
- [69] R. Dechend, F. Hirano, K. Lehmann et al., "The Bcl-3 oncprotein acts as a bridging factor between NF-kappaB/Rel and nuclear co-regulators," *Oncogene*, vol. 18, no. 22, pp. 3316–3323, 1999.
- [70] S. B. Hooks, S. P. Ragan, and K. R. Lynch, "Identification of a novel human phosphatidic acid phosphatase type 2 isoform," *FEBS Letters*, vol. 427, no. 2, pp. 188–192, 1998.
- [71] N. Sahni, S. Yi, M. Taipale et al., "Widespread macromolecular interaction perturbations in human genetic disorders," *Cell*, vol. 161, no. 3, pp. 647–660, 2015.
- [72] X. Zhao, K. Southwick, H. L. Cardasis et al., "Peptidomic profiling of human cerebrospinal fluid identifies YPRPIHPA as a novel substrate for prolylcarboxypeptidase," *Proteomics*, vol. 10, no. 15, pp. 2882–2886, 2010.
- [73] G. V. Pusapatı, T. Eiseler, A. Rykx et al., "Protein kinase D regulates RhoA activity via rhotekin phosphorylation," *The Journal of Biological Chemistry*, vol. 287, no. 12, pp. 9473–9483, 2012.
- [74] G. Lu, J. Chen, L. A. Espinoza et al., "Protein kinase D 3 is localized in vesicular structures and interacts with vesicle-associated membrane protein 2," *Cellular Signalling*, vol. 19, no. 4, pp. 867–879, 2007.
- [75] Q. Yang, D. Co, J. Sommercorn, and N. K. Tonks, "Cloning and expression of PTP-PEST. A novel, human, nontransmembrane protein tyrosine phosphatase," *The Journal of Biological Chemistry*, vol. 268, no. 9, pp. 6622–6628, 1993.
- [76] H. Li, F. Yang, C. Liu et al., "Crystal structure and substrate specificity of PTPN12," *Cell Reports*, vol. 15, no. 6, pp. 1345–1358, 2016.
- [77] T. Sun, N. Aceto, K. L. Meerbrey et al., "Activation of multiple proto-oncogenic tyrosine kinases in breast cancer via loss of the PTPN12 phosphatase," *Cell*, vol. 144, no. 5, pp. 703–718, 2011.
- [78] A. J. Garton, M. R. Burnham, A. H. Bouton, and N. K. Tonks, "Association of PTP-PEST with the SH3 domain of p130cas; a novel mechanism of protein tyrosine phosphatase substrate recognition," *Oncogene*, vol. 15, no. 8, pp. 877–885, 1997.
- [79] R. Arroyo, M. Duran-Frigola, C. Berenguer, M. Soler-López, and P. Aloy, "Charting the molecular links between driver and susceptibility genes in colorectal cancer," *Biochemical and Biophysical Research Communications*, vol. 445, no. 4, pp. 734–738, 2014.
- [80] J. Petschnigg, B. Groisman, M. Kotlyar et al., "The mammalian-membrane two-hybrid assay (MaMTH) for probing membrane-protein interactions in human cells," *Nature Methods*, vol. 11, no. 5, pp. 585–592, 2014.
- [81] Y. Katoh, H. Imakagura, M. Futatsumori, and K. Nakayama, "Recruitment of clathrin onto endosomes by the Tom1-Tollip complex," *Biochemical and Biophysical Research Communications*, vol. 341, no. 1, pp. 143–149, 2006.
- [82] M. Franco, O. Furstoss, V. Simon, C. Benistant, W. J. Hong, and S. Roche, "The adaptor protein Tom1L1 is a negative regulator of Src mitogenic signaling induced by growth factors," *Molecular and Cellular Biology*, vol. 26, no. 5, pp. 1932–1947, 2006.
- [83] A. Hegele, A. Kamburov, A. Grossmann et al., "Dynamic protein-protein interaction wiring of the human spliceosome," *Molecular Cell*, vol. 45, no. 4, pp. 567–580, 2012.
- [84] D. Cerecedo, R. Mondragon, B. Cisneros, F. Martinez-Perez, D. Martinez-Rojas, and A. Rendon, "Role of dystrophins and utrophins in platelet adhesion process," *British Journal of Haematology*, vol. 134, no. 1, pp. 83–91, 2006.
- [85] J. L. Costantini, S. M. Cheung, S. Hou et al., "TAPP2 links phosphoinositide 3-kinase signaling to B-cell adhesion through interaction with the cytoskeletal protein utrophin: expression of a novel cell adhesion-promoting complex in B-cell leukemia," *Blood*, vol. 114, no. 21, pp. 4703–4712, 2009.

- [86] H. Inagaki, Y. Matsushima, K. Nakamura, M. Ohshima, T. Kadokawa, and Y. Kitagawa, "A large DNA-binding nuclear protein with RNA recognition motif and serine/arginine-rich domain," *The Journal of Biological Chemistry*, vol. 271, no. 21, pp. 12525–12531, 1996.
- [87] M. Ehsan, H. Jiang, K. L. Thomson, and K. Gehmlich, "When signalling goes wrong: pathogenic variants in structural and signalling proteins causing cardiomyopathies," *Journal of Muscle Research and Cell Motility*, vol. 38, no. 3-4, pp. 303–316, 2017.
- [88] S. Onishi, E. Adnan, J. Ishizaki et al., "Novel autoantigens associated with lupus nephritis," *PLoS One*, vol. 10, no. 6, article e0126564, 2015.
- [89] J. Ren and A. J. Davidoff, " α 2-Heremans Schmid glycoprotein, a putative inhibitor of tyrosine kinase, prevents glucose toxicity associated with cardiomyocyte dysfunction," *Diabetes/Metabolism Research and Reviews*, vol. 18, no. 4, pp. 305–310, 2002.
- [90] Z. Yan, K. Cui, D. M. Murray et al., "PBAF chromatin-remodeling complex requires a novel specificity subunit, BAF200, to regulate expression of selective interferon-responsive genes," *Genes & Development*, vol. 19, no. 14, pp. 1662–1667, 2005.
- [91] V. Legendre-Guillemain, M. Metzler, M. Charbonneau et al., "HIP1 and HIP12 display differential binding to F-actin, AP2, and clathrin. Identification of a novel interaction with clathrin light chain," *Journal of Biological Chemistry*, vol. 277, no. 22, pp. 19897–19904, 2002.
- [92] C. Wu, M. H. Ma, K. R. Brown et al., "Systematic identification of SH3 domain-mediated human protein-protein interactions by peptide array target screening," *Proteomics*, vol. 7, no. 11, pp. 1775–1785, 2007.
- [93] T. Rolland, M. Tasan, B. Charlotteaux et al., "A proteome-scale map of the human interactome network," *Cell*, vol. 159, no. 5, pp. 1212–1226, 2014.
- [94] Y. Maiweilidan, I. Klauza, and E. Kordeli, "Novel interactions of ankyrins-G at the costameres: the muscle-specific obscurin/titin-binding-related domain (OTBD) binds plectin and filamin C," *Experimental Cell Research*, vol. 317, no. 6, pp. 724–736, 2011.
- [95] W. B. Holmes and C. L. Moncman, "Nebulette interacts with filamin C," *Cell Motility and the Cytoskeleton*, vol. 65, no. 2, pp. 130–142, 2008.
- [96] G. Blandin, S. Marchand, K. Charton et al., "A human skeletal muscle interactome centered on proteins involved in muscular dystrophies: LGMD interactome," *Skeletal Muscle*, vol. 3, no. 1, p. 3, 2013.
- [97] C. A. Sargent, C. Young, S. Marsh, M. A. Ferguson-Smith, and N. A. Affara, "The glycerol kinase gene family: structure of the Xp gene, and related intronless retroposons," *Human Molecular Genetics*, vol. 3, no. 8, pp. 1317–1324, 1994.
- [98] J. H. Yik, R. Chen, R. Nishimura, J. L. Jennings, A. J. Link, and Q. Zhou, "Inhibition of P-TEFb (CDK9/cyclin T) kinase and RNA polymerase II transcription by the coordinated actions of HEXIM1 and 7SK snRNA," *Molecular Cell*, vol. 12, no. 4, pp. 971–982, 2003.
- [99] N. Czudnochowski, F. Vollmuth, S. Baumann, K. Vogel-Bachmayr, and M. Geyer, "Specificity of Hexim1 and Hexim2 complex formation with cyclin T1/T2, importin alpha and 7SK snRNA," *Journal of Molecular Biology*, vol. 395, no. 1, pp. 28–41, 2010.
- [100] M. Varjosalo, R. Sacco, A. Stukalov et al., "Interlaboratory reproducibility of large-scale human protein-complex analysis by standardized AP-MS," *Nature Methods*, vol. 10, no. 4, pp. 307–314, 2013.
- [101] W. Liu, Q. Ma, K. Wong et al., "Brd4 and JMJD6-associated anti-pause enhancers in regulation of transcriptional pause release," *Cell*, vol. 155, no. 7, pp. 1581–1595, 2013.
- [102] Z. Chen, A. J. Kastaniotis, I. J. Miinalainen, V. Rajaram, R. K. Wierenga, and J. K. Hiltunen, "17beta-hydroxysteroid dehydrogenase type 8 and carbonyl reductase type 4 assemble as a ketoacyl reductase of human mitochondrial FAS," *The FASEB Journal*, vol. 23, no. 11, pp. 3682–3691, 2009.
- [103] R. Venkatesan, S. K. Sah-Teli, L. O. Awoniyi et al., "Insights into mitochondrial fatty acid synthesis from the structure of heterotetrameric 3-ketoacyl-ACP reductase/3R-hydroxyacyl-CoA dehydrogenase," *Nature Communications*, vol. 5, p. 4805, 2014.
- [104] S. Ohno, K. Nishikawa, Y. Honda, and S. Nakajin, "Expression in *E. coli* and tissue distribution of the human homologue of the mouse Ke 6 gene, 17 β -hydroxysteroid dehydrogenase type 8," *Molecular and Cellular Biochemistry*, vol. 309, no. 1-2, pp. 209–215, 2008.
- [105] V. Slama and J. Dudek, "Stability of diluted solutions of peracetic acid," *Ceskoslovenská Farmacie*, vol. 17, no. 1, pp. 40–41, 1968.
- [106] K. K. Leung, R. J. Hause Jr., J. L. Barking, M. F. Ciaccio, C. P. Chuu, and R. B. Jones, "Enhanced prediction of Src homology 2 (SH2) domain binding potentials using a fluorescence polarization-derived c-Met, c-Kit, ErbB, and androgen receptor interactome," *Molecular & Cellular Proteomics*, vol. 13, no. 7, pp. 1705–1723, 2014.
- [107] X. Huang, Z. Shi, W. Wang et al., "Identification and characterization of a novel protein ISOC2 that interacts with p16INK4a," *Biochemical and Biophysical Research Communications*, vol. 361, no. 2, pp. 287–293, 2007.
- [108] A. Ando, Y. Y. Kikuti, H. Kawata et al., "Cloning of a new kinesin-related gene located at the centromeric end of the human MHC region," *Immunogenetics*, vol. 39, no. 3, pp. 194–200, 1994.
- [109] S. Li, L. Wang, M. Berman, Y. Y. Kong, and M. E. Dorf, "Mapping a dynamic innate immunity protein interaction network regulating type I interferon production," *Immunity*, vol. 35, no. 3, pp. 426–440, 2011.
- [110] S. Eskildsen, J. Justesen, M. H. Schierup, and R. Hartmann, "Characterization of the 2'-5'-oligoadenylate synthetase ubiquitin-like family," *Nucleic Acids Research*, vol. 31, no. 12, pp. 3166–3173, 2003.
- [111] H. Kristiansen, H. H. Gad, S. Eskildsen-Larsen, P. Despres, and R. Hartmann, "The oligoadenylylate synthetase family: an ancient protein family with multiple antiviral activities," *Journal of Interferon & Cytokine Research*, vol. 31, no. 1, pp. 41–47, 2011.
- [112] J. Donovan, M. Dufner, and A. Korennykh, "Structural basis for cytosolic double-stranded RNA surveillance by human oligoadenylylate synthetase 1," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 110, no. 5, pp. 1652–1657, 2013.
- [113] M. Y. Hein, N. C. Hubner, I. Poser et al., "A human interactome in three quantitative dimensions organized by stoichiometries and abundances," *Cell*, vol. 163, no. 3, pp. 712–723, 2015.


- [114] Y. Chao, Y. Xing, Y. Chen et al., "Structure and mechanism of the phosphotyrosyl phosphatase activator," *Molecular Cell*, vol. 23, no. 4, pp. 535–546, 2006.
- [115] E. Ogris, X. Du, K. C. Nelson et al., "A protein phosphatase methylesterase (PME-1) is one of several novel proteins stably associating with two inactive mutants of protein phosphatase 2A," *The Journal of Biological Chemistry*, vol. 274, no. 20, pp. 14382–14391, 1999.
- [116] S. O. Marx, S. Reiken, Y. Hisamatsu et al., "PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts," *Cell*, vol. 101, no. 4, pp. 365–376, 2000.
- [117] A. Murayama, K. Ohmori, A. Fujimura et al., "Epigenetic control of rDNA loci in response to intracellular energy status," *Cell*, vol. 133, no. 4, pp. 627–639, 2008.
- [118] R. Kekuda, P. D. Prasad, Y. J. Fei, F. H. Leibach, and V. Ganapathy, "Cloning and functional expression of the human type 1 sigma receptor (hSigmaR1)," *Biochemical and Biophysical Research Communications*, vol. 229, no. 2, pp. 553–558, 1996.
- [119] S. Figaro, N. Scrima, R. H. Buckingham, and V. Heurgué-Hamard, "HemK2 protein, encoded on human chromosome 21, methylates translation termination factor eRF1," *FEBS Letters*, vol. 582, no. 16, pp. 2352–2356, 2008.
- [120] L. Songe-Moller, E. van den Born, V. Leihne et al., "Mammalian ALKBH8 possesses tRNA methyltransferase activity required for the biogenesis of multiple wobble uridine modifications implicated in translational decoding," *Molecular and Cellular Biology*, vol. 30, no. 7, pp. 1814–1827, 2010.
- [121] D. Fu, J. A. Brophy, C. T. Chan et al., "Human AlkB homolog ABH8 is a tRNA methyltransferase required for wobble uridine modification and DNA damage survival," *Molecular and Cellular Biology*, vol. 30, no. 10, pp. 2449–2459, 2010.
- [122] T. Stuven, E. Hartmann, and D. Gorlich, "Exportin 6: a novel nuclear export receptor that is specific for profilin.actin complexes," *The EMBO Journal*, vol. 22, no. 21, pp. 5928–5940, 2003.



**The Scientific
World Journal**




Gastroenterology
Research and Practice



MEDIATORS
of
INFLAMMATION



Journal of
Diabetes Research



Disease Markers



Journal of
Immunology Research



PPAR Research




Hindawi

Submit your manuscripts at
www.hindawi.com




International Journal of
Endocrinology




BioMed
Research International




Journal of
Ophthalmology



Stem Cells
International




eCAM
Evidence-Based
Complementary and
Alternative Medicine




Journal of
Obesity



Journal of
Oncology




Computational and
Mathematical Methods
in Medicine




Behavioural
Neurology



Parkinson's
Disease



AIDS
Research and Treatment



Oxidative Medicine and
Cellular Longevity