

Review Article

Immunomodulatory Effects of 17β -Estradiol on Epithelial Cells during Bacterial Infections

Ivan Medina-Estrada,¹ Nayeli Alva-Murillo ,² Joel E. López-Meza ,³
and Alejandra Ochoa-Zarzosa ³

¹Trayectoria en Genómica Alimentaria, Universidad de la Ciénega del Estado de Michoacán de Ocampo, Sahuayo De Morelos, MICH, Mexico

²Departamento de Biología, División de Ciencias Naturales y Exactas, Universidad de Guanajuato, Guanajuato, GTO, Mexico

³Centro Multidisciplinario de Estudios en Biotecnología, Facultad de Medicina Veterinaria y Zootecnia, Universidad Michoacana de San Nicolás de Hidalgo, Morelia, MICH, Mexico

Correspondence should be addressed to Alejandra Ochoa-Zarzosa; ochoaz@umich.mx

Received 21 March 2018; Revised 26 June 2018; Accepted 24 July 2018; Published 29 August 2018

Academic Editor: Martin Holland

Copyright © 2018 Ivan Medina-Estrada et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The innate immune system can function under hormonal control. 17β -Estradiol (E2) is an important sexual hormone for the reproductive cycle of mammals, and it has immunomodulatory effects on epithelial cells, which are the first line of defense against incoming bacteria. E2 regulates various pathophysiological processes, including the response to infection in epithelial cells, and its effects involve the regulation of innate immune signaling pathways, which are mediated through estrogen receptors (ERs). E2 modulates the expression of inflammatory and antimicrobial elements such as cytokines and antimicrobial peptides. The E2 effects on epithelial cells during bacterial infections are characterized by an increase in the production of antimicrobial peptides and by the diminution of the inflammatory response to abrogate proinflammatory cytokine induction by bacteria. Here, we review several novel molecular mechanisms through which E2 regulates the innate immune response of epithelial cells against bacterial infections.

1. Introduction

17β -Estradiol, also known as E2 (due to its two hydroxyl groups), is a steroidal hormone derived from cholesterol and is the most predominant and potent sexual hormone during the reproductive stage of females [1]. The principal effects of E2 are associated with reproductive and sexual functions, although it is also involved in the development of different pathologies, such as cancer, autoimmune diseases, and infectious processes, where the innate immune response (IIR) can be altered by this hormone [2].

The main immunomodulatory effects of E2 have been described in the epithelium of the reproductive tract of females, where it modulates the IIR helping to maintain the microbiota in addition to favoring an anti-inflammatory environment [3–5]. E2 can also regulate the differentiation and proliferation of epithelial cells in several tissues (e.g., epithelial mammary cells) [6, 7]. The anti-inflammatory

properties of E2 have been demonstrated in different epithelial models, such as the human uterus, rat oviduct, and mouse intestine, and it is well known that E2 can reverse the inflammatory response induced by bacterial components or whole pathogens. In addition, in some models, it has been reported that E2 improves the antimicrobial response against different pathogens inducing the production of antimicrobial peptides. To date, there is little information concerning the immunomodulatory effects of E2 in epithelial cells during infections [3, 8, 9]. Thus, the objective of this review was to provide the current knowledge on the pathways and elements that participate in the immunomodulatory effects of E2 in epithelial cells during bacterial infections.

2. General Aspects of E2

E2 is the most abundant and potent natural estrogen in all vertebrates. E2 is a C18 steroid derived from cholesterol

and is the main estrogen present in the serum of mammal females at reproductive age. In humans and other vertebrates, estrogens are made primarily in the female ovaries, but small amounts of this hormone may be produced in the male testes as well as in the adrenal glands, brain, and fat tissue of both sexes. E2 synthesis involves several chemical intermediates, such as progestagens and androgens (e.g., testosterone), which are derived from cholesterol (Figure 1). Testosterone undergoes conversion to estradiol by aromatase (P450aro = CYP19), resulting in the loss of one carbon (C19) and the aromatization of the A-ring, thereby producing E2. An alternative pathway to synthesize E2 occurs through the conversion of androstenedione to estrone (via aromatase), which can be enzymatically converted to E2 [10].

The main targets of E2 are the reproductive female organs (ovary, uterus, and mammary gland). E2 acts in a vast variety of cell populations; however, its main targets are epithelial cells. This hormone regulates epithelial cell differentiation, proliferation, and apoptosis, which are processes intimately linked to the female reproductive cycle such as lactation or ovulation [11–13]. E2 functions are diverse and play important roles in the regulation of metabolism (body fat distribution), hemodynamics (vascular tone), reproduction (ovulation and development of primary and secondary sexual characters, including growth and involution of the mammary gland), and pathological processes, such as cancer and diseases, where the immune response is compromised [11, 14, 15].

2.1. Estrogen Receptors. The effects of E2 are mediated by estrogen receptors (ERs), which are classified as ER α (565 aa) and ER β (530 aa), and these receptors were characterized in 1973 for the first time in a rat uterus [16]. ERs belong to the superfamily of nuclear receptors, functioning as transcriptional factors or associating with other transcriptional factors to regulate the expression of many genes related to the functions described for E2 [17]. In addition to nuclear receptors, there is a third ER, namely GPR30, which belongs to the G protein-coupled receptor family. The GPR30 receptor is located in the endoplasmic reticulum and plasma membrane and is mainly associated with nongenomic responses [16, 18].

The nuclear receptors ER α and ER β have well defined the functional and structural subunits. Both ERs remain inactive when they are bound to the heat shock protein complexes Hsp70 and Hsp90; however, binding with their ligands (e.g., E2) induces a conformational change, which favors the dissociation of the ER complex and promotes ER activation and translocation to the nucleus. In the nucleus, ER interacts with estrogen response elements (EREs) present in the DNA sequence (the minimum consensus sequence is a palindromic repeat sequence of 5'-GGTCAnnTGACC-3'). EREs are present in many gene promoters such as oxytocin, *c-fos*, *c-myc*, TGF- α , prolactin, complement proteins, and lactoferrin [19]. ERs can also be activated independent of estrogen binding (nonclassical signaling). This kind of ER activation is the consequence of several posttranslational modifications (phosphorylation, acetylation, and methylation) induced by different enzymes [20].

E2 levels vary depending on the species and its reproductive stage. The range in mammal females oscillates between 50 pg/ml (standard for basal condition) and 1500 pg/ml (standard for the preovulatory phase), which is associated with the effects that E2 have on reproduction and is the reason why sexual hormone levels are varied in mammals [21, 22]. ERs as well as E2 have been associated with some pathologies, such as cancer, being breast cancer the principal disease in which ERs are involved [23, 24]. Nevertheless, ERs and E2 also play an important role in the IIR to different stimuli. For example, females are more susceptible to develop inflammatory lung conditions and sex hormones have been implicated in this process [25]. However, the immune mechanisms responsible for sex-based disparity are unknown. The data suggest that specific sex hormone influences the IIR [1]. Accordingly, ERs can modulate the activation of transcriptional factors associated with the IIR, such as NF- κ B, SP1, AP-1, and C/EBP β . As a consequence, ERs regulate the transcription of elements of the inflammatory response, such as proinflammatory cytokines, anti-inflammatory cytokines, antimicrobial peptides, and chemokines. In addition, in response to their ligands, ERs regulate different signaling pathways, which can also be associated with the IIR [26].

GPR30 is associated with cellular estrogen functions, including canonical genomic signaling and novel nongenomic responses. The nongenomic events include production of second messengers (Ca²⁺ and cAMP), activation of tyrosine kinase receptors (such as EGFR and IGF-IR), and activation of different protein/lipid kinase pathways (PI3K, Akt, MAPK, Scr, and PKA/PKC) [16]. Thus, GPR30 signaling pathways are also related to the regulation of the IIR. However, the information concerning these functions is scarce.

3. Hormonal Regulation of the Innate Immune Response

Diverse hormones have been reported to participate in the modulation of the IIR, which is activated by different stimuli such as those occurring in infectious diseases or stress conditions, which can modulate the inflammatory response. This response during infections is mainly characterized by the production of proinflammatory cytokines, namely tumor necrosis factor alpha (TNF- α), interleukin-1 beta (IL-1 β), interleukin-2 (IL-2), and interleukin-6 (IL-6), as well as chemokines. These molecules are secreted by the injured tissue as well as by innate cells such as macrophages, dendritic cells, and mast cells in order to recruit neutrophils and monocytes to the affected tissue. Through the release of antimicrobial molecules and active phagocytosis, innate immune cells act to eliminate the foreign agent and recover tissue homeostasis [27]. Sexual hormones play an important role during the IIR to infection in particular estradiol, progesterone, and androgens [28].

On the other hand, the stress response stimulates the hypothalamic-pituitary-adrenal (HPA) axis, inducing the release of the corticotrophin-releasing hormone (CRH) from the hypothalamus to the pituitary portal system. Corticotrophin is then released from the pituitary gland to the




FIGURE 1: Estradiol synthesis. The production of estradiol can occur in different cell types. Estrogen is a product derived from cholesterol by a series of reactions throughout estrogen biosynthesis. The aromatase enzyme participates in the last step in E2 synthesis. Aromatase is a member of the cytochrome P450 superfamily and is widely expressed in many cell types. As a result of these reactions, an 18-carbon molecule is produced with two hydroxyl groups in its molecular structure. The resulting molecule also contains an aromatic ring with a hydroxyl group on carbon 3 and a β -hydroxyl group or ketone at position 17 of ring D. The phenolic ring A is the main structural feature in which selective and high-affinity binding to estrogen receptors occurs [10, 23].

systemic circulation, stimulating the suprarenal cortex to liberate glucocorticoids [29]. At the same time, stress induces the activation of the sympathetic nervous system, which leads to catecholamine release from the autonomic nervous terminals and the adrenal medulla [30]. Glucocorticoids and catecholamines have well-known immunoregulatory functions, namely, reducing the inflammatory response activated by the infection, which leads to the production of anti-inflammatory cytokines [29].

In addition to the HPA axis, other hormones have been implicated in the regulation of the IIR. Indeed, receptors for many hormones are expressed in cells from the innate and adaptive immune systems, which is the case for prolactin, growth hormone, thyroid hormone, or estrogen. The immunomodulatory effects of the hormones depend on the type of tissue, the hormone concentration, the physiological context, or the activated signaling pathway [31].

With regard to estradiol, its immunomodulatory properties comprise both the development as well as the function of innate immune cells. In this sense, ERs are expressed in immune cells triggering different signaling pathways. Estradiol exhibits both proinflammatory and anti-inflammatory responses related to hormone concentrations [26].

The transition to menopause, characterized by declining estradiol levels, has an impact on immunosenescence increasing the susceptibility to infection diseases and decreasing the efficacy of vaccination. These effects in postmenopausal women are characterized by an upregulation in the production of the cytokines TNF- α , IL-1 β , IL-10, and IL-6, the stimulation of the cytotoxic activity of NK cells, and a diminished phagocytic capacity of dendritic cells (DCs), which leads to impaired antigen presentation and activation of the adaptive immune system. Upon hormone replacement therapy, these age-related changes can be partially




FIGURE 2: Signal transduction pathways of epithelial cells involved in the recognition of pathogenic bacteria. Epithelial cells are part of the innate immune system, which senses the presence of microbial pathogen-associated molecular patterns (PAMPs) present in bacteria. These elements activate inflammatory pathways mainly through the activation of Toll-like receptors (TLRs). TLR proteins are coupled to the scaffold proteins, myeloid differentiation primary response 88 (MyD88), or TIR domain-containing adapter-inducing interferon- β (TRIF), which are associated with interleukin-1 receptor-associated kinase 1/4 (IRAK1/4). This kinase is coupled to TNF receptor-associated factor (TRAF) [38]. TLR signaling results in the downstream activation of the following three major families of proteins important in activating inflammatory gene expression: interferon regulatory factors (IRFs); mitogen-activated protein kinase (MAPK) pathway, such as c-Jun N-terminal kinase (JNK), protein 38 (P38), and extracellular signal-regulated kinases (ERKs); and the canonical inflammatory pathway, namely, nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B). However, other transcriptional factors, such as activator protein 1 (AP-1), can also be activated. Bacteria induces the expression of many inflammatory genes and cytokine secretion, such as interleukin 1-beta (IL-1 β), interleukin-6 (IL-6), and tumor necrosis factor-alpha (TNF- α) [36]. Intracellular bacteria can be recognized in epithelial cells by TLRs (endosome) or Nod-like receptors (NODs), which can activate the inflammasome. This is a multiprotein complex activated by caspases, and its activation and composition depend on the activator, which initiates inflammasome assembly, resulting in the processing of inflammatory cytokines such as IL-1 β and interleukin 18 (IL-18) [63].

prostaglandin release, glucose metabolism, eosinophil infiltration, and lipid and protein synthesis [41]. In addition, E2 induces ER α recruitment to ERE sites in target genes of mouse uterus, which leads to RNA and DNA syntheses, epithelial cell proliferation, and their differentiation toward a columnar secretory epithelium [42]. These effects are achieved at long time periods (after 24–72 h).

Otherwise, vaginal epithelial cells respond to E2 by undergoing cornification (production of keratins and involucrin), a process that involves both proliferation and differentiation. These effects are mediated by ER α in a direct way as well as through a paracrine route (involving stroma cells) [43].

E2 also modulates the permeability of the lower female reproductive tract (vagina and ectocervix). Epithelial cells

are linked by tight junction proteins, regulating the traffic of molecules across the epithelium. In the lower female reproductive tract, the stratified squamous epithelium shows tight junctions between basal epithelial cells. E2 increases the relaxation of epithelial tight junctions, which induces the flux across the epithelium. These effects are mediated by the expression of claudin and occludin [9, 44].

E2 also promotes lactobacillus growth in vaginal epithelial cells by increasing the storage of glycogen in the suprabasal and apical layers [45]. Glycogen is a substrate for acid production by these bacteria maintaining a low-pH environment [7]. Epithelial cells from the urinary tract (bladder epithelium) are also influenced by E2, and these cells play an important role of protection from infectious diseases where E2 has a relevant function, increasing the production

of antimicrobial peptides and tightening intercellular connection, thereby preventing bacteria to reach the cells where they can hide and cause infection [7]. *In vitro* and *in vivo* studies indicate that the production of different antimicrobial peptides increases in bladder epithelial cells after E2 treatment (see below).

In addition, it has been shown in rabbit bladders that E2 increases the volume of smooth muscle cells as well as vascularization, promoting bladder contraction [46]. In the absence of E2, the residual urine increases, whereas the urine flow is reduced, impairing the mechanical clearance of bacteria [47].

4.1. Immunomodulatory Effects of E2 in Epithelial Cells. The epithelial IIR is fast and acts directly upon the pathogen attack or tissue damage [48]. Several studies have demonstrated the regulation of the IIR of epithelia by E2. In all of the epithelial models where E2 plays an immunomodulatory role, E2 has also been demonstrated to have anti-inflammatory effects. In human epithelial cells from the endometrium, E2 (10^{-7} M) shows an anti-inflammatory action reversing the effects caused by PAMPs, such as Poly I:C [49]. In addition, E2 can block the inflammatory effects caused by IL-1 β and lipopolysaccharide (LPS) from *Escherichia coli* in human and rat uterus [6, 49]. In bovine oviduct, LPS stimulation increases the secretion of IL-1 β and TNF- α and it induces the expression of TLR2 and TLR4. However, E2 (1 ng/ml) reverses these inflammatory effects [44]. In the same way, the production of nitric oxide induced via mechanical damage or LPS stimulation in rat intestinal epithelium is diminished by E2 [3, 50].

Despite the aforementioned evidence, the relation of E2 with the IIR during infectious processes has not been properly analyzed because epithelial cells are challenged with PAMPs or proinflammatory cytokines instead of whole bacteria, in most of the models.

Experimental evidence suggests that E2 performs its immunomodulatory effects through the regulation of TLRs because they are the main PRRs that trigger the IIR signaling pathways. E2 (10^{-8} M) through ER α modulates the signaling of TLR3 (which recognizes double-stranded RNA) in the human reproductive tract during the beginning of the gestational cycle. Interestingly, the increase of E2 reduces the abundance of TLR3 favoring an anti-inflammatory environment [51].

E2 also regulates different inflammatory transcriptional factors. For example, in rat aortic smooth muscle cells, E2 inhibits the activation of NF- κ B, thus avoiding its dissociation from the I κ B inhibitor [52]. In addition, it is known that in both human uterine and urinary epithelial cells, E2 can modulate the activation of other transcriptional factors such as AP-1, SP1, STAT, and vitamin D receptor (VDR), which also participate in the regulation of the IIR. In this sense, E2 inhibits STAT phosphorylation and promotes the activation of VDR, modulating the MAPK signaling pathway (Figure 3) [6, 7, 53]. Finally, in human endometrium, E2 (10^{-8} M or 10^{-9} M) inhibits the protein levels of the IL-1 β receptor, thus reversing the stimulatory effects of IL-1 β on mRNA expression of TNF- α , IL-8, and NF- κ B [6]. A

summary of the effects of E2 on the IIR of epithelial cells is shown in Figure 3.

4.2. Antimicrobial Activity of E2 against Bacterial Infections in Epithelial Cells. Antimicrobial peptides (AMPs) are a diverse group of molecules with a broad range of activities, including cytotoxic, antifungal, antibacterial, or immunomodulatory effects. In mammals, these molecules are produced by different types of cells, highlighting those that are in intimate contact with pathogens such as epithelial cells [54]. In different models of infection, E2 exerts antibacterial activity against many pathogens because it favors the expression and secretion of AMPs. Human beta defensin-2 (HBD-2) is one of the most relevant AMPs in the human reproductive tract, and its secretion is regulated by E2. For example, human uterine epithelium exposed to LPS in the presence of E2 (10^{-3} M) shows increased secretion of the AMPs, calcineurin-like metallophosphoesterase superfamily protein (SLP1), and HBD-2, as well as decreased production of pro-inflammatory cytokines via NF- κ B inhibition [3]. Moreover, in vaginal epithelium, E2 (2 nM) induces HBD-2 secretion in response to LPS [55]. In addition, growth inhibition of *S. aureus* in the presence of E2 has been reported in a rat uterus and this inhibition is attributed to AMP secretion in the uterine lumen [8]. In the urinary tracts of women treated with E2 (10^{-3} M), a lesser recurrence of *Escherichia coli* infections has been observed, which is associated with the production of AMPs, such as defensins (HBD-1 and HBD-2), and the homeostasis of the common microbial flora [3]. In similar models (postmenopausal women treated with estrogens), a diminished risk of bacterial infections has been observed due to the increase of mRNA coding for the AMPs HBD-1 and HBD-3, as well as psoriasin and cAMP [3, 6, 8].

During the menstrual cycle, the fluctuation of hormonal concentrations leads to immune changes in the female reproductive tract (FRT). These changes vary depending on the location (upper or lower FRT) and the hormonal concentrations, which fluctuate differentially across the tissues in relation to their levels in circulation. At mid-cycle when E2 levels are high, there is a window of vulnerability for the establishment of infections in the lower FRT (viral, bacterial, or fungal) as a result of the inhibition in the production of AMPs (such as HBD-2). By contrast, in the upper FRT (uterus) persists a reduced activity of cytotoxic cell at a time when the production of AMPs is enhanced optimizing the conditions for a successful implantation [9]. In addition, during the proliferative stage of the menstrual cycle, cervical/vaginal secretion contains higher levels of other AMPs in relation with other stages of the cycle (medium and secretory phases), for example, the secretory leukocyte protease inhibitor-1 (SLP1), HBD-1, human neutrophil proteins 1–3 (HNP1–3), or lactoferrin [5, 6, 56, 57].

In animal models, immunomodulatory effects of E2 also comprise the production of AMPs. For example, SBD-1 (sheep beta-defensin-1) defensin is induced in ovine oviduct epithelial cells by E2 (10^{-8} M) [5]. In male rabbits exposed to *S. aureus* (a strain causing toxic shock), E2 treatment protects against the infection [58]. Similarly, in rat uterus using *in vivo* and *in vitro* experiments, higher levels of E2 induce




FIGURE 3: Anti-inflammatory and antibacterial effects of E2 in epithelial cells. E2 mediates its effects in epithelium by binding to intracellular receptors (ERs), leading to the translocation of hormone-ER (ER phosphorylate) complexes into the nucleus where they modulate gene transcription through binding to estrogen-responsive elements (EREs) on DNA (5'GGTCAnnnTGACC-3' consensus sequence). There is another type of ER, namely, G protein-coupled receptor 30 (GPR30), which binds E2, tamoxifen, and ICI 182780 (Fulvestrant). This receptor promotes the activation of epidermal growth factor receptor (EGFR) and mediates signaling in breast, endometrial, and thyroid cancer cells. This receptor activates signaling pathways in the cytosol and at the plasma membrane [18]. However, most of the molecular targets activated by GPR30 located at the plasma membrane during infections are unknown. GPR30 also has effects on the IIR associated with nongenomic responses (e.g., nitric oxide production). When cells are infected by bacteria, they trigger the activation of proinflammatory signaling pathways, mainly depending on TLRs. These pathways (described above) promote the production of proinflammatory elements [16]. However, E2 can modulate these pathways, influencing the IIR and decreasing the proinflammatory elements by blocking the activation of transcriptional factors, such as NF- κ B (canonical inflammatory pathway), ultimately resulting in the downregulation of IL-1 β , IL-6, and TNF- α gene expression. In contrast, the E2-ER complex mainly induces the gene expression of AMPs, such as defensins, and promotes an anti-inflammatory environment by inducing interleukin-10 (IL-10) gene expression. Gene expression regulated by E2 depends on the concentrations of the hormone and the interaction with pathogenic bacteria. It is well known that E2 reverses the inflammatory effects (association with bacteria and proinflammatory cytokines) and promotes an anti-inflammatory environment [8, 60, 64, 65].

resistance against *E. coli* infection and this effect is attributed to the secretion of antimicrobial molecules [8]. In human bladder epithelial cells, E2 induces the production of defensins HBD1-3, ribonuclease (RNase) 7, psoriasin, and cathelicidin; these effects can be associated with a reduced *E. coli* infection [59]. These findings indicate that AMP production constitutes a relevant defense mechanism induced by E2 in epithelia.

Evidence from our group indicates that bovine mammary epithelial cells (bMECs) treated with E2 produce antimicrobial molecules (probably AMPs) as the conditioned medium of bMECs treated with E2 inhibits the viability and growth of *S. aureus*. In addition, bMECs treated with E2 show upregulated gene expression of antimicrobial molecules, such as DEFB1 (defensin beta 1), BNBD5 (bovine neutrophil beta-defensin 5), and S100A7 (psoriasin) [60]. These effects are achieved through the activation of transcriptional factors,

such as AP-1, NF1, or ER (data unpublished) (Figure 4). These events lead to the inhibition of *S. aureus* internalization, presumably through the inhibition of focal adhesion kinase (FAK) (Figure 4) [60].

Considering the roles of epithelial cells during the IIR (Section 3.1), the combination of an anti-inflammatory environment and the antimicrobial response is essential for the maintenance of homeostatic epithelial flora. During infections, the host initiates an inflammatory response, sometimes aggravated, which results in damage to the tissue. Nevertheless, E2 in epithelial cells from all of the models described (using PAMPs or whole pathogens) inactivates inflammatory elements but activates the production of antimicrobial molecules, presumably induced via AP-1 or SP-1, thereby favoring elimination of the pathogen. In addition, the production of AMPs by the host can enhance the epithelial inflammatory response accordingly to reports that




FIGURE 4: Effects of E2 on bovine mammary epithelial cells during *Staphylococcus aureus* infection. During *S. aureus* infection in bovine mammary epithelial cells (bMECs), E2 inhibits the mitogen-activated protein kinases (MAPKs), resulting in anti-inflammatory effects (decreasing gene expression of IL-1 β and IL-6 as well as increasing gene expression of IL-10). E2 also activates protein kinase B (Akt), which in turn activates transcriptional factors, such as activator protein 1 (AP-1), NF1, or ER, that are related to IIR gene expression, such as AMPs, IL-10, or TNF- α . Through the zipper mechanism, *S. aureus* is internalized into epithelial cells, preventing it from being detected by the defense system of the host cell. However, E2 can block the signaling pathway of the zipper mechanism through the inhibition of focal adhesion kinase (FAK), resulting in the reduction of *S. aureus* internalization. The production of antimicrobial molecules (e.g., antimicrobial peptides) by bMECs treated with E2 also reduces *S. aureus* viability. However, the identity of the AMPs secreted by bMECs remains unknown. By RT-qPCR, the upregulation in the gene expression of DEFB1 (defensin beta 1), BNBD5 (bovine neutrophil beta-defensin 5), and S100A7 (psoriasin) has been shown. E2 induces the expression of TLR2, a key receptor in the recognition of *S. aureus*, as well as upregulates the expression of its receptors (ER α/β) [54, 58, 60].

show the immunomodulatory properties of these molecules through the induction of chemotaxis, enhancing the phagocytic activity of macrophages and dendritic cells, upregulating the production of inflammatory cytokines, chemokines, and so on [54, 61]. Furthermore, it is important to highlight that the production of AMPs is not enough to counteract an infection, in particular infections caused by intracellular pathogens, which require of a correct inflammatory response [62].

5. Concluding Remarks and Future Directions

Epithelial cells constitute a fundamental part of the local inflammatory response and host defense against pathogens. Epithelium is a target of E2 effects where the hormone induces the production of antimicrobial molecules, thereby reinforcing its role as a defense barrier against microorganisms besides its immunomodulatory properties. In addition, AMPs can also enhance the IIR through the regulation of other immunomodulatory effects.

Antimicrobial and anti-inflammatory properties of E2 are highly relevant along the reproductive life of women. For example, in the middle of the menstrual cycle when E2

levels are high, women are more likely to experience infections in the lower FRT (viral, bacterial, or fungal) as consequence of reducing the production of AMPs by epithelial cells. These effects of E2 are selective in ways that vary according to its concentrations and the FRT site. Based on epidemiological data, women appear to lose their immunological advantage after menopause, since in aging, they are more susceptible to infections and to develop inflammatory diseases. However, the participation at these stages of the E2-induced antimicrobial response requires further research.

Results discussed in this review may lead to the development of new hormonal strategies that would improve and enhance the IIR of epithelial cells against infections at the different reproductive stages in women.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Acknowledgments

Ivan Medina-Estrada was supported by a scholarship from Consejo Nacional de Ciencia y Tecnología. This work was

supported by grants from CIC14.1 to Alejandra Ochoa-Zarzosa and Consejo Nacional de Ciencia y Tecnología CB2016 287210.

References

- [1] S. Abid, S. Xie, M. Bose et al., "17 β -estradiol dysregulates innate immune responses to *Pseudomonas aeruginosa* respiratory infection and is modulated by estrogen receptor antagonism," *Infection and Immunity*, vol. 85, no. 10, 2017.
- [2] S. Nadkarni and S. McArthur, "Oestrogen and immunomodulation: new mechanisms that impact on peripheral and central immunity," *Current Opinion in Pharmacology*, vol. 13, no. 4, pp. 576–581, 2013.
- [3] M. V. Patel, J. V. Fahey, R. M. Rossoll, and C. R. Wira, "Innate immunity in the vagina (part I): estradiol inhibits HBD2 and elafin secretion by human vaginal epithelial cells," *American Journal of Reproductive Immunology*, vol. 69, no. 5, pp. 463–474, 2013.
- [4] M. L. Turner, J. G. Cronin, G. D. Healey, and I. M. Sheldon, "Epithelial and stromal cells of bovine endometrium have roles in innate immunity and initiate inflammatory responses to bacterial lipopeptides in vitro via Toll-like receptors TLR2, TLR1, and TLR6," *Endocrinology*, vol. 155, no. 4, pp. 1453–1465, 2014.
- [5] S. Wen, G. Cao, T. Bao et al., "Modulation of ovine SBD-1 expression by 17 β -estradiol in ovine oviduct epithelial cells," *BMC Veterinary Research*, vol. 8, no. 1, p. 143, 2012.
- [6] J. V. Fahey, J. A. Wright, L. Shen et al., "Estradiol selectively regulates innate immune function by polarized human uterine epithelial cells in culture," *Mucosal Immunology*, vol. 1, no. 4, pp. 317–325, 2008.
- [7] P. L uthje, A. Lind en Hirschberg, and A. Brauner, "Estrogenic action on innate defense mechanisms in the urinary tract," *Maturitas*, vol. 77, no. 1, pp. 32–36, 2014.
- [8] J. V. Fahey, R. M. Rossoll, and C. R. Wira, "Sex hormone regulation of anti-bacterial activity in rat uterine secretions and apical release of anti-bacterial factor(s) by uterine epithelial cells in culture," *The Journal of Steroid Biochemistry and Molecular Biology*, vol. 93, no. 1, pp. 59–66, 2005.
- [9] C. R. Wira, M. Rodr iguez-Garc a, and M. V. Patel, "The role of sex hormones in immune protection of the female reproductive tract," *Nature Reviews Immunology*, vol. 15, no. 4, pp. 217–230, 2015.
- [10] C. R. Nishida, S. Everett, and P. R. Ortiz de Montellano, "Specificity determinants of CYP1B1 estradiol hydroxylation," *Molecular Pharmacology*, vol. 84, no. 3, pp. 451–458, 2013.
- [11] A. Frank, L. M. Brown, and D. J. Clegg, "The role of hypothalamic estrogen receptors in metabolic regulation," *Frontiers in Neuroendocrinology*, vol. 35, no. 4, pp. 550–557, 2014.
- [12] I. Lamote, E. Meyer, A. M. Massart-Le en, and C. Burvenich, "Sex steroids and growth factors in the regulation of mammary gland proliferation, differentiation, and involution," *Steroids*, vol. 69, no. 3, pp. 145–159, 2004.
- [13] L. Yart, L. Finot, V. Lollivier, and F. Dessauge, "Oestradiol enhances apoptosis in bovine mammary epithelial cells in vitro," *Journal of Dairy Research*, vol. 80, no. 01, pp. 113–121, 2013.
- [14] J. F. Arnal, C. Fontaine, A. Billon-Gal es et al., "Estrogen receptors and endothelium," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 30, no. 8, pp. 1506–1512, 2010.
- [15] E. E. Connor, M. J. Meyer, R. W. Li, M. E. Van Amburgh, Y. R. Boisclair, and A. V. Capuco, "Regulation of gene expression in the bovine mammary gland by ovarian steroids," *Journal of Dairy Science*, vol. 90, pp. E55–E65, 2007.
- [16] E. R. Prossnitz, J. B. Arterburn, and L. A. Sklar, "GPR30: a G protein-coupled receptor for estrogen," *Molecular and Cellular Endocrinology*, vol. 265–266, pp. 138–142, 2007.
- [17] K. Yakimchuk, M. Jondal, and S. Okret, "Estrogen receptor α and β in the normal immune system and in lymphoid malignancies," *Molecular and Cellular Endocrinology*, vol. 375, no. 1–2, pp. 121–129, 2013.
- [18] S. N. Romano and D. A. Gorelick, "Crosstalk between nuclear and G protein-coupled estrogen receptors," *General and Comparative Endocrinology*, vol. 261, pp. 190–197, 2018.
- [19] S. Farzaneh and A. Zarghi, "Estrogen receptor ligands: a review (2013–2015)," *Scientia Pharmaceutica*, vol. 84, no. 3, pp. 409–427, 2016.
- [20] I. Barone, L. Brusco, and S. A. W. Fuqua, "Estrogen receptor mutations and changes in downstream gene expression and signaling," *Clinical Cancer Research*, vol. 16, no. 10, pp. 2702–2708, 2010.
- [21] A. W. Bell, "Regulation of organic nutrient metabolism during transition from late pregnancy to early lactation," *Journal of Animal Science*, vol. 73, no. 9, pp. 2804–2819, 1995.
- [22] I. Lamote, E. Meyer, A. De Ketelaere, L. Duchateau, and C. Burvenich, "Influence of sex steroids on the viability and CD11b, CD18 and CD47 expression of blood neutrophils from dairy cows in the last month of gestation," *Veterinary Research*, vol. 37, no. 1, pp. 61–74, 2006.
- [23] N. Heldring, A. Pike, S. Andersson et al., "Estrogen receptors: how do they signal and what are their targets," *Physiological Reviews*, vol. 87, no. 3, pp. 905–931, 2007.
- [24] C. Jiang, J. Guo, Z. Wang et al., "Decursin and decursinol angelate inhibit estrogen-stimulated and estrogen-independent growth and survival of breast cancer cells," *Breast Cancer Research*, vol. 9, no. 6, p. R77, 2007.
- [25] A. Tam, S. Wadsworth, D. Dorscheid, S.-F. P. Man, and D. D. Sin, "Estradiol increases mucus synthesis in bronchial epithelial cells," *PLoS One*, vol. 9, no. 6, article e100633, 2014.
- [26] S. Kovats, "Estrogen receptors regulate innate immune cells and signaling pathways," *Cellular Immunology*, vol. 294, no. 2, pp. 63–69, 2015.
- [27] G. M. Barton, "A calculated response: control of inflammation by the innate immune system," *Journal of Clinical Investigation*, vol. 118, no. 2, pp. 413–420, 2008.
- [28] S. L. Klein and K. L. Flanagan, "Sex differences in immune responses," *Nature Reviews Immunology*, vol. 16, no. 10, pp. 626–638, 2016.
- [29] E. Calcagni and I. Elenkov, "Stress system activity, innate and T helper cytokines, and susceptibility to immune-related diseases," *Annals of the New York Academy of Sciences*, vol. 1069, no. 1, pp. 62–76, 2006.
- [30] J. Roved, H. Westerdahl, and D. Hasselquist, "Sex differences in immune responses: hormonal effects, antagonistic selection, and evolutionary consequences," *Hormones and Behavior*, vol. 88, pp. 95–105, 2017.
- [31] Z. Spolarics, G. Pe a, Y. Qin, R. J. Donnelly, and D. H. Livingston, "Inherent X-linked genetic variability and cellular mosaicism unique to females contribute to sex-related differences in the innate immune response," *Frontiers in Immunology*, vol. 8, p. 1455, 2017.

- [32] C. Giefing-Kröll, P. Berger, G. Lepperdinger, and B. Grubeck-Loebenstein, "How sex and age affect immune responses, susceptibility to infections, and response to vaccination," *Aging Cell*, vol. 14, no. 3, pp. 309–321, 2015.
- [33] F. Abdi, H. Mobedi, N. Mosaffa, M. Dolatian, and F. Ramezani Tehrani, "Effects of hormone replacement therapy on immunological factors in the postmenopausal period," *Climacteric*, vol. 19, no. 3, pp. 234–239, 2016.
- [34] P. J. Sansonetti, "To be or not to be a pathogen: that is the mucosally relevant question," *Mucosal Immunology*, vol. 4, no. 1, pp. 8–14, 2011.
- [35] K. F. Ross and M. C. Herzberg, "Autonomous immunity in mucosal epithelial cells: fortifying the barrier against infection," *Microbes and Infection*, vol. 18, no. 6, pp. 387–398, 2016.
- [36] S. B. Rasmussen, L. S. Reinert, and S. R. Paludan, "Innate recognition of intracellular pathogens: detection and activation of the first line of defense," *APMIS*, vol. 117, no. 5–6, pp. 323–337, 2009.
- [37] R. P. Schleimer, A. Kato, R. Kern, D. Kuperman, and P. C. Avila, "Epithelium: at the interface of innate and adaptive immune responses," *The Journal of Allergy and Clinical Immunology*, vol. 120, no. 6, pp. 1279–1284, 2007.
- [38] X. Hu, J. Chen, L. Wang, and L. B. Ivashkiv, "Crosstalk among Jak-STAT, Toll-like receptor, and ITAM-dependent pathways in macrophage activation," *Journal of Leukocyte Biology*, vol. 82, no. 2, pp. 237–243, 2007.
- [39] D. Schams, S. Kohlenberg, W. Amselgruber, B. Berisha, M. W. Pfaffl, and F. Sinowatz, "Expression and localisation of oestrogen and progesterone receptors in the bovine mammary gland during development, function and involution," *Journal of Endocrinology*, vol. 177, no. 2, pp. 305–317, 2003.
- [40] A. Sobolewska, M. Gajewska, J. Zarzynska, B. Gajkowska, and T. Motyl, "IGF-I, EGF, and sex steroids regulate autophagy in bovine mammary epithelial cells via the mTOR pathway," *European Journal of Cell Biology*, vol. 88, no. 2, pp. 117–130, 2009.
- [41] W. Winuthayanon, S. L. Lierz, K. C. Delarosa et al., "Juxtacrine activity of estrogen receptor α in uterine stromal cells is necessary for estrogen-induced epithelial cell proliferation," *Scientific Reports*, vol. 7, no. 1, p. 8377, 2017.
- [42] S. C. Hewitt, L. Li, S. A. Grimm et al., "Research resource: whole-genome estrogen receptor α binding in mouse uterine tissue revealed by ChIP-seq," *Molecular Endocrinology*, vol. 26, no. 5, pp. 887–898, 2012.
- [43] G. R. Cunha, P. S. Cooke, and T. Kurita, "Role of stromal-epithelial interactions in hormonal responses," *Archives of Histology and Cytology*, vol. 67, no. 5, pp. 417–434, 2004.
- [44] R. Kowsar, N. Hambruch, J. Liu, T. Shimizu, C. Pfarrer, and A. Miyamoto, "Regulation of innate immune function in bovine oviduct epithelial cells in culture: the homeostatic role of epithelial cells in balancing Th1/Th2 response," *Journal of Reproduction and Development*, vol. 59, no. 5, pp. 470–478, 2013.
- [45] S. Ayehunie, A. Islam, C. Cannon et al., "Characterization of a hormone-responsive organotypic human vaginal tissue model: morphologic and immunologic effects," *Reproductive Sciences*, vol. 22, no. 8, pp. 980–990, 2015.
- [46] K. Aikawa, T. Sugino, S. Matsumoto, P. Chichester, C. Whitbeck, and R. M. Levin, "The effect of ovariectomy and estradiol on rabbit bladder smooth muscle contraction and morphology," *The Journal of Urology*, vol. 170, no. 2, pp. 634–637, 2003.
- [47] R. Raz, Y. Gennesin, J. Wasser et al., "Recurrent urinary tract infections in postmenopausal women," *Clinical Infectious Diseases*, vol. 30, no. 1, pp. 152–156, 2000.
- [48] D. D. Bannerman, M. J. Paape, J.-W. Lee, X. Zhao, J. C. Hope, and P. Rainard, "*Escherichia coli* and *Staphylococcus aureus* elicit differential innate immune responses following intramammary infection," *Clinical and Vaccine Immunology*, vol. 11, no. 3, pp. 463–472, 2004.
- [49] M. A. Crane-Godreau and C. R. Wira, "Effects of estradiol on lipopolysaccharide and Pam₃Cys stimulation of CCL20/macrophage inflammatory protein 3 alpha and tumor necrosis factor alpha production by uterine epithelial cells in culture," *Infection and Immunity*, vol. 73, no. 7, pp. 4231–4237, 2005.
- [50] T. Hirata, Y. Osuga, K. Hamasaki et al., "Expression of toll-like receptors 2, 3, 4, and 9 genes in the human endometrium during the menstrual cycle," *Journal of Reproductive Immunology*, vol. 74, no. 1–2, pp. 53–60, 2007.
- [51] M. J. Lesmeister, R. L. Jorgenson, S. L. Young, and M. L. Mifseldt, "17Beta-estradiol suppresses TLR3-induced cytokine and chemokine production in endometrial epithelial cells," *Reproductive Biology and Endocrinology*, vol. 3, no. 1, p. 74, 2005.
- [52] D. Xing, S. Oparil, H. Yu et al., "Estrogen modulates NF κ B signaling by enhancing I κ B α levels and blocking p65 binding at the promoters of inflammatory genes via estrogen receptor- β ," *PLoS One*, vol. 7, no. 6, article e36890, 2012.
- [53] B. Villaggio, S. Soldano, and M. Cutolo, "1,25-dihydroxyvitamin D3 downregulates aromatase expression and inflammatory cytokines in human macrophages," *Clinical and Experimental Rheumatology*, vol. 30, no. 6, pp. 934–938, 2012.
- [54] V. Díaz-Murillo, I. Medina-Estrada, J. E. López-Meza, and A. Ochoa-Zarzosa, "Defensin γ -thionin from *Capsicum chinense* has immunomodulatory effects on bovine mammary epithelial cells during *Staphylococcus aureus* internalization," *Peptides*, vol. 78, pp. 109–118, 2016.
- [55] J. H. Han, M. S. Kim, M. Y. Lee et al., "Modulation of human β -defensin-2 expression by 17 β -estradiol and progesterone in vaginal epithelial cells," *Cytokine*, vol. 49, no. 2, pp. 209–214, 2010.
- [56] L. Schaefer, A. Babelova, E. Kiss et al., "The matrix component biglycan is proinflammatory and signals through Toll-like receptors 4 and 2 in macrophages," *The Journal of Clinical Investigation*, vol. 115, no. 8, pp. 2223–2233, 2005.
- [57] G. Soboll, M. A. Crane-Godreau, M. A. Lyimo, and C. R. Wira, "Effect of oestradiol on PAMP-mediated CCL20/MIP-3 α production by mouse uterine epithelial cells in culture," *Immunology*, vol. 118, no. 2, pp. 185–194, 2006.
- [58] G. K. Best, T. O. Abney, J. M. Kling, J. J. Kirkland, and D. F. Scott, "Hormonal influence on experimental infections by a toxic shock strain of *Staphylococcus aureus*," *Infection and Immunity*, vol. 52, no. 1, pp. 331–333, 1986.
- [59] P. Lüthje, H. Brauner, N. L. Ramos et al., "Estrogen supports urothelial defense mechanisms," *Science Translational Medicine*, vol. 5, no. 190, p. 190ra80, 2013.
- [60] I. Medina-Estrada, J. E. López-Meza, and A. Ochoa-Zarzosa, "Anti-inflammatory and antimicrobial effects of estradiol in bovine mammary epithelial cells during *Staphylococcus aureus* internalization," *Mediators of Inflammation*, vol. 2016, Article ID 6120509, 16 pages, 2016.

- [61] E. F. Haney and R. E. W. Hancock, "Peptide design for antimicrobial and immunomodulatory applications," *Biopolymers*, vol. 100, no. 6, pp. 572–583, 2013.
- [62] B. Schitteck, "The antimicrobial skin barrier in patients with atopic dermatitis," in *Current Problems in Dermatology*, vol. 41, pp. 54–67, 2011.
- [63] K. Ellwanger, E. Becker, I. Kienes et al., "The NLR family pyrin domain-containing 11 protein contributes to the regulation of inflammatory signaling," *Journal of Biological Chemistry*, vol. 293, no. 8, pp. 2701–2710, 2018.
- [64] M. Gürsoy, U. K. Gürsoy, A. Liukkonen, T. Kauko, S. Penkkala, and E. Könönen, "Salivary antimicrobial defensins in pregnancy," *Journal of Clinical Periodontology*, vol. 43, no. 10, pp. 807–815, 2016.
- [65] E. Vegeto, S. Belcredito, S. Etteri et al., "Estrogen receptor- α mediates the brain antiinflammatory activity of estradiol," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 16, pp. 9614–9619, 2003.



Hindawi

Submit your manuscripts at
www.hindawi.com

