

Review Article

Regulatory T Cells in Human Ovarian Cancer

Dong-Jun Peng,¹ Rebecca Liu,² and Weiping Zou^{1,3,4}

¹ Department of Surgery, University of Michigan, Ann Arbor, MI 48109, USA

² Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI 48109, USA

³ Graduate Programs in Immunology and Cancer Biology, University of Michigan, Ann Arbor, MI 48109, USA

⁴ University of Michigan School of Medicine, C560B MSRB II, 1150 W. Medical Center Dr., Ann Arbor, MI 48109-0669, USA

Correspondence should be addressed to Weiping Zou, wzou@med.umich.edu

Received 15 September 2011; Accepted 26 November 2011

Academic Editor: Kentaro Nakayama

Copyright © 2012 Dong-Jun Peng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Multiple layers of suppressive components including regulatory T (T_{Reg}) cells, suppressive antigen-presenting cells, and inhibitory cytokines form suppressive networks in the ovarian cancer microenvironment. It has been demonstrated that as a major suppressive element, T_{Reg} cells infiltrate tumor, interact with several types of immune cells, and mediate immune suppression through different molecular and cellular mechanisms. In this paper, we focus on human ovarian cancer and will discuss the nature of T_{Reg} cells including their subsets, trafficking, expansion, and function. We will briefly review the development of manipulation of T_{Reg} cells in preclinical and clinical settings.

1. Introduction

Ovarian cancer is one of the most common and deadliest gynecologic cancers. In 2010, 21880 new cases were diagnosed, and such cancer caused nearly 13850 deaths in the United States alone [1]. Ovarian cancer usually has poor prognosis, and most patients were diagnosed at advanced stages. The five-year survival rate for all stages of ovarian cancer is 46% in 2010 [1]. It has been well documented that patients' clinical outcome and five-year survival rate are positively associated with the number of tumor-infiltrating lymphocytes (TILs) [2], and the ratio of intraepithelial $CD8^+$ TILs to T_{Reg} cells [3], or negatively associated with tumor-infiltrating T_{Reg} cells [4].

T_{Reg} cells are also known as suppressor T cells which consist of a specific subpopulation of cells that functionally suppress the activation of immune system and maintain immune tolerance to self-antigens. T_{Reg} cells contain two major subsets known as natural T_{Reg} cells (nT_{Reg}) and adaptive or induced T_{Reg} cells (iT_{Reg}). nT_{Reg} cells derived from thymus are considered as classic T_{Reg} cells, by contrast, iT_{Reg} cells develop in the periphery in response to self- or tumor antigens by converting naive $CD4^+$ T cells into T_{Reg} cells [5].

Because most tumors express self-antigens, T_{Reg} cells-mediated immunosuppression is believed to be one of the major contributors to immune evasion by tumors and becomes the main obstacle toward successful tumor immunotherapy [6]. In this paper, we will focus on human ovarian cancer and discuss the nature of T_{Reg} cells including their subsets, trafficking, differentiation, and proliferation and the clinical application of manipulation of T_{Reg} cells.

2. Regulatory T-Cell Subsets

In early 1970s, Gershon and Kondo first described the existence of thymus-derived suppressive T cells (later termed as T_{Reg} cell) *in vivo* [7, 8]. After more than a decade, Sakaguchi et al. demonstrated that $CD4^+$ T cells expressing interleukin-2 (IL-2) receptor alpha-chain (CD25) can be defined as the population of T_{Reg} cells with immune-suppressive activities and maintaining immune tolerance to self-antigen [9]. Later in 2003, Hori et al. found that the transcription factor forkhead box P3 (Foxp3) controls the development of T_{Reg} cells and is crucial for maintaining the immune-suppressive function of T_{Reg} cells [10].

Natural T_{Reg} cells differentiate in the thymus and migrate to periphery, which constitute 5–10% of CD4⁺ T cells [11–13]. In addition, there are several subsets of T_{Reg} cells other than CD4⁺CD25⁺FOXP3⁺ T_{Reg} cells. Groux et al. identified another subset of T_{Reg} cells, CD4⁺ T_R 1 cells, that suppress antigen-specific immune responses by producing high levels of IL-10 [14]. In addition to CD4⁺ T_{Reg}, CD8⁺ suppressive T cells have been found playing an important role in the regulation of autoimmune disease [7, 15]. CD8⁺ suppressive T cells now referred to as CD8⁺ T_{Reg} cells are characterized as CD8⁺CD25⁺, CD8⁺CD122⁺, or CD8⁺CD45RC^{low} T_{Reg} cells, which comprise less than 1% of peripheral CD8⁺ T cells [15]. Th3 T_{Reg} cells have similar immune-suppressive function; however, in contrast to natural T_{Reg} cells, Th3 exerts its suppressive capacity independent of cell membrane contact but mainly bases on the action of self-produced cytokine TGFβ [16].

3. Regulatory T-Cell Trafficking

T_{Reg} cells consist of ~10% of peripheral CD4⁺ T cells characterized as CD4⁺CD25⁺FOXP3⁺ T cells, which is important for the control of autoimmune reaction [9, 11]. Dysregulation of T_{Reg} can cause autoimmune diseases [17] and may contribute to tumor-initiated immune evasion [18]. As demonstrated by *in vivo* mouse model, the deletion of T_{Reg} cells results in tumor rejection [19]. However, the suppressive capacity of T_{Reg} cells is also determined by the ratio of T_{Reg} cells to effector T cells [3]. A high CD8⁺/T_{Reg} ratio is associated with favorable prognosis and improved survival [3, 20]. It has been reported that many human cancers are associated with high frequency of T_{Reg} cells in the circulation or in the tumor tissues, including ovarian cancer [4], lung cancer [21], breast cancer [22], liver cancer [23], head and neck cancer [24], and lymphoma [25]. These increased levels of T_{Reg} cells are linked to high death hazard and poor survival, while the depletion of tumor-infiltrated T_{Reg} cells and the blockade of T_{Reg} trafficking to tumors enhance anti-tumor immune response [4, 26].

CCR4 and its binding partners CCL22 and CCL17 are believed to be the most predominant axis in chemokine-mediated selective T_{Reg} trafficking to the tumors. Iellem et al. have profiled chemotactic responses and chemokine receptors expression of human T_{Reg} cells and found that T_{Reg} cells specifically express chemokine receptors CCR4 and CCR8 [27]. Chemokine CCL22, the ligand for CCR4, preferentially attracts activated-antigen-specific T cells to dendritic cells [28, 29]. It has also been shown that human ovarian cancer cells and tumor-associated macrophages produce chemokine CCL22, which mediates T_{Reg} cells trafficking to tumor [4]. Blockade of CCL22 *in vivo* significantly reduces human T_{Reg} cells trafficking to tumors in ovarian carcinoma [4]. This chemokine-mediated T_{Reg} trafficking has been also observed in other types of cancer, such as gastric cancer [30], Hodgkin's lymphoma [31], and breast cancer [32]. Interestingly, in gastric cancer, CCL22 and CCL17 seem both important to recruit T_{Reg} cells to the tumors as demonstrated by *in vivo* study as well as *in vitro* migration assay, and

the levels of CCL22 and CCL17 within tumors are correlated to the increased levels of T_{Reg} cells in early gastric cancer [33].

Besides CCR4 chemokine axis, CCR5/CCL5 axis may also selectively recruit T_{Reg} cells to the tumors. Using human pancreatic adenocarcinoma and murine pancreatic tumor model, it has been found that CCR5 is highly expressed in T_{Reg} cells, while tumor cells produce elevated amount of CCL5, and disruption of CCR5/CCL5 chemokine axis blocks T_{Reg} cells migration and reduces tumor growth [34]. In addition, CCL20 chemokine shows high affinity to CCR6 and can also mediate selective CCR6⁺ T_{Reg} cells trafficking [35].

4. Regulatory T-Cell Differentiation and Proliferation

CD4⁺CD25⁺ T_{Reg} cells are generated in the thymus. Papiernik et al. found that peripheral T_{Reg} migrates from the thymus and appears in the periphery as early as 10th day of life [36]. They also found that CD4⁺CD25⁺ T_{Reg} cells differentiation is totally dependent on IL-2, because IL-2 knockout mice do not develop CD4⁺CD25⁺ T_{Reg} *in vivo* [36]. Further evidences have been provided from the studies on irradiated rat model [37]. In this study, autoimmune diseases were induced in rats by thymectomy and irradiation; however the xenograft transfer of CD4⁺ T cells from normal rats can abrogate the autoimmune responses. These observations suggest that normal thymus-derived T cells have immune suppressive functions and thus prevent autoimmunity [37]. In another model system, adoptive transfer of thymocytes or peripheral T cells depleted of CD4⁺CD25⁺ T_{Reg} cells causes autoimmune diseases in mice, which provides further evidences of thymic origin of T_{Reg} cells and their peripheral existence [38].

However, there is little known about the comprehensive requirements for thymic T_{Reg} development. Although there are several arguments about how and what stromal components are involved in thymic T_{Reg} cell differentiation, thymic stromal cells, including cortical and medullary thymic epithelial cells and dendritic cells (DCs), contribute to T_{Reg} cells differentiation and selection [38]. Jordan et al. used TCR-transgenic mice which express the receptor recognizing specific self-antigen and found that thymocytes bearing a TCR with high affinity to a specific self-antigen undergo selection and become CD4⁺CD25⁺ T_{Reg} cells when interacting with a single self-antigen, but thymocytes bearing TCR with low affinity do not undergo selection [39].

In addition to thymus, T_{Reg} can also be generated in the periphery. For instance, tumor microenvironment favors the induction and differentiation of T_{Reg} cells, and that has been extensively studied for several years [40]. In the tumor microenvironment, DC differentiation and function were suppressed by tumor-associated factors IL-10, VEGF, and TGFβ, resulting in immature/dysfunctional DC [6]. Dysfunctional DC directly contributed to the induction of IL-10-producing T_{Reg} cells *in vivo* in human and *in vitro* [41, 42]. Tumor-associated plasmacytoid DC also induced IL-10⁺ T_{Reg} generation [43, 44]. Tumor can convert DC into TGFβ-producing

immature DC, which selectively promotes T_{Reg} proliferation in TGF β -dependent manner [45].

$CD4^+CD25^+$ T_{Reg} cells can also be converted from peripheral naïve $CD4^+CD25^-$ T cells by the action of TGF β . Tumor microenvironment contains high levels of TGF β which might mediate tumor-associated T_{Reg} cells conversion [46].

5. Targeting Regulatory T Cells

5.1. T_{Reg} Cell Depletion. In the mouse model, depletion of $CD4^+CD25^+$ T_{Reg} cells using anti-CD25 antibody causes tumor regression, which correlated to the reduced number of T_{Reg} cells [18, 47]. Using the recombinant IL-2 diphtheria toxin conjugate DAB(389)IL-2 (also known as denileukin diftitox and ONTAK), Dannull et al. demonstrated that DAB(389)IL-2 was capable of selectively eliminating $CD25^+$ T_{Reg} cells from the PBMCs of cancer patients without inducing toxicity on other cellular subsets, and DAB(389)IL-2-mediated T_{Reg} depletion enhanced anti-tumor immune responses and significantly reduced the number of T_{Reg} cells present in the blood of cancer patients [48]. Daclizumab (also known as Zenapex) and Basiliximab (also called Simulect) are monoclonal antibodies against CD25 [49, 50], and the administration of Daclizumab in patient with metastatic breast cancer enhanced anti-tumor immunity [51].

Cytotoxic T-lymphocyte-associated antigen 4 (CTLA4) is constitutively expressed and restricted to $CD4^+CD25^+$ T_{Reg} cells among all $CD4^+$ cells, and the immune-suppressive function of T_{Reg} is mediated by CTLA4 signaling [52, 53]. CTLA4 binds to inhibitory B7 members on APC and transmits an inhibitory signal to T cells. *In vivo* administration of anti-CTLA4 antibody resulted in tumor rejection including preestablished tumors [54]. Periodic infusions of anti-CTLA4 antibody in previously vaccinated patients with cancer created clinically effective antitumor immune response [55]. Patients with metastatic melanoma showed improved antitumor immunity and tumor regression by blockade of CTLA-4 together with peptide vaccination [56].

Glucocorticoid-induced tumor necrosis factor (TNF) receptor family-related protein (GITR or DTA-1) is predominantly expressed on the surface of T_{Reg} cells. An agonistic anti-GITR antibody administration in mice can abrogate T_{Reg} -mediated immune suppression and enhance effective anti-tumor immunity *in vivo* [57, 58]. In addition, treatment with anti-GITR antibody in B16 mice elicited immune response and rejected tumor [59]. However GITR is not exclusively expressed on T_{Reg} cell; it is also expressed by various $CD4^+$ T cells and others. Therefore, the clinical therapeutic relevance of GITR blockade and its side effects on potential deficits of other effective immune cells remain to be determined.

OX40 (CD134) also belongs to TNF receptor family and expressed on activated T cells. Both naïve and activated T_{Reg} express OX40. Similar to GITR, triggering OX40 by an agonistic antibody against OX40 reduces T_{Reg} -mediated immune suppression and restores effector T-cell function both *in vivo* and *in vitro* [60]. It has been also shown that OX40 is

necessary for T_{Reg} development, homeostasis, and immune-suppressive activity. However, stimulation of OX40 signal in naïve T cells can abrogate T_{Reg} -mediated suppression [61].

Clinical relevance of the depletion of T_{Reg} cells has been further confirmed by the treatment of cyclophosphamide (CY) in the patients bearing tumor. Cyclophosphamide is a nitrogen mustard alkylating agent that mediates DNA crosslinking. Low dose of CY administration improved patients' immune responses by reducing the number of T_{Reg} cells and by decreasing the suppressive activity of T_{Reg} cells [62]. Effects of T_{Reg} depletion on anti-tumor immune responses were further investigated by the study on B16 melanomas mouse model [63]. Other immunosuppressants like cyclosporine A (CSA) and azathioprine might also inhibit T_{Reg} cells generation [64, 65]. For instance, high dose of CSA abrogates T_{Reg} cell generation; by contrast, low dose of CSA can promote T_{Reg} cell development [64]. It is therefore important to determine whether lowdose of those agents can improve antitumor immunity in patients.

5.2. Targeting T_{Reg} Trafficking. Our group has demonstrated that human ovarian cancer cells and tumor-associated macrophage (TAM) produced chemokine CCL22, the ligand for CCR4 which functionally expressed on tumor T_{Reg} cells, mediating T_{Reg} cells trafficking to the tumor and ascites, and the blockade of CCL22 abrogated T_{Reg} cells migration [4]. It has been demonstrated that chemokine receptor CCR4 is selectively expressed by T_{Reg} cells, and the CCR4 and CCR4-associate chemokines axis is one of the most described tumor T_{Reg} recruitment axes [66]. The administration of anti-CCR4 antibody effectively depletes CCR4 $^+$ T cells and inhibits T_{Reg} cells migration in Hodgkin lymphoma [31]. Furthermore, the significant correlation between CCL17 or CCL22 chemokines and the number of tumor-infiltrating T_{Reg} cells was found in patients with neoplastic meningitis and gastric cancer [30, 33]. CCL5 and CCL20 chemokines are also involved in T_{Reg} trafficking, and that blockade of those chemokines reduces T_{Reg} cells trafficking and inhibits tumor growth [34, 35]. We have shown that CXCL12/CXCR4 axis mediated T_{Reg} trafficking to bone marrow [67]. Recently, a study has demonstrated that blockade of CXCR4 by a selective antagonist resulted in the significant reduction of intratumoral T_{Reg} cells, which was associated with greatly increased antitumor immunity and an improved survival in an immunocompetent mouse model of ovarian cancer [68].

5.3. Targeting TGF β Signaling Pathway. TGF β is implicated in T_{Reg} differentiation, conversion, and function. It is thought that blockade of TGF β signaling pathway may alter T_{Reg} phenotype and function and in turn enhances antitumor immunity [6]. In addition to T_{Reg} cells, ovarian carcinoma cells can also produce TGF β [69]. Notably, TGF β is not only important for T_{Reg} cell functional integrity, but also inhibits the proliferation and functional differentiation of T lymphocytes, NK cells, and macrophages [46, 70]. This may induce T-cell unresponsiveness to TCR stimulation, failure to produce Th1 cytokines, and production of additional TGF β [46]. TGF β signaling may also be crucial for tumor cell

transformation. Therefore, targeting TGF β signaling may be therapeutically meaningful. TGF β inhibitor AP 12009 was tested in a Phase I/II clinical trial for advanced pancreatic cancer and other malignancies [71]. LY2109761, an inhibitor of TGF β I/II receptors, can suppress pancreatic cancer metastases [72]. In a preclinical model, we have shown that anti-TGF β can reduce T_{Reg} cells in tumors and tumor-draining lymph nodes. This effect is enhanced by B7-H1 blockade [73]. Nonetheless, it is clear that blocking TGF β signaling may affect T_{Reg} compartment. However, as TGF β is implicated in multiple layers of biological activities, the ultimate clinical therapeutic efficiency and side effects of TGF β signaling blockade remain to be investigated.

5.4. Targeting Inhibitory B7 Family Members. The expression, regulation, functional, and clinical relevance of inhibitory B7 family members have been reviewed elsewhere [74]. Human ovarian cancer and cancer-associated myeloid antigen-presenting cells express high levels of B7-H1 (PD-L1), which are negatively associated with patient survival [74, 75]. Patients with high expression of B7-H1 had a significantly poor prognosis compared to the patients with low expression of B7H1 [76]. B7-H1 expression was also found inversely correlated to the intraepithelial CD8⁺ T lymphocyte count, indicating that B7-H1 on tumor cells may suppress antitumor CD8⁺ T cells [76]. The receptor, programmed death 1 (PD-1), is expressed on activated T-cell subsets, antigen-specific CD8⁺ T cells [77], and T_{Reg} [78]. Interestingly, B7-H1/PD-1 has been reported to be involved in the development of induced T_{Reg} cells [79]. Therefore, targeting B7-H1/PD-1 signaling pathway may reduce T_{Reg} development and function. As anti-PD-1 is in clinical application to treat patients with melanoma, renal cell carcinoma, and other cancers, further mechanistic studies on these patients will determine if the effects of anti-PD-1 on T_{Reg} cells are mechanistically and clinically relevant.

In addition to B7-H1, human ovarian cancer and cancer-associated myeloid antigen-presenting cells also express high levels of B7-H4 (B7x, B7s1), which are negatively associated with patient survival [74, 80, 81]. Interestingly, T_{Reg} cells can induce IL-10 expression by APCs and indirectly stimulate B7-H4 expression on APCs and convey suppressive activity to APCs [74, 80, 81]. Thus, it is tempting to speculate that blocking B7-H4 signaling pathway may disable the suppressive effects of T_{Reg} cells on APCs. Notably, as the receptor for B7-H4 has not been identified, B7-H4 signaling is much less understood in both mouse and human system. Nonetheless, studies on ovarian cancer patients and preclinical cancer models suggest that interruption of B7-H4 signaling may lead to improved antitumor T-cell response and decreased T_{Reg} suppressive function.

6. Conclusions

T_{Reg} cells infiltrate tumor including ovarian cancer. Their phenotype, trafficking mechanism, suppressive activity, and clinical relevance have been defined in human cancer. However, recent evidence indicates that T_{Reg} cells may not be

stable and are subject to environmental regulation. In this regard, it remains poorly understood how T_{Reg} cells evolve in human tumor microenvironment. Although their action mode of mechanisms has been investigated in many different physiological and pathological scenarios, the key suppressive mechanisms may be differed in different tumors or/and in different stages. Therefore, further patient-oriented studies are essential for dissecting T_{Reg} cell biology. Nonetheless, targeting T_{Reg} cells or/and reprogramming T_{Reg} cells is an important strategy to treat patients with cancer. It is suggested that combinatorial therapy by incorporating T_{Reg} manipulation may be ideal direction to develop novel therapeutic regimen to efficiently treat patients with cancer.

References

- [1] A. Jemal, R. Siegel, J. Xu, and E. Ward, "Cancer statistics, 2010," *CA Cancer Journal for Clinicians*, vol. 60, no. 5, pp. 277–300, 2010.
- [2] L. Zhang, J. R. Conejo-Garcia, D. Katsaros et al., "Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer," *New England Journal of Medicine*, vol. 348, no. 3, pp. 203–213, 2003.
- [3] E. Sato, S. H. Olson, J. Ahn et al., "Intraepithelial CD8⁺ tumor-infiltrating lymphocytes and a high CD8⁺/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 51, pp. 18538–18543, 2005.
- [4] T. J. Curiel, G. Coukos, L. Zou et al., "Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival," *Nature Medicine*, vol. 10, no. 9, pp. 942–949, 2004.
- [5] M. A. Curotto de Lafaille and J. J. Lafaille, "Natural and adaptive Foxp3⁺ regulatory T cells: more of the same or a division of labor?" *Immunity*, vol. 30, no. 5, pp. 626–635, 2009.
- [6] W. Zou, "Regulatory T cells, tumour immunity and immunotherapy," *Nature Reviews Immunology*, vol. 6, no. 4, pp. 295–307, 2006.
- [7] R. K. Gershon and K. Kondo, "Cell interactions in the induction of tolerance: the role of thymic lymphocytes," *Immunology*, vol. 18, no. 5, pp. 723–737, 1970.
- [8] R. K. Gershon and K. Kondo, "Infectious immunological tolerance," *Immunology*, vol. 21, no. 6, pp. 903–914, 1971.
- [9] S. Sakaguchi, N. Sakaguchi, M. Asano, M. Itoh, and M. Toda, "Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor α -chains (CD25): breakdown of a single mechanism of self-tolerance causes various autoimmune diseases," *Journal of Immunology*, vol. 155, no. 3, pp. 1151–1164, 1995.
- [10] S. Hori, T. Nomura, and S. Sakaguchi, "Control of regulatory T cell development by the transcription factor Foxp3," *Science*, vol. 299, no. 5609, pp. 1057–1061, 2003.
- [11] E. M. Shevach, "CD4⁺CD25⁺ suppressor T cells: more questions than answers," *Nature Reviews Immunology*, vol. 2, no. 6, pp. 389–400, 2002.
- [12] S. Sakaguchi, "Naturally arising Foxp3-expressing CD25⁺ CD4⁺ regulatory T cells in immunological tolerance to self and non-self," *Nature Immunology*, vol. 6, no. 4, pp. 345–352, 2005.

- [13] L.A. Stephens, C. Mottet, and D. Mason, "Human CD4⁺ CD25⁺ thymocytes and peripheral T cells have immune suppressive activity in vitro," *Eur J Immunol*, vol. 31, no. 4, pp. 1247–1254, 2001.
- [14] H. Groux, A. O'Garra, M. Bigler et al., "A CD4⁺ T-cell subset inhibits antigen-specific T-cell responses and prevents colitis," *Nature*, vol. 389, no. 6652, pp. 737–742, 1997.
- [15] T. R. F. Smith and V. Kumar, "Revival of CD8⁺ Treg-mediated suppression," *Trends in Immunology*, vol. 29, no. 7, pp. 337–342, 2008.
- [16] H. Jonuleit and E. Schmitt, "The regulator T cell family: distinct subsets and their interrelations," *Journal of Immunology*, vol. 171, no. 12, pp. 6323–6327, 2003.
- [17] E. Jones, M. Dahm-Vicker, A. K. Simon et al., "Depletion of CD25⁺ regulatory cells results in suppression of melanoma growth and induction of autoreactivity in mice," *Cancer Immunity*, vol. 2, p. 1, 2002.
- [18] S. Onizuka, I. Tawara, J. Shimizu, S. Sakaguchi, T. Fujita, and E. Nakayama, "Tumor rejection by in vivo administration of anti-CD25 (interleukin-2 receptor α) monoclonal antibody," *Cancer Research*, vol. 59, no. 13, pp. 3128–3133, 1999.
- [19] D. Golgher, E. Jones, F. Powrie, T. Elliott, and A. Gallimore, "Depletion of CD25⁺ regulatory cells uncovers immune responses to shared murine tumor rejection antigens," *European Journal of Immunology*, vol. 32, no. 11, pp. 3267–3275, 2002.
- [20] F. Ichihara, K. Kono, A. Takahashi, H. Kawaida, H. Sugai, and H. Fujii, "Increased populations of regulatory T cells in peripheral blood and tumor-infiltrating lymphocytes in patients with gastric and esophageal cancers," *Clinical Cancer Research*, vol. 9, no. 12, pp. 4404–4408, 2003.
- [21] E. Y. Woo, C. S. Chu, T. J. Goletz et al., "Regulatory CD4⁺ CD25⁺ T cells in tumors from patients with early-stage non-small cell lung cancer and late-stage ovarian cancer," *Cancer Research*, vol. 61, no. 12, pp. 4766–4772, 2001.
- [22] U. K. Liyanage, T. T. Moore, H. G. Joo et al., "Prevalence of regulatory T cells is increased in peripheral blood and tumor microenvironment of patients with pancreas or breast adenocarcinoma," *Journal of Immunology*, vol. 169, no. 5, pp. 2756–2761, 2002.
- [23] L. Ormandy, T. Hillemann, H. Wedemeyer, M. P. Manns, T. F. Greten, and F. Korangy, "Increased populations of regulatory T cells in peripheral blood of patients with hepatocellular carcinoma," *Cancer Research*, vol. 65, no. 6, pp. 2457–2464, 2005.
- [24] A. K. Schott, R. Pries, and B. Wollenberg, "Permanent up-regulation of regulatory T-lymphocytes in patients with head and neck cancer," *International Journal of Molecular Medicine*, vol. 26, no. 1, pp. 67–75, 2010.
- [25] T. Álvaro, M. Lejeune, M. T. Salvadó et al., "Outcome in Hodgkin's lymphoma can be predicted from the presence of accompanying cytotoxic and regulatory T cells," *Clinical Cancer Research*, vol. 11, no. 4, pp. 1467–1473, 2005.
- [26] J. Steitz, J. Brück, J. Lenz, J. Knop, and T. Tüting, "Depletion of CD25⁺ CD4⁺ T cells and treatment with tyrosinase-related protein 2-transduced dendritic cells enhance the interferon α -induced, CD8⁺ T-cell-dependent immune defense of B16 melanoma," *Cancer Research*, vol. 61, no. 24, pp. 8643–8646, 2001.
- [27] A. Illem, M. Mariani, R. Lang et al., "Unique chemotactic response profile and specific expression of chemokine receptors CCR4 and CCR8 by CD4⁺ CD25⁺ regulatory T cells," *Journal of Experimental Medicine*, vol. 194, no. 6, pp. 847–853, 2001.
- [28] M. T. Wu, H. Fang, and S. T. Hwang, "Cutting edge: CCR4 mediates antigen-primed T cell binding to activated dendritic cells," *Journal of Immunology*, vol. 167, no. 9, pp. 4791–4795, 2001.
- [29] H. L. Tang and J. G. Cyster, "Chemokine up-regulation and activated T cell attraction by maturing dendritic cells," *Science*, vol. 284, no. 5415, pp. 819–822, 1999.
- [30] J. Haas, L. Schopp, B. Storch-Hagenlocher et al., "Specific recruitment of regulatory T cells into the CSF in lymphomatous and carcinomatous meningitis," *Blood*, vol. 111, no. 2, pp. 761–766, 2008.
- [31] T. Ishida, T. Ishii, A. Inagaki et al., "Specific recruitment of CC chemokine receptor 4-positive regulatory T cells in Hodgkin lymphoma fosters immune privilege," *Cancer Research*, vol. 66, no. 11, pp. 5716–5722, 2006.
- [32] C. Ménétrier-Caux, M. Gobert, and C. Caux, "Differences in tumor regulatory T-cell localization and activation status impact patient outcome," *Cancer Research*, vol. 69, no. 20, pp. 7895–7898, 2009.
- [33] Y. Mizukami, K. Kono, Y. Kawaguchi et al., "CCL17 and CCL22 chemokines within tumor microenvironment are related to accumulation of Foxp3⁺ regulatory T cells in gastric cancer," *International Journal of Cancer*, vol. 122, no. 10, pp. 2286–2293, 2008.
- [34] M. C. B. Tan, P. S. Goedegebuure, B. A. Belt et al., "Disruption of CCR5-dependent homing of regulatory T cells inhibits tumor growth in a murine model of pancreatic cancer," *Journal of Immunology*, vol. 182, no. 3, pp. 1746–1755, 2009.
- [35] M. Kleinewietfeld, F. Puentes, G. Borsellino, L. Battistini, O. Rötzschke, and K. Falk, "CCR6 expression defines regulatory effector/memory-like cells within the CD25⁺ CD4⁺ T-cell subset," *Blood*, vol. 105, no. 7, pp. 2877–2886, 2005.
- [36] M. Papiernik, M. L. De Moraes, C. Pontoux, F. Vasseur, and C. Pénit, "Regulatory CD4 T cells: expression of IL-2R α chain, resistance to clonal deletion and IL-2 dependency," *International Immunology*, vol. 10, no. 4, pp. 371–378, 1998.
- [37] W. J. Penhale, W. J. Irvine, J. R. Inglis, and A. Farmer, "Thyroiditis in T cell depleted rats: suppression of the autoallergic response by reconstitution with normal lymphoid cells," *Clinical and Experimental Immunology*, vol. 25, no. 1, pp. 6–16, 1976.
- [38] S. Sakaguchi, M. Miyara, C. M. Costantino, and D. A. Hafler, "FOXP3⁺ regulatory T cells in the human immune system," *Nature Reviews Immunology*, vol. 10, no. 7, pp. 490–500, 2010.
- [39] M. S. Jordan, A. Boesteanu, A. J. Reed et al., "Thymic selection of CD4⁺ CD25⁺ regulatory T cells induced by an agonist self-peptide," *Nature Immunology*, vol. 2, no. 4, pp. 301–306, 2001.
- [40] W. Zou, "Immunosuppressive networks in the tumour environment and their therapeutic relevance," *Nature Reviews Cancer*, vol. 5, no. 4, pp. 263–274, 2005.
- [41] H. Jonuleit, E. Schmitt, G. Schuler, J. Knop, and A. H. Enk, "Induction of interleukin 10-producing, nonproliferating CD4⁺ T cells with regulatory properties by repetitive stimulation with allogeneic immature human dendritic cells," *Journal of Experimental Medicine*, vol. 192, no. 9, pp. 1213–1222, 2000.
- [42] M. V. Dhodapkar, R. M. Steinman, J. Krasovskiy, C. Munz, and N. Bhardwaj, "Antigen-specific inhibition of effector T cell function in humans after injection of immature dendritic cells," *Journal of Experimental Medicine*, vol. 193, no. 2, pp. 233–238, 2001.
- [43] W. Zou, V. Machelon, A. Coulomb-L'Hermin et al., "Stromal-derived factor-1 in human tumors recruits and alters the function of plasmacytoid precursor dendritic cells," *Nature Medicine*, vol. 7, no. 12, pp. 1339–1346, 2001.

- [44] S. Wei, I. Kryczek, L. Zou et al., "Plasmacytoid dendritic cells induce CD8⁺ regulatory T cells in human ovarian carcinoma," *Cancer Research*, vol. 65, no. 12, pp. 5020–5026, 2005.
- [45] F. Ghiringhelli, P. E. Puig, S. Roux et al., "Tumor cells convert immature myeloid dendritic cells into TGF- β -secreting cells inducing CD4⁺CD25⁺ regulatory T cell proliferation," *Journal of Experimental Medicine*, vol. 202, no. 7, pp. 919–929, 2005.
- [46] W. Chen, W. Jin, N. Hardegen et al., "Conversion of peripheral CD4⁺CD25⁻ naive T cells to CD4⁺CD25⁺ regulatory T cells by TGF- β induction of transcription factor Foxp3," *Journal of Experimental Medicine*, vol. 198, no. 12, pp. 1875–1886, 2003.
- [47] J. Shimizu, S. Yamazaki, and S. Sakaguchi, "Induction of tumor immunity by removing CD25⁺CD4⁺ T cells: a common basis between tumor immunity and autoimmunity," *Journal of Immunology*, vol. 163, no. 10, pp. 5211–5218, 1999.
- [48] J. Dannull, Z. Su, D. Rizzieri et al., "Enhancement of vaccine-mediated antitumor immunity in cancer patients after depletion of regulatory T cells," *Journal of Clinical Investigation*, vol. 115, no. 12, pp. 3623–3633, 2005.
- [49] F. Vincenti, R. Kirkman, S. Light et al., "Interleukin-2-receptor blockade with daclizumab to prevent acute rejection in renal transplantation," *New England Journal of Medicine*, vol. 338, no. 3, pp. 161–165, 1998.
- [50] B. Nashan, R. Moore, P. Amlot, A.-G. Schmidt, K. Abeywickrama, and J.-P. Soulillou, "Randomised trial of basiliximab versus placebo for control of acute cellular rejection in renal allograft recipients," *The Lancet*, vol. 350, no. 9086, pp. 1193–1198, 1997.
- [51] A. J. Rech and R. H. Vonderheide, "Clinical use of anti-CD25 antibody daclizumab to enhance immune responses to tumor antigen vaccination by targeting regulatory T cells," *Annals of the New York Academy of Sciences*, vol. 1174, pp. 99–106, 2009.
- [52] T. Takahashi, T. Tagami, S. Yamazaki et al., "Immunologic self-tolerance maintained by CD25⁺CD4⁺ regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4," *Journal of Experimental Medicine*, vol. 192, no. 2, pp. 303–310, 2000.
- [53] S. Read, V. Malmström, and F. Powrie, "Cytotoxic T lymphocyte-associated antigen 4 plays an essential role in the function of CD25⁺CD4⁺ regulatory cells that control intestinal inflammation," *Journal of Experimental Medicine*, vol. 192, no. 2, pp. 295–302, 2000.
- [54] D. R. Leach, M. F. Krummel, and J. P. Allison, "Enhancement of antitumor immunity by CTLA-4 blockade," *Science*, vol. 271, no. 5256, pp. 1734–1736, 1996.
- [55] F. S. Hodi, M. Butler, D. A. Oble et al., "Immunologic and clinical effects of antibody blockade of cytotoxic T lymphocyte-associated antigen 4 in previously vaccinated cancer patients," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 8, pp. 3005–3010, 2008.
- [56] G. Q. Phan, J. C. Yang, R. M. Sherry et al., "Cancer regression and autoimmunity induced by cytotoxic T lymphocyte-associated antigen 4 blockade in patients with metastatic melanoma," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 14, pp. 8372–8377, 2003.
- [57] J. Shimizu, S. Yamazaki, T. Takahashi, Y. Ishida, and S. Sakaguchi, "Stimulation of CD25⁺CD4⁺ regulatory T cells through GITR breaks immunological self-tolerance," *Nature Immunology*, vol. 3, no. 2, pp. 135–142, 2002.
- [58] K. Ko, S. Yamazaki, K. Nakamura et al., "Treatment of advanced tumors with agonistic anti-GITR mAb and its effects on tumor-infiltrating Foxp3⁺CD25⁺CD4⁺ regulatory T cells," *Journal of Experimental Medicine*, vol. 202, no. 7, pp. 885–891, 2005.
- [59] M. J. Turk, J. A. Guevara-Patiño, G. A. Rizzuto, M. E. Engelhorn, and A. N. Houghton, "Concomitant tumor immunity to a poorly immunogenic melanoma is prevented by regulatory T cells," *Journal of Experimental Medicine*, vol. 200, no. 6, pp. 771–782, 2004.
- [60] B. Valzasina, C. Guiducci, H. Dislich, N. Killeen, A. D. Weinberg, and M. P. Colombo, "Triggering of OX40 (CD134) on CD4⁺CD25⁺ T cells blocks their inhibitory activity: a novel regulatory role for OX40 and its comparison with GITR," *Blood*, vol. 105, no. 7, pp. 2845–2851, 2005.
- [61] I. Takeda, S. Ine, N. Killeen et al., "Distinct roles for the OX40-OX40 ligand interaction in regulatory and nonregulatory T cells," *Journal of Immunology*, vol. 172, no. 6, pp. 3580–3589, 2004.
- [62] M. E. C. Lutsiak, R. T. Semnani, R. De Pascalis, S. V. S. Kashmiri, J. Schlom, and H. Sabzevari, "Inhibition of CD4⁺25⁺ T regulatory cell function implicated in enhanced immune response by low-dose cyclophosphamide," *Blood*, vol. 105, no. 7, pp. 2862–2868, 2005.
- [63] M. J. Turk, J. A. Guevara-Patiño, G. A. Rizzuto, M. E. Engelhorn, and A. N. Houghton, "Concomitant tumor immunity to a poorly immunogenic melanoma is prevented by regulatory T cells," *Journal of Experimental Medicine*, vol. 200, no. 6, pp. 771–782, 2004.
- [64] M. Kawai, H. Kitade, C. Mathieu, M. Waer, and J. Pirenne, "Inhibitory and stimulatory effects of cyclosporine A on the development of regulatory T cells in vivo," *Transplantation*, vol. 79, no. 9, pp. 1073–1077, 2005.
- [65] S. Shibutani, F. Inoue, O. Aramaki et al., "Effects of immunosuppressants on induction of regulatory cells after intratracheal delivery of alloantigen," *Transplantation*, vol. 79, no. 8, pp. 904–913, 2005.
- [66] A. W. Mailloux and M. R. I. Young, "Regulatory T-cell trafficking: from thymic development to tumor-induced immune suppression," *Critical Reviews in Immunology*, vol. 30, no. 5, pp. 435–447, 2010.
- [67] L. Zou, B. Barnett, H. Safah et al., "Bone marrow is a reservoir for CD4⁺CD25⁺ regulatory T cells that traffic through CXCL12/CXCR4 signals," *Cancer Research*, vol. 64, no. 22, pp. 8451–8455, 2004.
- [68] E. Righi, S. Kashiwagi, J. Yuan et al., "CXCL12/CXCR4 blockade induces multimodal antitumor effects that prolong survival in an immunocompetent mouse model of ovarian cancer," *Cancer Research*, vol. 71, no. 16, pp. 5522–5534, 2011.
- [69] J. M. Bartlett, S. P. Langdon, W. N. Scott et al., "Transforming growth factor- β isoform expression in human ovarian tumours," *European Journal of Cancer*, vol. 33, no. 14, pp. 2397–2403, 1997.
- [70] B. B. Cazac and J. Roes, "TGF- β receptor controls B cell responsiveness and induction of IgA in vivo," *Immunity*, vol. 13, no. 4, pp. 443–451, 2000.
- [71] K. H. Schlingensiefen, B. Fischer-Blass, S. Schmaus, and S. Ludwig, "Antisense therapeutics for tumor treatment: the TGF- β 2 inhibitor AP 12009 in clinical development against malignant tumors," *Recent Results in Cancer Research*, vol. 177, pp. 137–150, 2008.
- [72] D. Melisi, S. Ishiyama, G. M. Scabas et al., "LY2109761, a novel transforming growth factor β receptor type I and type II dual inhibitor, as a therapeutic approach to suppressing pancreatic cancer metastasis," *Molecular Cancer Therapeutics*, vol. 7, no. 4, pp. 829–840, 2008.

- [73] S. Wei, A. B. Shreiner, N. Takeshita, L. Chen, W. Zou, and A. E. Chang, "Tumor-induced immune suppression of in vivo effector T-cell priming is mediated by the B7-H1/PD-1 axis and transforming growth factor β ," *Cancer Research*, vol. 68, no. 13, pp. 5432–5438, 2008.
- [74] W. Zou and L. Chen, "Inhibitory B7-family molecules in the tumour microenvironment," *Nature Reviews Immunology*, vol. 8, no. 6, pp. 467–477, 2008.
- [75] T. J. Curiel, S. Wei, H. Dong et al., "Blockade of B7-H1 improves myeloid dendritic cell-mediated antitumor immunity," *Nature Medicine*, vol. 9, no. 5, pp. 562–567, 2003.
- [76] J. Hamanishi, M. Mandai, M. Iwasaki et al., "Programmed cell death 1 ligand 1 and tumor-infiltrating CD8⁺ T lymphocytes are prognostic factors of human ovarian cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 9, pp. 3360–3365, 2007.
- [77] W. Ke, I. Kryczek, L. Chen, W. Zou, and T. H. Welling, "Kupffer cell suppression of CD8⁺ T cells in human hepatocellular carcinoma is mediated by B7-H1/programmed death-1 interactions," *Cancer Research*, vol. 69, no. 20, pp. 8067–8075, 2009.
- [78] W. Wang, R. Lau, D. Yu, W. Zhu, A. Korman, and J. Weber, "PD1 blockade reverses the suppression of melanoma antigen-specific CTL by CD4⁺ CD25^{Hi} regulatory T cells," *International Immunology*, vol. 21, no. 9, pp. 1065–1077, 2009.
- [79] L. M. Francisco, V. H. Salinas, K. E. Brown et al., "PD-L1 regulates the development, maintenance, and function of induced regulatory T cells," *Journal of Experimental Medicine*, vol. 206, no. 13, pp. 3015–3029, 2009.
- [80] I. Kryczek, S. Wei, G. Zhu et al., "Relationship between B7-H4, regulatory T cells, and patient outcome in human ovarian carcinoma," *Cancer Research*, vol. 67, no. 18, pp. 8900–8905, 2007.
- [81] I. Kryczek, L. Zou, P. Rodriguez et al., "B7-H4 expression identifies a novel suppressive macrophage population in human ovarian carcinoma," *Journal of Experimental Medicine*, vol. 203, no. 4, pp. 871–881, 2006.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

