Clinical Study

Endoscopic Revision (StomaphyX) versus Formal Surgical Revision (Gastric Bypass) for Failed Vertical Band Gastroplasty

Johan Bolton,¹ Richdeep S. Gill,¹ Akram Al-Jahdali,² Simon Byrns,³ Xinzhe Shi,⁴ Daniel W. Birch,⁴ and Shahzeer Karmali⁴

¹ Department of Surgery, University of Alberta, Edmonton, AB, Canada T6G 2B7

² Department of Surgery, University of British Columbia, Vancouver, British Columbia, Canada

³ Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB, Canada T6G 2B7

⁴ Center for the Advancement of Minimally Invasive Surgery (CAMIS), Community Services Center, Royal Alexandra Hospital, Room 405, 10240 Kingsway, Edmonton, AB, Canada T5H 3V9

Correspondence should be addressed to Shahzeer Karmali; shahzeer.karmali@albertahealthservices.ca

Received 30 September 2012; Revised 4 December 2012; Accepted 18 December 2012

Academic Editor: Francesco Saverio Papadia

Copyright © 2013 Johan Bolton et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Weight regain secondary to VBG pouch dilation is a typical referral for Bariatric surgeons. In this study we compare an endoluminal pouch reduction (Stomaphyx) to RYGB for revision. **Methods.** A retrospective review was completed for patients with a previous VBG presenting with weight regain between 2003–2010. **Results.** Thirty patients were identified for study 23 RYGB, 14 StomaphyX. Significant post procedure BMI loss was seen in each cohort (RYGB, $47.7 \pm 7 \text{ kg/m}^2$ to $35 \pm 7 \text{ kg/m}^2$; StomaphyX $43 \pm 10 \text{ kg/m}^2$ to $40 \pm 9 \text{ kg/m}^2$, $P = 0.0007$). Whereas nausea and headache were the only complications observed in StomaphyX patients, the RYGB group had a 43.5% complication rate and 1 mortality. Complications following RYGB include: incisional hernia (13%), anastomotic leak (8.7%), respiratory failure (8.7%), fistula (8.7%), and perforation (4.35%). The median length of stay following RYGB was 6 days compared to 1.5 ± 0.5 days following StomaphyX. **Conclusion.** This study suggests that while RYGB revision may achieve greater weight loss, the complication rates and severity is discouraging. StomaphyX may be a safe alternative. Further technical modifications of the device and longer follow-up may clarify the role of this approach.

1. Introduction

According to the World Health Organization, there are over 500 million obese individuals worldwide [1]. Bariatric surgery has been shown to be an effective treatment strategy to produce marked weight loss in patients with moderate to severe obesity ($\text{BMI} > 35 \text{ kg/m}^2$) [2]. Bariatric surgery includes both primarily restrictive and malabsorptive procedures. Specifically, vertical banded gastroplasty (VBG), a primarily restrictive bariatric surgical procedure, was first described by Mason in 1982 [3]. Despite initial optimism with VBG, long-term results have been disappointing. According to Balsiger et al., only 20% of patients maintained 50% excess weight loss (EWL) at 10-year followup [4]. Weight regain following VBG may be related to staple-line dehiscence and stomal pouch dilation [5]. In addition, reoperation and revision of VBG are needed in 20% to 30% of patients [4, 6].

Roux-en-Y gastric bypass (RYGB) remains the most common revisional procedure following failed VBG. RYGB had been shown to produce marked weight loss as a revisional procedure for previously failed restrictive bariatric surgical procedures [7–9]. However, some studies have suggested increased complication rates with revisional bariatric operations as high as 12% to 41% ([10] Schwartz RW, *Obes Surg* 2002; [11]). Recently, an endoscopic treatment strategy involving the StomaphyX device (Endogastric Solutions Inc., Redmond, WA, USA) has been used to revise gastric pouch dilatations. Endoscopic revision via the StomaphyX has been reported in patients following VBG and RYGB successfully [12, 13]. Our objective was to retrospectively compare weight loss and complication rates, following revision of failed VBG with either the StomaphyX device or formal surgical revision to RYGB at our institution.

2. Materials and Methods

A retrospective review was completed for all patients with a previous VBG presenting to a comprehensive adult weight management clinic (Weight Wise) with weight regain between 2003 and 2010. The multidisciplinary team at Weight Wise including physicians, nurses, physiotherapy, and dietitians assessed these patients. Patients with previous VBG that presented with persistent weight gain despite conservative measures and meeting the Canadian Guidelines for Surgical Intervention were considered for revision by StomaphyX or open conversion to RYGB [14]. VBG revision endoscopically via the Stomaphyx device was performed as a part of a clinical trial, with the results previously reported by Manouchehri et al. [12].

2.1. StomaphyX Revision of VBG. StomaphyX is an endoluminal device that has recently been developed as an alternative to revisional surgery. A gastroscope is inserted through the internal lumen of the device and full-thickness gastric tissue is suctioned into the device allowing for the application of a polypropylene fastener which if repeated circumferentially creates a circular pleat thus downsizing the gastric pouch. The StomaphyX device has since been used and studied at our institution as a minimally invasive revisional option for patients with failed VBG [12].

2.2. Preoperative Characteristics. Patient demographics were collected retrospectively, including age, sex, and mean preoperative weight and body mass index (BMI) following the initial VBG.

2.3. Outcomes. The primary outcomes of interest were complication rates after revisional procedure weight loss. This included perioperative and postoperative complications such as anastomotic leakage, intra-abdominal abscess, dehiscence, respiratory or cardiac complications, and incisional hernia. Mortality was also recorded. Weight loss following bariatric revisional surgery was recorded. Also, the operative time to complete either VBG revision with StomaphyX or formal conversion to RYGB was recorded, along with length of hospital stay (LOS).

2.4. Statistical Analysis. Descriptive statistics were reported as mean $\pm$ standard deviation. Comparison of pre- and postbariatric procedure outcomes was performed using a paired Student *t*-test. Statistical significance was defined as $P < 0.05$.

3. Results

There were a total of thirty-seven patients from 2003 to 2010 that were included in the analysis. The preoperative characteristics are detailed in Table 1. Twenty-three patients were identified that had previously failed VBG and conversion to RYGB. Fourteen patients were identified with previously failed VBG and endoscopic revision via StomaphyX. The preoperative BMI in the RYGB group was $47.7 \pm 7 \text{ kg/m}^2$

TABLE 1: Patient characteristics of failed VBG patients following StomaphyX or RYGB ($N = 37$).

Characteristic	StomaphyX ($N = 14$)	RYGB ($N = 23$)
Age (years)	46.4 (6.7)	49.0 (8.0)
Sex (% female)	93	78
Preoperative weight (kg)	119.5 (25.9)	140.3 (32.1)
Preoperative BMI (kg/m^2)	43.4 (9.7)	49.8 (8.4)

RYGB: Roux-en-Y gastric bypass; Standard Deviation indicated in brackets.

FIGURE 1: Change in body mass index (BMI) following formal conversion of VBG to RYGB. * $P < 0.05$ versus preoperative body mass index.

compared to $43 \pm 10 \text{ kg/m}^2$ in the StomaphyX group. As seen in Figure 1, patients with previously failed VBG had a significant decrease in BMI following RYGB ($47.7 \pm 7 \text{ kg/m}^2$ to $35 \pm 7 \text{ kg/m}^2$ at 24-month followup, $P = 0.0007$). In patients following StomaphyX endoscopic revision, there was also a significant decrease in BMI ($43 \pm 10 \text{ kg/m}^2$ to $40 \pm 9 \text{ kg/m}^2$ at 6-month followup, $P = 0.0007$). Figure 2 demonstrates the changes in BMI in both groups at 6-months followup.

The only complications observed in the StomaphyX group were short-term nausea and headache (Table 2). On the other hand, the RYGB group had an overall 43.5% complication rate with a postoperative mortality. Major complications in the RYGB group include anastomotic leak (8.7%), incisional hernia (13%), fistula (8.7%), respiratory failure (8.7%), and perforation (4.4%). The median LOS following RYGB was six days compared to 1.5 ± 0.5 days following StomaphyX endoscopic revision.

4. Discussion

This study retrospectively compares the use of two distinct treatment strategies in patients with failed VBG at a single institution. Formal conversion of the VBG to a RYGB in morbidly obese patients results in marked weight loss, however, with considerable morbidity. On the other hand, endoscopic revision with the StomaphyX device in morbidly obese patient produced less weight loss, with minimal morbidity.

FIGURE 2: Comparison of initial body mass index and at 6-month followup in VBG patients revised by RYGB or endoscopically (StomaphyX). * $P < 0.05$ versus preoperative body mass index.

TABLE 2: Complications following VBG revision by either StomaphyX or RYGB.

Type of complication	StomaphyX (N = 14)	RYGB (N = 23)
Death	0	1
Anastomotic leakage	N/A	2
Fistula	0	2
Wound infection/abscess	N/A	3
Cardiopulmonary failure	0	2
Minor complications	4*	6
Total	4	16

N/A: not applicable. *Short-term nausea and headache.

Similar to our findings, Gagné et al. reported that 38% of patients converted to RYGB had either early or late complications, including anastomotic leak, abscess, stricture, bleeding, and respiratory failure. Other studies have also suggested increased complication rates ranging from 12% to 41%, following revisional surgery in previously failed restrictive bariatric procedures [10, 11].

Endoscopic revision of failed VBG via StomaphyX device is a novel treatment strategy. Endoscopic treatment avoids the need for operative revision and related intra-abdominal complications, such as anastomotic leakage. Two common reasons for VBG failure are enlargement of the gastric pouch and staple line dehiscence [14]. The StomaphyX device is used to decrease the size of the gastric pouch by approximating and immobilizing two or more serosal surfaces through tissue fastening [13]. Mikami et al. reported successful use of the StomaphyX device in 39 patients to reduce the size of the gastric pouch following RYGB [13]. These authors reported a 10 kg weight loss at one-year followup with no major complications and no mortalities. Leitman et al. also reported a 7.3 kg weight loss at one-year followup and no major complications, following endoscopic revision with the StomaphyX device [15].

The two main limitations of our study are duration of followup and a small sample size. There are also other potentially

viable options such as revision with an adjustable gastric band that were not included. With these major limitations its not possible to make any definitive recommendations about the use of StomaphyX. It does, however, present an interesting debate about the relative importance of less complications versus more weight loss.

5. Conclusion

In conclusion, endoscopic revision via the StomaphyX device is a safe revisional treatment strategy in morbidly obese patients that have failed VBG. In addition, endoscopic revision may be a reasonable initial approach to failed VBG, with low complication rates. Further studies are needed to clarify the role of the StomaphyX pouch reduction in patients with failed VBG.

Authors' Contributin

J. Bolton and R. S. Gill both serve as first authors of this paper.

References

- [1] WHO, "Obesity and overweight," Fact Sheet 311, 2011.
- [2] R. Padwal, S. Klarenbach, N. Wiebe et al., "Bariatric surgery: a systematic review and network meta-analysis of randomized trials," *Obesity Reviews*, vol. 12, no. 8, pp. 602–621, 2011.
- [3] E. E. Mason, "Vertical banded gastroplasty for obesity," *Archives of Surgery*, vol. 117, no. 5, pp. 701–706, 1982.
- [4] B. M. Balsiger, J. L. Poggio, J. Mai, K. A. Kelly, and M. G. Sarr, "Ten and more years after vertical banded gastroplasty as primary operation for morbid obesity," *Journal of Gastrointestinal Surgery*, vol. 4, no. 6, pp. 598–605, 2000.
- [5] E. E. Mason and J. J. Cullen, "Management of complications in vertical banded gastroplasty," *Current Surgery*, vol. 60, no. 1, pp. 33–37, 2003.
- [6] D. A. del Amo, M. M. Díez, M. E. Guedeá, and V. A. Diago, "Vertical banded gastroplasty: is it a durable operation for morbid obesity?" *Obesity Surgery*, vol. 14, no. 4, pp. 536–538, 2004.
- [7] H. J. Sugerman, J. M. Kellum Jr., E. J. DeMaria, and H. D. Reines, "Conversion of failed or complicated vertical banded gastroplasty to gastric bypass in morbid obesity," *American Journal of Surgery*, vol. 171, no. 2, pp. 263–269, 1996.
- [8] J. Hedberg, S. Gustavsson, and M. Sundom, "Long-term follow-up in patients undergoing open gastric bypass as a revisional operation for previous failed restrictive procedures," *Surgery for Obesity and Related Diseases*, vol. 8, no. 6, pp. 696–701, 2012.
- [9] D. J. Gagné, E. Dovec, and J. E. Urbandt, "Laparoscopic revision of vertical banded gastroplasty to Roux-en-Y gastric bypass: outcomes of 105 patients," *Surgery for Obesity and Related Diseases*, vol. 7, no. 4, pp. 493–499, 2011.
- [10] R. W. Schwartz, W. E. Strodel, W. S. Simpson, and W. O. Griffen, "Gastric bypass revision: lessons learned from 920 cases," *Surgery*, vol. 104, no. 4, pp. 806–812, 1988.
- [11] J. de Csepe, R. Nahouraii, and M. Gagner, "Laparoscopic gastric bypass as a reoperative bariatric surgery for failed open restrictive procedures: initial experience in seven patients," *Surgical Endoscopy*, vol. 15, no. 4, pp. 393–397, 2001.

- [12] N. Manouchehri, D. Birch, C. Menzes, X. Shi, and S. Karmali, "Natural orifice surgery: endoluminal pouch reduction following failed vertical banded gastroplasty," *Obesity Surgery*, vol. 21, no. 11, pp. 1787–1791, 2011.
- [13] D. Mikami, B. Needleman, V. Narula, J. Durant, and W. S. Melvin, "Natural orifice surgery: initial US experience utilizing the StomaphyX™ device to reduce gastric pouches after Roux-en-Y gastric bypass," *Surgical Endoscopy and Other Interventional Techniques*, vol. 24, no. 1, pp. 223–228, 2010.
- [14] D. C. Lau, J. D. Douketis, K. M. Morrison, I. M. Hramiak, A. M. Sharma, and E. Ur, "2006 Canadian clinical practice guidelines on the management and prevention of obesity in adults and children," *Canadian Medical Association Journal*, vol. 176, no. 8, pp. S1–S13, 2007.
- [15] I. M. Leitman, C. S. Virk, D. V. Avgerinos et al., "Early results of trans-oral endoscopic plication and revision of the gastric pouch and stoma following Roux-en-Y gastric bypass surgery," *Journal of the Society of Laparoendoscopic Surgeons*, vol. 14, no. 2, pp. 217–220, 2010.