

Clinical Study

Calculation of Unknown Preoperative K Readings in Postrefractive Surgery Patients

Nicola Rosa ¹, Maddalena De Bernardo ¹ and Maria Borrelli²

¹Department of Medicine and Surgery, University of Salerno, Salerno, Italy

²Department of Ophthalmology, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

Correspondence should be addressed to Maddalena De Bernardo; mdebernardo@unisa.it

Received 11 July 2017; Accepted 20 December 2017; Published 11 February 2018

Academic Editor: Ciro Costagliola

Copyright © 2018 Nicola Rosa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. To determine the unknown preoperative K readings (Kpre) to be used in history-based methods, for intraocular lens (IOL) power calculation in patients who have undergone myopic photorefractive keratectomy (PRK). **Methods.** A regression formula generated from the left eyes of 174 patients who had undergone PRK for myopia or for myopic astigmatism was compared with other methods in 168 right eyes. The Pearson index and paired *t*-test were utilized for statistical analysis. **Results.** The differences between Kpre and those obtained with the other methods were as follows: 0.61 ± 0.94 D (range: -3.94 to 2.05 D, $p < 0.01$) subtracting the effective treatment, 0.01 ± 0.86 D (range: -2.61 to 2.34 D, $p = 0.82$) with Rosa's formula, -0.02 ± 1.31 D (range: -3.43 to 3.68 D, $p = 0.82$) with the current study formula, and -0.43 ± 1.40 D (range: -3.98 to 3.12 D, $p < 0.01$) utilizing a mean K (Km) of 43.5 D. **Conclusions.** These formulas may permit the utilization of history-based methods, that is, the double-K method in calculating the IOL power following PRK when Kpre are unknown.

1. Introduction

After refractive surgery for myopia, both keratometry and corneal topography tend to overestimate the corneal power, and consequently the calculated IOL power is underestimated, with a high risk of postoperative hyperopia, which may lead to IOL exchange or a piggyback lens [1–9].

This is an important problem because a large number of patients have undergone this surgery with excellent results. Consequently, in the case of a cataract developing, they will expect the same excellent uncorrected visual acuity that they had after refractive surgery, before the cataract onset.

The inaccuracy of the IOL power calculation, excluding an incorrect estimation of axial length [10], is due to several reasons. Among these are pupil width [11], inaccurate measurement of anterior corneal curvature by automated and manual keratometry (K) or computerized videokeratography [12, 13], inaccurate value of the keratometric index resulting from the modified relationship between the anterior and posterior corneal surface, and incorrect estimation of the

effective lens position (ELP) resulting from these modifications. To overcome the last problem, Aramberri [14] and Rosa et al. [15] described the so-called double-K method, in which the preoperative K readings (Kpre) are used to estimate the effective lens position. Unfortunately, in most cases, this value is unknown.

The purpose of our study was to find a new method that permits the calculation of the Kpre, when they are not available, and to compare and discuss the results of those obtained by other methods described in the literature [13–16].

2. Materials and Methods

This retrospective study was comprised of consecutive patients who had undergone photorefractive keratectomy (PRK) for myopia or for myopic astigmatism. The institutional review board (IRB) approved the retrospective review of records of analyzed data, which had been collected as part of standard of care. Patients gave their informed consent for the surgery.




FIGURE 1: Group B: correlation between the postoperative mean keratometry (Km) and the difference between postoperative and preoperative mean keratometry (ΔK_m).

Patients were asked to discontinue wearing contact lenses for at least 1 month before the last refractive evaluation, which occurred on the day the patients underwent PRK.

Patients with systemic and ocular diseases that might interfere with the corneal healing process [17–22] or with the refractive outcome, such as diabetes, connective tissue disorders, dry eye, uveitis, corneal and lens opacities, and glaucoma, were excluded from the treatment.

All PRK treatments were performed using topical anesthesia (oxybuprocaine eye drops). The lids were opened with a speculum, and the epithelium was debrided by mechanical brush epithelial removal. All treatments were performed with a 193 nm ESIRIS excimer laser (Schwind, Kleinostheim, Germany). Immediately after surgery, a bandage contact lens was applied to the treated eye under sterile conditions; the bandage was not removed until complete reepithelialization. During this period, operated eyes received diclofenac sodium 0.1% eye drops twice a day for the first 2 days, netilmicin preservative-free eye drops until reepithelialization, and preservative-free artificial tears for 1 month after reepithelialization. All patients received clobetasone eye drops for 1 month in a tapered dose as follows: 1 drop 4 times a day for the first week, 1 drop 3 times a day for the second week, 1 drop twice a day for the third week, and 1 drop once a day for the last week.

Before and 6 months after PRK, all patients had a complete ophthalmic examination, including automatic K readings with an IOLMaster (version 4.08.0002; Zeiss, Jena, Germany). The mean of 3 consecutive good-quality measurements for the keratometry was used.

The 6-month postoperative K readings (K_{post}) were related to the difference in K readings ($K_{post} - K_{pre}$) in an attempt to find a correlation formula that could be utilized by subtracting the calculated differences of the K_{post} to obtain the K_{pre} .

A total of 174 consecutive patients (75 males, 99 females) with a mean age of 32 ± 9 years (range: 18 to 56 years) who had undergone myopic PRK were analyzed.

The left eyes of these patients were utilized to identify a regression formula that correlated the K_{post} with the difference in K readings, evaluated with the IOLMaster: This

formula was used to calculate the K_{pre} in the right eyes of the same patients, and the differences between the K_{pre} obtained with this formula and those obtained preoperatively were compared with the differences obtained from the following methods:

- (1) The effective treatment calculated at the corneal plane (taking into account the preoperative and postoperative spectacle refraction, converted at the corneal plane with the formula [23] SE corneal plane = SE spectacle plane/[1 - (0.012 × SE spectacle plane)]) was added to the K_{post} .
- (2) In 2004, in an attempt to overcome the problem of the underestimation of the IOL power in patients who had undergone PRK, one of the authors of the current study published the formula $y = 0.7615x - 0.6773$, where x is the difference in refraction at the corneal plane and y is the keratometric difference evaluated with the IOLMaster [13], to be used when the effective treatment was known. In the current paper, the authors utilized this formula to calculate a modified difference in K readings to add to the postoperative ones in order to calculate the preoperative readings.
- (3) A mean K (K_m) of 43.5 [16] was used as standard K_{pre} , and the differences with the preoperative ones were calculated.

Statistical analysis was performed with a Student paired t -test, and $p < 0.01$ was considered to be statistically significant.

3. Results

The correlation between the K_{post} and the difference between K_{post} and K_{pre} , obtained by analyzing the 174 left eyes with a preoperative refraction of -4.87 ± 2.5 D (range: -10.0 to -0.5 D) and a 6-month postoperative refraction of 0.57 ± 0.81 D (range: -1.88 to $+2.5$ D), gave the following formula (Figure 1): $y = 0.8197x - 36.907$ (where $x = K_{post}$ and $y = K_{post} - K_{pre}$).

TABLE 1: Differences (in diopters) in 168 right eyes between preoperative keratometry readings and the calculated ones with the knowledge of the effective treatment.

	Effective treatment calculated at the corneal plane	Rosa et al. 2004
Mean	-0.61	0.01
SD	0.94	0.86
Min	-3.94	-2.61
Max	2.05	2.34
<i>p</i>	<0.01	0.82

TABLE 2: Comparison between the two methods to calculate the preoperative keratometry readings with the knowledge of the effective treatment.

Range	Effective treatment calculated at the corneal plane		Rosa et al. 2004	
	<i>n</i>	%	<i>n</i>	%
±0.5 D	65	38.7	94	56
±1 D	146	86.9	150	89.3
±2 D	160	95.2	165	98.2
>±2 D	8	4.8	3	1.8

n = number of eyes; % = percentage of eyes; D = diopters.

168 right eyes of the same patients (6 of them did not undergo surgery because they were emmetropic) with a preoperative refraction of -5.38 ± 2.86 D (range: -9.33 to -0.5 D) and a 6-month postoperative refraction of 0.49 ± 0.71 D (range: -2 to $+1.5$ D) were used to test the difference between the preoperative K readings and those calculated with the formula obtained from the left eyes and also with the other above-mentioned methods [5, 13, 16, 23].

Tables 1–4 show that the best results can be obtained mainly with the formula previously described [13] and with the one calculated in the current study, as there is no statistically significant difference with the preoperative K readings (Tables 1 and 3).

In particular, the comparison between these two methods shows that the formula based on the knowledge of preoperative parameters achieves higher percentages of eyes closer to the real preoperative K readings (Tables 2 and 4).

4. Discussion

With the third-generation IOL power theoretic formulas (e.g., SRK-T), the AL and K are used to calculate the ELP, which would be automatically underestimated after refractive surgery, as a consequence of K value underestimation [14]. The double-K method described by Aramberri requires a knowledge of the Kpre to estimate the effective lens position in the case of cataract surgery in patients who had previously undergone corneal refractive surgery.

This is an easy task if the Kpre have been recorded, but regrettably, this rarely happens.

TABLE 3: Differences (in diopters) in 168 right eyes between preoperative keratometry readings and the calculated ones without the knowledge of preoperative parameters.

	Current study	Preoperative mean keratometry = 43.5
Mean	-0.02	-0.43
SD	1.31	1.40
Min	-3.43	-3.98
Max	3.68	3.12
<i>p</i>	0.82	<0.01

TABLE 4: Comparison between the two methods to calculate the preoperative keratometry readings without the knowledge of preoperative parameters.

Range	Current study		Preoperative mean keratometry = 43.5	
	<i>n</i>	%	<i>n</i>	%
±0.5 D	54	32.1	39	23.2
±1 D	123	73.2	120	71.4
±2 D	160	95.2	156	92.9
>±2 D	8	4.8	12	7.1

n = number of eyes; % = percentage of eyes; D = diopters.

Theoretically, if the achieved correction is known, it should be possible to calculate Kpre by adding this value to the postoperative ones. This is not always correct in practice, because it has been shown that the effective treatment does not correspond to the changes detected by the machines; in particular, postoperatively, there is an overestimation of the corneal power, resulting in an underestimation of the effective treatment [24, 25].

Few methods have been described to calculate the preoperative Kpre [13, 16, 26, 27].

In 2004, Rosa et al. [13] described a formula which was not designed with the purpose of calculating the Kpre, but tested in the present paper for such a purpose, and have shown that the results are better than those obtained when the pure effective treatment is used.

In the case that the effective treatment is unknown, we may use a standard mean K [16], or we may use the formula obtained from this study. Although the results of both methods appear to be less accurate than the ones obtained when the effective treatment is known, the results obtained with this new formula are better than the ones obtained utilizing a standard value of 43.5 D.

Recently Saiki et al. [26] described a method to calculate the Kpre, starting from the assumption that it is possible to predict the preoperative anterior corneal power from the postoperative posterior one in patients who had undergone LASIK surgery.

Similar to Saiki, in a recent paper, De Bernardo et al. [27] found no significant changes in the posterior corneal surface after PRK with a good correlation between preoperative anterior Km and postoperative posterior Km. This not only confirms the findings of previously published studies [28–31] but also confirms the possibility of predicting the preoperative

anterior Km by utilizing the postoperative posterior one, as stated by Saiki et al. [26].

The results of our present study indicate that it is possible to calculate the Kpre without utilizing the posterior postoperative K readings. Unfortunately, we were not able to compare our data with those of Saiki and De Bernardo because to make such a calculation, the Pentacam is needed, and these data were not available for the patients examined in the present study.

Another limitation of our study is that the formula published in 2004 has been proven to be effective only for the IOLMaster, and at present, we do not know if it can be utilized with other devices. Nevertheless, as the IOLMaster is the most widely utilized device, we believe that this formula can be easily applied.

5. Conclusions

In conclusion, we suggest that to calculate the Kpre, if the effective treatment is known, the formula obtained from the study published in 2004 [13] is the best option. If, on the other hand, the effective treatment is not available, both standard mean K and the formula reported in this study may be utilized.

Conflicts of Interest

No conflicting relationship exists for any author.

References

- [1] D. S. Siganos, I. G. Pallikaris, J. E. Lambropoulos, and C. J. Koufala, "Keratometric readings after photorefractive keratectomy are unreliable for calculating IOL power," *Journal of Refractive Surgery*, vol. 12, no. 2, pp. S278–S279, 1996.
- [2] N. Rosa, L. Capasso, M. Lanza, and M. Borrelli, "Clinical results of a corneal radius correcting factor in calculating intraocular lens power after corneal refractive surgery," *Journal of Refractive Surgery*, vol. 25, no. 7, pp. 599–603, 2009.
- [3] B. Seitz and A. Langenbacher, "Intraocular lens power calculation in eyes after corneal refractive surgery," *Journal of Refractive Surgery*, vol. 16, no. 3, pp. 349–361, 2000.
- [4] J. G. Ladas, B. S. Boxer Wachler, J. D. Hunkeler, and D. S. Durrie, "Intraocular lens power calculations using corneal topography after photorefractive keratectomy," *American Journal of Ophthalmology*, vol. 132, no. 2, pp. 254–255, 2001.
- [5] S. Chen and F.-R. Hu, "Correlation between refractive and measured corneal power changes after myopic excimer laser photorefractive surgery," *Journal of Cataract & Refractive Surgery*, vol. 28, no. 4, pp. 603–610, 2002.
- [6] M. De Bernardo, P. Cornetta, and N. Rosa, "Intraocular lens calculation adjustment after laser refractive surgery," *Journal of Cataract & Refractive Surgery*, vol. 43, no. 4, p. 578, 2017.
- [7] V. Feiz, M. Moshirfar, M. J. Mannis et al., "Nomogram-based intraocular lens power adjustment after myopic photorefractive keratectomy and LASIK: a new approach," *Ophthalmology*, vol. 112, no. 8, pp. 1381–1387, 2005.
- [8] E. Borasio, J. Stevens, and G. T. Smith, "Estimation of true corneal power after keratorefractive surgery in eyes requiring cataract surgery: BESSt formula," *Journal of Cataract & Refractive Surgery*, vol. 32, no. 12, pp. 2004–2014, 2006.
- [9] W. Haigis, "Intraocular lens calculation after refractive surgery for myopia: Haigis-L formula," *Journal of Cataract & Refractive Surgery*, vol. 34, no. 10, pp. 1658–1663, 2008.
- [10] N. Rosa, L. Capasso, M. Lanza, and A. Romano, "Axial eye length evaluation before and after myopic photorefractive keratectomy," *Journal of Refractive Surgery*, vol. 21, no. 3, pp. 281–287, 2005.
- [11] P.-R. Preussner, J. Wahl, H. Lahdo, B. Dick, and O. Findl, "Ray tracing for intraocular lens calculation," *Journal of Cataract & Refractive Surgery*, vol. 28, no. 8, pp. 1412–1419, 2002.
- [12] N. Rosa, M. De Bernardo, M. Borrelli, M. L. Filosa, E. Minutillo, and M. Lanza, "Reliability of the IOLMaster in measuring corneal power changes after hyperopic photorefractive keratectomy," *Journal of Refractive Surgery*, vol. 27, no. 4, pp. 293–298, 2011.
- [13] N. Rosa, L. Capasso, M. Lanza, D. Furgiuele, and A. Romano, "Reliability of the IOLMaster in measuring corneal power changes after photorefractive keratectomy," *Journal of Cataract & Refractive Surgery*, vol. 30, no. 2, pp. 409–413, 2004.
- [14] J. Aramberri, "Intraocular lens power calculation after corneal refractive surgery: double-K method," *Journal of Cataract & Refractive Surgery*, vol. 29, no. 11, pp. 2063–2068, 2003.
- [15] N. Rosa, L. Capasso, and M. Lanza, "Double-K method to calculate IOL power after refractive surgery," *Journal of Cataract & Refractive Surgery*, vol. 31, no. 2, pp. 254–255, 2005.
- [16] K. J. Hoffer, "Intraocular lens power calculation after previous laser refractive surgery," *Journal of Cataract and Refractive Surgery*, vol. 35, no. 4, pp. 759–765, 2009.
- [17] M. De Bernardo, L. Capasso, M. Lanza et al., "Long-term results of corneal collagen crosslinking for progressive keratoconus," *Journal of Optometry*, vol. 8, no. 3, pp. 180–186, 2015.
- [18] N. Rosa, M. Lanza, M. Borrelli et al., "Corneal thickness and endothelial cell characteristics in patients with myotonic dystrophy," *Ophthalmology*, vol. 117, no. 2, pp. 223–225, 2010.
- [19] N. Rosa, M. Lanza, M. De Bernardo, G. Signoriello, and P. Chiodini, "Relationship between corneal hysteresis and corneal resistance factor with other ocular parameters," *Seminars in Ophthalmology*, vol. 30, no. 5–6, pp. 335–339, 2015.
- [20] M. De Bernardo and N. Rosa, "Central corneal thickness after cross-linking using high-definition optical coherence tomography, ultrasound, and dual Scheimpflug tomography: a comparative study over one year," *American Journal of Ophthalmology*, vol. 176, p. 254, 2017.
- [21] M. De Bernardo, P. Cornetta, and N. Rosa, "Safety and efficacy of sequential intracorneal ring segment implantation and cross-linking in pediatric keratoconus," *American Journal of Ophthalmology*, vol. 181, pp. 182–183, 2017.
- [22] M. De Bernardo and N. Rosa, "Fuchs' endothelial and myotonic dystrophies: corneal dystrophy in myotonic patients," *Investigative Ophthalmology & Visual Science*, vol. 58, no. 13, p. 5838, 2017.
- [23] B. Seitz, A. Langenbacher, N. X. Nguyen, M. M. Kus, and M. Kuchle, "Underestimation of intraocular lens power for cataract surgery after myopic photorefractive keratectomy," *Ophthalmology*, vol. 106, no. 4, pp. 693–702, 1999.
- [24] M. De Bernardo, L. Zeppa, M. Cennamo, S. Iaccarino, L. Zeppa, and N. Rosa, "Prevalence of corneal astigmatism before cataract surgery in Caucasian patients," *European Journal of Ophthalmology*, vol. 24, no. 4, pp. 494–500, 2014.
- [25] N. Rosa, M. De Bernardo, M. Borrelli, and M. Lanza, "New factor to improve reliability of the clinical history method for

- intraocular lens power calculation after refractive surgery,” *Journal of Cataract & Refractive Surgery*, vol. 36, no. 12, pp. 2123–2128, 2010.
- [26] M. Saiki, K. Negishi, N. Kato et al., “Modified double-K method for intraocular lens power calculation after excimer laser corneal refractive surgery,” *Journal of Cataract & Refractive Surgery*, vol. 39, no. 4, pp. 556–562, 2013.
- [27] M. De Bernardo, S. Iaccarino, M. Cennamo, L. Caliendo, and N. Rosa, “Corneal anterior power calculation for an IOL in post-PRK patients,” *Optometry and Vision Science*, vol. 92, no. 2, pp. 190–195, 2015.
- [28] J. B. Ciolino and M. W. Belin, “Changes in the posterior cornea after laser in situ keratomileusis and photorefractive keratectomy,” *Journal of Cataract & Refractive Surgery*, vol. 32, no. 9, pp. 1426–1431, 2006.
- [29] B. J. Ha, S. W. Kim, S. W. Kim, E. K. Kim, and T. I. Kim, “Pentacam and Orbscan II measurements of posterior corneal elevation before and after photorefractive keratectomy,” *Journal of Refractive Surgery*, vol. 25, no. 3, pp. 290–295, 2009.
- [30] N. Rosa, M. De Bernardo, S. Iaccarino, and M. Cennamo, “Intraocular lens power calculation: a challenging case,” *Optometry and Vision Science*, vol. 91, no. 2, pp. e29–e31, 2014.
- [31] M. De Bernardo, M. Borrelli, M. Mariniello, M. Lanza, and N. Rosa, “Pentacam vs SP3000P specular microscopy in measuring corneal thickness,” *Contact Lens & Anterior Eye*, vol. 38, no. 1, pp. 21–27, 2015.



Hindawi

Submit your manuscripts at www.hindawi.com

