Research Article

On the One Dimensional Poisson Random Geometric Graph

L. Decreusefond and E. Ferraz

Institut Télécom, Télécom ParisTech, CNRS LTCI, 75634 Paris, France

Correspondence should be addressed to E. Ferraz, eduardo.ferraz@telecom-paristech.fr

Received 24 March 2011; Revised 20 July 2011; Accepted 22 July 2011

Academic Editor: Kelvin K. W. Yau

Copyright © 2011 L. Decreusefond and E. Ferraz. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Given a Poisson process on a bounded interval, its random geometric graph is the graph whose vertices are the points of the Poisson process, and edges exist between two points if and only if their distance is less than a fixed given threshold. We compute explicitly the distribution of the number of connected components of this graph. The proof relies on inverting some Laplace transforms.

1. Motivation

As technology goes on [1–3], one can expect a wide expansion of the so-called *sensor networks*. Such networks represent the next evolutionary step in building, utilities, industry, home, agriculture, defense, and many other contexts [4].

These networks are built upon a multitude of small and cheap sensors which are devices with limited transmission capabilities. Each sensor monitors a region around itself by measuring some environmental quantities (e.g., temperature, humidity), detecting intrusion and so forth, and broadcasts its collected information to other sensors or to a central node. The question of whether information can be shared among the whole network is then of crucial importance. Mathematically speaking, sensors can be abstracted as points in $\mathbf{R}^2$, $\mathbf{R}^3$, or a manifold. The region a sensor monitors is represented by a circle centered at the location of the sensor. In what follows, it is assumed that the broadcast radius, that is, the distance at which a sensor can communicate with another sensor, is equal to the monitoring radius. Two questions are then of interest: can any two sensors communicate using others as hopping relays and is the whole region covered by all the sensors? The recent works of Ghrist and his collaborators [5, 6] show how, in any dimension, algebraic topology can be used to answer these questions. Their method consists in building the so-called simplicial complex associated to the configuration of points and the radius of communication. Then, simple algebraic computations yield the Betti numbers: the first Betti number, usually denoted as β_0 , is the

number of connected components; the second number β_1 is the number of coverage holes. Thus, we have a satisfactory deployment whenever $\beta_0 = 1$ and $\beta_1 = 0$. Trying to pursue their work for random settings, we quickly realized that the dimension of the ambient space played a key role. We then first began by the analysis of dimension 1, which appeared to be the simplest situation. In this case, there is no need of algebraic topology so we will not go further in the description of this line of thought even if it was our first motivation.

In dimension 1, the only question of interest is that of the connexity but it can take different forms. Imagine we are given $[0, 1]$ as a domain in which n points $\{x_1, \dots, x_n\}$ are drawn. For a radius r , one can wonder whether $[0, 1] \subset \cup_{i=1, \dots, n} [x_i - r, x_i + r]$, or one can investigate whether $[x_i - r, x_i + r] \cap [x_{i+1} - r, x_{i+1} + r] \neq \emptyset$ for all $i = 1, \dots, n - 1$. The second situation is less restrictive since we do not impose that the frontier of the interval is to be covered. Depending on the application, we have in mind, both questions are sensible. A slightly different but somehow close problem is that of the circle: consider now that the points are dispatched along a circle of unit perimeter C_1 and ask again whether $C_1 \subset \cup_{i=1, \dots, n} B(x_i, r)$ where $B(x, r)$ is the 2-dimensional ball of center x and radius r . Several years ago, this problem has been thoroughly analyzed ([7] and references therein) for a fixed number of i.i.d. arcs over the circle. A closed form formula can be given for the probability of coverage as a function of the number and of the common law of the arcs length. Some variations of this problem have been investigated since then, see, for instance, [8]. More recently, in [9], algorithms are devised to determine whether a domain can be protected from intrusion by a “belt” of sensors (namely, a ring or the border of a rectangle). There is no performance analysis in this work which is focused on algorithmic solutions for this special problem of coverage. Still motivated by applications to sensor networks, [10] considers the situation where sensors are actually placed in a plan, have a fixed radius of observation, and analyse the connectivity of the trace of the covered region over a line. Some recent results of Kahle [11, 12] are actually hardly linked to our results: the motivation is the same, studying the Betti numbers of some random simplicial complexes, but the results are only asymptotic and valid in dimension greater than 2.

Our main result is the distribution of the number of connected components for a Poisson distribution of sensors in a bounded interval. We could not use the method of [7] since the number of gaps does not determine the connectivity of the domain. For instance, one may have only one gap at the “beginning” which means that all the points are pairwise within the threshold distance and, thus, that the network is connected, or one may have only one gap in the “middle” which means that there is a true hole of connectivity.

Actually, our method is very much related to the queueing theory. Indeed, clusters, that is, sequence of neighboring points, are the strict analogous of busy periods—see Section 2. As will appear below, our analysis turns down to be that of an M/D/1/1 queue with a preemption: when a customer arrives during a service, it preempts the server, and, since there is no buffer, the customer who was in service is removed from the queueing system. This analogy led us to use standard tools of queueing theory: Laplace transform and renewal processes—see, for instance, [13, 14]. This works perfectly, and, with a bit of calculus, we can compute all the characteristics we are interested in. It is worthwhile to note that a queueing model (namely the M/G/ ∞) also appears in [10].

The paper is organized as follows: Section 2 presents the model and defines the relevant quantities to be calculated. The calculations and analytical results are presented in Section 3. For our situation, we find results analogous to that of [7]. In Section 4, two other scenarios are presented, considering the number of incomplete clusters and clusters placed in a circle. In Section 5, numerical examples are presented and analyzed.

2. Problem Formulation

Let $L > 0$, we assume that we are given a Poisson process, denoted by N , of intensity λ on $[0, L]$. Let $(X_i, i \geq 1)$ be the atoms of N . We, thus, know that the random variables, $\Delta X_i = X_{i+1} - X_i$ are i.i.d. and exponentially distributed. We fix $\epsilon > 0$. Two points, located, respectively, at x and y , are said to be *directly connected* whenever $|x - y| \leq \epsilon$. For $i < j$, two points of N , say X_i and X_j , are indirectly connected if X_i and X_{l+1} are directly connected for any $l = i, \dots, j-1$. A set of points directly or indirectly connected is called a *cluster*, a complete cluster is a cluster which begins and ends within $[0, L]$. The connectivity of the whole network is measured by the number of clusters.

The number of points in the interval $[0, x]$ is denoted by $N_x = \sum_{n=0}^{\infty} \mathbf{1}_{\{X_n \leq x\}}$. The random variable A_i given by

$$A_i = \begin{cases} X_1 & \text{if } i = 1, \\ \inf\{X_j : X_j > A_{i-1}, X_j - X_{j-1} > \epsilon\} & \text{if } i > 1, \end{cases} \quad (2.1)$$

represents the beginning of the i th cluster, denoted by C_i . In the same way, the end of this same cluster, E_i , is defined by

$$E_i = \inf\{X_j + \epsilon : X_j > A_i, X_{j+1} - X_j > \epsilon\}. \quad (2.2)$$

So, the i th cluster, C_i , has a number of points given by $N_{E_i} - N_{A_i}$. We define the length B_i of C_i as $E_i - A_i$. The intercluster size, D_i , is the distance between the end of C_i and the beginning of C_{i+1} , which means that $D_i = A_{i+1} - E_i$, and ΔA_i is the distance between the first points of two consecutive clusters C_i , given by $\Delta A_i = A_{i+1} - A_i = B_i + D_i$.

Remark 2.1. With this set of assumptions and definitions, we can see our problem as an $M/D/1/1$ preemptive queue, see Figure 1. In this nonconservative system, the service time is deterministic and given by ϵ . When a customer arrives during a service, the served customer is removed from the system and replaced by the arriving customer. Within this framework, a cluster corresponds to what is called a busy period, the intercluster size is an idle time, and $A_i + D_i$ is the length of the i th cycle.

The number of complete clusters in $[0, L]$ corresponds to the number of connected components $\beta_0(L)$ (since, in dimension 1, it coincides with the Euler characteristics of the union of intervals, see [5]) of the network. The distance between the beginning of the first cluster and the beginning of the $(i+1)$ th one is defined as $U_i = \sum_{k=1}^i \Delta A_k$. We also define $\Delta X_0 = D_0 = X_1$. Figure 2 illustrates these definitions.

For the sake of completeness, we recall the essentials of Markov's process theory needed to go along, for further details we refer, for instance, to [13, 14]. In what follows, for a process X , $(\mathcal{F}_t^X, t \geq 0)$ is the filtration generated by the sample-paths of X :

$$\mathcal{F}_t^X = \sigma\{X(s), s \geq t\}. \quad (2.3)$$

Figure 1: Queueing representation of the model. A down arrow denotes that user i starts his service. An up arrow indicates that user i leaves the system without having finished the service. A double up arrow illustrates that the service of user i finishes. Beginning and end of the i th busy period, respectively, A_i and E_i , are also shown.

Figure 2: Definitions of the relevant quantities of the network. Distance between points, distance between clusters, clusters size, interclusters size, beginning and ends of clusters.

Definition 2.2. A process $(X(t), t \geq 0)$ with values in a denumerable space E is said to be Markov whenever

$$\mathbb{E}[F(X(t+s)) \mid \mathcal{F}_t^X] = \mathbb{E}[F(X(t+s)) \mid X(t)], \quad (2.4)$$

for any bounded function F from E to $\mathbf{R}$, any $t \geq 0$ and $s \geq 0$.

Equivalently, a process X is Markov if and only if, given the present (i.e., given $X(t)$), the past (i.e., the sample-path of X before time t) and the future (i.e., the sample-path of X after time t) of the process are independent.

Definition 2.3. A random variable τ with values in $\mathbf{R}^+ \cup \{+\infty\}$ is an $\mathcal{F}_t^X$ -stopping time whenever, for any $t \geq 0$, the event $\{\tau \leq t\}$ belongs to $\mathcal{F}_t^X$.

The point is that (2.4) still holds when t is replaced by a stopping time τ : given $X(\tau)$, the past and the future of X are independent. X is then said to be strong Markov. This property always holds for the Markov processes with values in a denumerable space but is not necessarily true for the Markov processes with values in an arbitrary space.

From now on, the Markov process under consideration is N , the Poisson process of intensity λ over $[0, L]$.

Lemma 2.4. *For any $i \geq 1$, A_i and E_i are stopping times.*

Proof. Let us consider the filtration $\mathcal{F}_t^N = \sigma\{N_a, a \leq t\}$. For $i = 1$, we have

$$\{A_1 \leq t\} \iff \{X_1 \leq t\} \iff \{N_t \geq 1\} \in \mathcal{F}_t^N. \quad (2.5)$$

Thus, A_1 is a stopping time. For A_2 , we have

$$\{A_2 > t\} \iff \bigcup_{n \geq 1} \left\{ N_t = n, \bigcup_{j=1}^n \left\{ \Delta X_j \geq \epsilon, \bigcup_{k=j+1}^n \{\Delta X_k \leq \epsilon\} \right\} \right\} \in \mathcal{F}_t^N, \quad (2.6)$$

so A_2 is also a stopping time. We proceed along the same line for others A_i and as well for E_i to prove that they are stopping times. $\square$

Since N is a (strong) Markov's process, the next corollary is immediate.

Corollary 2.5. *The set $\{B_i, D_i, i \geq 1\}$ is a set of independent random variables. Moreover, D_i is distributed as an exponential random variable with mean $1/\lambda$, and the random variables $\{B_i, i \geq 1\}$ are i.i.d.*

3. Calculations

Theorem 3.1. *The Laplace transform of the distribution of B_i is given by*

$$\mathbb{E}\left[e^{-sB_i}\right] = \frac{\lambda + s}{\lambda + se^{(\lambda+s)\epsilon}}. \quad (3.1)$$

Proof. Since ΔX_j is an exponentially distributed random variable,

$$\mathbb{E}\left[e^{-s\Delta X_j} \mathbf{1}_{\{\Delta X_j \leq \epsilon\}}\right] = \int_0^\epsilon e^{-st} \lambda e^{-\lambda t} dt = \frac{\lambda}{s + \lambda} \left(1 - e^{-(s+\lambda)\epsilon}\right). \quad (3.2)$$

Hence, the Laplace transform of the distribution of B_1 is given by

$$\begin{aligned} \mathbb{E}\left[e^{-sB_1}\right] &= \sum_{n=1}^{\infty} \mathbb{E}\left[e^{-sB_1}, N_{E_1} = n\right] \\ &= \sum_{n=1}^{\infty} \mathbb{E}\left[e^{-s(\sum_{j=1}^{n-1} \Delta X_j + \epsilon)} \mathbf{1}_{\{\Delta X_n > \epsilon\}} \prod_{j=1}^{n-1} \mathbf{1}_{\{\Delta X_j \leq \epsilon\}}\right] \\ &= \sum_{n=1}^{\infty} \left(\mathbb{E}\left[e^{-s\Delta X_1} \mathbf{1}_{\{\Delta X_1 \leq \epsilon\}}\right]\right)^{n-1} \mathbb{E}\left[e^{-s\Delta X_n} \mathbf{1}_{\{\Delta X_n > \epsilon\}}\right] e^{-s\epsilon} \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \left(\frac{\lambda}{s+\lambda} \left(1 - e^{-(s+\lambda)\epsilon} \right) \right)^n e^{-s\lambda} e^{-s\epsilon} \\
&= \frac{\lambda + s}{se^{\lambda\epsilon} e^{s\epsilon} + \lambda}.
\end{aligned} \tag{3.3}$$

Using Corollary 2.5, we have $\mathbb{E}[e^{-sB_1}] = \mathbb{E}[e^{-sB_i}]$, which concludes the proof. $\square$

From this result, we can immediately calculate the Laplace transform of the distribution of ΔA_i . Since $\Delta A_i = B_i + D_i$, we have $\mathbb{E}[e^{-s\Delta A_i}] = \mathbb{E}[e^{-s(B_i+D_i)}]$, and using Corollary 2.5:

$$\mathbb{E}[e^{-s\Delta A_i}] = \mathbb{E}[e^{-sB_i}] \mathbb{E}[e^{-sD_i}] = \frac{\lambda}{\lambda + se^{(\lambda+s)\epsilon}}. \tag{3.4}$$

Corollary 3.2. *The Laplace transform of the distribution of U_n , for $n \geq 0$, is given by*

$$\mathbb{E}[e^{-sU_n}] = \frac{\lambda^n}{(\lambda + se^{(\lambda+s)\epsilon})^n}. \tag{3.5}$$

Proof. We use Corollary 2.5 and Theorem 3.1 to calculate the Laplace transform of the distribution of U_n since $U_n = \sum_{i=1}^n (B_i + D_i)$:

$$\begin{aligned}
\mathbb{E}[e^{-sU_n}] &= \prod_{i=1}^n \mathbb{E}[e^{-sB_i}] \mathbb{E}[e^{-sD_i}] \\
&= \left(\frac{\lambda + s}{\lambda + se^{(\lambda+s)\epsilon}} \right)^n \left(\frac{\lambda}{\lambda + s} \right)^n,
\end{aligned} \tag{3.6}$$

hence, the result. $\square$

Let us define the function p_n as

$$p_n : x \in \mathbf{R}^+ \mapsto p_n(x) = \Pr(\beta_0(x) = n), \tag{3.7}$$

that is, $p_n(x)$ is the probability of having n clusters in the interval $[0, x]$. Since, for all $x \in \mathbf{R}_+$, $0 \leq p_n(x) \leq 1$, the Laplace transform of p_n with respect to x ,

$$\mathcal{L}\{p_n\}(s) = \int_0^{\infty} e^{-sx} p_n(x) dx, \tag{3.8}$$

is well defined.

Theorem 3.3. *For any $n \geq 0$, the Laplace transform of p_n is given by*

$$\mathcal{L}\{p_n\}(s) = \frac{\lambda^n e^{(\lambda+s)\epsilon}}{(se^{(\lambda+s)\epsilon} + \lambda)^n}. \tag{3.9}$$

Proof. We note that; see Figure 3,

$$\{\beta_0(x) \geq n\} \iff \begin{cases} \{\Delta X_0 + U_{n-1} + B_n \leq L\} & \text{if } n \geq 1, \\ \{\Delta X_0 < \infty\} & \text{if } n = 0. \end{cases} \quad (3.10)$$

Hence,

$$\Pr(\beta_0(x) = 0) = 1 - \Pr(\Delta X_0 + B_1 \leq x), \quad (3.11)$$

$$\Pr(\beta_0(x) = n) = \Pr(\Delta X_0 + U_{n-1} + B_n \leq x) - \Pr(\Delta X_0 + U_n + B_{n+1} \leq x). \quad (3.12)$$

Let

$$Y_n = \begin{cases} \Delta X_0 + U_{n-1} + B_n, & \text{if } n \geq 1, \\ 0, & \text{if } n = 0, \end{cases} \quad (3.13)$$

then we have

$$\begin{aligned} \mathcal{L}\{\Pr(Y_n \leq \cdot)\}(s) &= \int_0^\infty \Pr(Y_n \leq x) e^{-sx} dx \\ &= \int_0^\infty \int_0^x dP_{Y_n}(y) e^{-sx} dx \\ &= \frac{1}{s} \mathbb{E}[e^{-sY_n}] \\ &= \frac{1}{s} \mathbb{E}[e^{-s\Delta X_0}] \mathbb{E}[e^{-sU_{n-1}}] \mathbb{E}[e^{-sB_n}] \\ &= \frac{1}{s} \frac{\lambda^n}{(e^{\lambda\epsilon} s e^{\epsilon s} + \lambda)^n}, \end{aligned} \quad (3.14)$$

for $n \geq 1$, where we used Corollary 2.5 in the third line. For $n = 0$, the Laplace transform is trivial and given by $\mathcal{L}\{\Pr(Y_0 \leq \cdot)\}(s) = 1/s$. Substituting (3.14) in the Laplace transform of both sides of (3.12) yields

$$\begin{aligned} \mathcal{L}\{p_n\}(s) &= \mathcal{L}\{\Pr(Y_n \leq \cdot)\}(s) - \mathcal{L}\{\Pr(Y_{n+1} \leq \cdot)\}(s) \\ &= \frac{e^{\epsilon\lambda} e^{\epsilon s} \lambda^n}{(e^{\epsilon\lambda} s e^{\epsilon s} + \lambda)^{n+1}}, \quad n \geq 0. \end{aligned} \quad (3.15)$$

The proof is, thus, complete. □

Lemma 3.4. *Let m be an positive integer. For any $x > 0$, when $\epsilon \rightarrow 0$, $\mathbb{E}[\beta_0^m] \rightarrow \mathbb{E}[N_L^m]$.*

Figure 3: Illustration of the condition equivalent to $\beta_0 \geq n$.

Proof. Since there is almost surely a finite number of points in $[0, x]$, for almost all sample-paths, there exists $\eta > 0$ such that $\Delta X_j \geq \eta$ for any $j = 1, \dots, N_x$. Hence, for $\epsilon < \eta$, $\beta_0(x) = N_x$. This implies that $\beta_0(x)$ tends almost surely to N_x as ϵ goes to 0. Moreover, it is immediate by the very definition of $\beta_0(x)$ that $\beta_0(x) \leq N_x$. Since, for any m , $\mathbb{E}[N_x^m]$ is finite, the proof follows by dominated convergence. $\square$

Let $\text{Li}_t(z)$, $z, t \in \mathbf{R}$, $z < 1$, be the polylogarithm function with parameter t , defined by

$$\text{Li}_t(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^t}. \quad (3.16)$$

For m a positive integer, consider the function of x

$$M_{\beta_0}^m : x \mapsto \mathbb{E}[\beta_0^m(x)] = \sum_{i=0}^{\infty} i^m p_i(x). \quad (3.17)$$

Its Laplace transform is given by

$$\mathcal{L}\{M_{\beta_0}^m\}(s) = \int_0^{\infty} \mathbb{E}[\beta_0(x)^m] e^{-sL} dx. \quad (3.18)$$

Corollary 3.5. Let α be defined as follows:

$$\alpha = \frac{e^{\epsilon\lambda}}{\lambda} s e^{\epsilon s}. \quad (3.19)$$

The Laplace transform of the m th moment of $\beta_0(L)$ is

$$\mathcal{L}\{M_{\beta_0}^m\}(s) = \frac{\alpha}{s(\alpha+1)} \text{Li}_{-m}\left(\frac{1}{\alpha+1}\right), \quad (3.20)$$

which converges, provided that $\alpha > 0$.

Proof. Applying the Laplace transform of both sides of (3.17), we get

$$\begin{aligned}\mathcal{L}\{M_{\beta_0}^m\}(s) &= \sum_{i=1}^{\infty} i^m \mathcal{L}\{p_i\}(s) \\ &= \frac{((e^{\epsilon\lambda}/\lambda)e^{\epsilon s})}{((e^{\epsilon\lambda}/\lambda)se^{\epsilon s} + 1)} \sum_{i=1}^{\infty} \frac{i^m}{((e^{\epsilon\lambda}/\lambda)se^{\epsilon s} + 1)^i} \\ &= \frac{\alpha}{s(\alpha + 1)} \text{Li}_{-m}\left(\frac{1}{\alpha + 1}\right),\end{aligned}\tag{3.21}$$

concluding the proof. $\square$

We define $\left\{ \begin{smallmatrix} m \\ k \end{smallmatrix} \right\}$ as the Stirling number of second kind [15]; that is, $\left\{ \begin{smallmatrix} m \\ k \end{smallmatrix} \right\}$ is the number of ways to partition a set of m objects into k groups. They are intimately related to polylogarithm by the following identity (see [16]) valid for any positive integer m ,

$$\text{Li}_{-m}(z) = \sum_{k=0}^m \frac{(-1)^{m+k} k! \left\{ \begin{smallmatrix} m+1 \\ k+1 \end{smallmatrix} \right\}}{(1-z)^{k+1}}.\tag{3.22}$$

Corollary 3.6. *The m th moment of the number of clusters on the interval $[0, L]$ is given by*

$$M_{\beta_0}^m(L) = \sum_{k=1}^m \left\{ \begin{smallmatrix} m \\ k \end{smallmatrix} \right\} \left(\frac{L}{\epsilon} - k \right)^k \left(\lambda \epsilon e^{-\epsilon\lambda} \right)^k \mathbf{1}_{\{L/\epsilon > k\}}.\tag{3.23}$$

Proof. Using (3.22) in the result of Corollary 3.5, we get

$$\begin{aligned}\mathcal{L}\{M_{\beta_0}^m\}(s) &= \frac{\alpha}{s} \sum_{k=0}^m \frac{(-1)^{m+k} k! \left\{ \begin{smallmatrix} m+1 \\ k+1 \end{smallmatrix} \right\} (1+\alpha)^k}{\alpha^{k+1} (\alpha + 1)} \\ &= \frac{1}{s} \sum_{k=0}^m c_{k,m} \frac{1}{\alpha^k},\end{aligned}\tag{3.24}$$

where the coefficients $c_{k,m}$ are integers given by

$$c_{k,m} = \sum_{j=k}^m (-1)^j j! \left\{ \begin{smallmatrix} m+1 \\ j+1 \end{smallmatrix} \right\} \binom{j}{k}.\tag{3.25}$$

Using the following identity of the Stirling numbers [17],

$$\sum_{j=0}^m (-1)^j j! \left\{ \begin{smallmatrix} m+1 \\ j+1 \end{smallmatrix} \right\} = 0,\tag{3.26}$$

we find that $c_{0,m} = 0$ for m a positive integer. So, we can write the Laplace transform of the moments as

$$\mathcal{L}\{M_{\beta_0}^m\}(s) = \sum_{k=1}^m c_{k,m} \frac{(\lambda e^{-\epsilon\lambda})^k}{s^{k+1} e^{k s \epsilon}}, \quad (3.27)$$

and apply the inverse of the Laplace transform in both sides of (3.12) to obtain

$$\begin{aligned} M_{\beta_0}^m(L) &= \mathcal{L}^{-1} \left\{ \sum_{k=1}^m c_{k,m} \frac{(\lambda e^{-\epsilon\lambda})^k}{s^{k+1} e^{k s \epsilon}} \right\} (L) \\ &= \sum_{k=1}^m c_{k,m} (\lambda e^{-\epsilon\lambda})^k \mathcal{L}^{-1} \left\{ \frac{1}{s^{k+1} e^{k s \epsilon}} \right\} (L) \\ &= \sum_{k=1}^m \frac{c_{k,m}}{k!} (L - k\epsilon)^k (\lambda e^{-\epsilon\lambda})^k \mathbf{1}_{\{L > k\epsilon\}}. \end{aligned} \quad (3.28)$$

According to Lemma 3.4, when $\epsilon \rightarrow 0$, we obtain

$$M_{\beta_0}^m(L) = \mathbb{E}[N_L^m] = \sum_{k=1}^m \frac{c_{k,m}}{k!} (L\lambda)^k \mathbf{1}_{\{L > 0\}}. \quad (3.29)$$

Hence, for any $\lambda > 0$,

$$\sum_{k=1}^m \frac{c_{k,m}}{k!} (L\lambda)^k \mathbf{1}_{\{L > 0\}} = \sum_{k=1}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} (L\lambda)^k \mathbf{1}_{\{L > 0\}}, \quad (3.30)$$

which shows that

$$c_{k,m} = \left\{ \begin{matrix} m \\ k \end{matrix} \right\} k!. \quad (3.31)$$

Thus, we have proved (3.23) for any positive integer m . $\square$

Theorem 3.7. For any n, L, λ , and ϵ , we have

$$\Pr(\beta_0(L) = n) = \frac{1}{n!} \sum_{i=0}^{\lfloor L/\epsilon \rfloor - n} \frac{(-1)^i}{i!} \left((L - (n+i)\epsilon) \lambda e^{-\lambda\epsilon} \right)^{n+i}. \quad (3.32)$$

Proof. Since $\beta_0(L) \leq N_L$ and since $\mathbb{E}[e^{sN_L}]$ is finite for any $s \in \mathbf{R}$, we have, for any $s \geq 0$,

$$\mathbb{E}[e^{-s\beta_0(L)}] = \sum_{k=0}^{\infty} (-1)^k \frac{s^k}{k!} \mathbb{E}[\beta_0^k(L)]. \quad (3.33)$$

Rearranging the terms of the right-hand side, and substituting $M_{\beta_0}^m(L)$, by the result of (3.23), we obtain

$$\mathbb{E}\left[e^{-s\beta_0(L)}\right] = \sum_{k=0}^{\infty} \left((L - k\epsilon)^k (\lambda e^{-\lambda\epsilon})^k \mathbf{1}_{\{L > k\epsilon\}} \sum_{j=k}^{\infty} \frac{(-s)^j}{j!} \left\{ \begin{matrix} j \\ k \end{matrix} \right\} \right). \quad (3.34)$$

Furthermore, it is known (see [17]) that

$$\sum_{j=k}^{\infty} \frac{x^j}{j!} \left\{ \begin{matrix} j \\ k \end{matrix} \right\} = \frac{1}{k!} (e^x - 1)^k. \quad (3.35)$$

Hence,

$$\mathbb{E}\left[e^{-s\beta_0(L)}\right] = \sum_{k=0}^{\infty} (L - k\epsilon)^k (\lambda e^{-\lambda\epsilon})^k \mathbf{1}_{\{L > k\epsilon\}} \frac{(e^{-s} - 1)^k}{k!}. \quad (3.36)$$

By inverting the Laplace transforms, we get

$$\sum_{k=0}^{\infty} \sum_{i=k}^{\infty} \frac{(-1)^i}{i!} \binom{i}{n} \delta_{(k-n)} (k\epsilon - L)^k (\lambda e^{-\lambda\epsilon})^k \mathbf{1}_{\{L > k\epsilon\}}, \quad (3.37)$$

where δ_a is the Dirac measure at point a . After some simple algebra, we find the expression of the probability that an interval contains n complete clusters:

$$\Pr(\beta_0(L) = n) = p_n(L) = \frac{1}{n!} \sum_{i=0}^{\lfloor L/\epsilon \rfloor - n} \frac{(-1)^i}{i!} \left([L - (n+i)\epsilon] \lambda e^{-\lambda\epsilon} \right)^{n+i}, \quad (3.38)$$

concluding the proof. $\square$

Lemma 3.8. For $x \geq 0$, $p_n(x)$ has the three following properties.

- (i) $p_n(x)$ is differentiable.
- (ii) $\lim_{x \rightarrow \infty} p_n(x) = 0$.
- (iii) $\lim_{x \rightarrow \infty} dp_n(x)/dx = 0$.

Proof. Let j be a nonnegative integer. The function is obviously differentiable when $x/\epsilon \neq j$. Besides, we have

$$\lim_{x \rightarrow \epsilon j^+} p_n(x) - \lim_{x \rightarrow \epsilon j^-} p_n(x) = \lim_{x \rightarrow \epsilon j^+} \frac{(-1)^j}{j!} \left((x - (n+j)\epsilon) \frac{1}{a} \right)^{n+j}. \quad (3.39)$$

Since the right-hand term function of x is zero as well as its derivative for all j , the function is also derivable when $x/\epsilon = j$, which proves (i). Items (ii) and (iii) are direct consequences of the Final Value theorem in the Laplace transform of p_n and its derivative. $\square$

The expression of p_n gives us a Laplace pair between the x and s domains:

$$\frac{\mathbf{1}_{\{x \geq 0\}}}{n!} \sum_{i=0}^{\lfloor x/\epsilon \rfloor - n} \frac{(-1)^i}{i!} \left((x - (n+i)\epsilon) \frac{1}{a} \right)^{n+i} \xLeftrightarrow{\mathcal{L}} \frac{ae^{\epsilon s}}{(ase^{\epsilon s} + 1)^{n+1}}. \quad (3.40)$$

We can use this relation to find the distributions of B_i and U_n .

Theorem 3.9. *The probability density functions of B_i and U_n , denoted, respectively, by $f_{B_i}(x)$ and $f_{U_n}(x)$, are given:*

$$f_{B_i}(x) = \left[\lambda e^{-\epsilon \lambda} p_0(x - \epsilon) + e^{-\epsilon \lambda} \frac{d}{dx} p_0(x - \epsilon) \right] \mathbf{1}_{\{x > \epsilon\}}, \quad (3.41)$$

$$f_{U_n}(x) = \lambda e^{-\epsilon \lambda} p_{n-1}(x - \epsilon) \mathbf{1}_{\{x > \epsilon\}}, \quad (3.42)$$

where the expressions of $p_0(x - \epsilon)$ and $(d/dx)p_0(x - \epsilon)$ are straightforwardly obtained from (3.32).

Proof. According to Theorem 3.1,

$$\begin{aligned} \mathbb{E}[e^{-sB_i}] &= \frac{1}{\lambda} \frac{(\lambda + s)}{(e^{\epsilon \lambda} / \lambda) s e^{\epsilon s} + 1} \\ &= \lambda e^{-\epsilon \lambda} \frac{e^{\epsilon \lambda}}{\lambda} \frac{e^{\epsilon s}}{(e^{\epsilon \lambda} / \lambda) s e^{\epsilon s} + 1} e^{-\epsilon s} + e^{-\epsilon \lambda} s \frac{e^{\epsilon \lambda}}{\lambda} \frac{e^{\epsilon s}}{(e^{\epsilon \lambda} / \lambda) s e^{\epsilon s} + 1} e^{-\epsilon s} \\ &= \lambda e^{-\epsilon \lambda} e^{-\epsilon s} \mathcal{L}\{p_0(\cdot)\}(s) + e^{-\epsilon \lambda} e^{-\epsilon s} s \mathcal{L}\{p_0(\cdot)\}(s). \end{aligned} \quad (3.43)$$

Here, using the inverse Laplace transform established in (3.40) and remembering that $p_0(x^-) = 0$, we get an analytical expression for $f_{B_i}(x)$, proving (3.41).

Proceeding in a similar fashion, we can find the distribution of U_n by inverting its Laplace transform given by Corollary 3.2 as follows:

$$\begin{aligned} \mathbb{E}[e^{-sU_n}] &= \frac{1}{((e^{\epsilon \lambda} / \lambda) s e^{\epsilon s} + 1)^n} \\ &= \lambda e^{-\epsilon \lambda} \frac{e^{\epsilon \lambda}}{\lambda} \frac{e^{\epsilon s}}{((e^{\epsilon \lambda} / \lambda) s e^{\epsilon s} + 1)^n} e^{-\epsilon s} \\ &= \lambda e^{-\epsilon \lambda} e^{-\epsilon s} \mathcal{L}\{p_{n-1}(\cdot)\}(s). \end{aligned} \quad (3.44)$$

We, thus, have (3.42). □

We can also obtain the probability that the segment $[0, L]$ is completely covered by the sensors. To do this, we remember that the first point (if there is one) is capable to cover the interval $[X_1 - \epsilon, X_1 + \epsilon]$.

Theorem 3.10. Let $R_{m,n}(x)$ be defined as follows:

$$R_{m,n}(x) = \sum_{i=m}^{\lfloor x/\epsilon \rfloor - 1} \left[\left(e^{-\lambda\epsilon} \right)^{i+n} \sum_{j=0}^{i+n} \frac{(\lambda[(1-i)\epsilon - x])^j}{j!} \right]. \quad (3.45)$$

Then,

$$\Pr([0, L] \text{ is covered}) = R_{0,1}(L) - e^{-\lambda\epsilon} R_{0,1}(L - \epsilon) - e^{-\lambda\epsilon} R_{1,0}(L) + e^{-2\lambda\epsilon} R_{1,0}(L - \epsilon). \quad (3.46)$$

Proof. The condition of total coverage is the same as

$$\{\forall x \in [0, L], \exists X_i \in [0, L] \mid x \in [X_i - \epsilon, X_i + \epsilon]\}, \quad (3.47)$$

which means that

$$\{[0, L] \text{ is covered}\} \iff \{B_1 \geq L - X_1\} \cap \{X_1 \leq \epsilon\}. \quad (3.48)$$

Hence,

$$\Pr([0, L] \text{ is covered}) = \int_0^\epsilon \Pr(B_1 \geq L - X_1 \mid X_1 = x) dP_{X_1}(x), \quad (3.49)$$

and since B_1 and X_1 are independent

$$\Pr([0, L] \text{ is covered}) = \int_0^\epsilon \int_{L-x}^\infty f_{B_1}(u) \lambda e^{-x\lambda} du dx. \quad (3.50)$$

The result then follows from Lemma 3.8 and some tedious but straightforward algebra. $\square$

4. Other Scenarios

The method can be used to calculate p_n for other definitions of the number of clusters. We consider two other definitions: the number of incomplete clusters and the number of clusters in a circle.

4.1. Number of Incomplete Clusters

The major difference with Section 3 is that a cluster is now taken into account as soon as one of the point of the cluster is inside the interval $[0, L]$. So, for instance, in Figure 3, we count actually $n + 1$ incomplete clusters. We define $\beta'_0(L)$ as the number of incomplete clusters on an interval $[0, L]$.

Theorem 4.1. Let $G(k)$ be defined as

$$G(k) = (-1)^k \left(e^{-k\lambda\epsilon} \sum_{j=0}^k \frac{[\lambda(k\epsilon - L)]^j}{j!} - e^{-\lambda L} \right) \mathbf{1}_{\{T > k\epsilon\}}, \quad (4.1)$$

for $k \in \mathbb{N}_+$ and $G(-1) = e^{-\lambda L}$. Then,

$$\Pr(\beta'_0(L) = n) = \sum_{i=n}^{\lfloor L/\epsilon \rfloor + 1} (-1)^{i+n} \binom{i}{n} (G(i-1) + G(i)), \quad \text{for } n \geq 0. \quad (4.2)$$

Proof. The condition of $\beta'_0(L) \geq n$ is now given by

$$\{\beta'_0 \geq n\} \iff \begin{cases} \{\Delta X_0 + U_{n-1} \leq L\} & \text{if } n \geq 1, \\ \{\Delta X_0 < \infty\} & \text{if } n = 0. \end{cases} \quad (4.3)$$

We define Y_n as

$$Y_n = \begin{cases} \Delta X_0 + U_{n-1} & \text{if } n \geq 1, \\ 0 & \text{if } n = 0. \end{cases} \quad (4.4)$$

Repeating the same calculations, we find the Laplace transform of $\Pr(\beta'_0(\cdot) = n)$:

$$\mathcal{L}\{\Pr(\beta'_0(\cdot) = n)\}(s) = \begin{cases} \frac{\lambda}{s + \lambda} \frac{e^{\epsilon\lambda}}{\lambda} \frac{e^{\epsilon s}}{((e^{\epsilon\lambda}/\lambda)se^{\epsilon s} + 1)^n} & \text{if } n \geq 1, \\ \frac{1}{\lambda + s} & \text{if } n = 0. \end{cases} \quad (4.5)$$

With this expression, following the lines of Lemma 3.4, we obtain

$$\mathcal{L}\{\mathbb{E}[\beta'_0(\cdot)^m]\}(s) = \sum_{k=1}^{m+1} \binom{m+1}{k} (k-1)! \frac{1}{s^k} \frac{\lambda}{\lambda + s} \left(\frac{\lambda e^{-\lambda\epsilon}}{e^{s\epsilon}} \right)^{k-1}. \quad (4.6)$$

Then, we write

$$\frac{\lambda}{\lambda + s} \frac{1}{s^k} = \frac{(-1)^k}{\lambda^{k-1}} \frac{1}{\lambda + s} + \sum_{i=1}^k \frac{1}{s^i} \left(\frac{-1}{\lambda} \right)^{k-i}, \quad (4.7)$$

to find an expression with a well-known Laplace transform inverse, and, after inverting it, we obtain

$$\mathbb{E}[\beta'_0{}^m] = \sum_{k=0}^m \binom{m+1}{k+1} k! G(k). \quad (4.8)$$

Expanding the Laplace transform of the distribution of $\beta'_0(L)$ in a Taylor series and rearranging terms, we get

$$\mathbb{E}\left[e^{-s\beta'_0(L)}\right] = 1 + G(0) \sum_{j=1}^{\infty} \frac{(-s)^j}{j!} \left\{ \begin{matrix} j \\ 1 \end{matrix} \right\} + \left(\sum_{k=1}^{\infty} G(k) \sum_{j=k}^{\infty} \frac{(-s)^j}{j!} \left\{ \begin{matrix} j+1 \\ k+1 \end{matrix} \right\} \right). \quad (4.9)$$

Now, we use another recurrence that the Stirling numbers obey [17],

$$\left\{ \begin{matrix} j+1 \\ k+1 \end{matrix} \right\} = \left\{ \begin{matrix} j \\ k \end{matrix} \right\} + (k+1) \left\{ \begin{matrix} j \\ k+1 \end{matrix} \right\}, \quad (4.10)$$

to get

$$\begin{aligned} \sum_{j=k}^{\infty} \frac{x^j}{j!} \left\{ \begin{matrix} j+1 \\ k+1 \end{matrix} \right\} &= \sum_{j=k}^{\infty} \frac{x^j}{j!} \left(\left\{ \begin{matrix} j \\ k \end{matrix} \right\} + (k+1) \left\{ \begin{matrix} j \\ k+1 \end{matrix} \right\} \right) \\ &= \frac{1}{k!} (e^x - 1)^k + \frac{1}{k!} (e^x - 1)^{k+1}. \end{aligned} \quad (4.11)$$

Hence,

$$\mathbb{E}\left[e^{-s\beta'_0(L)}\right] = 1 + \sum_{k=1}^{\infty} (G(k-1) + G(k)) (e^{-s} - 1)^k. \quad (4.12)$$

Inverting this expression for any nonnegative integer n , we have the searched distribution. $\square$

4.2. Number of Clusters in a Circle

We investigate now the case where the points of the process are deployed over a circumference, and we want to count the number of complete clusters, which corresponds to calculate the Euler's Characteristic of the total coverage, so we call this quantity χ . Without loss of generality, we can choose an arbitrary point to be the origin.

Theorem 4.2. *The distribution of the Euler's Characteristic, $\chi(L)$, when the points are deployed over a circumference of length L is given by*

$$\begin{aligned} \Pr(\chi(L) = n) &= e^{-\lambda L} \mathbf{1}_{\{n=0\}} + \left(1 - e^{-\lambda L}\right) \frac{\lambda e^{-\epsilon \lambda}}{n!} \\ &\times \sum_{i=0}^{\lfloor L/\epsilon \rfloor - n} \left[\frac{(-1)^i}{i!} \left([L - (n+i)\epsilon] \lambda e^{-\epsilon \lambda} \right)^{n+i-1} \left(L + (n+i) \left(\frac{1}{\lambda} - \epsilon \right) \right) \right] \quad \text{for } n \geq 0. \end{aligned} \quad (4.13)$$

Proof. If there is no points on the circle, $\chi(L) = 0$. Otherwise, if there is at least one point, we choose the origin at this point, and we have equivalence between the events:

$$\{\chi(L) \geq n\} \iff \begin{cases} \{U_{n-1} + B_n \leq L\} \cap \{N_L > 0\} & \text{if } n \geq 1, \\ \{\Delta X_0 < \infty\} & \text{if } n = 0. \end{cases} \quad (4.14)$$

In Figure 4, we present an example of this equivalence.

We can define Y_n as

$$Y_n = \begin{cases} U_{n-1} + B_n & \text{if } n \geq 1, \\ 0 & \text{if } n = 0, \end{cases} \quad (4.15)$$

to find the Laplace transform or $\Pr(\chi(L) = n)$:

$$\mathcal{L}\{\Pr(\chi(\cdot) = n)\}(s) = (1 - e^{-\lambda L}) \frac{\lambda + s}{\lambda} \frac{e^{\epsilon s}}{((e^{\epsilon \lambda}/\lambda) s e^{\epsilon s} + 1)^n}. \quad (4.16)$$

The number of clusters is almost surely equal to the number of points when $\epsilon \rightarrow 0$, so

$$\mathbb{E}[\chi(L)^m] = (1 - e^{-\lambda L}) \lambda e^{-\epsilon \lambda} \sum_{k=1}^m \left[\left\{ \begin{matrix} m \\ k \end{matrix} \right\} ([L - k\epsilon] \lambda e^{-\epsilon \lambda})^{k-1} \left(L + k \left(\frac{1}{\lambda} - \epsilon \right) \right) \mathbf{1}_{\{L > k\epsilon\}} \right]. \quad (4.17)$$

Expanding the Laplace transform in a Taylor series and rearranging terms, as we did previously, yields

$$\mathbb{E}[e^{-s\chi(L)}] = (1 - e^{-\lambda L}) \lambda e^{-\epsilon \lambda} \sum_{k=0}^{\infty} \left[([L - k\epsilon] \lambda e^{-\epsilon \lambda})^{k-1} \left(L + k \left(\frac{1}{\lambda} - \epsilon \right) \right) \mathbf{1}_{\{L > k\epsilon\}} \sum_{j=k}^{\infty} \frac{(-s)^j}{j!} \left\{ \begin{matrix} j \\ k \end{matrix} \right\} \right]. \quad (4.18)$$

Since

$$\sum_{j=k}^{\infty} \frac{(-s)^j}{j!} \left\{ \begin{matrix} j \\ k \end{matrix} \right\} = \frac{(e^{-s} - 1)^k}{k!}, \quad (4.19)$$

we can directly invert this Laplace transform, add the case where there are no points for $\chi(L) = 0$, and the theorem is proved. $\square$

5. Examples

We consider some examples to illustrate the results of the paper. Here, the behavior of the mean and the variance of $\beta_0(L)$ as well as $\Pr(\beta_0(L) = n)$ are presented.

Figure 4: Illustration of the condition equivalent to $\chi(L) \geq n$. Since the coverage of the last point on $[0, L]$ overlaps the cluster with a point in zero, they are actually contained in the same cluster.

From (3.23), we have that $\mathbb{E}[\beta_0(L)]$ is given by

$$\mathbb{E}[\beta_0(L)] = (L - \epsilon)\lambda e^{-\epsilon\lambda} \mathbf{1}_{\{L > \epsilon\}}. \quad (5.1)$$

This expression agrees with the intuition in that there are three typical regions given a fixed ϵ . When λ is much smaller than $1/\epsilon$, the number of clusters is approximatively the number of sensors, since the connections with few sensors will unlikely happen, which can be seen from the fact that $\mathbb{E}[\beta_0(L)] \rightarrow L\lambda$ when $\lambda \rightarrow 0$. As we increase λ , the mean number of direct connections overcomes the mean number of sensors, and, at some value of λ , we expect that $\mathbb{E}[\beta_0(L)]$ decreases, when adding a point is likely to connect disconnected clusters. We remark that the maximum occurs exactly for $\epsilon = 1/\lambda$, that is, when the mean distance between two sensors equals the threshold distance for them to be connected. At this maximum, $\mathbb{E}[\beta_0(L)]$ takes the value of $(L/\epsilon - 1)e^{-1}$. Finally, when λ is too large, all sensors tend to be connected,

Figure 5: Variation of the mean number of clusters in function of λ when $L = 4$ and $\epsilon = 1$.

and there is only one cluster which even goes beyond L , so there are no complete clusters into the interval $[0, L]$. This is trivial when we make $\lambda \rightarrow \infty$ in the last equation. Figure 5 shows this behavior when $L = 4$ and $\epsilon = 1$.

The variance can be obtained also by (3.23) as follows:

$$\text{Var}(\beta_0(L)) = (L - \epsilon)\lambda e^{-\epsilon\lambda} \mathbf{1}_{\{L > \epsilon\}} + (L - 2\epsilon)\lambda^2 e^{-2\epsilon\lambda} \mathbf{1}_{\{L > 2\epsilon\}} - (L - \epsilon)^2 \lambda^2 e^{-2\epsilon\lambda} \mathbf{1}_{\{L > \epsilon\}}, \quad (5.2)$$

and, under the condition that $L > 2\epsilon$

$$\text{Var}(\beta_0(L)) = (L - \epsilon)\lambda e^{-\epsilon\lambda} + \epsilon(3\epsilon - 2L)\lambda^2 e^{-2\epsilon\lambda}. \quad (5.3)$$

Figure 6 shows a plot of $\text{Var}(\beta_0(L))$ in function of λ for $L = 4$ and $\epsilon = 1$. We can expect that, when λ is small compared to ϵ , the plot should be approximatively linear, since there would not be too much connections in the network, and the variance of the number of clusters should be close to the variance of the number of sensors given by λL . Since $\beta_0(L)$ tends almost surely to 0 when λ goes to infinity, $\text{Var}(\beta_0(L))$ should also tend to 0 in this case. Those two properties are observed in the plot. Besides, we find the critical points of this function, and again, $\lambda = 1/\epsilon$ is one of them, and at this value $\text{Var}(\beta_0(L)) = (L/\epsilon)e^{-1} + (3 - 2L/\epsilon)e^{-1}$. The other two are the ones satisfying the transcendent equation:

$$\lambda e^{-\lambda\epsilon} = \frac{L - \epsilon}{2\epsilon(2L - 3\epsilon)}. \quad (5.4)$$

By using the second derivative, we realize that $1/\epsilon$ is actually a minimum. Besides, if $L \leq 2\epsilon$, there is just one critical point, a maximum, at $\lambda = 1/\epsilon$.

Figure 6: Behavior of the variance of the number of clusters in function of λ when $L = 4$ and $\epsilon = 1$.

Figure 7: Probabilities of connectiveness, $\Pr(\beta_0(L) = n)$, for $n = 0, 1, 2, 3$, in function of λ when $L = 4$ and $\epsilon = 1$.

The last example in the section is performed with the result obtained in Theorem 3.7. We consider again $L = 4$ and $\epsilon = 1$ to obtain the following distributions:

$$\begin{aligned}
 \Pr(\beta_0(L) = 0) &= 1 - 3\lambda e^{-\lambda} + 2\lambda^2 e^{-2\lambda} - 1/6\lambda^3 e^{-3\lambda}, \\
 \Pr(\beta_0(L) = 1) &= 3\lambda e^{-\lambda} - 4\lambda^2 e^{-2\lambda} + 1/2\lambda^3 e^{-3\lambda}, \\
 \Pr(\beta_0(L) = 2) &= 2\lambda^2 e^{-2\lambda} - 1/2\lambda^3 e^{-3\lambda}, \\
 \Pr(\beta_0(L) = 3) &= 1/6\lambda^3 e^{-3\lambda}, \\
 \Pr(\beta_0(L) > 3) &= 0.
 \end{aligned} \tag{5.5}$$

Those expressions are simple, and they have at most four terms, since $L = 4\epsilon$. We plot these functions in Figure 7. The critical points on those plots at $\lambda = 1/\epsilon$ are confirmed for the fact that, in function of λ for every n , $\Pr(\chi(L) = n)$ can be represented as a sum:

$$\sum_{i=0}^j q_{i,j} (\lambda e^{-\lambda\epsilon})^i, \quad (5.6)$$

where the coefficients $q_{i,j}$ are constant in relation to λ . However, $(\lambda e^{-\lambda\epsilon})^i$ has a critical point at $\lambda = 1/\epsilon$ for all $i > 0$, so this should be also a critical point of $\Pr(\chi(L) = n)$. If λ is small, we should expect that $\Pr(\chi(L) = 0)$ is close to one, since it is likely to N have no points. For this reason, in this region, $\Pr(\chi(L) = n)$ for $n > 0$ is small. When λ is large, we expect to have very large clusters, likely to be larger than L , so it is unlikely to have a complete cluster in the interval, and, again, $\Pr(\chi(L) = 0)$ approaches to the unity, while $\Pr(\chi(L) = n)$ for $n > 0$ become again small.

Acknowledgment

The authors would like to thank the anonymous referee whose constructive remarks helped us to improve the presentation of this paper.

References

- [1] J. M. Kahn, R. H. Katz, and K. S. J. Pister, "Mobile networking for smart dust," in *Proceedings of the International Conference on Mobile Computing and Networking (MOBICOM '99)*, pp. 271–278, Seattle, Wash, USA, August 1999.
- [2] F. L. Lewis, *Wireless Sensor Networks*, chapter 2, John Wiley, New York, NY, USA, 2004.
- [3] G. J. Pottie and W. J. Kaiser, "Wireless integrated network sensors," *Communications of the ACM*, vol. 43, no. 5, pp. 51–58, 2000.
- [4] C. Y. Chong and S. P. Kumar, "Sensor networks: evolution, opportunities, and challenges," *Proceedings of the IEEE*, vol. 91, no. 8, pp. 1247–1256, 2003.
- [5] R. Ghrist and A. Muhammad, "Coverage and hole-detection in sensor networks via homology," in *Proceedings of the 4th International Symposium on Information Processing in Sensor Networks (IPSN '05)*, pp. 254–260, April 2005.
- [6] V. de Silva and R. Ghrist, "Coordinate-free coverage in sensor networks with controlled boundaries via homology," *International Journal of Robotics Research*, vol. 25, no. 12, pp. 1205–1222, 2006.
- [7] A. F. Siegel and L. Holst, "Covering the circle with random arcs of random sizes," *Journal of Applied Probability*, vol. 19, no. 2, pp. 373–381, 1982.
- [8] L. Holst, "On multiple covering of a circle with random arcs," *Journal of Applied Probability*, vol. 17, no. 1, pp. 284–290, 1980.
- [9] S. Kumar, T. H. Lai, and A. Arora, "Barrier coverage with wireless sensors," in *Proceedings of the 11th Annual International Conference on Mobile Computing and Networking (MOBICOM '05)*, pp. 284–298, Cologne, Germany, September 2005.
- [10] P. Manohar, S. S. Ram, and D. Manjunath, "Path coverage by a sensor field: the nonhomogeneous case," *ACM Transactions on Sensor Networks*, vol. 5, no. 2, pp. 17:1–17:26, 2009.
- [11] M. Kahle, "Random geometric complexes," *Discrete & Computational Geometry*, vol. 45, no. 3, pp. 553–573, 2011.
- [12] M. Kahle and E. Meckes, "Limit theorems for Betti numbers of random simplicial complexes," *Probability*. Preprint, <http://arxiv.org/abs/1009.4130>.
- [13] S. Asmussen, *Applied Probability and Queues*, vol. 51 of *Stochastic Modelling and Applied Probability*, Springer, New York, NY, USA, 2nd edition, 2003.

- [14] R. B. Cooper, *Introduction to Queueing Theory*, North-Holland, New York, NY, USA, 2nd edition, 1981.
- [15] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics: A Foundation for Computer Scienc*, chapter 6.1, Addison-Wesley, Reading, Mass, USA, 2nd edition, 1994.
- [16] D. C. Wood, "Technical report 15-92," Tech. Rep., University of Kent computing Laboratory, University of Kent, Canterbury, UK, 1992.
- [17] S. Roman, *The Umbral Calculus*, Pure and Applied Mathematics, Academic Press, New York, NY, USA, 1984.