Research Article

Interval Estimations of the Two-Parameter Exponential Distribution

Lai Jiang and Augustine C. M. Wong

*Department of Mathematics and Statistics, York University, 4700 Keele Street,
Toronto, ON, Canada M3J 1P3*

Correspondence should be addressed to Augustine C. M. Wong, august@yorku.ca

Received 11 November 2011; Revised 24 January 2012; Accepted 31 January 2012

Academic Editor: A. Thavaneswaran

Copyright © 2012 L. Jiang and A. C. M. Wong. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In applied work, the two-parameter exponential distribution gives useful representations of many physical situations. Confidence interval for the scale parameter and predictive interval for a future independent observation have been studied by many, including Petropoulos (2011) and Lawless (1977), respectively. However, interval estimates for the threshold parameter have not been widely examined in statistical literature. The aim of this paper is to, first, obtain the exact significance function of the scale parameter by renormalizing the p^* -formula. Then the approximate Studentization method is applied to obtain the significance function of the threshold parameter. Finally, a predictive density function of the two-parameter exponential distribution is derived. A real-life data set is used to show the implementation of the method. Simulation studies are then carried out to illustrate the accuracy of the proposed methods.

1. Introduction

The two-parameter exponential distribution with density:

$$f(x; \mu, \sigma) = \frac{1}{\sigma} \exp\left\{-\frac{x - \mu}{\sigma}\right\}, \quad (1.1)$$

where $\mu < x$ is the threshold parameter, and $\sigma > 0$ is the scale parameter, is widely used in applied statistics. For example, Lawless [1] applied the two-parameter exponential distribution to analyze lifetime data, and Baten and Kamil [2] applied the distribution to analyze inventory management systems with hazardous items. Petropoulos [3] proposed two new classes of confidence interval for the scale parameter σ . Lawless [1] obtained a prediction interval for a future observation from the two-parameter exponential distribution.

In this paper, we consider a sample $(x_1, \dots, x_n)$ from the two-parameter exponential distribution with density (1.1). In Section 2, we show that by renormalizing the p^* -formula, the exact significance function of the scale parameter σ can be obtained. In Section 3, the approximate Studentization method, based on the significance function of σ obtained in Section 2, is applied to obtain the significance function of the threshold parameter μ . In Section 4, we combine the results of the previous two sections and derive a predictive density for a future observation from the two-parameter exponential distribution. Some concluding remarks are given in Section 5. Throughout this paper, a real-life data set is used to show the implementation of the proposed methods, and simulation results are presented to illustrate the accuracy of the proposed methods.

2. Confidence Interval for the Scale Parameter

For the two-parameter exponential distribution with density (1.1), it can be shown that the marginal density of $X_{(1)} = \min(X_1, \dots, X_n)$ is

$$f_m(x_{(1)}; \mu, \sigma) = \frac{n}{\sigma} \exp\left\{-\frac{n}{\sigma}(x_{(1)} - \mu)\right\} \quad x_{(1)} > \mu. \quad (2.1)$$

With an observed sample $(x_1, \dots, x_n)$, the log conditional likelihood function that depends only on σ , can be written as

$$\begin{aligned} \ell_c(\sigma) &= \ell(\mu, \sigma) - \ell_m(\mu, \sigma) \\ &= a - (n-1) \log \sigma - \frac{n}{\sigma}(\bar{x} - x_{(1)}), \end{aligned} \quad (2.2)$$

where a is an additive constant and $\bar{x} = \sum_{i=1}^n x_i$. Without loss of generality, in this paper, a is set as 0. Moreover, denote $s = (\bar{x} - x_{(1)})$ and $\psi = 1/\sigma$, then the log conditional likelihood function can be rewritten as

$$\ell_c(\psi) = (n-1) \log \psi - ns\psi. \quad (2.3)$$

Note that (2.3) has the same form as a log likelihood function of an exponential family model with canonical parameter ψ . The maximum likelihood estimate of ψ , $\hat{\psi} = (n-1)/ns$ is obtained by solving $(\partial \ell_c(\psi))/\partial \psi|_{\psi=\hat{\psi}} = 0$. Furthermore, an estimate of the variance of $\hat{\psi}$ is $\hat{j}^{-1}$, where $\hat{j} = -(\partial^2 \ell_c(\psi))/\partial \psi^2|_{\psi=\hat{\psi}} = (n^2 s^2)/(n-1)$ is the observed information.

Under regularity conditions as stated in DiCiccio et al. [4], both the standardized maximum likelihood estimate

$$q = q(\psi) = (\hat{\psi} - \psi)\hat{j}^{1/2}, \quad (2.4)$$

and the signed log likelihood ratio statistic:

$$r = r(\psi) = \text{sign}(\hat{\psi} - \psi) \sqrt{2[\ell_c(\hat{\psi}) - \ell_c(\psi)]} \quad (2.5)$$

have limiting standard normal distribution with rate of convergence $O(n^{-1/2})$. Hence the significance function of σ can be approximated by

$$p(\sigma) = \Phi\left(q\left(\frac{1}{\psi}\right)\right), \quad (2.6)$$

or

$$p(\sigma) = \Phi\left(r\left(\frac{1}{\psi}\right)\right), \quad (2.7)$$

where $\Phi(\cdot)$ is the cumulative distribution function of the standard normal distribution.

Since the conditional log likelihood function given in (2.3) in exponential family form with ψ being the canonical parameter, the modified signed log likelihood statistic by Barndorff-Nielsen [5, 6] can be simplified into

$$r^* = r^*(\psi) = r - \frac{1}{r} \log \frac{r}{q}, \quad (2.8)$$

where q and r are defined in (2.4) and (2.5) respectively, and r^* has limiting standard normal distribution with rate of convergence $O(n^{-3/2})$. Hence the significance function of σ can be approximated by

$$p(\sigma) = \Phi\left(r^*\left(\frac{1}{\psi}\right)\right). \quad (2.9)$$

Barndorff-Nielsen [7] derived the p^* -formula—an approximate density for the maximum likelihood estimator. In this case, we have

$$p^*(\hat{\sigma}; \sigma) = k \hat{j}^{1/2} \exp\{\ell_c(\sigma) - \ell_c(\hat{\sigma})\} = k \frac{\hat{\sigma}^{n-2}}{\sigma^{n-1}} \exp\left\{-\frac{(n-1)\hat{\sigma}}{\sigma}\right\}, \quad (2.10)$$

where k is the renormalizing constant. Since $\hat{\sigma} = ns/(n-1) = 1/\hat{\psi}$, by change of variable and renormalization, we have

$$p^*(s; \sigma) = \Gamma^{-1}(n-1) \left(\frac{ns}{\sigma}\right)^{n-1} \frac{1}{s} \exp\left\{-\frac{ns}{\sigma}\right\}, \quad (2.11)$$

which is the exact density of $S = \bar{X} - X_{(1)}$, and it is free of μ . Therefore, inference concerning σ can be based on the distribution of S . Thus, the exact confidence interval for σ can be obtained using $p^*(s; \sigma)$ and is

$$p(\sigma) = \int_0^s p^*(t; \sigma) dt. \quad (2.12)$$

Grubbs [8] reported the following data set (see Table 1).

Table 1

162	200	271	302	393	508	539	629	706	777
884	1008	1101	1182	1463	1603	1984	2355	2880	

We have

$$n = 19, \quad x_{(1)} = 162, \quad s = 835.21. \quad (2.13)$$

Table 2 recorded the 90%, 95%, and 99% confidence interval for σ . The proposed method (r^*) and the exact method (exact) give approximately the same confidence intervals, whereas the results obtained by the standardized maximum likelihood estimate method (mle) and the signed log likelihood ratio method (r) are quite different. Moreover, from Table 2, it is clear that r gives the narrowest confidence intervals and mle gives the widest confidence intervals. The significance functions of σ obtained from the four methods are plotted in Figure 1.

In order to compare the accuracies of the four methods, Monte Carlo simulation studies with 10,000 replicates are performed. For each simulation study, we generate sample of size n from the two-parameter exponential distribution with scale parameter σ and threshold parameter μ . Then for each sample, the 95% confidence intervals for σ is calculated from the four methods discussed in this section.

Table 3 recorded the results of the simulation studies for some combinations of n , μ , and σ . The “Lower Error” is the proportion of the true σ that falls outside the lower limit of the 95% confidence intervals while the “Upper Error” is the proportion of the true σ that falls outside the upper limit of the 95% confidence intervals, and $(1 - \text{Lower Error} - \text{Upper Error})$ is recorded as “Central Coverage”. The nominal values for the “Lower Error”, “Upper Error”, and “Central Coverage” are 0.025, 0.025, and 0.95, respectively. Moreover, for 10,000 Monte Carlo simulations, the standard errors for the “Lower Error” and the “Upper Error” are the same and are $\sqrt{0.025(1 - 0.025)/10000} = 0.0016$. Similarly, the corresponding standard error for the “Central Coverage” is 0.0022.

From Table 3, even for the smallest possible sample size ($n = 2$), the proposed method and the exact method give almost identical results and have excellent coverage properties. The results obtained by the other two methods are not as satisfactory especially when the sample size is small.

Since mle method gives the poorest coverage, we excluded it from further investigation. Table 4 recorded the average width of the confidence interval for σ for the simulation study with $(n, \sigma, \mu) = (10, 0.5, 5)$. We can note that even though r has the shortest average width of the confidence interval for σ , it also has the poorest coverage properties as demonstrated in Table 3. For inference purpose, coverage properties are more important than width of the confidence interval. Hence the proposed method is recommended for this problem.

3. Confidence Interval for the Threshold Parameter

For the two-parameter exponential distribution, Petropoulos [3] showed that $(X_{(1)}, S)$ is a sufficient statistic. Note that μ is completely contained in the marginal density of $X_{(1)}$, but it also depends σ . In Section 2, the significance function of σ , $p(\sigma)$, was obtained. Hence, we can apply the approximate Studentization method, which is discussed in Fraser and Wong

Figure 1: The significance function of σ .

Table 2: Confidence intervals for σ .

Method	90% CI	95% CI	99% CI
mle	(635.30, 1439.83)	(603.03, 1638.59)	(548.56, 2244.03)
r	(512.51, 1334.35)	(574.08, 1454.18)	(507.92, 1732.03)
r^*	(622.32, 1363.95)	(583.01, 1487.51)	(515.37, 1774.35)
exact	(622.33, 1363.98)	(583.02, 1487.54)	(515.38, 1774.39)

Table 3: Results from simulation studies for the parameter of interest σ .

n	σ	μ	Method	Lower error	Upper error	Central coverage
2	3	1	mle	0.0505	0.0000	0.9495
			r	0.0117	0.0526	0.9357
			r^*	0.0245	0.0227	0.9528
			exact	0.0249	0.0231	0.9520
5	1	2	mle	0.0447	0.0000	0.9553
			r	0.0175	0.0400	0.9425
			r^*	0.0248	0.0266	0.9486
			exact	0.0249	0.0266	0.9485
8	6	4	mle	0.0388	0.0021	0.9591
			r	0.0167	0.0339	0.9494
			r^*	0.0217	0.0256	0.9527
			exact	0.0217	0.0256	0.9527
10	0.5	5	mle	0.0415	0.0042	0.9543
			r	0.0220	0.0331	0.9449
			r^*	0.0274	0.0248	0.9478
			exact	0.0275	0.0248	0.9477

Table 4: Average width of the confidence interval for σ .

Method	Confidence level		
	90%	95%	99%
r	0.6180	0.7714	0.1.1372
r^*	0.6529	0.8160	1.2064
exact	0.6530	0.8161	1.2065

[9], to eliminate the dependence of σ from the marginal density of $X_{(1)}$. More specifically, the approximate Studentized marginal density of $X_{(1)}$ is

$$\begin{aligned}
 f^S(x_{(1)}; \mu) &= c \int_0^\infty f_m(x_{(1)}; \mu, \sigma) |dp(\sigma)| \\
 &= c \int_0^\infty \Gamma^{-1}(n-1) \frac{n^n s^{n-1}}{\sigma^{n+1}} \exp\left\{-\frac{n}{\sigma}(x_{(1)} - \mu + s)\right\} d\sigma,
 \end{aligned} \tag{3.1}$$

where $c = \int_\mu^\infty \int_0^\infty f_m(x_{(1)}; \mu, \sigma) |dp(\sigma)| dx_{(1)}$ is the normalizing constant. Hence, we have

$$f^S(x_{(1)}; \mu) = \frac{n-1}{s} \left(1 + \frac{x_{(1)} - \mu}{s}\right)^{-n} \quad x_{(1)} > \mu, \tag{3.2}$$

and the corresponding significance function of μ is

$$p(\mu) = \int_\mu^{x_{(1)}} f^S(v; \mu) dv = 1 - \left(1 + \frac{x_{(1)} - \mu}{s}\right)^{-(n-1)}. \tag{3.3}$$

Thus, the explicit $(1 - \alpha)100\%$ confidence interval for μ obtained by the approximate Studentization method is

$$\left(x_{(1)} - s \left[\left(\frac{\alpha}{2}\right)^{-1/(n-1)} - 1\right], x_{(1)} - s \left[\left(1 - \frac{\alpha}{2}\right)^{-1/(n-1)} - 1\right]\right). \tag{3.4}$$

Applying the approximate Studentization method to the Grubbs [8] data set, the 90%, 95%, and 99% confidence intervals for the threshold parameter μ are $(-27.97, 160.82)$, $(-68.22, 161.42)$, and $(-157.87, 161.88)$, respectively.

To illustrate the accuracy of the proposed method, we performed a Monte Carlo simulation study. Table 5 records the results from this study. The proposed approximate Studentization method gives extremely good coverage properties even when the sample size is extremely small.

4. Prediction Interval for a Future Observation

The density of a future observation from the two-parameter exponential distribution is given in (1.1), which depends on both parameters μ and σ . With observed data $(x_1, \dots, x_n)$, or equivalently, observed sufficient statistic (s, t) , $p(\sigma)$ was obtained by the method in Section 2,

Table 5: Results from simulation studies and the parameter of interest are μ using the approximate Studentization method.

n	σ	μ	Lower error	Upper error	Central coverage
2	3	1	0.0239	0.0258	0.9503
5	1	2	0.0277	0.0254	0.9469
8	6	4	0.0249	0.0241	0.9510
10	0.5	5	0.0251	0.0248	0.9501

Figure 2: The predictive cumulative distribution function, $F^P(x)$.

and it gives us as much information about σ as we can extract from the observed data in the absence of knowledge of μ . Moreover, $p(\mu)$ was obtained by the approximate Studentization method in Section 3 gives us as much information about μ as we can extract from the observed data after averaging out the effect of σ . Therefore, to eliminate μ and σ from the density of a future observation, we apply the approximate Studentization method to obtain a predictive density of X :

$$\begin{aligned}
 f^P(x) &= \int_{-\infty}^{x_{(1)}} \int_0^{\infty} f(x; \mu, \sigma) |dp(\sigma)| |dp(\mu)| \\
 &= \int_{-\infty}^{x_{(1)}} \left[\int_0^{\infty} \frac{1}{\sigma} \exp\left\{-\frac{x-\mu}{\sigma}\right\} \Gamma^{-1}(n-1) \frac{(ns)^{n-1}}{\sigma^n} \exp\left\{-\frac{ns}{\sigma}\right\} d\sigma \right] |dp(\mu)| \\
 &= (n-1)^2 n^{n-1} s^{2n-2} \int_{-\infty}^{x_{(1),1}} \left(\frac{1}{(x-\mu+ns)(x_{(1)}-\mu+s)} \right)^n d\mu.
 \end{aligned} \tag{4.1}$$

Hence, the corresponding predictive cumulative distribution function is

$$F^P(x) = \int_{-\infty}^x f^P(u) du. \tag{4.2}$$

Although the explicit form of the predictive interval is not available, it can be obtained numerically from softwares like *Maple* or *Matlab*.

Applying the proposed method to the Grubbs [8] data set, the predictive cumulative distribution function obtained in (4.2) is plotted in Figure 2, and the corresponding 90%, 95% and 99% predictive intervals are (161, 2980), (128, 3714), and (47, 5530), respectively. The corresponding intervals obtained by the method discussed in Lawless [1] are (161.00, 2982.23), (129.21, 3715.96), and (48.02, 5532.63), respectively. The two methods give almost identical results. Lawless's method is easy to apply but the derivation is more difficult. The derivation of the proposed method is easy to follow but it requires good numerical integration methods to carry out the calculation.

5. Conclusion

In this paper, by renormalizing the p^* -formula, the exact significance function of the scale parameter of the two-parameter exponential distribution is obtained. This significance function is then used in the approximate Studentization method to obtain the significance function of the threshold parameter. Simulation results illustrated that these two significance functions have excellent coverage properties even when the sample size is extremely small. Finally, these two significance functions are used in the approximate Studentization method to obtain a predictive density and hence a predictive cumulative distribution function, of a future observation from the two-parameter exponential distribution.

References

- [1] J. F. Lawless, "Prediction intervals for the two parameter exponential distribution," *Technometrics*, vol. 19, no. 4, pp. 469–472, 1977.
- [2] A. Baten and A. Kamil, "Inventory management systems with hazardous items of two-parameter exponential distribution," *Journal of Social Sciences*, vol. 5, pp. 183–187, 2009.
- [3] C. Petropoulos, "New classes of improved confidence intervals for the scale parameter of a two-parameter exponential distribution," *Statistical Methodology*, vol. 8, no. 4, pp. 401–410, 2011.
- [4] T. J. DiCiccio, C. A. Field, and D. A. S. Fraser, "Approximations of marginal tail probabilities and inference for scalar parameters," *Biometrika*, vol. 77, no. 1, pp. 77–95, 1990.
- [5] O. E. Barndorff-Nielsen, "Inference on full or partial parameters based on the standardized signed log likelihood ratio," *Biometrika*, vol. 73, no. 2, pp. 307–322, 1986.
- [6] O. E. Barndorff-Nielsen, "Modified signed log likelihood ratio," *Biometrika*, vol. 78, no. 3, pp. 557–563, 1991.
- [7] O. E. Barndorff-Nielsen, "On a formula for the distribution of the maximum likelihood estimator," *Biometrika*, vol. 70, no. 2, pp. 343–365, 1983.
- [8] F. E. Grubbs, "Approximate fiducial bounds on reliability for the two parameter negative exponential distribution," *Technometrics*, vol. 13, pp. 873–876, 1971.
- [9] D. A. S. Fraser and A. C. M. Wong, "On the accuracy of approximate Studentization," *Statistical Papers*, vol. 38, no. 3, pp. 351–356, 1997.