Research Article

Increasing Trend of Resistance to Penicillin, Tetracycline, and Fluoroquinolone Resistance in *Neisseria gonorrhoeae* from Pakistan (1992–2009)

Kausar Jabeen, Summiya Nizamuddin, Seema Irfan, Erum Khan, Faisal Malik, and Afia Zafar

Department of Pathology and Microbiology, Aga Khan University, Karachi 74800, Pakistan

Correspondence should be addressed to Kausar Jabeen, kausar.jabeen@aku.edu

Received 4 April 2011; Revised 12 July 2011; Accepted 12 July 2011

Academic Editor: Thomas R. Unnasch

Copyright © 2011 Kausar Jabeen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Emergence and spread of drug resistant *Neisseria gonorrhoeae* is global concern. We evaluated trends of antimicrobial resistance in *Neisseria gonorrhoeae* over years 1992–2009 in Pakistan. Resistance rates were compared between years (2007–2009) and (1992–2006). Antimicrobial susceptibility testing was performed and interpreted according to Clinical Laboratory Standards Institute (CLSI) criteria using the disk diffusion methodology against penicillin, ceftriaxone, tetracycline and ofloxacin. Additional antibiotics tested in 100 strains isolated during 2007–2009, included cefotaxime, ceftazidime, ceftazidime, cefixime, cefpodoxime, spectinomycin and azithromycin. *Neisseria gonorrhoeae* ATCC 49226 was used as control. Chi-square for trend analysis was conducted to assess resistance trend over the study period. During study period significant increase in combined resistance to penicillin, tetracycline and ofloxacin was observed (P value < 0.01). Resistance rates during the two study period also increased significantly (P value < 0.01). Ceftriaxone resistance was not observed. None of the isolates were found to be resistant or with intermediate sensitivity to additional antibiotics. Our findings suggest that penicillin, ciprofloxacin, tetracycline should not be used in the empirical treatment of gonorrhea in Pakistan. Ceftriaxone and cefixime should be the first line therapy; however periodic MICs should be determined to identify emergence of strains with reduced susceptibility.

1. Introduction

Gonorrhea continues to be a public health problem with the emergence of multidrug resistant strains [1]. Accurate diagnosis with effective treatment to prevent further transmission is one of the essential elements for control of gonococcal infections [2]. Treatment strategies should be devised to utilize appropriate and preferably single-dose therapy that could be conveniently administered at the time of diagnosis [3]. Surveillance of *Neisseria gonorrhoeae* antimicrobial resistance is crucial in guiding empirical therapy in any individual setting as resistance may vary in different countries [4].

Neisseria gonorrhoeae infections have been reported from Pakistan as a cause of sexually transmitted infections, but unfortunately national surveillance data is not available to assess its prevalence or antimicrobial resistance [5, 6]. Due to limited resources and lack of trained technologist and microbiologists, most of the laboratories do not report *Neisseria*

gonorrhoeae. Therefore, existing antimicrobial resistance data from the country is limited and mainly laboratory based. Previously high resistance rates to penicillin, tetracycline, and quinolone have already been reported from the country [7, 8]. Thus ongoing surveillance of resistance is important considering the public health importance of gonococcal infections. In this study, we have evaluated antimicrobial resistance in *Neisseria gonorrhoeae* for the years (2007–2009) and compared it with previously reported data (1992–2006). Additionally, 100 strains from 2007–2009 were further analyzed for additional treatment options.

2. Materials and Methods

This study was conducted in the clinical laboratory of the Aga Khan University Hospital (AKUH), Karachi, Pakistan. The AKUH laboratory is accredited with the Joint Commission of

TABLE 1: Antimicrobial resistance trends in *Neisseria gonorrhoeae* strains over the years 1992–2009 showing significant increase in resistance to penicillin, tetracycline, and ofloxacin. There is also significant increase in combined resistance to all 3 antibiotics. *n* is the total number of resistance strains for that particular year. Percentage is shown in parenthesis (%).

Years	Total number of strains	CRO <i>n</i> (%)	OFX <i>n</i> (%)	P <i>n</i> (%)	TE <i>n</i> (%)	MDR <i>n</i> (%)
1992	15	0	0	1 (7.7)	0	0
1993	11	0	0	0	0	0
1994	16	0	0	4 (25)	0	0
1995	28	0	0	1 (3.6)	0	0
1996	20	0	0	0	0	0
1997	25	0	0	8 (32)	5 (20)	0
1998	26	0	0	3 (11.5)	1 (3.8)	0
1999	33	0	6 (28.8)	6 (19.4)	3 (9.7)	0
2000	25	0	6 (24)	2 (8)	0	0
2001	26	0	6 (23.1)	5 (19.2)	1 (3.8)	0
2002	19	0	8 (42.1)	1 (5.3)	2 (10.5)	0
2003	51	0	37 (72.5)	15 (29.4)	13 (25.5)	7 (13.7)
2004	56	0	47 (83.9)	29 (51.8)	26 (46.4)	17 (30.4)
2005	61	0	48 (78.7)	18 (29.5)	36 (59)	13 (21.3)
2006	90	0	71 (78.9)	39 (43.3)	49 (54.4)	25 (27.8)
2007	95	0	79 (83.2)	57 (60)	62 (65.3)	39 (41.1)
2008	116	0	107 (92.2)	94 (81)	93 (80.2)	79 (68.1)
2009	106	0	98 (92.5)	92 (86.8)	82 (77.6)	75 (70.8)
<i>P</i> -value for trends		NA	<0.01	<0.01	<0.01	<0.01

P: Penicillin; TE: Tetracycline; OFX: Ofloxacin; CRO: Ceftriaxone; MDR: strains resistant to penicillin, tetracycline, and ofloxacin.

TABLE 2: *Neisseria gonorrhoeae* strains with intermediate susceptibility over the study period 1992–2009. *n* is the total number of intermediate strains for that particular year. Percentage is shown in parenthesis (%).

Years	<i>n</i>	OFX	P	TE
1992	15	0	1 (7.7)	0
1993	11	0	1 (9.1)	0
1994	16	0	5 (31.3)	3 (23.1)
1995	28	0	9 (32.1)	8 (28.6)
1996	20	0	1 (5.3)	2 (10)
1997	25	0	7 (28)	7 (28)
1998	26	0	3 (11.5)	4 (15.4)
1999	33	1 (4.8)	3 (9.7)	8 (25.8)
2000	25	0 (0)	4 (16)	3 (12)
2001	26	2 (7.7)	2 (7.7)	3 (11.5)
2002	19	1 (5.3)	10 (52.6)	2 (10.5)
2003	51	5 (9.8)	22 (43.1)	6 (11.8)
2004	56	0 (0)	13 (23.2)	3 (5.4)
2005	61	2 (3.3)	36 (59)	5 (8.2)
2006	90	9 (10.1)	42 (46.7)	15 (16.7)
2007	95	4 (4.2)	31 (32.6)	23 (24.2)
2008	116	5 (4.3)	16 (13.8)	17 (14.7)
2009	106	6 (5.7)	12 (11.3)	16 (15.1)

P: Penicillin; TE: Tetracycline; OFX: Ofloxacin.

appropriate antibiotics for empirical treatment of gonorrhea is challenging.

This 10-year analysis clearly showed that the level of resistance to traditional antibiotics used in the treatment

of gonorrhea continued to increase in local isolates. This trend follows the global pattern of antibiotic resistance in *N. gonorrhoeae* and leaves third-generation cephalosporins as the recommended antibiotic for its treatment [13]. However emergence and spread of strains with reduced susceptibility to ceftriaxone is another emerging issue. Unfortunately, these strains are simultaneously resistance to multiple drug classes, including quinolones, macrolides, penicillins, and tetracyclines [14]. We did not perform ceftriaxone MIC, required to detect reduced susceptibility, and there is a possibility to miss their occurrence as the frequency of MDR strains constitutes a large proportion of our current *N. gonorrhoeae* isolates circulating in the community. On the other hand, treatment failure with ceftriaxone has not been documented from the country. The best strategy in this scenario would be periodic determination of ceftriaxone MICs to detect existence of these strains and close followup with clinicians to identify treatment failure.

A high quinolone resistance approaching almost 100% in recent years was demonstrated in local isolates. This exponential rise in quinolone resistance has also been reported from the region and raises several concerns [15, 16]. We had not evaluated the exact reason for this finding, but probable reasons could be its continuous use despite higher rate of resistance and poor disease control in the community. Increased resistance rates to quinolones have been reported from the country in other community acquired pathogens highlighting the overall misuse of quinolones as the major reason of increased resistance [17, 18].

In view of high resistance to the currently tested antimicrobials in our setting, we tested 8 cephalosporins including

3 oral options: cefixime, cefpodoxime, and cefuroxime. The efficacy of cefixime has been shown to be similar to ceftriaxone [19]. Although cefpodoxime and cefuroxime have shown a cure rate of more than 95%, their pharmacodynamic parameters are less favorable than those of cefixime and ceftriaxone [20, 21]. Nevertheless, they could be used as oral alternatives in the treatment of uncomplicated gonorrhea in our setting [22]. Cefotaxime, cefoxitin, cefepime, ceftazidime, and ceftizoxime are the other single-dose cephalosporins included in the updated treatment regimens for gonococcal infections by the Centers for Disease Control and Prevention [23]. Although studies have reported emergence of resistance to these agents, we were not able to detect resistance against any of them in this study.

Local isolates were also uniformly susceptible to spectinomycin, which is a good option especially in those patients who cannot tolerate cephalosporins [24]. The most likely reason for its consistent susceptibility is its less-frequent use in the community. Rapid emergence of resistance has been reported with the wider use of spectinomycin mandating its rationale use.

Though azithromycin is so far not recommended as a single agent for the treatment of gonorrhea, it has a potential role in the management of these infections in combination with third-generation cephalosporin. We did not detect resistance to this drug although strains with high level resistance have emerged as another threat [25, 26].

Although our dataset included samples from all over Pakistan, sampling limitations prevent us from generalizing our results to the entire population of the country. However, it is important to note that only a few laboratories in the country perform antimicrobial susceptibility testing in *Neisseria gonorrhoeae*, and our data is the largest ever from the country reporting resistance over time. Another limitation is lack of MICs determination for cephalosporins with possibility of missing strains with reduced susceptibility.

In conclusion, we recommend that penicillin, ciprofloxacin, and tetracycline should not be used in the empirical treatment of gonorrhoeae. Ceftriaxone and cefixime should be the first line therapy; however, periodic MICs should be determined to identify emergence of strains with reduced susceptibility. Optimized, standardized, and quality assured antibiotic susceptibility testing needs to be established in laboratories in Pakistan. Simultaneously prevention strategies should be strengthened to maximize the clinical utility of these antimicrobials. Such strategies could include the use of a limited number of treatment regimens and overall avoiding a general misuse of the remaining yet effective antibiotics. In addition, a national policy for management of gonococcal infections should be developed and disseminated to clinicians in the country.

References

- [1] J. W. Tapsall, F. Ndowa, D. A. Lewis, and M. Unemo, "Meeting the public health challenge of multidrug- and extensively drug-resistant *Neisseria gonorrhoeae*," *Expert Review of Anti-Infective Therapy*, vol. 7, no. 7, pp. 821–834, 2009.
- [2] J. W. Tapsall, "Monitoring antimicrobial resistance for public health action," *Communicable Diseases Intelligence*, supplement 27, pp. s70–s74, 2003.
- [3] G. R. Burstein and K. A. Workowski, "Sexually transmitted diseases treatment guidelines," *Current Opinion in Pediatrics*, vol. 15, no. 4, pp. 391–397, 2003.
- [4] J. Tapsall, "Current concepts in the management of gonorrhoea," *Expert Opinion on Pharmacotherapy*, vol. 3, no. 2, pp. 147–157, 2002.
- [5] S. Hawkes, M. Collumbien, L. Platt et al., "HIV and other sexually transmitted infections among men, transgenders and women selling sex in two cities in Pakistan: a cross-sectional prevalence survey," *Sexually Transmitted Infections*, vol. 85, pp. ii8–ii16, 2009.
- [6] A. Altaf, N. Z. Janjua, S. Kristensen et al., "High-risk behaviours among juvenile prison inmates in Pakistan," *Public Health*, vol. 123, no. 7, pp. 470–475, 2009.
- [7] K. Jabeen, E. Khan, and R. Hasan, "Emergence of quinolone-resistant *Neisseria gonorrhoeae* in Pakistan," *International Journal of STD and AIDS*, vol. 17, no. 1, pp. 30–33, 2006.
- [8] A. Zafar and K. Jabeen, "Antimicrobial resistance in *Neisseria gonorrhoeae* and limited treatment options," *Journal of the Pakistan Medical Association*, vol. 57, no. 7, pp. 333–334, 2007.
- [9] E. W. Koneman, S. D. Allen, W. M. Janda, P. C. Schreckengaber, and J. W. C. Winn, *Color Atlas and Text Book of Diagnostic Microbiology*, Lippincott, Philadelphia, Pa, USA, 5th edition, 1997.
- [10] Clinical Laboratory Standards Institute, "Performance standards for antimicrobial susceptibility testing," Sixteen informational supplement, M100-S16 (M7-A7), 2006.
- [11] "British Society of Antimicrobial Chemotherapy Methods for Antimicrobial Susceptibility Testing," Version 8, January 2009.
- [12] J. W. Tapsall, *Antibiotic Resistance in Neisseria Gonorrhoeae*, World Health Organization, Geneva, Switzerland, 2001.
- [13] J. W. Tapsall, "Antibiotic resistance in *Neisseria gonorrhoeae*," *Clinical Infectious Diseases*, vol. 41, no. 4, pp. S263–S268, 2005.
- [14] J. W. Tapsall, "Implications of current recommendations for third-generation cephalosporin use in the WHO Western Pacific Region following the emergence of multiresistant gonococci," *Sexually Transmitted Infections*, vol. 85, no. 4, pp. 256–258, 2009.
- [15] A. Sutrisna, O. Soebjakto, F. S. Wignall et al., "Increasing resistance to ciprofloxacin and other antibiotics in *Neisseria gonorrhoeae* from East Java and Papua, Indonesia, in 2004—Implications for treatment," *International Journal of STD and AIDS*, vol. 17, no. 12, pp. 810–812, 2006.
- [16] WHO Western Pacific Programme, South East Asian Gonococcal Antimicrobial Surveillance Programme, J. W. Tapsall et al., "Surveillance of antibiotic resistance in *Neisseria gonorrhoeae* in the WHO Western Pacific and South East Asian regions, 2007–2008," *Communicable Diseases Intelligence*, vol. 34, no. 1, pp. 1–7, 2010.
- [17] K. Jabeen, A. Zafar, S. Irfan, E. Khan, V. Mehraj, and R. Hasan, "Increase in isolation of extended spectrum beta lactamase producing multidrug resistant non typhoidal *Salmonellae* in Pakistan," *BMC Infectious Diseases*, vol. 10, article 101, 2010.
- [18] E. Khan, K. Jabeen, M. Ejaz, J. Siddiqui, M. F. Shezad, and A. Zafar, "Trends in antimicrobial resistance in *Shigella* species in Karachi, Pakistan," *Journal of Infection in Developing Countries*, vol. 3, no. 10, pp. 798–802, 2009.
- [19] R. M. Ramus, J. S. Sheffield, J. A. Mayfield, and G. D. Wendel Jr., "A randomized trial that compared oral cefixime and intramuscular ceftriaxone for the treatment of gonorrhea in

- pregnancy," *American Journal of Obstetrics and Gynecology*, vol. 185, no. 3, pp. 629–632, 2001.
- [20] R. P. Das, K. Jones, A. J. Robinson, and D. J. Timmins, "Cefuroxime axetil to treat gonorrhea," *Genitourinary Medicine*, vol. 64, no. 6, p. 394, 1988.
- [21] G. R. Kinghorn, R. C. Spencer, T. K. Smith, P. D. Woolley, R. Patel, and A. J. Robinson, "Comparative study of cefuroxime axetil and procaine penicillin in the treatment of uncomplicated gonorrhoea," *International Journal of STD and AIDS*, vol. 1, no. 4, pp. 285–287, 1990.
- [22] Centers for Disease Control and Prevention, "Oral alternatives to cefixime for the treatment of uncomplicated *Neisseria gonorrhoeae* urogenital infections," <http://www.cdc.gov/std/treatment/cefixime.htm>.
- [23] Centers for Disease Control and Prevention, "Updated recommended treatment regimens for gonococcal infections and associated conditions," <http://www.cdc.gov/std/treatment/2006/updated-regimens.htm>.
- [24] K. A. Workowski and S. M. Berman, "Sexually transmitted diseases treatment guidelines," *The MMWR Recommendations and Reports*, vol. 55, no. 11, pp. 1–94, 2006.
- [25] S. Starnino and P. Stefanelli, "Azithromycin-resistant *Neisseria gonorrhoeae* strains recently isolated in Italy," *Journal of Antimicrobial Chemotherapy*, vol. 63, no. 6, pp. 1200–1204, 2009.
- [26] P. G. Galarza, B. Alcalá, C. Salcedo et al., "Emergence of high level azithromycin-resistant *Neisseria gonorrhoeae* strain isolated in Argentina," *Sexually Transmitted Diseases*, vol. 36, no. 12, pp. 787–788, 2009.