CYSTIC fibrosis (CF) lung disease is characterized by persistent inflammation. Antiinflammatory drugs, such as corticosteroids and ibuprofene, have proved to slow the decline of pulmonary function although their use is limited because of frequent adverse events. We hypothesized that colchicine could be an alternative treatment because of its antiinflammatory properties and upregulatory effect on cystic fibrosis transmembrane regulator (CFTR) closely related proteins. We herein present results obtained in an open study of eight CF children treated with colchicine for at least 6 months. Clinical status was better in all patients and respiratory function tests significantly improved in five. Median duration of antibiotherapy decreased significantly. These preliminary results support our hypothesis of a beneficial effect of colchicine in CF patients and stress the need for a controlled therapeutic trial.

Key words: Cystic Fibrosis, Inflammation, Cystic fibrosis transmembrane regulator, ABC protein, Colchicine

Interest of colchicine for the treatment of cystic fibrosis patients. Preliminary report

I. Sermet-Gaudelus,¹ V. Stoven,² J.P. Annereau,² V. Witko-Sarsat,³ P. Reinert,⁴ M. Guyot,⁵ B. Descamps-Latscha,³ J.Y. Lallemand² and G. Lenoir^{1,CA}

¹Service de Pédiatrie II, Hôpital Necker-Enfants Malades, 149 rue de Sèvres, 75015 Paris;

²Laboratoire de Résonnance Magnétique Nucléaire, Ecole Polytechnique, Massy Palaiseau; ³INSERM U507, Hôpital Necker, Paris; ⁴Pediatric Department, Hôpital Intercommunal, Créteil; ⁵Pediatric Department, Hôpital de Lisieux, France

^{CA}Corresponding Author

Tel: (+33) 1 44 49 48 83

Fax: (+33) 1 47 83 32 26

Email: gérard.lenoir@nck.ap-hop-paris.fr

Introduction

Lung disease in cystic fibrosis (CF) patients is characterized by persistent bacterial infection leading to chronic bronchitis and bronchiectasis.¹ Vigorous antibiotherapy, clearance of mucus and adequate nutritional support have been the bases of conventional therapy. Recently, attention has been focused on the lung inflammation which not only damages the parenchyma directly but also impairs local defence against pathogens.^{2,3} Blunting the inflammatory response by using antiinflammatory drugs such as corticosteroids or ibuprofen has been proven to slow the decline of pulmonary function, although the high incidence of adverse events outweighs beneficial effect.^{4,5}

Colchicine should be considered as an alternative treatment because of its antiinflammatory properties notably by interference with PNN margination, chemotaxis and phagocytosis.⁶ Interestingly, colchicine has also been shown to upregulate the expression of the multi-drug-resistance (MDR) gene.⁷ This gene encodes for the P-glycoprotein or P-gp which belongs to the ATP binding cassette (ABC) protein family and shares remarkable homology with cystic fibrosis transmembrane regulator (CFTR).⁸ *In vitro* experiments have

shown that closely related ABC proteins can complement each other.⁹ Taken together, these data led us to propose that colchicine might induce complementation of CFTR by P-gp.

Recently, we reported improvement of lung function in a CF patient after antitumoral therapy and suggested that this could be due to the complementation of CFTR.¹⁰ However the toxicity of these drugs precludes their wide use in CF patients. We herein present preliminary results showing that colchicine may exert a beneficial effect on CF clinical status.

Materials and methods

Patients

Eight patients with CF diagnosed according to conventional criteria entered this open study conducted from 1 March 1997 to 31 December 1997.¹² All these patients had end-stage CF lung disease or chronic airflow limitation unresponsive to conventional therapy. Colchicine (Houdé, Paris, France) was administered orally (1 mg daily), during at least 6 months without other antiinflammatory treatment (inhaled or oral steroid therapy, nonsteroidal antiinflammatory drugs) or recombinant human DNase.

The following parameters were considered every 3 months during the duration of the assay, i.e. at least 6 months: body weight (BW) and height (expressed as standard deviation from ideal weight and height for age); respiratory function tests (forced expiratory volume- FEV1, and forced vital capacity-FVC); number of antibiotic courses in relation to respiratory tract infections. Concomitant therapies and adverse clinical events were documented.

Statistics

Results are presented as median (range) values. The average values in the 6 months before the beginning of treatment were compared with their post-treatment average after completion of at least 6 months of treatment. Statistical analysis was done using Wilcoxon's signed rank test. *P* value of less than 0.05 was considered significant.

Results

Patient characteristics

Eight patients (six boys) were studied. The characteristics of the patients at beginning of treatment are shown in Table 1. Median age was 13.5 years (range 5–28 years). Six patients were ΔF508 homozygous. Median FEV1 and FVC were respectively 54% (range 21–63%) and 62% (37–64%). Patient 7 was a 27-year-old patient with secondary nephropathic amyloidosis. The occurrence of renal failure justified the start of colchicine therapy. Patient 8 was a 13-year-old girl on the lung transplant waiting list because of a dramatic decline of respiratory function and nutritional status (Fig. 1). Patient 2 and patient 8 both also had concomitant long-term azithromycin started 2 years before.

FIG. 1. Evolution of FEV1 and FVC. *Wilcoxon's signed rank test.

Effect of colchicine on patient clinical status

Median duration of treatment was 7 months (range 6–12 months). No child had change in the baseline treatment during the study period. For all the patients there was a subjective improvement in clinical status. Median duration of antibiotherapy significantly decreased from 5 to 1.2 days per month (*P* < 0.05). Median weight gain was + 0.25 SD (*P* = 0.1).

The respiratory function tests could be performed in seven patients. FEV1 substantially improved in six patients, rising from 54% (21–63%) predicted to 57% (37–64%) predicted (*P* = 0.14). In contrast, FVC remained unchanged during the duration of colchicine therapy. Patient 8, on the lung transplant waiting list, had a marked improvement of FEV1 from 21% to 37% predicted, and FVC from 32% to 44% which prompted his removal from this list. Concomitantly, weight gain (4 kg) and reduced need for antibiotics (15 days during the 6 months after beginning of treatment versus 30 days during the 6 months before) were observed. The child had recovered good daily activity. It is worth noting that there had been no concomitant change in the usual treatment of this patient.

Adverse effects

The drug was well tolerated except in two patients who had mild diarrhoea. However, this mild side

Table 1. Baseline characteristics of patients before colchicine treatment

Patient (No.)	Age (years)	BW*	CF mutation	FVC**	FEV1**	Lung colonization
1	10	−0.5	DF508/DF508	66%	54%	<i>P. aeruginosa</i>
2	16	−1	DF508/DF508	65%	60%	<i>P. aeruginosa</i> <i>Staph. aureus</i> <i>B. Cepacia</i>
3	14	−0.5	DF508/DF508	81%	78%	<i>Staph. aureus</i>
4	5	−1.5	?	ND	ND	<i>P. aeruginosa</i> <i>Staph. aureus</i>
5	6	−2	DF508/DF508	66%	54%	<i>H. influenzae</i>
6	13	0.5	DF508/DF508	90%	92%	<i>P. aeruginosa</i>
7	28	−1	DF508/DF508	72%	61%	<i>P. aeruginosa</i>
8	14	−2	DF508/?	32%	32%	<i>P. aeruginosa</i> <i>B. Cepacia</i>
Median	13.5	−1		54	62	
Range	5–28	−2–0.5		21–63	32–68	

* Expressed as SD for theoretical weight for age;
** Expressed as percentages of theoretical value for age and sex.

effect did not lead to discontinuation of the treatment.

Discussion

We report preliminary results showing that colchicine treatment in CF patients is associated with a significant decrease of antibiotherapy, and a potential improvement of respiratory status. To our knowledge, colchicine has not yet been proposed in CF but interestingly, it has already been used for the treatment of idiopathic pulmonary fibrosis¹² and IgE-mediated asthma.¹³

The beneficial effect of colchicine may be due to two different mechanisms: (i) colchicine has anti-inflammatory properties due to interference with PNN margination, chemotaxis and phagocytosis¹⁴ and inhibition of leukotriene synthesis.¹⁵ It also inhibits fibroblast proliferation and total collagen synthesis;¹⁶ (ii) colchicine also promotes expression of ABC proteins such as P-gp which shares great structural homology with CFTR.^{7,8,17,18} Several recent data also suggest functional homology. CFTR overexpression leads to a multidrug-resistance phenotype similar to P-gp.¹⁹ CFTR exports glutathione and glucuronate anions and may play a role in cell detoxification similar to MRP.^{20,21} As closely related ABC proteins can complement each other, upregulation of P-gp and/or MRP may hence lead to functional complementation of defective CFTR.^{9,10} This hypothesis is also supported by the expression of MDR in nasal epithelial cells of a CF patient treated with antitumoural drugs but not in a control CF patient who never had chemotherapy.¹⁰ This function of upregulation of ABC proteins could be emphasized by the adjunction of other ABC protein inducers such as azithromycin.²² Interestingly, two patients were treated by both drugs and experienced an improvement in clinical status.

In conclusion, colchicine should be considered in the treatment in CF patients, because of its anti-inflammatory role and its property of inducing ABC proteins. Although preliminary, the potential importance of these results in this severe disease seems to us to justify their early report and has formed the basis of a controlled trial which is now being undertaken by our group.

References

- Koch C, Hoiby N. Pathogenesis of cystic fibrosis. *Lancet* 1993; **341**: 1065–1069.
- Khan TZ, Wagener JS, Bost T, *et al.* Early pulmonary inflammation in infants with cystic fibrosis. *Am J Respir Crit Care Med* 1995; **151**: 1075–1082.
- Descamps-Latscha B, Döring G, Galanaud P, Lenoir G, Navarro J, Schaffar L. Inflammation in Cystic Fibrosis. *Med Inflamm* 1996; **5**: 121–143.
- Auerbach HS, Williams M, Kirkpatrick JA, Cotten HR. Alternate day prednisolone reduces the morbidity and improves pulmonary function in cystic fibrosis. *Lancet* 1985; **ii**: 686–688.
- Konstan MW, Byard PJ, Hoppel CL, Davis PB. Effect of high-dose ibuprofen in patients with cystic fibrosis. *N Engl J Med* 1995; **331**: 848–854.
- Levy M, Spino M, Read E. Colchicine: a state of the art review. *Pharmacother* 1991; **11**(3): 196–211.
- Ueda K, Pastan I, Gottesman MM. Isolation and sequence of the promoter region of the human multidrug-resistance (P-glycoprotein) gene. *J Biol Chem* 1987; **262**: 17432–17436.
- Higgins CF. ABC transporters: from microorganisms to man. *Annu Rev Biol* 1992; **8**: 67–113.
- Tommasini R, Evers R, Vogt E, *et al.* The human multidrug resistance-associated protein functionally complements the yeast cadmium resistance factor 1. *Proc Natl Acad Sci USA* 1996; **93**: 6743–6748.
- Lallemant JY, Stoven V, Annereau JP, Boucher J, Blanquet S, Barthe J, Lenoir G. Induction by antitumoural drugs of proteins that functionally complement CFTR: a novel therapy for cystic fibrosis? *Lancet* 1997; **350**: 711–712.
- Stern RC. The diagnosis of cystic fibrosis. *N Engl J Med* 1997; **336**: 487–491.
- Douglas WW, Ryu JH, Swensen SJ, *et al.* Colchicine versus prednisone in the treatment of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1998; **158**: 220–225.
- Kelly SJ, Uri AJ, Freeland HS, *et al.* Effects of colchicine on IgE-mediated early and late airway reactions. *Chest* 1995; **107**: 985–991.
- Dallaverde E, Fan PT, Chang YH. Mechanism of action of colchicine. V. Neutrophil adherence and phagocytosis in patients with acute gout treated with colchicine. *J Pharmacol Exp Ther* 1982; **223**: 197–202.
- Peters-Golden M, McNish RW, Davis JA, *et al.* Colchicine inhibits arachidonate release and 5-lipoxygenase action in alveolar macrophages. *Am J Physiol* 1996; **271**: 6 Pt 1, L1004–1013.
- Entzian P, Schlaak M, Seitzer U, *et al.* Antiinflammatory and antifibrotic properties of colchicine: implications for idiopathic pulmonary fibrosis. *Lung* 1997; **175**: 41–51.
- Vollrath V, Wielandt AM, Acuna C, *et al.* Effect of colchicine and heat shock on multidrug resistance gene and P-glycoprotein expression in rat liver. *J Hepatol* 1994; **21**: 754–763.
- Eliseenkova AV, Kakpakova ES, Abdjashitov RI, Stavrovskaya AA. Colchicine-resistance and enhancement of P-glycoprotein activity after co-cultivation of drug-sensitive cells with multidrug resistant variants. *Cell Biol Int* 1995; **19**(2): 113–119.
- Wei LY, Stutts MJ, Hoffman MM, Roepe PD. Overexpression of the CFTR in NIH 3T3 cells lowers membrane potential and intracellular pH and confers a multidrug resistance phenotype. *Biophys J* 1995; **69**: 883–895.
- Lindsell P, Hanrahan JW. Glutathione permeability of CFTR. *Am J Cell Physiol* 1998; **275**(44): C323–326.
- Lindsell P, Hanrahan JW. Adenosine triphosphate dependant asymmetry of anion permeation in cystic fibrosis transmembrane conductance regulator chloride channel. *J Gen Physiol* 1998; **111**: 601–614.
- Altschuler EL. Azithromycin, the multidrug-resistant protein, and cystic fibrosis. *Lancet* 1998; **351**: 1286.

ACKNOWLEDGEMENT. This work was supported by the association 'ABCF Protéine'.

Received 17 November 1998;
accepted in revised form 18 December 1998