

MACROPHAGE migration inhibitory factor (MIF) has recently been forwarded as a critical regulator of inflammatory conditions, and it has been hypothesized that MIF may have a role in the pathogenesis of asthma and chronic obstructive pulmonary disease (COPD). Hence, we examined effects of MIF immunoneutralization on the development of allergen-induced eosinophilic inflammation as well as on lipopolysaccharide (LPS)-induced neutrophilic inflammation in lungs of mice. Anti-MIF serum validated with respect to MIF neutralizing capacity or normal rabbit serum (NRS) was administered i.p. repeatedly during allergen aerosol exposure of ovalbumin (OVA)-immunized mice in an established model of allergic asthma, or once before instillation of a minimal dose of LPS into the airways of mice, a tentative model of COPD. Anti-MIF treatment did not affect the induced lung tissue eosinophilia or the cellular composition of bronchoalveolar lavage fluid (BALF) in the asthma model. Likewise, anti-MIF treatment did not affect the LPS-induced neutrophilia in lung tissue, BALF, or blood, nor did it reduce BALF levels of tumor necrosis factor- α (TNF- α) and macrophage inflammatory protein-1 α (MIP-1 α). The present data suggest that MIF is not critically important for allergen-induced eosinophilic, and LPS-induced neutrophilic responses in lungs of mice. These findings do not support a role of MIF inhibition in the treatment of inflammatory respiratory diseases.

Key words: Macrophage migration inhibitory factor (MIF), Allergen-induced eosinophilia, LPS-induced neutrophilia, Mouse models of inflammatory lung diseases

Role of macrophage migration inhibitory factor (MIF) in allergic and endotoxin-induced airway inflammation in mice

M. Korsgren,^{1,CA} L. Källström,³ L. Uller,¹ T. Bjerke,³ F. Sundler,¹ C. G. A. Persson² and O. Korsgren⁴

Departments of ¹Physiological Sciences and ²Clinical Pharmacology, Lund University Hospital, 221 85 Lund; ³Department of Inflammation Pharmacology, AstraZeneca R&D Lund, 221 87 Lund; ⁴Department of Clinical Immunology and Transfusion Medicine, Uppsala University Hospital, 751 85 Uppsala, Sweden

^{CA} Corresponding Author

Tel: (+46) 46 177714

Fax: (+46) 46 177720

Email: Magnus.Korsgren@mphy.lu.se

Introduction

Macrophage migration inhibitory factor (MIF) is considered to be a critical regulator of various inflammatory conditions. For example, MIF is a pivotal mediator in the host response to endotoxic shock,¹ and plays an important role in the development of the delayed type hypersensitivity reaction and collagen-induced arthritis in mice.^{2,3} MIF may further have a proinflammatory role in the development of human acute respiratory distress syndrome.⁴ MIF was described originally to be a T-cell product, which inhibited the random migration of macrophages.^{5,6} Interestingly, recent data indicate that MIF is predominantly expressed by Th2-like T-cells.⁷ However, MIF also exists preformed in monocytes/macrophages, eosinophils, B-cells, airway epithelial cells, and corticotrophic cells within the anterior pituitary gland, and is released in response to various proinflammatory stimuli.^{1,8–12} MIF shares with other cytokines sensitivity to inhibition by therapeutic concentrations of glucocorticoids. However, at low physiological glucocorticoid concentrations MIF

expression is induced.^{13,14} MIF has the property of counteracting anti-inflammatory and immunosuppressive actions of glucocorticoids.^{13,14} Anti-MIF therapeutic strategies are thus under development with the aim to increase the immunosuppressive and anti-inflammatory properties of endogenously released glucocorticoids, thereby reducing the requirement for steroid therapy in a variety of inflammatory conditions. It has further been suggested that drugs inhibiting MIF would be effective also in inflammatory conditions that exhibit steroid resistance.¹⁴

Airway mucosal inflammation in allergic asthma is characterized by infiltration and/or activation of eosinophils, macrophages, T-lymphocytes, and mast cells.¹⁵ Since all these immune cells can produce MIF in significant quantities, and since elevated levels of MIF have been detected in BALF from asthmatic patients compared with controls, it has been forwarded that MIF may have a role in eosinophilic airway disease.⁹

Recent studies have shown that genetically MIF-deficient mice are resistant to the lethal effects of a high systemic dose of LPS, but susceptible to a

combination of a low-dose of LPS and D-galactosamine.¹⁶ These mice exhibit no impairment of neutrophil migration to the peritoneum elicited by thioglycollate; despite diminished neutrophil accumulation in BALF they clear *Pseudomonas aeruginosa* instilled into the trachea better than wild type mice.¹⁶ Makita *et al.*¹¹ recently demonstrated that immunoneutralization of MIF attenuates pulmonary neutrophil influx and acute lung injury induced by intraperitoneal (i.p.) administration of LPS in rats. These effects were associated with reduced BALF levels of macrophage inflammatory protein-2, a powerful neutrophil chemokine.¹¹ Moreover, it has been demonstrated that MIF induces alveolar cells/macrophages to secrete TNF- α and IL-8,^{4,17} two cytokines widely thought to be critically important for neutrophil infiltration in pulmonary inflammatory conditions such as chronic obstructive pulmonary disease (COPD).^{18,19}

In an attempt to further explore the hypothesis that MIF may have a role in bronchopulmonary eosinophilic and neutrophilic inflammation, we examined effects of MIF inhibition both in an established murine model of allergic asthma, and in a model involving LPS-induced neutrophilic inflammation in the lungs of mice. Specifically, anti-MIF serum was administered repeatedly during allergen aerosol exposure of immunized mice, or given once before instillation of a low dose of LPS into the lower airways of mice.

Materials and methods

Animals

Male C57BL/6 mice ($n=184$, 6–8 weeks of age), were purchased from Bomholtgaard, Denmark. All mice were kept in well-controlled animal housing facilities and had free access to tap water and pelleted food throughout the experimental period. All animals were used under protocols approved by the Ethics Committee of the Faculty of Medicine at the University of Lund.

Induction of allergic eosinophil-rich airway inflammation

We have used a protocol slightly modified from that developed by Brusselle and colleagues.²⁰ On the first day of the experiment (Day 0), all mice were actively immunized by i.p. injection of 10 μ g chicken OVA (Grade III, Sigma, St Louis, MO, USA), adsorbed to 1 mg of alum adjuvant. Starting 14–16 days after immunization the mice were exposed once daily during 7 days to aerosolized saline (SAL) or OVA over a 30-min period by placing groups of 5–10 awake mice in an exposition chamber. The aerosol was generated into the chamber using a nebulizer (Bird 500 ml Inline Micronebulizer

driven at 4 bar, Bird Co., Palm Springs, CA). The concentration of OVA in the nebulizer was 1% w/v. Animals were sacrificed by i.p. injection of pentobarbital 8 h after the last aerosol exposure.

Induction of neutrophil-rich airway inflammation

Groups of mice received one intratracheal instillation of a low dose of *Escherichia coli* LPS (Difco Lab., MI, USA, 4 μ g/kg, i.e. ~ 0.08 μ g/animal, diluted in ~ 20 μ l saline) or SAL. The present LPS dose is comparable with occupational levels.²¹ For example, it has been estimated that cotton mill workers are exposed to 60 μ g endotoxin per day.²² In preliminary dose-response experiments our selected dose of LPS was shown to induce submaximal responses regarding TNF- α levels and total cell numbers in BALF. For the instillation procedure, animals were anaesthetized with enflurane, and a blunted cannula was introduced perorally into the trachea. Animals were sacrificed by i.p. injection of pentobarbital 4 or 24 h after LPS or SAL administration.

In vivo neutralization of MIF

Starting the day before the first allergen aerosol challenge, mice ($n = 20$) were injected i.p. with 200 μ l of rabbit anti-murine MIF serum; this treatment was then repeated every 3 days until termination of the experiment. Control mice ($n = 20$) were injected with a similar volume (200 μ l) of NRS. Other groups received no treatment at all. Groups of mice subjected to intratracheal LPS instillation received one i.p. injection of anti-MIF serum (200 μ l), NRS, or SAL 12 h before LPS challenge ($n = 8$ in each group). This anti-MIF treatment has previously been shown to neutralize MIF *in vivo* using the same dose level and administration route in mice as in this study.^{1,23} Components in serum may have some capacity to downregulate inflammatory responses, underscoring the need for proper control groups in the evaluation of experiments using anti-serum. Thus, in this study groups of mice treated with anti-MIF serum were compared with corresponding groups of mice treated with NRS.

In additional experiments we explored whether anti-MIF treatment might be effective via the local route. Groups of mice ($n = 4$ in each group) received 15 μ l anti-MIF serum or NRS administered intratracheally, alone or together with LPS solution (4 μ g/kg) in a total volume of ~ 20 μ l.

The present batch of anti-MIF serum was checked for bioactivity at our laboratory using a protocol, which has previously been employed to analyse the role of MIF in endotoxaemia.¹⁶ Groups of mice ($n = 5$ –6 in each group) received one i.p. injection of anti-MIF serum (200 μ l), NRS, or SAL 2 h before i.p. injection with a high dose of LPS (25 mg/kg). Ninety minutes after LPS challenge blood was collected by

cardiac puncture and placed into EDTA tubes. As an indicator of LPS-induced plasma extravasation, the haematocrit was determined by an automated haematology analyser (Sysmex K-4500, TOA Medical Electronics Co., Kobe, Japan).

Histological analysis of allergic airway inflammation

Lung tissue specimens obtained 8 h after the last OVA or SAL exposure were immersed overnight in Stefani's fixative (2% paraformaldehyde and 0.2% picric acid in 0.1 M phosphate buffer, pH 7.2), rinsed repeatedly in buffer (Tyrode's buffer supplemented with 10% sucrose), frozen in mounting medium (Tissue-Tek, Miles Inc, Elkhart, IN, USA), and stored at -80°C until sectioning. Eosinophils were detected by histochemical visualization of cyanide-resistant eosinophil peroxidase (EPO) activity.²⁴⁻²⁶ Briefly, cryosections (10 μm) were incubated for 8 min at room temperature in PBS buffer (pH 7.4) supplemented with 3,3-diaminobenzidine tetrahydrochloride (60 mg/100 ml, Sigma), 30% H_2O_2 (0.3 ml/100 ml), and NaCN (120 mg/100 ml). Slides were then rinsed in water and mounted in Kaiser's medium (Merck, Darmstadt, Germany). Eosinophils were identified by their dark-brown reaction product. For evaluation of the number of eosinophils in pulmonary tissue, 40 randomly selected areas (0.04 mm^2 each) in one lung section from each animal were examined. The number of eosinophils in the 40 areas was counted at a magnification of 400, and the mean was expressed as eosinophils/unit area. Cell counts were made in a blinded fashion. For assessment of general airway morphology sections were stained with haematoxylin and erythrosin. Lung tissue specimens and tracheobronchial lymph nodes immersed in buffered 4% paraformaldehyde (pH 7.2), dehydrated, and embedded in paraffin, were used for immunohistochemical visualization of MIF-expressing cells. Sections were incubated overnight in 4°C in a moist chamber with a 1:800 dilution of the rabbit anti-murine MIF serum. The anti-MIF serum used in the present study has previously been used for immunohistochemistry.^{1,2} NRS at a dilution of 1:800 or PBS were used in control sections. The site of the antigen-antibody reaction was revealed by application of fluorescein isothiocyanate-conjugated swine antisera directed at rabbit immunoglobulins (DAKO, Glostrup, Denmark) diluted 1:80 for 1 h at room temperature.

Histological analysis of LPS-induced airway inflammation

Lung tissue specimens obtained 4 h and 24 h after LPS or SAL challenge were immersed in buffered 4% paraformaldehyde (pH 7.2), dehydrated, and embedded in paraffin. One section (6 μm) per animal ($n = 3$ from each group) were stained with haematoxylin and

erythrosin to examine the extent of pulmonary inflammation. Other sections (one per animal, $n = 3$ from each group) were used for immunohistochemical demonstration of neutrophils in lung tissue. Sections were incubated overnight in 4°C in a moist chamber with a 1:200 dilution of rabbit antisera directed at human myeloperoxidase (MPO, DAKO, Glostrup, Denmark). The site of the antigen-antibody reaction was revealed by application of fluorescein isothiocyanate-conjugated swine antisera directed at rabbit immunoglobulins (DAKO, Glostrup, Denmark) for 1 h at room temperature. In control sections, omitting the primary antibody, only slight yellowish auto-fluorescence was found.

Analysis of cells in peripheral blood and BALF, and measurement of TNF- α and MIP-1 α levels in BALF and plasma

Animals were anaesthetized with an i.p. injection of pentobarbital. The chest was opened and a blood sample was collected via the still beating heart. A tracheal cannula was inserted via a midcervical incision. The airways were lavaged once (LPS-challenged mice) or twice (OVA-exposed mice) with 0.7 ml of PBS (Life Technologies, Paisley, UK). The BALF was immediately centrifuged (10 min, 4°C , $160 \times g$). Cell pellets were resuspended in 250 μl PBS for total and differential cell counting and the supernatants were rapidly frozen. Differential counting was performed on May-Grünwald-Giemsa stained cytopins and blood smears. Between 200 and 500 cells were counted on each cytopsin, and 100 cells were counted on each blood smear. Commercial ELISA kits (R&D systems, MN, USA) were used to measure levels of TNF- α and MIP-1 α in the BALF of LPS and SAL challenged mice. TNF- α levels were also measured in plasma obtained from LPS challenged mice. The limit of detection was 5.1 pg/ml for TNF- α , and 1.5 pg/ml for MIP-1 α .

Statistics

Data are expressed as mean \pm SEM. To calculate significance levels between treatment groups, the Student's t -test was used throughout the study. ELISA values below detection limits were assigned the value of the detection limit. Probabilities <0.05 were used as the generally accepted level of statistical significance for differences between mean values.

Results

Additional validation of the present anti-MIF sera

Exposure of a high dose of endotoxin is known to cause plasma extravasation and subsequent loss of circulating plasma volume.^{27,28} In order to ascertain a

preserved activity of the anti-MIF serum, the haematocrit was determined 90 min after i.p. LPS challenge as a measure of plasma extravasation. The haematocrit was significantly increased in LPS-challenged mice compared with SAL-challenged mice ($53.1 \pm 4.0\%$ vs. $45.2 \pm 6.2\%$, $P < 0.001$). This response was inhibited in mice treated with anti-MIF serum before LPS challenge ($45.7 \pm 2.1\%$ vs. $50.8 \pm 5.0\%$ in corresponding NRS-treated mice, $P < 0.01$).

Effect of anti-MIF treatment on allergen-induced airway inflammation

To assess the role of MIF in allergic airway inflammation, lung tissue eosinophilia and cellular composition of BALF were determined in anti-MIF-treated and NRS-treated mice 8 h after last allergen aerosol exposure. The number of eosinophils in lung tissue was similar in allergen aerosol exposed anti-MIF-treated and NRS-treated mice (Fig. 1). Likewise, total cellular content (data not shown) and the percentage of eosinophils, neutrophils, lymphocytes, and macrophages in BALF did not differ significantly between anti-MIF-treated and NRS-treated mice after allergen aerosol exposure (Fig. 2).

Untreated SAL and OVA exposed mice were also included in the study, to check that the present inflammation was specifically induced by OVA challenge. Histologic analysis of lungs taken from OVA-exposed mice revealed the presence of peribronchial and perivascular infiltrates, whereas lung tissue taken from SAL-exposed animals showed normal lung histology. Allergen aerosol exposures caused a marked eosinophilia in pulmonary tissue (12.1 ± 2.0 vs. 3.0 ± 0.5 eosinophils/unit area in SAL exposed mice, $P < 0.01$). A significant increase in the percentage of




FIG. 1. Effect of anti-MIF treatment on allergen-induced lung tissue eosinophilia. Mice were immunized and then challenged daily with aerosolized OVA. Data are from two independent experiments and are presented as mean \pm SEM. NRS = NRS-treated mice ($n=20$), anti-MIF = anti-MIF-treated mice ($n=20$). Lung tissue eosinophilia did not differ significantly between anti-MIF-treated and NRS-treated mice.




FIG. 2. Cellular composition of BALF in immunized and allergen challenged mice treated with anti-MIF serum or NRS. Data are presented as mean \pm SEM. NRS = NRS-treated mice ($n=10$), anti-MIF = anti-MIF-treated mice ($n=10$). Eos. = eosinophils, Neutro. = neutrophils, Lymph. = lymphocytes, Macro. = macrophages. OVA challenge induced similar changes of the cellular profile in BALF of anti-MIF-treated and NRS-treated mice.

eosinophils ($44.4 \pm 12.2\%$ vs. $2.2 \pm 1.6\%$ in SAL-exposed mice, $P < 0.05$), neutrophils ($1.7 \pm 0.5\%$ vs. $0.1 \pm 0.1\%$ in SAL-exposed mice, $P < 0.05$), and lymphocytes ($7.5 \pm 1.6\%$ vs. $0.8 \pm 0.3\%$ in SAL-exposed mice, $P < 0.01$) was demonstrated in response to OVA challenge.

Immunohistochemistry was used to visualize the MIF-expressing cells in the pulmonary infiltrates and tracheobronchial lymph nodes of OVA-challenged mice. A majority of the leukocytes in the perivascular and peribronchial pulmonary infiltrates were MIF-positive (Fig. 3a). Interestingly, large, intensely MIF-positive cells with dendritic shape were seen in the superficial cortex of the tracheobronchial lymph nodes in OVA-challenged mice (Fig. 3b). Lymphocytes, mainly located in the cortex, were also stained for MIF, although less intensely (Fig. 3c). The exact identity of the MIF-positive cells was not further evaluated in the present study. No staining except for a yellowish auto-fluorescence (compare Fig. 3a and d) was observed in control sections where NRS or PBS was used instead of the anti-MIF serum (not shown).

Characterization of the present 'low dose' LPS model

Intratracheal instillation of a minimal dose of LPS into the lower airways of mice induced a significant increase in total cellular content of BALF 4 h after challenge (51.9 ± 5.5 vs. 10.6 ± 2.6 BALF cells $\times 10^4$ /ml in SAL challenged controls, $P < 0.001$). This increase remained 24 h after LPS instillation (33.8 ± 4.6 vs. 12.6 ± 1.8 BALF cells $\times 10^4$ /ml in SAL challenged controls, $P < 0.01$). The percentage of




FIG. 3. Immunohistochemistry. Visualization of MIF-expressing cells in the pulmonary infiltrates (a) and tracheobronchial lymph nodes (b and c) of OVA challenged mice, and visualization of MPO-positive cells in the lungs of LPS-challenged mice (d). A majority of the leukocytes in the pulmonary infiltrates (arrows) of OVA-challenged mice were stained with anti-MIF serum (a). Interestingly, large, intensely MIF-positive cells with dendritic shape (arrow) were seen in the superficial cortex of the tracheobronchial lymph nodes in OVA-challenged mice (b). Lymphocytes, mainly located in the cortex, were also stained for MIF, although less intensely (c). In deeper portions of the lymph nodes many cells were unstained (c). A multifocal perivascular and peribronchial MPO-positive (neutrophilic) distribution (arrows) was seen at 4 h after intratracheal LPS-challenge (d). In (a) and (d) a slight yellowish auto-fluorescence, mainly located in the lung parenchyma, is observed. B = bronchus, V = blood vessel. Original magnification $\times 250$.

neutrophils in BALF was strikingly increased at both 4 h and 24 h after LPS challenge ($77.3 \pm 2.7\%$ and $53.1 \pm 5.9\%$, respectively, vs. $6.8 \pm 4.3\%$ and $8.3 \pm 5.9\%$, respectively, in SAL-challenged controls, $P < 0.001$). The percentage of lymphocytes remained low 4 h after LPS challenge ($0.8 \pm 0.2\%$ vs. $1.4 \pm 0.3\%$ in SAL-challenged controls). However, an increase of lymphocytes was observed 24 h after LPS instillation ($2.3 \pm 0.5\%$ vs. $0.9 \pm 0.3\%$ in SAL-challenged controls, $P < 0.05$).

The percentage of polymorphonuclear leukocytes (PMN) in peripheral blood increased rapidly after intratracheal LPS instillation ($48.8 \pm 3.1\%$ vs. $21.2 \pm 3.1\%$ in SAL-challenged controls at the 4 h time point, $P < 0.001$). At 24 h after LPS instillation the percentage of PMN in peripheral blood had returned to baseline levels ($23.1 \pm 1.6\%$ vs. $20.1 \pm 3.5\%$ in SAL-challenged controls).

Histological analysis of lungs taken 4 hours after LPS challenge demonstrated a moderate neutrophilia perivascularly and peribronchially (Fig. 3d). Neutrophils were also detected in alveolar walls and spaces. A similar, but reduced, distribution of neutrophils was observed 24 hours post LPS instillation (not shown).

High levels of TNF- α were detected in BALF 4 h after LPS challenge (3302.7 ± 437.9 pg/ml versus 17.0 ± 7.0 pg/ml in SAL-challenged controls, $P < 0.001$). A small increase in levels of TNF- α could still be detected 24 h after LPS challenge (16.3 ± 3.1 pg/ml vs. 5.3 ± 0.2 pg/ml in SAL-challenged controls, $P < 0.01$). Intratracheal instillation of LPS also induced a significant increase of MIP-1 α levels in BALF 4 h after challenge (3942.6 ± 280.4 pg/ml vs. 26.2 ± 14.9 pg/ml in SAL-challenged controls, $P < 0.001$).

Effect of anti-MIF treatment on LPS-induced airway inflammation

To examine the role of MIF in LPS-induced airway inflammation, cellular profile in BALF and neutrophilia in lung tissue and blood were determined in anti-MIF-treated and NRS-treated mice. At 4 h and 24 h after intratracheal instillation of a low dose LPS total cellular content (data not shown) and the percentage of neutrophils, lymphocytes, and macrophages in BALF were similar in LPS-challenged anti-MIF-treated and NRS-treated mice (Fig. 4a and b). Consistent with the findings in BALF, the number of PMN in blood did not differ significantly between anti-MIF-treated and NRS-treated animals at either time points (Fig. 4a and b). Also, no obvious difference in lung tissue neutrophilia was observed between anti-MIF-treated and NRS-treated animals at 4 h and 24 h after LPS challenge (data not shown).

Since MIF has been reported to modulate the expression of TNF- α and chemokines in models of




FIG. 4. Cellular composition of BALF and percentage of PMN in blood of anti-MIF-treated and NRS-treated mice 4 h (a) and 24 h (b) after LPS challenge. Data are presented as mean \pm SEM. NRS = NRS-treated mice ($n=8$), anti-MIF = anti-MIF-treated mice ($n=8$). Eos. = eosinophils, Neutro. = neutrophils, Lymph. = lymphocytes, Macro. = macrophages. At both 4 h (a) and 24 h (b) after LPS challenge, a similar cellular composition of BALF was demonstrated in anti-MIF-treated and NRS-treated mice. Consistent with the findings in BALF, the number of PMN in blood did not differ significantly between anti-MIF-treated and NRS-treated animals 4 h (a) and 24 h (b) after LPS challenge. (Epithelial cells and eosinophils in BALF are not included in the figures.)

endotoxaemia and acute lung injury,^{11,16} TNF- α and MIP-1 α levels were measured in LPS-challenged anti-MIF-treated and NRS-treated mice. At 4 h after LPS instillation mice treated with anti-MIF serum exhibited similarly increased levels of TNF- α and MIP-1 α in BALF as NRS-treated mice (Fig. 5). Equally low levels of TNF- α in BALF were observed in anti-MIF-treated and NRS-treated mice at 24 h after LPS challenge (12.5 ± 2.6 pg/ml and 15.9 ± 4.8 pg/ml, respectively). Plasma levels of TNF- α were below detection limit in both anti-MIF-treated and NRS-treated mice at 4 h after LPS challenge (data not shown).




FIG. 5. Levels of TNF- α and MIP-1 α in BALF 4 h after LPS instillation of mice treated with anti-MIF serum or NRS. Data are presented as mean \pm SEM. NRS = NRS-treated animals ($n=8$), anti-MIF = anti-MIF-treated animals ($n=8$). Mice treated with anti-MIF serum exhibited similar levels of TNF- α and MIP-1 α in BALF as NRS-treated mice.

Effect of intratracheal anti-MIF treatment on LPS-induced airway inflammation

LPS-challenged mice treated with topical intratracheal anti-MIF serum or NRS exhibited a similar neutrophilia in BALF at the 4-h time point ($86.5 \pm 1.4\%$ and $82.4 \pm 2.0\%$ neutrophils in BALF, respectively). Also 24 h after LPS challenge no significant difference in BALF neutrophilia was observed between the intratracheally treated mice. Anti-MIF-treated and NRS-treated mice exhibited $68.0 \pm 1.6\%$ and $71.0 \pm 4.8\%$ neutrophils in BALF, respectively. The BALF neutrophilia in mice receiving intratracheal doses of anti-MIF serum or NRS together with LPS was somewhat increased when compared with that observed in BALF from animals receiving LPS challenge only. In accord, SAL-challenged mice treated intratracheally with NRS demonstrated a mild neutrophilia at both the 4- and 24-h time points (35.9 ± 13.0 and $39.4 \pm 11.8\%$ neutrophils in BALF, respectively).

Discussion

This study demonstrates that anti-MIF treatment does not have any major effects on the eosinophil-rich airway inflammation occurring in a murine model of allergic asthma. Similarly, anti-MIF treatment did not change the neutrophilic inflammatory response seen after instillation of a low dose of LPS into the lower airways of mice. Although we cannot exclude the possibility that MIF may regulate other indices of pulmonary inflammation than measured in this study, the present data do not support the view that MIF is critically involved in pulmonary eosinophilic or neutrophilic inflammatory conditions.

Human asthma is characterized by peribronchial inflammatory infiltrates, mainly consisting of eosinophils, T-lymphocytes, and macrophages.^{15,29} Given that all these immune cells express MIF in significant quantities, and that MIF has proinflammatory effects, MIF has been implicated in development of asthma and other inflammatory airway diseases.^{9,14} Moreover, IL-5, a cytokine considered pivotal for the recruitment of eosinophils to the airways in both human asthma and allergic mice, induces MIF secretion by cultured eosinophils.⁹ Indeed, abnormally high levels of MIF have been detected in BALF from asthmatic subjects.⁹ In similarity to human asthma, the present allergic mice exhibit peribronchial and perivascular infiltrates of eosinophils, T-lymphocytes, and macrophages.^{25,26} The present study also showed the presence of MIF-positive cells in the pulmonary infiltrates. In addition, an intriguing distribution of MIF-positive cells was observed in the tracheobronchial lymph nodes of the allergic mice, a location where essential immune responses to antigens take place. Hence, this allergic model would be well suited for exploration of anti-inflammatory efficacy of anti-MIF active compounds.

It was hypothesized that anti-MIF treatment might inhibit the allergic responses in the present asthma model both by the enhancement of anti-inflammatory effects of endogenous corticosteroids no longer counter-regulated by MIF and by diminished production of MIF-inducible proinflammatory cytokines. However, immunoneutralization of MIF during the period of allergen aerosol challenge did not influence the magnitude of pulmonary eosinophilia and cellular composition of inflammatory cells obtained by BAL. Hence, the present study failed to support the view that MIF is a critical regulator of pulmonary eosinophilic inflammation.

Instillation of a low dose of LPS into the airways of mice proved sufficient for producing consistent, predominantly neutrophilic pulmonary inflammation with moderate neutrophilia perivascularly and peribronchially, but also involving alveolar walls and spaces. These features are reminiscent of the pathology of COPD.²⁹ In addition, chronic exposure to endotoxin, as a component of organic dust in occupational settings, has been related to development of COPD-like conditions.³⁰⁻³² The potential relevance of the present model to human COPD may further be supported by studies suggesting that inhaled LPS, as a constituent of cigarette smoke, is of importance for the development of COPD.³³ The present demonstration of LPS-induced increases in BALF TNF- α and MIP-1 α levels is also of interest in relation to COPD, since these two mediators are potentially important for pulmonary neutrophil recruitment.^{18,34,35} These findings together with an increase in the percentage of lymphocytes in BALF and lack of eosinophilia, support the possibility that

the present LPS challenge produced a potentially useful model of COPD.²⁹

In contrast to most cytokines, MIF mRNA and protein are expressed constitutively in a variety of cell types, such as monocytes/macrophages, T-cells, airway epithelial cells, and pituitary endocrine cells. Proinflammatory stimuli, including LPS, are known to increase MIF mRNA expression above the level present constitutively.^{8,12} LPS administered systemically (i.p.) in high doses has been used in earlier studies to elucidate the role of MIF in endotoxaemia and acute lung injury.^{1,11,12,16} For example, treatment of mice with anti-MIF serum conferred full protection to the lethal effects of LPS (17.5 mg/kg) administered i.p.¹ Consistent with these previous findings and confirming the validation of the employed anti-MIF serum, the present study showed that anti-MIF treatment inhibits plasma extravasation in response to a high systemic dose of LPS.

In a rat model of acute lung injury, Makita *et al.*¹¹ demonstrated that anti-MIF treatment reduced the number of neutrophils per alveolus and the BALF neutrophilia, induced by a high dose of LPS given systemically. In mice, i.p. administration of LPS (1–20 mg/kg) does not induce transpulmonary neutrophil migration and infiltration of neutrophils into the alveolar space, but only neutrophil sequestration within the lung vasculature.^{36–38} In this study, a fraction of the previously employed i.p. doses of LPS (250–5000-fold less) was given locally into the airways. Reflecting the low dose and route of administration, plasma levels of TNF- α were below detection limit at 4 h after LPS challenge. In apparent contrast to the important role of MIF in host responses to high systemic doses of LPS, anti-MIF treatment did not change the neutrophil-rich inflammatory response induced by this mode of LPS exposure. To explain these data it is suggested that the importance of MIF in different models of LPS-induced host reactions may vary depending on the dose and/or the administration route of LPS. In accord, it has previously been shown that the mechanisms behind host responses to LPS may be completely different in models using high or low doses of LPS.³⁹ Genetically MIF-deficient mice are also resistant to the lethal effects of a high systemic dose of LPS, but susceptible to a combination of a low dose of LPS and D-galactosamine.¹⁶ The demonstration that MIF is not involved in the present pulmonary neutrophilic inflammation may reduce the promise of anti-MIF compounds as future COPD drugs.

In conclusion, the present data suggest that MIF is not critically important for allergic eosinophilic, or LPS-induced neutrophilic inflammation, in airways of mice. If translatable to human disease conditions,⁴⁰ these findings do not support the notion that MIF inhibitors will be effective against eosinophilic or neutrophilic respiratory diseases, such as allergic asthma and COPD.

Acknowledgements

We thank Dr Thierry Calandra, Lausanne, Switzerland, for providing control and anti-MIF serum. This work was supported by the Swedish Medical Research Council (V1180, 06P-11813, 8308, 4499), Vårdalstiftelsen, AstraZeneca, the Swedish Heart and Lung Foundation, and the Swedish Society for Medical Research.

References

- Bernhagen J, Calandra T, Mitchell RA, Martin SB, Tracey KJ, Voelker W, Manogue KR, Cerami A, Bucala R. MIF is a pituitary-derived cytokine that potentiates lethal endotoxemia. *Nature* 1993; **365**: 756–759.
- Bernhagen J, Bacher M, Calandra T, Metz CN, Doty SB, Donnelly T, Bucala R. An essential role for macrophage migration inhibitory factor in the tuberculin delayed-type hypersensitivity reaction. *J Exp Med* 1996; **183**: 277–282.
- Mikulowska A, Metz CN, Bucala R, Holmdahl R. Macrophage migration inhibitory factor is involved in the pathogenesis of collagen type II-induced arthritis in mice. *J Immunol* 1997; **158**: 5514–5517.
- Donnelly SC, Haslett C, Reid PT, Grant IS, Wallace WAH, Metz CN, Bruce LJ, Bucala R. Regulatory role for macrophage migration inhibitory factor in acute respiratory distress syndrome. *Nature Med* 1997; **3**: 320–323.
- David J. Delayed hypersensitivity in vitro: its mediation by cell free substances formed by lymphoid cell-antigen interaction. *Proc Natl Acad Sci USA* 1966; **56**: 72–77.
- Bloom BR, Bennett B. Mechanism of a reaction in vitro associated with delayed type hypersensitivity. *Science* 1966; **153**: 80–82.
- Bacher M, Metz CN, Calandra T, Mayer K, Chesney J, Lohoff M, Gemsa D, Donnelly T, Bucala R. An essential regulatory role for macrophage migration inhibitory factor in T-cell activation. *Proc Natl Acad Sci USA* 1996; **93**: 7849–7854.
- Calandra T, Bernhagen J, Mitchell RA, Bucala R. The macrophage is an important and previously unrecognized source of macrophage migration inhibitory factor. *J Exp Med* 1994; **179**: 1895–1902.
- Rossi AG, Haslett C, Hirani N, Greening AP, Rahman I, Metz CN, Bucala R, Donnelly SC. Human circulating eosinophils secrete macrophage migration inhibitory factor (MIF). Potential role in asthma. *J Clin Invest* 1998; **101**: 2869–2874.
- Wymann D, Bluggel M, Kalbacher H, Blesken T, Akdis CA, Meyer HE, Blaser K. Human B cells secrete migration inhibition factor (MIF) and present a naturally processed MIF peptide on HLA-DRB1*0405 by a FXXL motif. *Immunology* 1999; **96**: 1–9.
- Makita H, Nishimura M, Miyamoto K, Nakano T, Tanino Y, Hirokawa J, Nishihira J, Kawakami Y. Effect of anti-macrophage migration inhibitory factor antibody on lipopolysaccharide-induced pulmonary neutrophil accumulation. *Am J Respir Crit Care Med* 1998; **158**: 573–579.
- Bacher M, Meinhardt A, Lan HY, Mu W, Metz CN, Chesney JA, Calandra T, Gemsa D, Donnelly T, Atkins RC, Bucala R. Macrophage inhibitory factor expression in experimentally induced endotoxemia. *Am J Pathol* 1997; **150**: 235–246.
- Calandra T, Bernhagen J, Metz CN, Spiegel IA, Bacher M, Donnelly T, Cerami A, Bucala R. MIF as a glucocorticoid-induced modulator of cytokine production. *Nature* 1995; **377**: 68–71.
- Donnelly SC, Bucala R. Macrophage migration inhibitory factor: a regulator of glucocorticoid activity with a critical role in inflammatory disease. *Mol Med Today* 1997; **3**: 502–507.
- Bradley BL, Azzawi M, Jacobson M, Assoufi B, Collins JV, Irani AMA, Schwartz LB, Durham SR, Jeffery PK, Kay AB. Eosinophil, T lymphocytes, mast cells, neutrophils, and macrophages in bronchial biopsy specimens from atopic subjects with asthma: comparison with biopsy specimens from atopic subjects without asthma and normal control subjects and relationship to bronchial hyperresponsiveness. *J Allergy Clin Immunol* 1991; **88**: 661–674.
- Bozza M, Satoskar AR, Lin G, Lu B, Humbles AA, Gerard C, David JR. Targeted disruption of migration inhibitory factor gene reveals its critical role in sepsis. *J Exp Med* 1999; **189**: 341–346.
- Bernhagen J, Mitchell RA, Calandra T, Voelker W, Cerami A, Bucala R. Purification, bioactivity, and secondary structure analysis of mouse and human macrophage migration inhibitory factor (MIF). *Biochemistry* 1994; **33**: 14144–14155.
- Barnes PJ. Chronic obstructive pulmonary disease: new opportunities for drug development. *Trends Pharmacol Sci* 1998; **19**: 415–423.
- Pesci A, Majori M, Cuomo A, Borciani N, Bertacco S, Cacciani G, Gabrielli M. Neutrophils infiltrating bronchial epithelium in chronic obstructive pulmonary disease. *Respir Med* 1998; **92**: 863–870.
- Brusselle GJ, Kips JC, Tavernier JH, Van der Heyden JG, Cuvelier CA, Pauwels RA, Bluethmann H. Attenuation of allergic airway inflammation in IL-4 deficient mice. *Clin Exp Allergy* 1994; **24**: 73–80.

21. Rylander R. Health effects of cotton dust exposures. *Am J Ind Med* 1990; **17**: 39–45.
22. Nightingale JA, Rogers DF, Hart LA, Kharitonow SA, Chung KF, Barnes PJ. Effect of inhaled endotoxin on induced sputum in normal, atopic, and atopic asthmatic subjects. *Thorax* 1998; **53**: 563–571.
23. Calandra T, Spiegel LA, Metz CN, Bucala R. Macrophage migration inhibitory factor is a critical mediator of the activation of immune cells by exotoxins of Gram-positive bacteria. *Proc Natl Acad Sci USA* 1998; **95**: 11383–11388.
24. Ten RM, Pease LR, McKean DJ, Bell MP, Gleich GJ. Molecular cloning of the eosinophil peroxidase: evidence for the existence of a peroxidase multigene family. *J Exp Med* 1989; **169**: 1757–1769.
25. Korsgren M, Erjefält JS, Korsgren O, Sundler F, Persson CGA. Allergic eosinophil-rich inflammation develops in lungs and airways of B cell-deficient mice. *J Exp Med* 1997; **185**: 885–892.
26. Korsgren M, Persson CGA, Sundler F, Bjerke T, Hansson T, Chambers BJ, Hong S, Van Kaer L, Ljunggren H-G, Korsgren O. Natural killer cells determine development of allergen-induced eosinophilic airway inflammation in mice. *J Exp Med* 1999; **189**: 553–562.
27. Parrillo JE. Pathogenetic mechanisms of septic shock. *N Engl J Med* 1993; **328**: 1471–1477.
28. Bauss F, Dröge W, Männel DN. Tumor necrosis factor mediates endotoxic effects in mice. *Infect Immun* 1987; **55**: 1622–1625.
29. Jeffery PK. Pathology of asthma and COPD: a synopsis. *Eur Respir Rev* 1997; **43**: 111–118.
30. Smid T, Heederik D, Houba R, Quanjer PH. Dust- and endotoxin-related respiratory effects in the animal feed industry. *Am Rev Respir Dis* 1992; **146**: 1474–1479.
31. Schwartz DA, Donham KJ, Olenchock SA, Popendorf WJ, Van Fossen DS, Burmeister LF, Merchant JA. Determinants of longitudinal changes in spirometric function among swine confinement operators and farmers. *Am J Respir Crit Care Med* 1995; **151**: 47–53.
32. Schwartz DA, Thorne PS, Yagla SJ, Burmeister LF, Olenchock SA, Watt JL, Quinn TJ. The role of endotoxin in grain dust-induced lung disease. *Am J Respir Crit Care Med* 1995; **152**: 603–608.
33. Hasday JD, Bascom R, Costa JJ, Fitzgerald T, Dubin W. Bacterial endotoxin is an active component of cigarette smoke. *Chest* 1999; **115**: 829–835.
34. Goncalves de Moraes VL, Vargaftig BB, Lefort J, Meager A, Chignard M. Effect of cyclo-oxygenase inhibitors and modulators of cyclic AMP formation on lipopolysaccharide-induced neutrophil infiltration in mouse lung. *Br J Pharmacol* 1996; **117**: 1792–1796.
35. Christman JW, Blackwell TR, Cowan HB, Shepard VL, Rinaldo JE. Endotoxin induces the expression of macrophage inflammatory protein 1 α mRNA by rat alveolar and bone marrow-derived macrophages. *Am J Respir Cell Mol Biol* 1992; **7**: 455–461.
36. Cardozo C, Edelman J, Jagirdar J, Lesser M. Lipopolysaccharide-induced pulmonary vascular sequestration of polymorphonuclear leukocytes is complement independent. *Am Rev Respir Dis* 1991; **144**: 173–178.
37. Hirano S. Migratory responses of PMN after intraperitoneal and intratracheal administration of lipopolysaccharide. *Am J Physiol* 1996; **270**: L836–L845.
38. Lefort J, Singer M, Leduc D, Renesto P, Nahori MA, Huerre M, Créminon C, Chigard M, Vargaftig BB. Systemic administration of endotoxin induces bronchopulmonary hyperreactivity dissociated from TNF- α formation and neutrophil sequestration into the murine lungs. *J Immunol* 1998; **161**: 474–480.
39. Gutierrez-Ramos JC, Bluethmann H. Molecules and mechanisms operating in septic shock: lessons from knockout mice. *Immunol Today* 1997; **18**: 329–334.
40. Persson CGA, Erjefält JS, Korsgren M, Sundler F. The mouse trap. *Trends Pharmacol Sci* 1997; **18**: 465–467.

**Received 13 February 2000;
accepted 1 March 2000**



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

