Maturation of a hypothesis

Fernando D. Martinez

Respiratory Sciences Center, The University of Arizona, 1501 N Campbell, Avenue, Suite 2349, Tucson, AZ 85724-5030, USA

Tel: +1 520 626 6387

Fax: +1 520 626 6970

E-mail: fernando@resp-sci.arizona.edu

Recent findings of associations between markers of exposure to microbial products that do not directly cause disease and the risk of having asthma and allergies have significantly changed and enriched our understanding of the so-called hygiene hypothesis.

This hypothesis was originally presented as the proposed increased burden of atopy that Western societies paid for a dramatic decrease in infectious diseases during the first years of life.¹ This enunciation, provocative and trail-blazing, as it was, suggested a more or less direct relation between the occurrence of acute episodes of infectious disease and the subsequent development of allergies. Many researchers thus studied associations between the incidence of such episodes and subsequent asthma and allergies, but the results were contradictory.²⁻⁵ It became clear that some acute infections in early life could be protective, but this may depend on other factors that went unmeasured.

More recently, studies performed in rural and urban areas of developed countries have considerably widened the scope of the original hygiene hypothesis. Surveys in rural communities have consistently found that children living on animal farms have considerably less burden of atopic diseases than those living in the same communities but away from farms.⁶⁻⁸ Moreover, children who frequently visit farms but do not live on them also have a lower prevalence of allergies.⁸ More in-depth studies of the indoor environment in animal-farm homes suggest that the endotoxin content in these homes is much higher than that in rural homes away from animal farms.⁹ A recent study in an urban environment in the USA suggests that exposure to indoor endotoxin (as measured in house dust samples) is inversely associated with the risk of developing allergic sensitization in infants and young children, and is also associated with a higher proportion of interferon- γ -producing T cells in circulation.

A broader approach is thus needed for our understanding of the environmental factors that may influence the development of the immune system. Endotoxin contains lipopolysaccharides (LPS) that form part of the cell wall of Gram-negative bacteria. A very sensitive mechanism that responds to LPS is present in vertebrate immune systems. This receptor system constitutes the first, non-adaptive response to

external microbial stimuli. It activates a complex intercellular signaling mechanism and triggers the production of a set of immune mediators by antigen presenting cells. It is possible that these mediators, when produced by co-stimulation with perennial exposure to endotoxin, may deviate or modify the original responses to allergens in such a way that drives them away from T helper cell (Th)-2-type responses mediated by immunoglobulin E (IgE).

Based on these premises, and in collaboration with Holt, we speculated that polymorphisms in the receptor system for LPS could influence the risk of allergies. We have thus screened the genes involved in this receptor system for LPS, starting with CD14 gene, which codes for one of the main components of the endotoxin receptor system.¹⁰ We reported a C $\rightarrow$ T variation at position -159 of the promoter region of the gene (CD14/-159). We found that, both in Caucasians and Hispanics, one-half of all chromosomes contain one or the other allele (C or T). Carriers of the T allele in homozygote form had significantly higher levels of the circulating soluble form of CD14. We also found that skin-test-positive homozygotes for the T allele had significantly lower levels of total serum IgE, and atopic carriers of the T allele had significantly lower numbers of positive skin tests than carriers of the C allele. Similar findings have been reported by two other research groups,^{11,12} although not all researchers have been able to confirm our findings.^{13,14} It thus appears that, at least in some populations and at some ages, polymorphisms that increase the expression of CD14 may be associated with lower levels of IgE. A potential explanation for this finding could be that increased sensitivity to endotoxin and other microbial products that interact with CD14 could increase interleukin-12-mediated and interleukin-10 responses to these products, with increased likelihood of development of Th1-type responses or modified Th-2 responses and thus with less likelihood of atopic diseases.

These findings may allow us to assess the role of genetic variants in genes of the receptor system for endotoxin as determinants of susceptibility allergies in individuals who are exposed to different levels of endotoxin. Our final goal is to design prevention strategies based on exposure to innocuous surrogates of endotoxin in individuals who may be more susceptible to the preventive effects of bacterial products because of their specific genetic background.

References

1. Gerrard JW, Geddes CA, Reggin PL, Gerrard CD, Horne S. Serum IgE levels in white and Metis communities in Saskatchewan. *Ann Allergy* 1976; 37: 91-100.
2. Matricardi PM, Rosmini F, Riondino S, Fortini M, Ferrigno L, Rapicetta M, Bonini S. Exposure to foodborne and orofecal microbes versus airborne viruses in relation to atopy and allergic asthma: epidemiological study. *Br Med J* 2000; 320: 412-417.

3. Martinez FD, Stern DA, Wright AL, Taussig LM, Halonen M, Associates GHM. Association of non-wheezing lower respiratory tract illnesses in early life with persistently diminished serum IgE levels. *Thorax* 1995; **50**: 1067-1072.
4. Shaheen SO, Aaby P, Hall AJ, Barker DJ, Heyes CB, Shiell AW, Goudiaby A. Measles and atopy in Guinea-Bissau. *Lancet* 1996; **347**: 1792-1796.
5. Paunio M, Heinonen OP, Virtanen M, Leinikki P, Patja A, Peltola H. Measles history and atopic diseases: a population-based cross-sectional study. *J Am Med Assoc* 2000; **283**: 343-346.
6. Von Ehrenstein OS, Von Mutius E, Illi S, Baumann L, Bohm O, von Kries R. Reduced risk of hay fever and asthma among children of farmers. *Clin Exp Allergy* 2000; **30**: 187-193.
7. Braun-Fahrlander C, Gassner M, Grize L, *et al*. Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. SCARPOL team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution. *Clin Exp Allergy* 1999; **29**: 28-34.
8. Riedler J, Eder W, Oberfeld G, Schreuer M. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin Exp Allergy* 2000; **30**: 194-200.
9. Von Mutius E, Braun-Fahrlander C, Schierl R, *et al*. Exposure to endotoxin or other bacterial components might protect against the development of atopy. *Clin Exp Allergy* 2000; **30**: 1230-1234.
10. Baldini M, Lohman IC, Halonen M, Erickson RP, Holt PG, Martinez FD. A polymorphism in the 5' flanking region of the CD14 gene is associated with circulating soluble CD14 levels and with total serum IgE. *Am J Respir Cell Mol Biol* 1999; **20**: 976-983.
11. Amelung PJ, Weisch DG, Xu J, Paigen B, Meyers DA, Bleecker ER. A polymorphism in CD14 is associated with high IgE levels in a population with laboratory animal allergy. *Am J Respirat Crit Care Med* 2000; **161**: A927.
12. O'Donnell AR, Hayden CM, Laing IA, *et al*. Association study of CC16 and CD14 polymorphisms in an unselected population assessed at age 8 and 25. *Am J Respirat Crit Care Med* 2000; **161**: A928.
13. Joos L, Zhu S, Becker A, *et al*. Polymorphisms of the CD14 and TGF β 1 genes in a cohort of infants at high risk of allergic disorders. *Am J Respirat Crit Care Med* 2000; **161**: A928.
14. Celedon JC, Soto-Quiros M, Drazen JM, *et al*. Lack of association between a polymorphism in the 5' flanking region of the CD14 gene and skin test reactivity to allergens in the central valley of Costa Rica. *Am J Respirat Crit Care Med* 2000; **161**: A927.

Edited by Dr. F. J. Zijlstra; accepted 24 September 2001