It has recently been reported that phycocyanin, a biliprotein found in the blue–green microalgae *Spirulina*, exerts anti-inflammatory effects in some animal models of inflammation. Taking into account these findings, we decided to elucidate whether phycocyanin might exert also inhibitory effects in the induced allergic inflammatory response and on histamine release from isolated rat mast cells. In *in vivo* experiments, phycocyanin (100, 200 and 300 mg/kg post-orally (p.o.)) was administered 1 h before the challenge with 1 µg of ovalbumin (OA) in the ear of mice previously sensitized with OA. One hour later, myeloperoxidase activity and ear edema were assessed. Phycocyanin significantly reduced both parameters. In separate experiments, phycocyanin (100 and 200 mg/kg p.o.) also reduced the blue spot area induced by intradermal injections of histamine, and the histamine releaser compound 48/80 in rat skin. In concordance with the former results, phycocyanin also significantly reduced histamine release induced by compound 48/80 from isolated peritoneal rat mast cells. The inhibitory effects of phycocyanin were dose dependent. Taken together, our results suggest that inhibition of allergic inflammatory response by phycocyanin is mediated, at least in part, by inhibition of histamine release from mast cells.

Key words: Phycocyanin, Inhibition, Histamine release, Allergic inflammatory response, Rat mast cells

Role of histamine in the inhibitory effects of phycocyanin in experimental models of allergic inflammatory response

D. Remirez, N. Ledón and R. González^{CA}

Ozone Research Center, National Center for Scientific Research, Post Box 6412, Havana, Cuba

^{CA}Corresponding Author

Fax: +53 7 210233

E-mail: ozono@infomed.sld.cu

Introduction

We recently reported that phycocyanin, a biliprotein found in the blue–green microalgae *Spirulina* exerts antioxidative and anti-inflammatory effects in various *in vitro* and *in vivo* experimental models.¹ These effects are mediated by the scavenging action of phycocyanin against reactive oxygen species (ROS) such as OH[•], RO[•] and RO₂, anti-lipoperoxidative activity, and inhibitory effects on prostaglandin and leukotriene biosynthesis.^{2,3} Recently, an inhibitory effect of *Spirulina* on mast cell-mediated immediate-type allergic reactions in rats has also been reported.⁴ Thus, taking into account the former findings and the fact that phycocyanin accounts for more than 20% of algae dry weight, we decided to elucidate whether or not both allergic inflammatory response and histamine release from rat mast cells are inhibited by phycocyanin.

Materials and methods

Chemicals

Ovalbumin, *o*-dianisidine and compound 48/80 were purchased from Sigma (St Louis, MO, USA).

Orthophthalaldehyde was obtained from Merck (Barcelona, Spain).

Other reagents of analytical grade were purchased from normal commercial sources.

Preparation of phycocyanin extract

Phycocyanin was extracted from microalgae *Arthrospira maxima* as described in a Cuban patent.⁵ The blue powder thus obtained showed a peak in the visible absorption spectrum of 617 nm, which is very close to that reported for c-phycocyanin.⁶

Animals

Male Balb/c mice (20–25 g) and male Sprague–Dawley rats (175–250 g) were obtained from the breeding colony of the National Center for Laboratory Animal Production (CENPALAB, Havana, Cuba). The animals were housed under a 12 h light–dark cycle with room temperature maintained at 25°C, humidity at 60%, and food and water available *ad libitum*. The experiments were conducted in accordance with the ethical guidelines for investigations in laboratory animals and were approved by the Ethical Committee for Animal Experimentation of the National Center for Scientific Research.

Sensitization of mice and ear swelling response to ovalbumin

This test was carried out as described by Hessel *et al.*⁷ Active sensitization was performed by seven intra-peritoneal injections of 10 µg of ovalbumin in 0.5 ml of saline on alternate days. A group of animals received injections with saline only.

Four weeks after the last injection, the mice were divided into groups of seven animals and treated by gavage with phycocyanin (100, 200 and 300 mg/kg), with triamcinolone (10 mg/kg) used as a reference drug or saline. One hour later, the animals were challenged intracutaneously with 1 µg of ovalbumin in 0.01 ml of saline on the right ear and saline only on the left ear. Inflammation was followed for 1 h and thereafter animals were killed by cervical dislocation. A 6 mm diameter disc from each ear was removed with a metal punch and weighed. The swelling induced by ovalbumin was assessed as the increase in the weight of the right ear punch biopsy over that of the left ear and was called the edema index.

Determination of myeloperoxidase activity

Tissue samples were assessed biochemically for the neutrophil marker enzyme myeloperoxidase (MPO), using the method of Bradley *et al.*⁸ All the ear tissue was homogenized in 50 mM $\text{K}_2\text{HPO}_4/\text{KH}_2\text{PO}_4$ buffer (pH 6) containing 0.5% hexadecyl trimethylammonium bromide using a polytron (Ultra turrax x-25) homogenizer. After freeze-thawing three times, the samples were centrifuged at $2500 \times g$ for 30 min at 4°C and aliquots of the supernatant assayed spectrophotometrically for MPO. In brief, 40 µl of sample was mixed with 960 µl of 50 mM phosphate buffer (pH 6) containing 0.167 mg/ml of *o*-dianisidine dihydrochloride and 0.0005% hydrogen peroxide. The change in absorbance at 460 nm was measured with a Spekol 220 spectrophotometer (Carl Zeiss, Germany). MPO activity data are presented as units per milligram of tissue. One unit of MPO activity was defined as that degrading one micromole of hydrogen peroxide to water per minute at 25°C.

Skin reactions to histamine and compound 48/80 in rats

The method described by Awouters *et al.*⁹ was used. Groups of five rats were pre-treated with phycocyanin (50, 100 and 200 mg/kg per-orally (p.o.)), saline and promethazine (25 mg/kg) as positive control.

One hour thereafter, the animals were injected by duplicate into the shaved dorsal skin with histamine (50 µg) and the histamine releaser compound 48/80 (50 µg), and immediately afterwards also injected intravenously with 1 ml of 0.4% Evans blue dye dissolved in saline.

After 30 min, the rats were euthanized, skinned and the positive blue skin reactions were measured to determine their area.

Histamine release from isolated rat peritoneal mast cells

Experiments were performed using mixed peritoneal cells of the rat (1–5% mast cells) prepared as described by Garland and Mongar.¹⁰

The peritoneal washings from at least five rats were pooled and kept on ice until used. The washings were centrifuged at $250 \times g$ for 6 min, the supernatant was discarded and the cell pellet resuspended in cold Tyrode solution (3 ml per rat) of the following composition: NaCl, 137 mM; NaHCO_3 , 12 mM; NaH_2PO_4 , 0.3 mM; KCl, 2.7 mM; MgCl_2 , 1 mM; CaCl_2 , 1.8 mM; dextrose, 5.6 mM.

The cell suspension (0.5 ml aliquots in duplicate) was pre-incubated at 37°C for 10 min with three concentrations of phycocyanin (3, 5 and 8 mg/ml) and stimulated for a further 5 min with compound 48/80 (1 µg/ml). The reaction was then stopped by placing the tubes in an ice bath. Each sample was centrifuged at $250 \times g$ for 6 min, the supernatant was decanted, and the pellets resuspended in 1 ml of Tyrode solution and boiled for 10 min to release residual histamine. Release and residual histamine were measured fluorimetrically.¹¹

The results are expressed as the percentage of histamine released from the total content of mast cells. Histamine release values were corrected for a spontaneous release of between 2 and 8%. Preliminary assays demonstrated that phycocyanin did not interfere with fluorimetric histamine determination.

Statistical analysis

Data are presented as mean $\pm$ SEM. Mean differences between groups were compared by one-way analysis of variance with the Duncan multi-range test. In experiments with rat peritoneal mast cells, Student's *t*-test was used to make a statistical comparison between the mean values. The level of statistical significance was taken as $p < 0.05$.

Results

Effects of phycocyanin on mouse ear swelling response to ovalbumin

One hour after ovalbumin challenge, a marked increase in both ear thickness (4.70 ± 0.16 mg) and MPO activity (3.13 ± 0.21 U/mg of tissue) were observed in ovalbumin-sensitized mice (Table 1). Pre-treatment with phycocyanin (100, 200 and 300 mg/kg p.o.) 1 h before ovalbumin challenge significantly inhibited ear edema and MPO activity in this model.

Table 1. Effect of phycocyanin on mouse ear swelling response to ovalbumin

Treatment	Edema index (mg)	Inhibition (%)	MPO (U/mg of tissue)	Inhibition (%)
Saline (non-sensitized)	0.15 ± 0.05*	–	0.14 ± 0.05*	–
Ovalbumin controls	4.70 ± 0.16*	–	3.13 ± 0.21	–
Triamcinolone (10 mg/kg)	0.25 ± 0.05*	98	0.23 ± 0.09*	93
Phycocyanin (100 mg/kg)	3.80 ± 0.43*	19	2.45 ± 0.46*	22
Phycocyanin (200 mg/kg)	2.28 ± 0.19*	52	1.51 ± 0.24*	52
Phycocyanin (300 mg/kg)	1.78 ± 0.40*	62	1.07 ± 0.14*	66

Values are expressed as mean ± SEM. Ten mice per group. The edema index was assessed as the increase in the weight of the right ear punch biopsy over that of the left ear. Edema was measured 1 h after ovalbumin challenge.

* $p < 0.05$, statistically significant with respect to ovalbumin control group.

The inhibitory effect of phycocyanin was dose dependent. Triamcinolone (10 mg/kg) also induced remarkable reduction on edema index and MPO activity (Table 1).

Effect of phycocyanin on skin reactions induced by intradermal injections of histamine and compound 48/80 in rats

Pre-treatment with phycocyanin (100 and 200 mg/kg p.o.) significantly inhibited, in a dose-dependent manner, the skin reactions induced by compound 48/80 and histamine. The lower dose of phycocyanin (50 mg/kg) did not induce significant inhibitory effect. The histamine H_1 -receptor antagonist promethazine (25 mg/kg) produced a strong inhibition of skin reaction induced by compound 48/80 and histamine (Fig. 1).

Effect of phycocyanin on histamine release from isolated rat peritoneal mast cells

Phycocyanin (3, 5, and 8 mg/ml) significantly inhibited compound 48/80-induced histamine release from rat peritoneal mast cells following 10 min pre-incubation.

FIG. 1. Effects of phycocyanin and promethazine on histamine and compound 48/80-induced cutaneous reactions in rats. Phycocyanin and promethazine were given orally 1 h before the challenge. Each value represents the mean of five animals. * $p < 0.05$, statistically significant with respect to the controls (histamine and compound 48/80).

The inhibitory effect of phycocyanin was concentration dependent (Fig. 2).

Discussion

Increased microvascular permeability to plasma proteins is one of the characteristic features of the allergic inflammatory response. Mast cells are recognized to play a crucial role in allergen-induced plasma leakage because they can release pre-formed vasoactive mediators, among which the role of histamine and, in the case of rats and mice also of serotonin, has been historically emphasized.^{12,13}

More recently, it has been demonstrated that, besides the former vasoactive amines, mast cells

FIG. 2. Effect of phycocyanin on compound 48/80-induced histamine release from rat peritoneal mast cells. Rat peritoneal mast cells were pre-incubated at 37°C for 10 min with three different concentrations of phycocyanin and stimulated for a further 5 min with compound 48/80 (1 µg/ml). Each value is the mean ± SEM of three experiments performed in duplicate. Significantly different from the control at $p < 0.05$.

contain pre-formed stores of cytokines, such as tumor necrosis factor- α (TNF- α) and vascular permeability factor/vascular endothelial cell growth factor among others, that can be released in immunoglobulin E (IgE)-dependent reactions and can provide a potential effector and immunoregulatory roles to mast cells in allergic inflammatory response.¹⁴ Also, it is important that the participation of the newly synthesized lipid mediators from mast cells such as prostaglandin D₂, leukotrienes (LTC₄, LTD₄, LTE₄, LTB₄) and platelet-activating factor (PAF), as well as ROS, are strongly involved in IgE-dependent allergic inflammation.^{14,15}

In previous studies, we have demonstrated the antioxidant and anti-inflammatory properties of phycocyanin in various *in vitro* and *in vivo* experimental models.^{1,16-18} It was thus also shown that phycocyanin extract reduces LTB₄ and prostaglandin E₂ levels in the arachidonic acid-induced mouse ear inflammation test.^{2,3}

In the present work, we have found that phycocyanin also exerts an anti-inflammatory effect in mouse ear swelling response to ovalbumin in which the various mediators quoted earlier are released in a IgE-mediated reaction in sensitized mice (Table 1).

In this test, ear edema and MPO activity were reduced by phycocyanin. MPO is an enzyme present in neutrophils and at a much lower concentration in monocyte and macrophages. It is well known that the level of MPO activity is directly proportional to the neutrophil concentration on the inflamed tissue,⁸ by which measurement of the enzyme activity has been considered a quantitative and sensitive marker of chemotaxis and neutrophil infiltration in the inflammatory process. Therefore, the inhibitory effect of phycocyanin on MPO activity in ovalbumin-induced ear edema in mice constitutes further evidence of the anti-chemoattractant action of the biliprotein because previously we demonstrated that, in acetic acid-induced colitis in the rat, phycocyanin also reduced MPO activity in colonic tissue, and the histological findings also showed a substantial decrease in neutrophil infiltration in colonic mucosal injury.¹⁸

In our opinion the mechanism by which phycocyanin exerts its anti-chemoattractant action is through inhibitory effects on the biosynthesis of the neutrophil chemotactic mediator LTB₄. This view is supported by our previous findings that phycocyanin reduced LTB₄ levels in the arachidonic acid-induced mouse ear inflammation test.³

To elucidate whether phycocyanin is able to inhibit histamine release, we used the rat skin reaction test to histamine and compound 48/80 as well as the *in vitro* method of histamine release induced by compound 48/80 from isolated rat mast cells.

Promethazine (an H₁-receptor antagonist), and to a lesser extent phycocyanin, inhibited skin reactions to histamine and compound 48/80, respectively (Fig. 1).

Through this observation, the possibility of phycocyanin or its metabolites interacting with H₁-receptors must be considered, although this cannot be elucidated with this kind of experiment and will require further studies with the use of other methods such as radioligands.

On the contrary, the inhibitory effect of phycocyanin on histamine release from isolated rat mast cells (Fig. 2) supports the involvement of this event in the mode of action of the biliprotein as an anti-inflammatory agent, and particularly in the allergic reactions of the immediate type triggered by IgE.

At the present time, evidence is available that ROS such as superoxide anion, hydrogen peroxide, hydroxyl and peroxy radicals may initiate the arachidonic acid cascade, PAF synthesis or histamine release.¹⁹⁻²² Thus, it has been demonstrated that ROS degranulate mast cells leading to release of histamine, serotonin, TNF- α and other mediators of inflammation. Furthermore, evidence has been provided that histamine is a marker of OH[•] radicals.²³ Phycocyanin is able to scavenge hydroxyl, alkoxy and peroxy radicals.^{1,24}

Taking together the presented results, it is conceivable that these ROS scavenging properties of the biliprotein may explain its inhibitory effects on histamine release and other mediators of inflammation. However, further experiments are required to confirm whether ROS and other intracellular messengers such as cAMP and intracellular calcium²⁵ might be involved in the inhibitory effect of phycocyanin on histamine release from rat mast cells.

References

1. Romay C, Armesto J, Ramirez D, González R. Antioxidant and anti-inflammatory properties of c-phycocyanin from blue-green algae. *Inflamm Res* 1998; **47**: 36-41.
2. Romay C, Ledón N, González R. Effects of phycocyanin extract on prostaglandin E₂ levels in mouse ear inflammation test. *Arzneim Forsch/Drug Res* 2000; **50**: 1106-1109.
3. Romay C, Ledón N, González R. Phycocyanin extract reduces leukotrienes B₄ levels in arachidonic acid-induced mouse ear inflammation test. *J Phar Pharmacol* 1999; **51**: 641-642.
4. Kim HM, Lee EH, Cho H, Moon YH. Inhibitory effect of mast cell-mediated immediate type allergic reactions in rats by Spirulina. *Biochem Pharmacol* 1998; **55**: 1071-1076.
5. Benítez F, Travieso L, Dupeyron R. Method for phycocyanin obtainment from microalgae. Cuban Patent Pending, RPI: 11/97.
6. Berns DS, MacColl R. Phycocyanin in physical-chemical studies. *Chem Rev* 1992; **89**: 807-825.
7. Hessel EM, Van Oosterhout AJM, Hofstra CL. Bronchoconstriction and airways hyperresponsiveness after ovalbumin inhalation in sensitized mice. *Eur J Pharmacol* 1995; **293**: 401-412.
8. Bradley PR, Priebe DA, Chritensen RD, Rothstein G. Measurement of cutaneous inflammation: estimation of neutrophils content with an enzyme marker. *J Invest Dermatol* 1982; **78**: 206-209.
9. Awouters F, Niemegeers CJE, Janssen PAJ. Interaction of astemizole and other drugs with PCA, histamine, serotonin and compound 48/80-induced skin reactions in the rat A procedure to determine anti-allergic effectiveness. *Drug Dev Res* 1985; **5**: 137-145.
10. Garland LG, Mongar JL. Inhibition by cromoglycate of histamine release from rat peritoneal mast cells induced by mixtures of dextran, phosphatidyl serine and calcium ions. *Br J Pharmacol* 1974; **50**: 137-143.
11. Shore PA, Burkhalter A, Cohn VN. A method for the fluorimetric assay of histamine in tissues. *J Pharmacol Exp Ther* 1959; **127**: 182-186.
12. Halpern BN, Neveu T, Spector S. On the nature of the chemical mediators involved in anaphylactic reaction in mice. *Br J Pharmacol* 1963; **20**: 389-398.

13. Ohuchi K, Hirasawa N, Watanabe M, Tsurufuji S. Pharmacological analysis of the vascular permeability response in the anaphylactic phase of allergic inflammation in rats. *Eur J Pharmacol* 1985; **117**: 337-345.
14. Williams CM, Galli SJ. The diverse potential effector and immunoregulatory roles of mast cells in allergic disease. *J Allergy Clin Immunol* 2000; **105**: 847-859.
15. Fantozzi R, Brunelleschi S, Griulattine L, *et al*. Mast cell and neutrophil interactions: a role for superoxide anion and histamine. *Agents Actions* 1985; **16**: 260-264.
16. Romay C, Ledón N, González R. Further studies on anti-inflammatory activity of phycocyanin in some animal models of inflammation. *Inflamm Res* 1998; **47**: 334-338.
17. Ramirez D, González A, Merino N, Gonzalez R, *et al*. Effect of phycocyanin in zymosan-induced arthritis in mice. Phycocyanin as an antiarthritic compound. *Drug Dev Res* 1999; **48**: 70-75.
18. González R, Rodriguez S, Romay C. Anti-inflammatory activity of phycocyanin extract in acetic acid-induced colitis in rats. *Pharmacol Res* 1999; **39**: 55-59.
19. Lewis MS, Whatley RE, Cain P. Hydrogen peroxide stimulates the synthesis of platelet activating factor by endothelium and induces endothelial cell-dependent neutrophil adhesion. *J Clin Invest* 1988; **82**: 2045-2055.
20. Mannaioni PF, Masini E. The release of histamine by free radicals. *Free Radic Biol Med* 1988; **5**: 177-197.
21. Masini E, Palmerani B, Gambassi F. Histamine release from rat mast cells induced by metabolic activation of polyunsaturated fatty acids into free radicals. *Biochem Pharmacol* 1990; **39**: 879-889.
22. Alanko J, Riutta A, Mucha I. Modulation of arachidonic acid metabolism by phenols: relation to positions of hydroxyl groups and peroxy radical scavenging properties. *Free Radic Biol Med* 1993; **14**: 19-24.
23. Ching TL, Van der Hee RM, Bhoelan NM. Histamine as a marker for hydroxyl radicals. *Mediat Inflamm* 1995; **4**: 339-343.
24. Lissi EA, Pizarro M, Aspee A. Kinetics of phycocyanin bilin groups destruction by peroxy radicals. *Free Radic Biol Med* 2000; **28**: 1051-1055.
25. Alfonso A, Cabado AG, Vieytes MR. Functional compartments in rat mast cells for cAMP and calcium on histamine release. *Cell Signal* 2000; **12**: 342-350.

Received 25 September 2001

Accepted 21 January 2002