

Background: Sodium nitroprusside (SNP) and molsidomine are used in the treatment of coronary heart disease. Since the neutrophils play a pathological role in ischaemic heart disease, it is important to understand the direct action of nitrovasodilators on their function.

Aim: We examined the effects of SNP and 3-morpholinylsindolizone (SIN-1, molsidomine metabolite) on the respiratory burst of human neutrophils and their adhesion *in vitro*. The influence of nitric oxide (NO) donors on the activity of protein kinases, which are involved in the NADPH oxidase activation, was also investigated.

Methods: The respiratory burst of neutrophils was determined by chemiluminescence and fluorescence methods, while the adhesion was assayed by adherence of neutrophils to the plastic surface.

Results: NO donors decreased the oxidative burst of activated neutrophils. However, the effects of SNP and SIN-1 strongly depended on the treatment time of neutrophils and on the stimulus employed to cells activation. Protein kinase C inhibitor did not prevent the inhibitory effect of SIN-1, but diminished the inhibitory effect of SNP on the neutrophils' respiratory burst. Protein tyrosine kinase inhibitor did not affect the action of SNP, but diminished the inhibitory effect of SIN-1 on fMLP-stimulated but not on PMA-stimulated oxidative burst of neutrophils. This suggests that SNP action is mainly associated with protein kinase C, while SIN-1 is associated with protein tyrosine kinase activity. We also found that SIN-1 but not SNP diminished the adhesive activity of neutrophils.

Conclusions: Our data show that SIN-1 biological effect on some neutrophils activity is different from both spermine NONOate and SNP, and mainly depends on ONOO⁻, while SNP action is mediated by NO.

Key words: Nitrovasodilators, Neutrophils, Respiratory burst, Protein kinases, Adherence

Oxidative and adhesive responses of human neutrophils to nitrovasodilators *in vitro*: the role of protein kinases

Magdalena Klink^{1,CA}, Henryk Tchorzewski^{1,2} and Zofia Sulowska¹

¹Microbiology and Virology Centre, Polish Academy of Sciences, Lodowa 106, 93-232 Lodz, Poland; and

²Department of Clinical Immunology, Institute of Polish Mother's Memorial Hospital, Lodz, Poland

^{CA}Corresponding author

Tel: +48 42 6771236

Fax: +48 42 6771230

E-mail: MKlink@poczta.onet.pl

Introduction

Neutrophils play a key role in the pathogenesis of ischaemic heart disease as the mediators of myocardial ischaemia reperfusion injury. Activated neutrophils produce and release a variety of powerful inflammatory mediators such as reactive oxygen species, enzymes and cytokines that are responsible for cell and tissue damage and serious complications associated with heart diseases.^{1–3}

The reactive oxygen species are produced by neutrophils as result of NADPH oxidase activation (reactions named 'respiratory burst'). Various protein kinases are involved in the NADPH oxidase activation including protein kinase C (PKC), mitogen-activated protein kinases, protein kinase A, and protein tyrosine kinase (PKT).^{4–10} NADPH oxidase can be

activated by various stimuli such as chemotactic peptide (formyl-methionyl-leucyl-phenylalanine (fMLP)), phorbol 12-myristate 13-acetate (PMA) or opsonized zymosan. Stimulation of neutrophils with fMLP, a receptor-dependent stimulator, activates G-proteins and various downstream proteins including PKC, PKT, and mitogen-activated protein kinase. PMA as a receptor-independent stimulator directly activates PKC without G-proteins participation. It also indirectly activates mitogen-activated protein kinase and probably PKT.^{6,9–11}

Nitric oxide (NO) donors such as sodium nitroprusside (SNP) and molsidomine are used as nitrovasodilators in the treatment of patients with coronary heart disease.^{12–14} Besides their effects on the vascular smooth muscle cells, NO-releasing compounds may affect the circulating white blood

cells; for example, inhibit the adhesion, aggregation and chemotaxis of neutrophils.¹⁵

The recognition of the pathway by which the nitrovasodilators affect neutrophils function seems to be significant. The aim of this study was to examine the effects of SNP and 3-morpholinonydnomine (SIN-1, molsidomine metabolite) on the respiratory burst of human neutrophils and their adhesion *in vitro*. The influence of these compounds on the activity of kinases (PKT and PKC) that are involved in the NADPH oxidase activation was investigated.

Materials and methods

Chemical reagents

Polymorphprep was obtained from Nycomed (Oslo, Norway). Hanks' balanced salt solution (HBSS), SNP, SIN-1, PMA, spermine NONOate, genistein, herbimicin A, 1-(5-isoquinolinylnsulfonyl)-2-methylpiperazine (H-7), NPC 15437, fMLP, tumour necrosis factor- α (TNF- α), human recombinant, 2',7'-dichlorofluorescein (DCFH) diacetate, 3-[4,5-dimethylthiazol-2-yl]-2,3-diphenyltetrazolium bromide (MTT), sodium nitrite, sulfanilamide, naphthylethylenediamine dihydrochloride, and phosphoric acid were purchased from Sigma (USA). Luminol was a product of Serva (Germany). Phosphate-buffered saline (PBS) was purchased from BIOMED-Lublin (Poland).

Isolation of neutrophils

Heparinized peripheral blood was obtained from healthy volunteers and the neutrophils were isolated by centrifugation of blood sample on Polymorphprep as described previously.¹⁶ The Regional Commission approved the protocol of these studies for Ethics in Research.

Measurement of the generation of nitric oxide from NO donor compounds

NO donors (10–1000 μ M) were added to HBSS (cell free system) or to neutrophils (1×10^5 cells) in HBSS for 30 or 60 min at 37°C. The generation of NO was measured using Griess reagent as described previously.¹⁶

Respiratory burst of neutrophils

Two assays were used to assess the respiratory burst because a single technique could be insufficient to determine the effect of different NO donors on the neutrophil function.¹⁶ Luminol-enhanced chemiluminescence measures the intracellularly and extracellu-

larly released NADPH-derived and myeloperoxidase-derived oxygen metabolites, while DCFH oxidation assay measures the NADPH-derived oxygen species generated mainly intracellularly.^{17,18}

Chemiluminescence (CL) was measured in a 96-well plate Fluoroscan Ascent FL fluorometer (Labsystem, Helsinki, Finland). Neutrophils (1×10^5 cells/well in HBSS) were untreated or pre-treated with 1, 10, 100 or 1000 μ M of SNP or SIN-1 for 1–2 min or 30 min at 37°C with 5% CO₂ as indicated in the figures. In control experiments, the neutrophils were pre-treated with 10 μ M or 100 μ M of spermine NONOate for 1–2 min or 30 min at 37°C with 5% CO₂. PMA (0.1 μ g/ml) or fMLP (10^{-6} M) to initiate the respiratory burst and luminol (10^{-5} M) to enhance CL were added to the cells. In some experiments the neutrophils were pre-treated with 100 μ M or 1000 μ M of SNP or SIN-1 for 30 min at 37°C with 5% CO₂, and NO donors were then removed. PMA, fMLP and luminol were then added as previously. The CL reading for all experiments was recorded for 30 min at 2 min intervals. The CL intensity was given in relative light units (RLU). The data were expressed as the area under the curve of CL versus time (RLU total).

The DCFH oxidation assay was measured in a 96-well plate Fluoroscan Ascent FL fluorometer (Labsystem). Briefly, the neutrophils (2×10^6 cells/ml) were loaded with DCFH diacetate (100 μ g/ml) in HBSS for 30 min at 37°C, with 5% CO₂. Cells were then washed twice with PBS, resuspended in fresh HBSS and distributed into a 96-well plate (1×10^5 cells/well). Neutrophils were untreated or pre-treated with SNP or SIN-1 at concentrations of 1, 10, 100 or 1000 μ M for 1–2 min or 30 min at 37°C with 5% CO₂. The plate was placed into fluorometer and the intensity of fluorescence was determined at 485 nm excitation and 530 nm emission wavelengths (T_0). Neutrophils were then stimulated with PMA (0.1 μ g/ml) or fMLP (10^{-6} M) for the next 30 min at 37°C with 5% CO₂. After that time, the intensity of fluorescence was measured once more as described earlier (T_{30}). In some experiments the neutrophils were untreated or pre-treated with 100 μ M or 1000 μ M of SNP or SIN-1 for 30 min at 37°C with 5% CO₂. Cells were then washed to remove NO-releasing drugs. The intensity of fluorescence at T_0 and after stimulation with PMA or fMLP at T_{30} was measured. The percentage of increase in the intensity of fluorescence was calculated from the formula $[(FT_{30} - FT_0)/FT_0] \times 100$.¹⁹

Determination of the role of PKT and PKC activity in the NO donor-mediated respiratory burst

Neutrophils (1×10^5 cells/well) were pre-treated with PKT inhibitors genistein (10 μ M) and herbimicin A (1 μ M) or PKC inhibitors H-7 (100 μ M) and NPC (1 μ M) for 15 min at 37°C with 5% CO₂. Neutrophils

were then incubated with SNP or SIN-1 (100 μ M and 1000 μ M) or with NONOate (10 μ M and 100 μ M) for 30 min. PMA, fMLP and luminol were then added as already described. The data were expressed as the mediated by NO donors' per cent inhibition of PMA-induced or fMLP-induced respiratory burst of neutrophils calculated by the following formulae:

- A. $[(\text{RLU}_{\text{total}}$ of cells treated with SNP or SIN-1 + stimulated with PMA or fMLP) - ($\text{RLU}_{\text{total}}$ of cells treated with SNP or SIN-1)/($\text{RLU}_{\text{total}}$ of cells stimulated with PMA or fMLP) - ($\text{RLU}_{\text{total}}$ of untreated (control) cells)] \times 100%
- B. $[(\text{RLU}_{\text{total}}$ of cells pretreated with PKC or PKT inhibitor and SNP or SIN-1 + stimulated with PMA or fMLP) - ($\text{RLU}_{\text{total}}$ of cells treated with SNP or SIN-1 + stimulated with PKC or PKT)/($\text{RLU}_{\text{total}}$ of cells treated with PKC or PKT inhibitor + stimulated with PMA or fMLP) - ($\text{RLU}_{\text{total}}$ of cells treated with PKC or PKT inhibitor)] \times 100%.

Measurement of neutrophil adhesion

The adhesion was assayed by adherence of neutrophils to the plastic surface. This is a simple quantitative technique to examine the ability of human neutrophils to active adherence *in vitro*.²⁰ Briefly, neutrophils were treated with SNP and SIN-1 or with 20 ng/ml of TNF- α (as indicated later in Table 2). Non-adherent cells were removed by carefully washing the wells' surface with PBS and adherent neutrophils were exposed to MTT (2 mg/ml of PBS) containing 10 ng/ml of PMA for 30 min. Formazan produced by adherent cells was dissolved in 500 μ l of 2-propanol. The absorbance at 560 nm and 630 nm dual wavelength was measured in an ELISA reader Multiscan RC (Labsystem). The absorption of formazan is directly related to the number of adherent neutrophils.²⁰ The amount of adherent cells was also controlled microscopically by independent re-

searcher. The loss of cells treated with SIN-1 was observed under the microscope.

Statistical analysis

Data are presented as the mean \pm standard deviation. Statistical analysis was performed with Wilcoxon's signed rank test. Statistical significance was defined as $p \leq 0.05$.

Results

Oxidative response of neutrophils to nitrovasodilators

The generation of nitrite (a stable metabolite of NO) from NO donors in the presence of neutrophils or in the cell free system is presented in Table 1.

In pilot experiments, we noticed that 1 min or 30 min pre-treatment of neutrophils with SNP or SIN-1 at 1 μ M and 10 μ M had no significant effects on oxygen radical generation by stimulated cells (data not shown). Therefore, the concentrations of 100 μ M and 1000 μ M were selected for further experiments.

We observed that 1–2 min pre-incubation with either SNP or SIN-1 decreased the respiratory burst of PMA-stimulated neutrophils as it was measured with CL and DCFH oxidation assays. The effect of pre-incubation with SNP and SIN-1 on the respiratory burst of fMLP-stimulated neutrophils was not seen when CL measurement was used. However, different effects of SNP and SIN-1 on fMLP-stimulated neutrophils were observed when DCFH oxidation assay was used. SNP increased, while SIN-1 decreased the fMLP-stimulated respiratory burst of neutrophils (Fig. 1). Figure 2 demonstrates that 30 min pre-incubation with both nitrovasodilators decreased the respiratory burst of PMA-stimulated and fMLP-stimulated neutrophils as it was measured with CL and DCFH oxidation assays. As shown in Fig. 3, the removal of SNP from the cells environment cancelled its down-regulating effect on the oxidative burst of

Table 1. Nitrite generation from NO donors in the presence of neutrophils and in the cell free system

Incubation (min)	SNP (cell free system)		SIN-1 (cell free system)		Spermine NONOate (cell free system)		
	1000 μ M	100 μ M	1000 μ M	100 μ M	1000 μ M	100 μ M	10 μ M
30	1.2 \pm 0.6	0.1 \pm 0.1	48.4 \pm 15.3	9.4 \pm 3.1	288.1 \pm 32.2	137.8 \pm 3.1	15.5 \pm 0.5
60	2.3 \pm 0.4	0.7 \pm 0.4	187 \pm 20.3	13.8 \pm 2.5	288.0 \pm 10.1	125.2 \pm 7.9	13.3 \pm 1.6
	SNP + neutrophils		SIN-1 + neutrophils		Spermine NONOate + neutrophils		
	1000 μ M	100 μ M	1000 μ M	100 μ M	1000 μ M	100 μ M	10 μ M
30	4.2 \pm 1.2	1.2 \pm 0.8	36.1 \pm 8.7	4.1 \pm 0.8	308.0 \pm 16.9	132.2 \pm 12.2	13.7 \pm 7.8
60	12.3 \pm 1.7	2.0 \pm 0.9	104.3 \pm 12.7	19.3 \pm 9.6	300.9 \pm 11.9	118.5 \pm 9.4	12.0 \pm 1.2

SNP (100 μ M and 1000 μ M) or SIN-1 (100 μ M and 1000 μ M) or spermine NONOate (10–1000 μ M) were added to HBSS (cell free system) or to neutrophils (1×10^5 cells) in HBSS for 30 or 60 min at 37°C with 5% CO₂. The generation of nitrite from NO donors was measured using Griess reagents. Data are expressed as mean \pm standard deviation of three independent experiments.




FIG. 1. Effect of brief pre-treatment with nitrovasodilators on the respiratory burst of stimulated neutrophils *in vitro*. Neutrophils (1×10^5 cells) were untreated or pre-treated with nitrovasodilators at the concentrations of 100 μM and 1000 μM for 1–2 min, and then stimulated with PMA (closed bars) or with fMLP (hatched bars) or were unstimulated (open bars). See details in Materials and methods. Data are expressed as means of RLU total or per cent increase of fluorescence \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. * Statistically significant decrease: neutrophils versus neutrophils with nitrovasodilators ($p \leq 0.02$). # Statistically significant increase: neutrophils versus neutrophils with SNP ($p \leq 0.02$).




FIG. 2. Effect of 30 min pre-treatment with nitrovasodilators on the respiratory burst of activated neutrophils *in vitro*. Neutrophils (1×10^5 cells) were untreated or pre-treated with nitrovasodilators at the concentrations of 100 μM and 1000 μM for 30 min and then stimulated with PMA (closed bars) or with fMLP (hatched bars) or were unstimulated (open bars). See details in Materials and methods. Data are expressed as means of RLU total or per cent increase of fluorescence \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. * Statistically significant decrease: neutrophils versus neutrophils with nitrovasodilators ($p \leq 0.02$).




FIG. 3. Effect of withdrawal of nitrovasodilators from neutrophil environment prior to stimulation of cells to respiratory burst *in vitro*. Neutrophils (1×10^5 cells) were untreated or pre-treated with nitrovasodilators at the concentrations of 100 μM and 1000 μM for 30 min, washed to remove SNP and SIN-1, and then stimulated with PMA (closed bars) or with fMLP (hatched bars) or were unstimulated (open bars). See details in Materials and methods. Data are expressed as means of RLU total or per cent increase of fluorescence \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. * Statistically significant decrease: neutrophils versus neutrophils with nitrovasodilators ($p \leq 0.02$).

fMLP-stimulated but not PMA-stimulated neutrophils, whereas the inhibitory effects of SIN-1 were not reversed after its withdrawal.

Data presented in Fig. 4 demonstrate that 1-2 min pre-incubation with spermine NONOate decreased the respiratory burst of PMA-stimulated but not of fMLP-stimulated neutrophils. Thirty minutes' pre-

incubation with this NO donor decreased the oxidative burst of PMA-activated and fMLP-activated cells. It was not possible to perform experiments with NONOate at concentrations higher than 100 μM because that caused cell damage.

It should be stressed that both nitrovasodilators at the concentrations used did not interfere with lumi-




FIG. 4. Effect of NONOate on the respiratory burst of stimulated neutrophils *in vitro*. Neutrophils (1×10^5 cells) were untreated or pre-treated with NONOate at the concentrations of 10 μM and 100 μM for 1-2 min or 30 min, and then stimulated with PMA (closed bars) or with fMLP (hatched bars) or were unstimulated (open bars). See details in Materials and methods. Data are expressed as means of RLU total \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. * Statistically significant decrease: neutrophils versus neutrophils with nitrovasodilators ($p \leq 0.02$).

nol or DCFH diacetate and did not produce any signal in the cell free system as was verified in both assays.

Involvement of PKC and PKT pathways in the NO donor-mediated decrease of respiratory burst of activated neutrophils

We examined the effect of blocking the PKC or PKT activity on the nitrovasodilator-mediated decrease of respiratory burst of stimulated neutrophils. As shown in Fig. 5, PKC inhibitors (H-7 and NPC) at the used doses had no effect on fMLP-induced neutrophil oxidative burst, but when added prior to the induction of this process by PMA they caused a significant reduction of it. These data indicate that fMLP activates neutrophils' oxidative metabolism through the PKC-independent mechanism. PKT inhibitors (genistein and herbimycin A) caused a significant reduction of respiratory burst of PMA-stimulated and fMLP-stimulated neutrophils (Fig. 5). These results indicate that PKT participates in the induction of PMA-activated neutrophils' respiratory burst. Data presented in Fig. 6 demonstrate that PKC inhibitor H-7, administered prior to the addition of SNP and PMA or fMLP, partially but significantly diminished the inhibitory effect of the NO donor, whereas H-7 did not affect the

action of SIN-1. The similar results we observed with NPC (data not shown). As shown in Fig. 7, the PKT inhibitor genistein did not affect the inhibitory effect of SNP. However, genistein attenuated the inhibitory effect of SIN-1 on fMLP-stimulated but not on PMA-stimulated neutrophils. Similar results were observed with herbimycin A (data not shown).

Genistein did not affect the inhibitory effect of spermine NONOate but H-7 significantly diminished the inhibitory effect of spermine NONOate on respiratory burst of activated neutrophils. The percentage of inhibition was as follow: neutrophils+NONOate+fMLP, 22.2 ± 11.2 versus neutrophils+H-7+NONOate+fMLP, 10.7 ± 9.4 ($p \leq 0.05$, $n = 5$); neutrophils+NONOate+PMA, 23.9 ± 11.4 versus neutrophils+H-7+NONOate+PMA, 15 ± 7.2 ($p \leq 0.05$, $n = 5$).

Adhesive activity of nitrovasodilator-treated neutrophils

TNF- α enhanced the adhesion of neutrophils after 30 min of incubation with cells (from optical density (OD) = 0.070 ± 0.029 in the control sample up to OD = 0.130 ± 0.052 in the presence of cytokine; $p \leq 0.04$, $n = 5$). The data in Table 2 demonstrate the different effect of SNP and SIN-1 on the neutrophil




FIG. 5. Effects of PKC (A) and PKT (B) inhibitors on PMA-induced and fMLP-induced respiratory burst of neutrophils. Neutrophils (1×10^5) were first treated with H-7 (100 μ M) or NPC (1 μ M) (A) or with genistein (10 μ M) or herbimycin A (1 μ M) (B) for 15 min. Then, the cells were stimulated with 0.1 μ g/ml of PMA or 10^{-6} M fMLP, respectively. The CL reading was recorded for 30 min at 2 min intervals. The data are expressed as the area under the obtained curve of CL versus time (RLU total). Data are the means \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. Statistical significance: # PMA versus PMA+PKC inhibitors, $p \leq 0.02$; * PMA versus PMA+PKT inhibitors, $p \leq 0.02$; ** fMLP versus fMLP+PKT inhibitors, $p \leq 0.02$.




FIG. 6. Effect of PKC inhibitor on the SNP-mediated and SIN-1-mediated oxidative burst of activated neutrophils. Neutrophils (1×10^5) were first treated with H-7 (100 μ M) for 15 min. Then, the cells were incubated with 1000 μ M of SNP or 1000 μ M of SIN-1 for 30 min and next stimulated with 0.1 μ g/ml of PMA or 10^{-6} M of fMLP. The CL reading was recorded for 30 min at 2 min intervals. The data are expressed as means of per cent inhibition of respiratory burst \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. Statistical significance: * SNP+PMA versus SNP+PMA+H-7, $p \leq 0.02$; ** SNP+fMLP versus SNP+fMLP+H-7, $p \leq 0.02$.




FIG. 7. Effect of PKT inhibitor on the SNP-mediated and SIN-1-mediated oxidative burst of activated neutrophils. Neutrophils (1×10^5) were first treated with genistein (10 μ M) for 15 min. Then, the cells were incubated with 1000 μ M of SNP or 1000 μ M of SIN-1 for 30 min and next stimulated with 0.1 μ g/ml of PMA or 10^{-6} M of fMLP. The CL reading was recorded for 30 min at 2 min intervals. The data are expressed as means of per cent inhibition of respiratory burst \pm standard deviation of six independent experiments carried out with neutrophils from different individuals. Statistical significance: * SIN-1+fMLP versus SIN-1+fMLP+genistein, $p \leq 0.02$.

Table 2. Effect of nitrovasodilators on the adhesive activity of neutrophils

Time (min)	OD (560 nm and 630 nm)				
	Control	SNP		SIN-1	
		100 μ M	1000 μ M	100 μ M	1000 μ M
30	0.070 \pm 0.023	0.068 \pm 0.010	0.075 \pm 0.018	0.064 \pm 0.019	0.051 \pm 0.019*
60	0.258 \pm 0.039	0.215 \pm 0.043	0.260 \pm 0.043	0.207 \pm 0.027*	0.189 \pm 0.049*

Neutrophils (1×10^6 cells) were distributed into 24-well plates and were incubated without or with nitrovasodilators (100 μ M or 1000 μ M) for 30 and 60 min at 37°C, with 5% of CO₂. Non-adherent cells were removed, and adherent cells were exposed for MTT solution as described in Materials and methods. Data are expressed as means of optical densities \pm standard deviation of eight independent experiments carried out with neutrophils from different individuals. * Statistical significance: neutrophils versus neutrophils with SIN-1 ($p \leq 0.05$).

adhesion. Neutrophil adhesion was reduced after treatment of cells with 100 μ M of SIN-1 for 60 min and with 1000 μ M of SIN-1 for 30 and 60 min. In contrast, SNP did not affect their adhesion to the plastic surface. Spermine NONOate did not influence on the adherence of neutrophils (data not shown).

Discussion

The mechanism of NO generation from SNP is controversial. Some authors have shown that NO could be spontaneously generated from SNP,^{13,21} but others have demonstrated that NO formation required the presence of vascular tissue¹² and/or the visible light.²² We have shown here that SNP released NO in the cell free system; however, the presence of neutrophils enhanced its generation (Table 1). SIN-1, the metabolite of molsidomine, produced NO spontaneously without any cofactors.^{12–14}

The results of our experiments are in accordance with observations of others^{21–24} that SNP and SIN-1 at concentrations above 50 μ M decrease the reactive oxygen species production by neutrophils. However, it should be also emphasized that biological effects of SNP and SIN-1 strongly depended on the treatment time of neutrophils and on the stimulus employed to activate the cells. Our data indicate that SNP and SIN-1 should act on neutrophils for longer than 1–2 min to exert their modulatory effects, when the receptor-dependent activator is used (Figs. 1 and 2). We may speculate that the different oxidative response of short-time nitrovasodilator pre-treated neutrophils arise from different kinetics of the respiratory burst induced by PMA and by fMLP. The kinetics of CL responses show that the peak of the PMA-induced oxidative response occurs 10–15 min after activation of cells while the peak of the fMLP-induced respiratory burst of neutrophils occurs 1–5 min after neutrophil activation.²³ Our results and those of others²¹ suggest that SNP and SIN-1 may affect the respiratory burst of neutrophils by various signalling pathways (Fig. 3). The inhibition of the PMA-dependent respiratory burst of neutrophils by SNP and SIN-1 indicates that NO donors exert effects independent

of G-proteins. However, SIN-1 can also exert some effect via or close to G-proteins since it decreased the fMLP-mediated neutrophils' oxidative burst.²⁵

In the course of NADPH oxidase activation distinct signal transduction pathways can be activated.^{4,7,11} The signal transduction through receptors with tyrosine kinase activity occurs upon interaction of fMLP with its receptor; however, the other signalling pathway involves PKC via activation of phospholipase C. PMA directly activates PKC, and indirectly PKT.^{11,26} The role of protein kinases in the action of NO donors on the respiratory burst of neutrophils is not clear and it was not examined intensively. In this paper we show that SNP action is mainly associated with PKC, while SIN-1 is associated with PKT activity (Figs. 6 and 7). We suppose that the influence of NO donors on PKC, PKT and/or G-protein activities can be responsible for the decrease of respiratory burst of activated neutrophils.

It was demonstrated by others that the effect of SNP and SIN-1 on the neutrophil functions was mediated by NO.^{21,23,24} Our results suggest that NO donors can also affect neutrophils in the NO-independent pathway. Feelisch¹³ maintained that a relatively small amount of NO released from SNP *in vitro* was not sufficient for activating soluble guanylate cyclase and that biological effects of SNP were largely independent of NO. It was also suggested that SNP might mediate effects by a NO-dependent and soluble guanylate cyclase-independent mechanism.¹⁴ It should be also considered that during decomposition of SNP a cyanide is released. However, Forslund and Sundqvist²³ demonstrated that the effect of SNP on the respiratory burst of neutrophils was mediated by NO not by cyanide. Our data show that SNP, similar to spermine NONOate, decreases PMA-induced respiratory burst of neutrophils and affects the PKC activity. It can suggest that the action of SNP may depend on NO. Spontaneous decomposition of SIN-1 leads to the generation of equimolar quantities of NO and O₂⁻, which interact almost instantly to produce peroxynitrite anion (ONOO⁻).¹² However, it should be emphasized that formation of ONOO⁻ during decomposition of SIN-1 *in vitro* strongly depends on the oxygen concentration in the reaction mixture.^{27,28}

Our data show that the SIN-1 biological effect on some neutrophils' activity is different from both spermine NONOate and SNP, and mainly depends on ONOO⁻, while SNP action is mediated by NO.

ACKNOWLEDGEMENTS. This research was supported by grant 4 P05A 034 18 from The State Committee for Scientific Research (KBN) Poland.

References

- Siminiak T, Ozawa T. Neutrophil mediated myocardial injury. *Int J Biochem* 1993; **25**: 147–156.
- Hansen PR. Role of neutrophils in myocardial ischemia and reperfusion. *Circulation* 1995; **91**: 1872–1885.
- Baxter GF. The neutrophil as a mediator of myocardial ischemia-reperfusion injury: time to move on. *Basic Res Cardiol* 2002; **97**: 268–275.
- Babior BM. NADPH oxidase: an update. *Blood* 1999; **93**: 1464–1476.
- Jones RD, Hancock JT, Morice AH. NADPH oxidase: a universal oxygen sensor? *Free Radic Biol Med* 2000; **29**: 416–424.
- Dang PMC, Fontayne A, Hakim J, El-Benna J, Perianin A. Protein kinase C ζ phosphorylates a subset of selective sites of the NADPH oxidase component p47^{phox} and participates in formyl peptide-mediated neutrophil respiratory burst. *J Immunol* 2001; **166**: 1206–1213.
- Reeves EP, Dekker LV, Forbes LV, Wientjes FB, Grogan A, Pappin DC, Segal AW. Direct interaction between p47^{phox} and protein kinase C: evidence for targeting of protein kinase C by p47^{phox} in neutrophils. *Biochem J* 1999; **344**: 859–866.
- Karlson A, Nixin JB, McPhail LC. Phorbol myristate acetate induces neutrophil NADPH-oxidase activity by two separate signal transduction pathways: dependent or independent of phosphatidylinositol 3-kinase. *J Leukoc Biol* 2000; **67**: 396–404.
- Watson F, Robinson J, Edwards SW. Protein kinase C-dependent and -independent activation of the NADPH oxidase of human neutrophils. *J Biol Chem* 1991; **266**: 7432–7439.
- Dewas CD, Fay M, Gougerot-Pocidalo MA, El-Benna J. The mitogen-activated protein kinase extracellular signal-regulated kinase 1/2 pathway is involved in formyl-methionyl-leucyl-phenylalanine-induced p47^{phox} phosphorylation in human neutrophils. *J Immunol* 2000; **165**: 5238–5244.
- Worthen GS, Avdi N, Buhl AM, Suzuki N, Johnson GL. fMLP activates Ras and Raf in human neutrophils. Potential role in activation of MAP kinase. *J Clin Invest* 1994; **94**: 815–823.
- Yamamoto T, Bing RJ. Nitric oxide donors. *Proc Soc Exp Biol Med* 2000; **225**: 200–206.
- Feelisch MJ. The biochemical pathways of nitric oxide formation from nitrovasodilators: appropriate choice of exogenous NO donors and aspects of preparation and handling of aqueous NO solutions. *J Cardiovasc Pharmacol* 1991; **17(Suppl 3)**: S25–S33.
- Megson IL. Nitric oxide donor drugs. *Drugs* 2000; **25**: 701–715.
- Armstrong R. The physiological role and pharmacological potential of nitric oxide in neutrophils activation. *Int Immunopharmacol* 2001; **1**: 1501–1512.
- Klink M, Cedzynski M, Swierzko Ast, Tchorzewski H, Sulowska Z. Involvement of nitric oxide donor compounds in the bactericidal activity of human neutrophils *in vitro*. *J Med Microbiol* 2003; **52**: 303–308.
- Dahlgren C, Karlsson A. Respiratory burst in human neutrophils. *J Immunol Methods* 1999; **232**: 3–14.
- DeLeo FR, Allen LH, Apicella M, Nauseef WM. NADPH oxidase activation assembly during phagocytosis. *J Immunol* 1999; **163**: 6732–6740.
- Wang H, Joseph JA. Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. *Free Radic Biol Med* 1991; **27**: 612–616.
- Oez S, Weltre K, Platzer E, Kalden JR. A simple assay for quantifying the inducible adherence of neutrophils. *Immunobiology* 1990; **180**: 308–315.
- Forslund T, Sundqvist T. Nitric oxide regulates the chemiluminescence from stimulated human neutrophils. *APMIS* 1995; **103**: 813–817.
- Bates JN, Baker MT, Guerra R Jr, Harrison DG. Nitric oxide generation from nitroprusside by vascular tissue. Evidence that reduction of the nitroprusside anion and cyanide loss are required. *Biochem Pharmacol* 1991; **42**: S157–S165.
- Forslund T, Sundqvist T. Nitric oxide reduces hydrogen peroxide production from human polymorphonuclear neutrophils. *Eur J Clin Invest* 1995; **25**: 9–14.
- Pieper GM, Clarke GA, Gross GJ. Stimulatory and inhibitory action of nitric oxide donor agents vs. nitrovasodilators on reactive oxygen production by isolated polymorphonuclear leukocytes. *J Pharmacol Exp Ther* 1994; **269**: 451–456.
- Clancy RM, Leszczynska-Piziak J, Abramson SB. Nitric oxide, an endothelial cell relaxation factor, inhibits neutrophil superoxide anion production via a direct action on the NADPH oxidase. *J Clin Invest* 1992; **90**: 1116–1121.
- Ptasznik A, Prossnitz ER, Yoshikawa D, Smrcka A, Traynor-Kaplan AE, Bokoch GM. A tyrosine kinase signaling pathway accounts for the majority of phosphatidylinositol 3,4,5-triphosphate formation in chemoattractant-stimulated human neutrophils. *J Biol Chem* 1996; **271**: 25204–25207.
- Brunelli L, Crow JP, Beckman JS. The comparative toxicity of nitric oxide and peroxynitrite to *Escherichia coli*. *Arch Biochem Biophys* 1995; **316**: 327–334.
- Doulias PT, Barbouti A, Galaris D, Ischiropoulos H. SIN-1 induced DNA damage in isolated human peripheral blood lymphocytes as assessed by single cell gel electrophoresis (comet assay). *Free Radic Biol Med* 2001; **30**: 679–685.

Received 18 July 2003

Accepted 29 September 2003



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

