Research Article

Protective Effects of *Salvia miltiorrhizae* on Multiple Organs of Rats with Obstructive Jaundice

Zhang Ling,¹ Zhang Xiping,² Qiu Fengmei,³ Yan Ping,³ and Cheng Qihui⁴

¹ The Cardiovascular Research Laboratory, Lakehead University, 290 Munro Street, Thunder Bay, ON, Canada P7A 7 T1

² Department of General Surgery, Hangzhou First People's Hospital, Hangzhou 310006, Zhejiang, China

³ Zhejiang University of Traditional Chinese Medicine, Hangzhou 310053, Zhejiang, China

⁴ Department of Gynaecology and Obstetrics, Hangzhou First People's Hospital, Hangzhou 310006, Zhejiang, China

Correspondence should be addressed to Zhang Xiping, zxp99688@vip.163.com

Received 9 May 2009; Accepted 10 June 2009

Recommended by Vera L. Petricevich

Purpose. we aim to explore the protective effects of *Salvia miltiorrhizae* injection on multiple organs of obstructive jaundice (OJ) rats through observing the impact of this injection on the pathological alterations in these organs and the contents of endotoxin, PLA₂, and TNF- α in the blood. **Methods.** A total of 90 mice were randomly divided into sham-operated group, model-control group, and *Salvia miltiorrhizae*-treated group ($n = 30$). According to the duration of postoperative administration, each group was further divided into two subgroups, namely, 21 d subgroup (consecutive administration for 21 d, $n = 15$) and 28 d subgroup (consecutive administration for 28 d, $n = 15$). After administration, the pathological alterations in multiple organs were observed and the contents of endotoxin, PLA₂, and TNF- α in the blood were determined. **Results.** Compared to model control group, the number of dead rats in treated group decreased though there was no statistical difference between the two groups. The pathological alterations in the liver, kidney, and spleen in treated group showed varying degrees of mitigation. At all time points, the contents of plasma endotoxin declined significantly. On day 28, plasma PLA₂ content in treated group was significantly lower than that in model-control group. **Conclusion.** *Salvia miltiorrhizae* injection is able to obviously reduce the contents of inflammatory mediators in the blood of OJ rats and exert some protective effects on multiple organs of these rats.

Copyright © 2009 Zhang Ling et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

It is currently believed that obstructive jaundice (OJ) can induce functional damage to the liver, spleen, lymphocytes, and gastrointestinal mucosa as well as cardiac hemodynamic dysfunction [1–4]. However, the mechanisms underlying OJ-induced multiple organ damage are still unclear, and there are many controversial hypotheses to explain this phenomenon. It is generally believed that endotoxin, inflammatory mediators, and increased bile acids contribute to OJ-induced liver or other organs injuries. Moreover, elevated intrabiliary duct pressure and portal vein pressure are also important factors contributing to this injury.

S. miltiorrhizae is a kind of traditional Chinese drug commonly used for activating blood circulation, the research on the effect of it in the treatment of OJ has been reported [5, 6]. *S. miltiorrhizae* injection is the extraction of wild

Salvia roots, the main active ingredients include danshensu, salvianolic acid as well as tanshinone, dihydrotanshinone, and cryptotanshinone, which are able to protect endothelial cells, fight against inflammation, and prevent lipid peroxidation and calcium overload [7, 8]. Some studies have shown that when *S. miltiorrhizae* is used to treat liver and kidney damage in OJ rats, it can exert some therapeutic effects against this damage through protecting hepatic cells, maintaining liver function, as well as reducing the damage to renal cortex and renal function [4, 6, 9, 10]. Moreover, *S. miltiorrhizae* also shows some therapeutic effects on OJ patients [11]. Although *S. miltiorrhizae* has been used to treat single organ injury complicating OJ in few studies [6, 11], a comprehensive study on the treatment of OJ-induced multiorgan damage with *S. miltiorrhizae* is lacking. For this reason, the purpose of this study is to make up for this deficiency. Through observing the impact of *S. miltiorrhizae*

injection on the pathological alterations in multiple organs and the contents of inflammatory mediators in the blood of OJ rats, we explored the protective effects of *S. miltiorrhizae* on these organs.

2. Materials and Methods

2.1. Materials. A total 90 healthy male SD rats of clean grade, weighing between 270 and 330 g, were provided by the Laboratory Animal Research Center of the Zhejiang University of Traditional Chinese Medicine (China); Sodium taurocholate and sodium pentobarbital were purchased from Sigma Corporation, USA. *S. miltiorrhizae* injection (each 10 mL vial contains active components equivalent to 15 g of the original medicine) was purchased from Chiatai Qingchunbao Pharmaceutical Co., Ltd. (China). Endotoxin ELISA Kit was purchased from Associates of Cape Cod (USA), the calculation unit for content is EU/mL. Serum TNF- α ELISA Kit was purchased from Shanghai Senxiong Technology Industry Co., Ltd. (China), the calculation unit for content is ng/L (pg/mL). Serum secretory phospholipase A₂ enzyme Assay ELA kit (PLA₂) was purchased from R&D system Ins (USA), the calculation unit for content is U/mL.

2.2. Methods

2.2.1. Animal Grouping. 90 rats were utilized for OJ-associated experiments and randomly divided into sham-operated, model-control, and treated group ($n = 30$), which were further randomly subdivided into 21 d and 28 d groups ($n = 15$) according to time duration after operation.

2.2.2. Preparation of OJ Models and Associated Therapeutic Regimen. After rats were anesthetized with an intraperitoneal injection of 2.5% sodium pentobarbital (0.2 mL/100 g), the abdominal cavity was opened to identify and dissociate common bile duct along the hepatoduodenal ligament. For rats in the model-control groups and the treated groups, the proximal end of common bile duct was double-ligated with surgical threads, common bile duct was cut off, and a layered suture of the abdominal wall was performed to close the abdominal cavity. For rats in the sham-operated groups, common bile duct was only dissociated but not ligated, and a layered suture of the abdominal wall was also performed to close the abdominal cavity. An intraperitoneal injection of *S. miltiorrhizae* injection at a dose of 0.2 mL/100 g/d [12–15] was given to rats in the treated groups while equal volume of physiological saline solution was used in the sham-operateds and the model-control groups. Injection was maintained until the end of the 21st day and 28th day observation period in the corresponding groups.

2.2.3. Specimen Collection. On the 21st day and 28th day after operation, all alive rats were anesthetized with 2.5% sodium pentobarbital and killed mercyfully. Blood samples and tissue specimens of liver, lung, spleen, kidney, thymus,

intestinal muscoa, and mesenteric lymph node were then collected, respectively.

2.2.4. Determination of Experimental Parameters

(1) Observation of Mortality Rate and Pathological Changes. The mortality rates of rats in various groups were recorded. The gross pathological changes and pathological changes under light microscopy of different tissues were observed, respectively.

(2) Determination of the Levels of Plasma Endotoxin, PLA₂, and Serum TNF- α . The determination of these parameters was conducted according to the instructions provided with the kits.

2.2.5. Statistical Analysis. The compiled data were first put into Excel sheet, and then read SPSS15.0 for further analysis. Normal data were expressed as means (standard deviation) while abnormal data were expressed as medians (interquartile range). Analysis of variance and pairwise comparisons were used for normal data, whereas abnormal data were subjected to non-parametric test, among which Kruskal-Wallis H test was used for pairwise comparisons and Mann-Whitney U test for multiple comparisons. Yates' chi-square test (χ^2) was used for intergroup comparisons of mortality rates.

3. Results

3.1. Comparison of Mortality Rate. 4 and 7 rats died in the model-control groups on 21st and 28th day, respectively; and 2 and 4 rats died in the treated groups on 21st and 28th day. On the 21st day, the mortality rates in the sham-operated groups were significantly lower than those in the model-control groups ($P = .032$); On 28 d, the mortality rate in the sham-operated group was significantly lower than those in both the model-control group ($P = .006$) and the treated group ($P = .032$), and the difference was statistically significant. Compared with model-control group, though the number of dead OJ rats in the treated group declined, no statistical difference was noted ($P > .05$).

3.2. Pathological Changes of the Multiple Organs Under Gross and Light Microscopy

3.2.1. Liver

(1) Sham-Operated Group

Gross. The liver was normal and showed no obvious pathological changes.

Under the Light Microscope. There was no marked difference among various time points after operation, and the liver was basically normal. The focal infiltrations of inflammatory cells were occasionally seen in few hepatic tissues, see Figure 1.

FIGURE 1: Sham-operated group 28 days; normal liver cells; HE $\times$ 200.

FIGURE 2: Model-control group 21 days; degeneration and swelling of hepatic cells as well as hyperplasia of bile ducts; HE $\times$ 200.

(2) Model-Control Group

Gross. The gross pathological changes manifested as an increase in size and hypertrophy. On 21st and 28th d, 80 percent of livers had a thickness of 0.8 cm. The section plane showed yellow plaques and was slightly oily. The texture of the liver became fragile, the bile ducts were dilated and enlarged, and even formed cysts (an average size of approximately $2 \times 2 \times 1 \text{ cm}^3$).

Under the Light Microscope. There were aggravated with the increase of postoperative duration. These pathological changes mainly manifested as swelling and obscure boundary in the majority of hepatic cells, the narrowing or partial disappearance of some hepatic sinusoids, as well as vasculature and dissolution of hepatic cells in occasional part of the liver. The number of apoptotic bodies increased. The connective tissue in the most part of the portal area showed stellated hyperplasia. The bile ducts, whose surroundings revealed acute and chronic inflammatory cell infiltration, were enlarged and hyperproliferated. The limiting plate of hepatocytes were destroyed and showed focal necrosis, see Figures 2 and 3.

(3) Treated Group

Gross. In rats treated with *S. miltiorrhizae*, the liver was mildly enlarged in size, showed yellowish-brown color, and

FIGURE 3: Model-control group 28 days; hyperplasia of small hepatic bile ducts and obvious degeneration of hepatic cells; HE $\times$ 200.

FIGURE 4: Sham-operated group 28 days; granular substances were seen in the space between glomerular capillaries as well as in the glomerulus and the cytoplasm of renal tubular epithelial cells, and patchy necrosis of renal tubular epithelium was also observed; HE $\times$ 200.

had a sharper edge than that in model-control group. The left and right hepatic ducts as well as the proximal common bile duct were dilated to form cysts that had a size smaller than those in model-control group.

Under the Light Microscope. The pathological changes showed no obvious difference between 21 and 28 days after operation, but there were mitigated to varying degrees when compared with those in the model-control group.

3.2.2. Kidney

(1) Sham-Operated Group

Gross. No obvious abnormality was seen.

Under the Light Microscope. No obvious changes were seen in kidney, and an edema of renal tubular epithelial cells could be seen occasionally.

(2) Model-Control Group

Gross. Diffuse yellow staining of renal capsule, cortex, and medulla was seen. The renal cortex in nearly half of

FIGURE 5: Treated group 21 days; obvious glomerular congestion and renal tubular degeneration; HE $\times$ 200.

FIGURE 6: Treated group 28 days; turbidity and degeneration of renal tubular epithelial cells as well as vascular congestion; HE $\times$ 200.

rats showed brownish-black color. The kidney showed no changes in size and texture.

Under the Light Microscope. Swelling and necrosis of proximal or distal tubules, bile pigment deposition, bile pigment casts within the lumen of renal tubules, patchy necrosis and bile pigment deposition in renal tubules, and inflammatory cell infiltration as well as hyperplasia and edema of connective tissue in renal interstitium were seen. These pathological changes were aggravated with the prolongation of time, see Figure 4.

(3) Treated Group

Gross. Renal cortex in 80% of rats in treated group showed dark brown color, which was lighter than that in model-control group; renal medulla showed yellow color. The kidney showed no changes in size and texture.

Under the Light Microscope. Mainly manifested as hyperplasia of glomerular capillary endothelial cells, epithelial cells, mesangial cells, and basement membrane were milder than those in model-control group, see Figures 5 and 6.

FIGURE 7: Sham-operated group 28 days; normal lung tissue; HE $\times$ 200.

3.2.3. Lung

(1) Sham-Operated Group

Gross. The color and shape of the lungs were normal, and there was no exudate in the thoracic cavity.

Under the Light Microscope. There was no obvious difference in pathological changes. The majority of lung tissues showed normal structure. Edema and infiltrated inflammatory cells were seen in a very small part of pulmonary interstitium and alveolar space. Occasional widening of alveolar septum, capillary distention, and congestion were observed.

(2) Model-Control Group

Gross. Dark red plaques were seen in the lungs of some rats.

Under the Light Microscope. The pathological alterations in model-control group were slightly aggravated with an increase in postoperative duration. On day 21, the lung of the majority of rats showed edema in the interstitium and alveolar wall as well as capillary distension in the alveolar wall while the lung of some rats showed widening of the alveolar septum. On day 28 in model-control group, the lung of the majority of rats showed hyperplasia and widening of interstitial fiber in the alveolar wall as well as edema and inflammatory cell infiltration in the interstitium and alveolar wall, see Figure 7.

(3) Treated Group

Gross. No marked difference in pathological changes was observed compared to those in model-control group.

Under the Light Microscope. On day 21, the lung of the majority of rats showed edema in the alveolar wall and inflammatory cell infiltration in the interstitium. The lung of some rats showed hyperplasia and widening of the alveolar wall as well as vascular distention. The lung of individual rats was normal. On day 28, the lung of the majority of rats showed edema and inflammatory cell infiltration in the interstitium and alveolar wall as well as hyperplasia and

FIGURE 8: Treated group 28 days; capillary distension and congestion in the alveolar wall as well as inflammatory cell infiltration in the alveolar septum; HE $\times 400$.

FIGURE 9: Sham-operated group 28 days; hyperplasia of cells in the splenic sinus, enlargement of blood sinus, inflammatory cell infiltration, and hemorrhage; HE $\times 200$.

widening of interstitial fiber while the lung of individual rats was normal. Overall, the pathological alterations in treated group were milder than those in model-control group, see Figure 8.

3.2.4. Intestinal Mucosa

(1) Sham-Operated Group

Gross. No obvious abnormality was seen.

Pathological Changes under Light Microscopy. No obvious difference in pathological changes were observed. The intestinal mucosa was normal in the majority of rats. The intestinal mucosa epithelium was not intact in very few rats. Inflammatory cell infiltration was seen in proper layer.

(2) Model-Control Group

Gross. Yellow staining of the intestinal wall and peritoneum were seen in all rats.

Pathological Changes under Light Microscopy. No obvious difference in pathological changes was observed. On day

FIGURE 10: Sham-operated group 28 days; enlargement of splenic sinusoid and hyperplasia of fibrous tissue; HE $\times 200$.

21 after operation, intestinal mucosa was not intact in the majority of rats, the edema of proper layer, submucous layer, and serosal layer was seen in the majority of rats, and very few rats showed no abnormality of the intestinal mucosa. On 28 d after operation, focal necrosis in intestinal mucosa epithelium, as well as the edema of proper layer, submucous layer, and serosal layer were seen in the majority of rats.

(3) Treated Group

Gross. No obvious difference was observed when compared to those in model-control group.

Pathological Changes under Light Microscopy. No obvious difference in pathological changes was observed among each time points after operation. Inflammatory cell infiltration was seen in proper layer, submucous layer, and serosal layer in the majority of rats. Intestinal mucosa was normal in some rats, and intestinal mucosa was not intact in very few rats. On 21 d after operation, some rats showed no abnormality of the intestinal mucosa; inflammatory cell infiltration was seen in proper layer in some rats; intestinal mucosa was not intact in very few rats. On 28 d after operation, some rats showed no abnormality of the intestinal mucosa; inflammatory cell infiltration was seen in proper layer, submucous layer, and serosal layer in some rats; and intestinal mucosa was not intact in very few rats.

3.2.5. Spleen

(1) Sham-Operated Group

Gross. The morphology of spleen was normal, and no obvious pathological changes were seen.

Pathological Changes under Light Microscopy. The spleen was roughly normal in all rats.

(2) Model-Control Group

Gross. The spleen became enormous, with an average size of $4 \times 1 \times 1$ cm. The texture of spleen became fragile, and the color of spleen became purple black.

FIGURE 11: Treated group 28 days; enlargement and congestion of splenic sinusoid; HE $\times$ 200.

Pathological Changes under Light Microscopy. The fusion, enlargement, or spotty necrosis of follicular germinal centers in the white pulp of spleen, the hyperplasia of the fibrous tissue in the sinus, as well as splenic arteriolar sclerosis were seen in few rats. The spleen was roughly normal in some rats, see Figures 9 and 10.

(3) Treated Group

Gross. The size of spleen increased, with an average size of $3 \times 1 \times 0.5$ cm.

Pathological changes under light microscopy. No significant difference in pathological changes was noted among each time point after operation. The spleen was roughly normal in the majority of rats. Splenic arteriolar sclerosis was seen in few rats, see Figure 11.

3.2.6. Thymus

(1) Sham-Operated Group

Gross. The morphology of thymus was normal, and no obvious pathological changes were seen.

Pathological changes under light microscopy. No significant difference in pathological changes was noted among each time point after operation. The thymic tissue was roughly normal in all rats.

(2) Model-Control Group

Gross. The thymus significantly shrank and became jaundice in all rats.

Pathological Changes under Light Microscopy. No significant difference in pathological changes was noted among each time point after operation. The thymic tissue was roughly normal in the majority of rats. Obscure boundary between thymic cortex and medulla was occasionally seen. The pathological changes were similar between 21 and 28 days after operation: the thymic tissue was roughly normal in

FIGURE 12: Sham-operated group 28 days; hyperplasia of sinus cell and inflammatory cell infiltration in lymph node; HE $\times$ 200.

FIGURE 13: Treated group 28 days; normal lymph node; HE $\times$ 100.

the majority of rats, and obscure boundary between thymic cortex and medulla was occasionally seen.

(3) Treated Group

Gross. The thymus became slightly jaundice but showed no obvious shrinkage.

Pathological Changes under Light Microscopy. No significant difference in pathological changes was noted among each time point after operation. The thymic tissue was normal. Obscure boundary between thymic cortex and medulla was occasionally seen. The thymic tissue was roughly normal in the majority of rats, and obscure boundary between thymic cortex and medulla of was seen in some rats.

3.2.7. Lymph Nodes

(1) Sham-Operated Group

Gross. The morphology of lymph nodes was normal.

Pathological changes under light microscopy. No marked difference in pathological changes was observed. The morphology and structure of lymph nodes were roughly normal. The enlargement of the follicular germinal centers and the hyperplasia of sinus cells were seen in few rats.

TABLE 1: Comparison of endotoxin, PLA₂, and TNF- α level.

Indexes	Sham-operated group		Model-control group		Treated group	
	21 d	28 d	21 d	28 d	21 d	28 d
Endotoxin (Eu/mL)	0.2 $\pm$ 0.05	0.3 $\pm$ 0.04	0.7 $\pm$ 0.1**	0.8 $\pm$ 0.1**	0.5 $\pm$ 0.1** ++	0.5 $\pm$ 0.1** ++
PLA ₂ (U/mL)	207.2 $\pm$ 38.6	203.0 $\pm$ 31.0	701.0 $\pm$ 151.7**	738.6 $\pm$ 60.2**	593.2 $\pm$ 146.3**	645.6 $\pm$ 95.1** ++
TNF- α (pg/mL)	14.2 $\pm$ 3.2	15.6 $\pm$ 4.9	31.5 $\pm$ 11.0**	57.70 $\pm$ 13.0**	24.4 $\pm$ 9.2**	44.2 $\pm$ 23.1**

Compare to sham-operated group, * P < .05, ** P < .01; compare to model-control group, + P < .05, ++ P < .01.

(2) Model-control group

Gross. Lymph nodes became yellow in half of the rats, the texture of lymph nodes showed no changes.

Pathological changes under light microscopy. No marked difference in pathological changes was observed. The enlargement of the follicular germinal centers and the hyperplasia of sinus cells were seen in the majority of rats, and spotty necrosis could be seen in the mantle zone and germinal centers, see Figure 12.

(3) Treated Group

Gross. The pathological changes showed no marked difference compared with those in model-control group.

Pathological Changes under Light Microscopy. No marked difference in pathological changes was observed. The boundary of the follicular germinal centers in lymph nodes was clear. The enlargement of the follicular germinal centers and the hyperplasia of sinus cells were seen in the majority of rats; and few rats showed no obvious pathological changes in lymph nodes, see Figure 13.

3.3. Comparison of the Content of Endotoxin in Plasma. On 21st and 28th day, the contents in the sham-operated group were significantly lower than those in model-control group and treated group (P < .01), and those of treated group were significantly lower than that in model-control group (P < .05), see Table 1.

3.4. Comparison of the Content of PLA₂ in Plasma. On 21 and 28 d, the contents in sham-operated group were significantly lower than those in model-control group and treated group (P < .01). On 28th day, the content in treated group was significantly lower than that in model-control group (P < .01), see Table 1.

3.5. Comparison of the Content of TNF- α in Serum. On 21 and 28 d, the contents of serum TNF- α in the sham-operated group were significantly lower than those in the model-control group and the treated group (P < .01), see Table 1.

4. Discussion

When obstructive jaundice (OJ) develops, the hepatic sinusoid is enlarged, endothelial cells swell, and many bacteria

and toxins enter into the systemic circulation via hepatic veins and induce endotoxemia. As a result, systemic infection and multiorgan damage were caused [16–18]. In the present study, varying degrees of pathological alterations in multiple organs of obstructive jaundice rats were observed. Under light microscopy, obvious biliary cirrhosis in the liver, congestion and edema in the lung, as well as obvious inflammatory reaction in the intestinal mucosa and kidney were seen. These pathological alterations were identical to those reported in literature. For this reason, it is of important significance to protect these organs in the treatment of OJ.

Some researchers have reported the therapeutic effect of *S. miltiorrhizae* injection against OJ [6, 19]. They found that *S. miltiorrhizae* could significantly improve liver damage in OJ animals, protect intestinal mucosal barrier function in patients with early OJ and promote perioperative recovery of renal function in OJ patients. In the present study, we observed the pathological alterations in multiple organs of OJ rats on days 21 and 28, which represented pathological manifestations of late-stage OJ. *S. miltiorrhizae* showed some protective effects on the majority of organs of rats with late-stage OJ, especially prominent on the liver, kidney and spleen. The protective effects of *S. miltiorrhizae* on the lung, lymph nodes and intestinal mucosa were relatively weak. The observation that *S. miltiorrhizae* had a weak protective effect on the intestinal mucosa may be due to reduced repair capacity for the damaged intestinal mucosa in the late stage of OJ.

During the development process of OJ, an important contributing factor to multiorgan damage is excessive inflammatory reaction mediated by inflammatory mediators such as endotoxin, TNF- α , and PLA₂. OJ is often complicated with intestinal endotoxemia to generate endotoxin that can activate Kupffer cells to produce a large number of inflammatory mediators such as NO, TNF- α , and oxygen-free radicals. These inflammatory mediators are involved in inducing functional damage to multiple organs such as the liver and kidney. TNF- α is thought to be one of the most critical inflammatory mediators that can mediate endotoxin-induced damage [20–22]. Beside directly acting upon its receptors to induce multiorgan damage, TNF- α can also induce the production of IL-1 and IL-6 that can amplify its biological effects to form a cascade reaction and thereby cause the damage to the lung, liver, and intestinal mucosa [16, 18, 23, 24]. An increase in endotoxin and TNF- α level can directly alter renal hemodynamics and induce the redistribution of renal blood flow, thereby leading to renal cortical ischemia as well as necrosis of renal tubules and

cortex [11, 13, 22]. In animal, OJ can also induce pulmonary edema as well as apoptosis and necrosis of splenic cells, whose extent is positively correlated with the contents of endotoxin and TNF- α in the serum [5, 16, 25–27]. PLA₂ is another important inflammatory mediator contributing to OJ-induced multiorgan damage. It has direct toxic effects on renal tubular epithelial cells and can increase pulmonary capillary permeability [28–30]. Therefore, its activity is correlated with the extent of lung injury [31, 32]. TNF- α is able to activate PLA₂ and, together with endotoxin, is involved in the development and progression of OJ-induced multiorgan damage. At present, it is known that *S. miltiorrhizae* is a traditional Chinese drug that has relatively unambiguous antiendotoxin effects and can directly neutralize and destroy endotoxin [33–35]. It has been pointed out in some studies that *S. miltiorrhizae* can protect hepatic cells and maintain liver function through reducing inflammatory mediator levels and improving microcirculation [6, 9]. *S. miltiorrhizae* is also able to significantly improve the liver function of OJ patients at the early postoperative stage [36]. In OJ, *S. miltiorrhizae* is able to reduce endotoxemia, regulate the production and secretion of vasoactive substances, and improve renal blood perfusion [37], thereby exerting protective effects against OJ-induced damage to kidney function [10].

In this study, we found that the mortality rates of OJ rats in model-control group were almost two times high as those in treated group, suggesting that *S. miltiorrhizae* injection can indeed reduce the mortality rate of rats. However, no statistically significant difference in the mortality rate of rats was observed, which may be due to the limited sample numbers. We also found that the contents of endotoxin and PLA₂ of treated group were significantly lower than those in model-control group. However, after careful analysis, we found that the absolute median values of the serum TNF- α content of treated group were actually reduced. We guessed that *S. miltiorrhizae* injection can indeed reduce the TNF- α content. The main reason for the absence of significant difference may be due to the lower number of experimental rats. In follow-up studies, we plan to increase the number of experimental rats so that the result, that is, consistent with the speculation could be obtained. Thus, we speculate that the protective effects of *S. miltiorrhizae* against OJ-induced multi-organ damage in rats may be associated with reducing the contents of endotoxin, TNF- α , and PLA₂.

Currently, the mechanism underlying the protective effects of *S. miltiorrhizae* against OJ-induced multi-organ damage has not been fully clarified and double-blind control studies of *S. miltiorrhizae* with a large sample size are lacking. Therefore, our future study will be aimed to further explore these relevant issues. This experiment results provided an experimental basis for treatment of OJ patients with *S. miltiorrhizae*.

Acknowledgments

This work was supported by technological foundation project of Traditional Chinese Medicine Science of Zhejiang province (no. 2003C130; no. 2004C142), foundation project for medical science and technology of Zhejiang province

(no. 2003B134), grave foundation project for technological and development of Hangzhou (no. 2003123B19), intensive foundation project for technology of Hangzhou (no. 2004Z006), foundation project for medical science and technology of Hangzhou (no. 2003A004), and foundation project for technology of Hangzhou (no. 2005224).

References

- [1] S. F. Assimakopoulos, C. D. Scopa, and C. E. Vagianos, "Pathophysiology of increased intestinal permeability in obstructive jaundice," *World Journal of Gastroenterology*, vol. 13, no. 48, pp. 6458–6464, 2007.
- [2] V. Papadopoulos, D. Filippou, E. Manolis, and K. Mimidis, "Haemostasis impairment in patients with obstructive jaundice," *Journal of Gastrointestinal and Liver Diseases*, vol. 16, no. 2, pp. 177–186, 2007.
- [3] S. F. Assimakopoulos, A. G. Mavrakis, K. Grintzalis, et al., "Superoxide radical formation in diverse organs of rats with experimentally induced obstructive jaundice," *Redox Report*, vol. 13, no. 4, pp. 179–184, 2008.
- [4] N. Bulbul, M. A. Akkus, Z. Cetinkaya, et al., "Effects of melatonin and lactulose on the liver and kidneys in rats with obstructive jaundice," *Pediatric Surgery International*, vol. 18, no. 8, pp. 677–680, 2002.
- [5] Z. Xiping, W. Dijiong, L. Jianfeng, et al., "Effects of *Salvia miltiorrhizae* on ICAM-1, TLR4, NF-kappaB and Bax proteins expression in multiple organs of rats with severe acute pancreatitis or obstructive jaundice," *Inflammation*, vol. 32, no. 4, pp. 218–232, 2009.
- [6] X. P. Zhang and D. R. Liu, "Advances in research on the mechanism of *Salvia miltiorrhizae* in the treatment of obstructive jaundice," *Chinese Journal of Surgery of Integrated Traditional and Western Medicine*, vol. 12, no. 1, pp. 69–70, 2006 (Chinese).
- [7] N. Zhao, Z. X. Guo, and X. Zhao, "Chemical composition and pharmacological effects of *Salvia miltiorrhizae*," *World Phytochemicals*, vol. 22, no. 4, pp. 155–160, 2007 (Chinese).
- [8] M. Ding, T.-X. Ye, G.-R. Zhao, Y.-J. Yuan, and Z.-X. Guo, "Aqueous extract of *Salvia miltiorrhiza* attenuates increased endothelial permeability induced by tumor necrosis factor- α ," *International Immunopharmacology*, vol. 5, no. 11, pp. 1641–1651, 2005.
- [9] X. H. Chen, F. Wang, and D. W. Han, "Effects of *salvia miltiorrhiza* solution on endotoxemia and hepatic CD14 expression after obstructive jaundice," *Chinese Journal of Experimental Surgery*, vol. 20, no. 9, pp. 802–804, 2003 (Chinese).
- [10] M. J. Zhang, W. B. Yuan, G. L. Zhang, and F. Cen, "Effect of *salvia miltiorrhiza* on renal function in perioperative patient with obstructive jaundice," *Chinese Journal of Surgery of Integrated Traditional and Western Medicine*, vol. 12, no. 6, pp. 526–528, 2006 (Chinese).
- [11] B. Peng, A.-Y. Qiao, Y.-L. Luo, Z.-Y. Li, S. He, and Y.-T. Wu, "The mechanism and significance of cellular signal transduction in kidney glomerular mesangial cells of rat with obstructive jaundice," *Sichuan Da Xue Xue Bao Yi Xue Bao*, vol. 38, no. 1, pp. 116–118, 2007 (Chinese).
- [12] Z. Xiping, L. Daren, W. Dijiong, et al., "Effect of *Salvia miltiorrhizae* on the expressions of TLR4 protein in the liver of rats with SAP or OJ," *Inflammation*, vol. 32, no. 3, pp. 151–162, 2009.
- [13] X.-P. Zhang, G.-H. Feng, J. Zhang, et al., "Protective effects of *Salvia miltiorrhizae* on the hearts of rats with severe

- acute pancreatitis or obstructive jaundice," *Journal of Zhejiang University: Science B*, vol. 10, no. 3, pp. 193–202, 2009.
- [14] X. Zhang, L. Chen, J. Zhang, et al., "Effect of *Salvia miltiorrhiza* on apoptosis and NF- κ B p65 expression in the liver of rats with severe acute pancreatitis or obstructive jaundice," *Journal of Gastroenterology and Hepatology*, vol. 24, no. 5, pp. 841–852, 2009.
 - [15] Z. Xiping, C. Yang, W. Dijiong, et al., "Effects of *Salvia miltiorrhiza* on intercellular adhesion molecule 1 protein expression in the lungs of rats with severe acute pancreatitis or obstructive jaundice," *Pancreas*, vol. 38, no. 3, pp. 309–317, 2009.
 - [16] E. A. Beierle, J.-N. Vauthey, L. L. Moldawer, and E. M. Copeland III, "Hepatic tumor necrosis factor- α production and distant organ dysfunction in a murine model of obstructive jaundice," *American Journal of Surgery*, vol. 171, no. 1, pp. 202–206, 1996.
 - [17] E. Bjornsson, J. Gustafsson, J. Borkman, and A. Kilander, "Fate of patients with obstructive jaundice," *Journal of Hospital Medicine*, vol. 3, no. 2, pp. 117–123, 2008.
 - [18] V. G. Margaritis, K. S. Filos, M. A. Michalaki, et al., "Effect of oral glutamine administration on bacterial translocation, endotoxemia, liver and ileal morphology, and apoptosis in rats with obstructive jaundice," *World Journal of Surgery*, vol. 29, no. 10, pp. 1329–1334, 2005.
 - [19] R. Xia and X. Chen, "Effects of Danshen injection on the malignant obstructive jaundice in the SD rat model," *Journal of Huazhong University of Science and Technology*, vol. 26, no. 6, pp. 686–689, 2006 (Chinese).
 - [20] J. Dawiskiba, P. Kornafel, D. Kwiatkowska, and M. Zimecki, "Alterations of tumor necrosis factor- α and interleukin 6 production and activity of the reticuloendothelial system in experimental obstructive jaundice in rats," *HPB*, vol. 4, no. 1, pp. 11–19, 2002.
 - [21] J.-Y. Hong, E. F. Sato, K. Hiramoto, M. Nishikawa, and M. Inoue, "Mechanism of liver injury during obstructive jaundice: role of nitric oxide, splenic cytokines, and intestinal flora," *Journal of Clinical Biochemistry and Nutrition*, vol. 40, no. 3, pp. 184–193, 2007.
 - [22] M. Ljungdahl, J. Osterberg, U. Ransjou, L. Engstrand, and U. Haglund, "Inflammatory response in patients with malignant obstructive jaundice," *Scandinavian Journal of Gastroenterology*, vol. 42, no. 1, pp. 94–102, 2007.
 - [23] Y. F. Tian, Y. Li, Q. Zhao, et al., "Effect of ulinastatin on intestinal mucosal barrier function of rats with obstructive jaundice," *Nan Fang Yi Ke Da Xue Xue Bao*, vol. 27, no. 7, pp. 987–990, 2007 (Chinese).
 - [24] T.-Z. Liu, K.-T. Lee, C.-L. Chern, J.-T. Cheng, A. Stern, and L.-Y. Tsai, "Free radical-triggered hepatic injury of experimental obstructive jaundice of rats involves overproduction of proinflammatory cytokines and enhanced activation of nuclear factor κ B," *Annals of Clinical and Laboratory Science*, vol. 31, no. 4, pp. 383–390, 2001.
 - [25] P. Sileri, S. Morini, G. S. Sica, et al., "Bacterial translocation and intestinal morphological findings in jaundiced rats," *Digestive Diseases and Sciences*, vol. 47, no. 4, pp. 929–934, 2002.
 - [26] M. E. Sewnath, T. van der Poll, C. J. F. van Noorden, F. J. W. ten Kate, and D. J. Gouma, "Cholestatic interleukin-6-deficient mice succumb to endotoxin-induced liver injury and pulmonary inflammation," *American Journal of Respiratory and Critical Care Medicine*, vol. 169, no. 3, pp. 413–420, 2004.
 - [27] G. Lazar Jr., A. Paszt, J. Kaszaki, et al., "Kupffer cell phagocytosis blockade decreases morbidity in endotoxemic rats with obstructive jaundice," *Inflammation Research*, vol. 51, no. 10, pp. 511–518, 2002.
 - [28] H. Ito, M. Miyazaki, I. Udagawa, H. Koshikawa, and K. Okui, "Changes in lysosomal enzymes and cell damage of the liver in obstructive jaundiced rats," *Nippon Geka Gakkai Zasshi*, vol. 93, no. 2, pp. 144–149, 1992.
 - [29] X.-P. Zhang, H. Tian, Y.-H. Lai, et al., "Protective effects and mechanisms of Baicalin and octreotide on renal injury of rats with severe acute pancreatitis," *World Journal of Gastroenterology*, vol. 13, no. 38, pp. 5079–5089, 2007.
 - [30] S. Goto, H. Nakamura, H. Morooka, Y. Terao, O. Shibata, and K. Sumikawa, "Role of reactive oxygen in phospholipase A_2 activation by ischemia/reperfusion of the rat kidney," *Journal of Anesthesia*, vol. 13, no. 2, pp. 90–93, 1999.
 - [31] H.-Q. Zhang, H.-D. Wang, D.-X. Lu, et al., "Berberine inhibits cytosolic phospholipase A_2 and protects against LPS-induced lung injury and lethality independent of the α_2 -adrenergic receptor in mice," *Shock*, vol. 29, no. 5, pp. 617–622, 2008.
 - [32] G. Kostopanagiotou, E. Avgerinos, C. Costopanagiotou, et al., "Acute lung injury in a rat model of intestinal ischemia-reperfusion: the potential time depended role of phospholipases A_2 ," *Journal of Surgical Research*, vol. 147, no. 1, pp. 108–116, 2008.
 - [33] Z. H. Li, B. L. Han, and J. You, "Anti-endotoxin treatment of patients with obstructive jaundice," *Chinese Journal of Experimental Surgery*, vol. 15, no. 2, p. 119, 1998 (Chinese).
 - [34] K. Hu, W. L. Shen, and Y. G. Luo, "Study on the anti-endotoxin effects of *salvia miltiorrhiza* solution (SMS) in vitro," *Journal of Shantou University Medical College*, vol. 13, no. 3, pp. 16–17, 2000 (Chinese).
 - [35] B. Peng, J. Du, Q. Jia, et al., "The effect of *salvia miltiorrhiza* and shengmai on inflammatory mediator and renal function of post-operative patients with obstructive jaundice," *Hua Xi Yi Ke Da Xue Xue Bao*, vol. 32, no. 4, pp. 587–589, 2001 (Chinese).
 - [36] J. M. He, Z. J. Tan, and X. S. Zhong, "Effect of post-operative treatment with *salvia miltiorrhiza* on liver function of patients with obstructive jaundice," *Chinese Journal of Modern Medicine*, vol. 17, no. 12, pp. 1478–1480, 2007 (Chinese).
 - [37] Y. Zhang, Z. Wang, L. Yu, S. Han, and L. Zheng, "Observation on apoptosis of renal cell in the rat with obstructive jaundice and *salvia miltiorrhiza*'s effect on apoptosis," *Journal of Xi'an Jiaotong University (Medical Sciences)*, vol. 27, no. 1, pp. 42–45, 2006 (Chinese).