Research Article

Analysis of Several PLA₂ mRNA in Human Meningiomas

Yves Denizot,¹ Rafael De Armas,² Karine Durand,² Sandrine Robert,² Jean-Jacques Moreau,³ François Caire,³ Nicolas Weinbreck,² and François Labrousse²

¹ UMR CNRS 6101, Centre National de la Recherche Scientifique, Faculté de Médecine, Université de Limoges, 2 rue Dr. Marcland, 87025 Limoges, France

² Service d'Anatomie Pathologique, CHU Dupuytren, 87045 Limoges, France

³ Service de Neurochirurgie, CHU Dupuytren, 87045 Limoges, France

Correspondence should be addressed to Yves Denizot, yves.denizot@unilim.fr

Received 7 September 2009; Revised 25 November 2009; Accepted 21 December 2009

Recommended by Muzamil Ahmad

In view of the important oncogenic action of phospholipase A₂ (PLA₂) we investigated PLA₂ transcripts in human meningiomas. Real-time PCR was used to investigate PLA₂ transcripts in 26 human meningioma tumors. Results indicated that three Ca²⁺-dependent high molecular weight PLA₂ (PLA₂-IVA, PLA₂-IVB, PLA₂-IVC), one Ca²⁺-independent high molecular weight PLA₂ (PLA₂-VI) and five low molecular weight secreted forms of PLA₂ (PLA₂-IB, PLA₂-IIA, PLA₂-III, PLA₂-V, and PLA₂-XII) are expressed with PLA₂-IVA, PLA₂-IVB, PLA₂-VI, and PLA₂-XIIA as the major expressed forms. PLA₂-IIE, PLA₂-IIF, PLA₂-IVD, and PLA₂-XIIB are not detected. Plasma (PLA₂-VIIA) and intracellular (PLA₂-VIIB) platelet-activating factor acetylhydrolase transcripts are expressed in human meningiomas. However no difference was found for PLA₂ transcript amounts in relation to the tumor grade, the subtype of meningiomas, the presence of inflammatory infiltrated cells, of an associated edema, mitosis, brain invasion, vascularisation or necrosis. In conclusion numerous genes encoding multiples forms of PLA₂ are expressed in meningiomas where they might act on the phospholipid remodeling and on the local eicosanoid and/or cytokine networks.

Copyright © 2009 Yves Denizot et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Meningiomas are the second most common primary intracranial tumor. Meningiomas present clinically by causing focal or generalized seizure disorders, focal neurological deficits or neuropsychological decline [1]. A large number of molecular and genetic pathways that are altered in brain tumor cells have been identified. Among them, a possible role for the eicosanoid cascade has been suggested in meningiomas [2]. Phospholipase A₂ (PLA₂) is the key enzyme involved in eicosanoid generation [3–5]. PLA₂ catalyzes the hydrolysis of the *sn*-2 position of membrane glycerophospholipids to liberate the eicosanoid precursor arachidonic acid (AA) [3–5]. Four distinct families have been documented: low molecular weight secreted forms of PLA₂ (sPLA₂), Ca²⁺-dependent high molecular weight PLA₂ (cPLA₂), Ca²⁺-independent high molecular weight PLA₂ (iPLA₂); and the platelet-activating factor acetylhydrolase (PAF-AH). The sPLA₂ family is implicated in several biological processes such as inflammation and host defense [3, 4].

Nine isoenzymes have been identified in human: PLA₂-IB, PLA₂-IIA, PLA₂-IID, PLA₂-IIE, PLA₂-IIF, PLA₂-III, PLA₂-V, PLA₂-X, PLA₂-XIIA, and PLA₂-XIIB. In addition to their function in digestion of dietary phospholipids, host defense against bacteria and AA release from cellular phospholipids for eicosanoid synthesis, two classes of receptors (M and N) and several extracellular binding proteins have been identified indicating that sPLA₂ might signal through receptor activation [3–5]. In human the cPLA₂ family consists of four members, PLA₂-IVA, PLA₂-IVB, PLA₂-IVC, and PLA₂-IVD; PLA₂-IVA being the central regulator of stimulus-coupled cellular AA release [3–5]. In human the iPLA₂ group consists of seven members, iPLA₂ (PLA₂-VIA-1) currently being the best known member and playing major role in phospholipids remodeling and cancer [3, 5]. Beside its important place for eicosanoid generation, PLA₂ is also the key enzyme for the generation of the pro-inflammatory lipid mediator PAF recently documented in human meningioma [6]. PAF-AH activity which hydrolysis PAF into the inactive PAF precursor, lyso-PAF is detected in meningioma [6].

However no results reported whether this enzymatic activity originated from PLA₂-VIIA and/or PLA₂-VIIB, the plasma PAF-AH and the intracellular PAF-AH forms, respectively. Currently the contribution of PLA₂ in meningiomas is poorly documented despite the fact that PLA₂ inhibition decreased the growth of cultured meningioma cells [7]. In view of the potentially important oncogenic action of the various PLA₂ species, we investigated, at the mRNA levels, which of them were expressed in intracranial human meningiomas.

2. Materials and Methods

2.1. Patients. The procedure of the present study followed the rules edited by the French National Ethics. Ethics approval was obtained from the ethics committee of our hospital (CHU Dupuytren, Limoges, France). Twenty six patients who underwent surgery for intracranial meningiomas (from 1998 to 2004) were investigated. Tumors were from the Service d'Anatomie Pathologique of the CHU Dupuytren (France). After undergoing the routine hospital analysis, the excess of sample was kept at -80°C until use and this in accordance with the regulations in force in France. No written or oral consent was obtained because it is a study of samples already collected and referred to research prior the French bioethical law (2004). Thus, ethics committee explicitly approved the waiver of consent. Normal meninges were not available in our institution in light of our ethic committee law. The low amounts (10–15 mg) of available tumors only permitted investigations at the mRNA level. Tumors were classified according to the WHO criteria [8]. There were 16 grade I meningiomas including 8 transitional (2 men, 6 women, mean age 60 years), 3 meningothelial (1 man, 2 women, mean age 60 years), and 5 fibrous (5 women, mean age 59 years). Height tumors were grade II meningiomas: 7 atypical (6 men, 1 woman, mean age 58 years) and 1 chordoid meningiomas (1 man, 54 years). Two tumors were classified as anaplastic grade III meningiomas (2 men, mean age 54 years). Necrosis was assessed using morphological criteria defined by the WHO classification of meningiomas: “foci of spontaneous or geographic necrosis”. The chronic inflammatory infiltrate was mainly lymphoplasmocytic.

2.2. RNA Extraction. Total RNA was extracted using the “RNeasy Lipid Tissue mini kit” (Qiagen, Courtaboeuf, France) from 10–15 mg of tumor tissue. Before RNA extraction, tumor fragments were incubated with 14mm ceramic beads “Lysing matrix D” (Bertin Technologies, Montigny-Le-Bretonneux, France) in 1 mL QIAzol lysis reagent and homogenized at 5500 rpm during 2-fold 40 sec in the automated mixer Precellys (Bertin Technologies). Then homogenized samples were used for RNA purification according to the manufacturer's protocol. A DNase I digestion step was included for each extraction to avoid RNA contamination by genomic DNA. RNA integrity was checked by capillary electrophoresis on the Bioanalyzer 2100 (Agilent Technologies, Massy, France). Only RNA with an RNA Integrity Number

(R.I.N) higher than 5.5 was used for reverse transcription. Total RNA concentration was determined by measuring absorbance at 260 nm with a spectrophotometre NanoDrop ND-1000 (Labtech, Paris, France).

2.3. Reverse Transcription. Total RNA was reverse transcribed in single strand cDNA using random hexamers and as described in the protocol of the “SuperScript III First-Strand Synthesis System for RT-PCR” (Invitrogen, Cergy-Pontoise, France). Briefly, 1 μg total RNA was incubated with 200 U M-MLV reverse transcriptase in the presence of 0.5 mM dNTPs, 50 ng random hexamers, 5 mM MgCl_2 , 10 mM dithiothreitol, and 40 U RNase inhibitor, in a final volume of 20 μL 1X RT buffer. Reverse transcription was performed as follows in a thermocycler Gold 9700 (Applied Biosystem): denaturation during 5 min at 65°C and chilling 1 min on ice, hybridization during 10 min at 25°C followed by cDNA synthesis during 50 min at 50°C ; enzyme was inactivated by a 5 min incubation at 85°C and chilling on ice; finally the destruction of the RNA portion of the RNA:cDNA hybrids was performed by 2 U RNase H during 20 min at 37°C . Reactions were frozen at -20°C until quantitative real-time PCR realisation.

2.4. Real-Time PCR Analysis. PLA₂-IB, PLA₂-IIA, PLA₂-IID, PLA₂-IIE, PLA₂-IIF, PLA₂-III, PLA₂-IVA, PLA₂-IVB, PLA₂-IVC, PLA₂-IVD, PLA₂-V, PLA₂-VI, PLA₂-X, PLA₂-XIIA, PLA₂-XIIB, PLA₂-VIIA (plasma PAF-AH), PLA₂-VIIB (intracellular PAF-AH), and PLA₂R transcripts were analyzed using real-time polymerase chain reaction (PCR). PCR was performed in duplicate by using TaqMan assay reagents (Applied Biosystems, Foster City, CA) [9, 10]. Product references were the following: PLA₂-IB: Hs00386701-m1; PLA₂-IIA: Hs00179898-m1; PLA₂-IID: Hs00173860-m1; PLA₂-IIE: Hs00173897-m1; PLA₂-IIF: Hs00224482-m1; PLA₂-III: Hs00210447-m1; PLA₂-IVA: Hs00233352-m1; PLA₂-IVB: Hs00979952-m1; PLA₂-IVC: Hs00234345-m1; PLA₂-IVD: Hs00603557-m1; PLA₂-V: Hs00173472-m1; PLA₂-VI: Hs001/85926-m1; PLA₂-X: Hs00358567-m1; PLA₂-XIIA: Hs00830106-s1; PLA₂-XIIB: Hs00261432-m1; PLA₂-VIIA: Hs00968593-m1; PLA₂-VIIB: Hs01042135-m1; PLA₂R: Hs00234853-m1. Real-time PCR were performed following the recommendations of the manufacturer in a final volume of 20 μL with 10 μL of 2X Universal PCR Master Mix, 20 ng of cDNA in a volume of 9 μL (the amount of cDNA is an equivalent based on the initial amount of RNA used for the RT reaction and 1 μL of a 20X TaqMan gene expression specific probe. PCR parameters were the following: 95°C for 10 min and forty cycles of $95^{\circ}\text{C}/15$ sec and $60^{\circ}\text{C}/60$ sec. Amplification products were analyzed on an ABI Prism 7000 system (Applied Biosystems) [9, 10]. Gene expression levels were normalized to 18S RNA (product reference: Hs99999901-s1) according to the manufacturer's recommendation. Amounts of various transcripts were compared to sample with the lowest level of transcripts (a patient who was arbitrary quoted 1). The relative quantification of gene expression was performed using the comparative C_T method ($\Delta\Delta C_T$) (Figure 1(a)). Experiments were made

FIGURE 1: Q-PCR analysis of PLA₂ transcripts in human meningioma tumors. (a) Representative tracing of 18S and PLA₂-IVA of tumors from 2 different patients (duplicate samples). (b) Detection (in duplicate) of 18S, PLA₂-IVA, PLA₂-IVB, PLA₂-IVC, PLA₂-IVD, and PLA₂-VI in a meningioma tumor.

in duplicate. Mean C_T values were used in the $\Delta\Delta C_T$ calculation by using the “relative quantitation calculation and analysis software for Applied Biosystems real-time PCR systems”. NonRT controls (only with RNA) and blank RT controls (RT without RNA) were run to make sure the amplifications were specific.

2.5. Data Analysis. Significance was assessed by using the Kruskal-Wallis test followed by a Mann-Whitney U -test.

3. Results and Discussion

In a first set of experiments we investigated if mRNAs derived from the five intracellular PLA₂ genes (four cPLA₂ and iPLA₂) were detected in meningiomas. As shown in Figure 1(b), mRNAs derived from four of these five cloned

PLA₂ genes are detected. Mean C_T values are reported in Table 1. PLA₂-IVD transcripts were not present at detectable levels. In contrast, PLA₂-IVA, PLA₂-IVB, PLA₂-IVC, and PLA₂-VI were detected in 96% (25/26), 100% (26/26), 92% (24/26), and 100% (26/26) of tumors, respectively (Figure 2). These results confirm a previous study highlighting PLA₂ activity in 100% of human meningiomas [6]. No difference ($P > .05$, Mann Whitney U -test) was found for PLA₂ transcript amounts in relation to the tumor grade (Figure 1), nor the subtype of meningiomas, the presence of inflammatory infiltrated cells, of an associated edema, mitosis, brain invasion, vascularisation or necrosis (data not shown). The analysis of twenty six patients indicated the following rank of magnitude in human meningiomas: PLA₂-VI = PLA₂-IVB > PLA₂-IVA > PLA₂-IVC (Table 1). PLA₂-IVB and PLA₂-IVC had little specificity for the *sn*-2 fatty acid as compared with PLA₂-IVA which preferentially hydrolyses

FIGURE 2: Cytosolic PLA₂ transcripts in human meningiomas. Sixteen-grade I and eleven-grade II+III meningiomas were investigated. Gene expression levels were normalized to 18S RNA. Amounts of transcripts were compared to sample with the lowest level of transcripts (a patient who was arbitrarily quoted 1). (○) indicates patients with no detectable transcripts. No significant differences were documented between groups.

phospholipids containing AA at the *sn*-2 position [3–5]. PLA₂-VI was originally reported to mediate phospholipid remodeling and, thus, to act as a housekeeping protein without significant role in cell growth [3, 4]. However several recent studies have demonstrated that PLA₂-VI exhibited roles in cell regulation, growth, and death. Especially, one mechanism by which PLA₂-VI mediates cell growth involves regulation of AA release, p53, and MAPK activation [11]. Of interest, involvements of p53 and MAPK kinase have been recently reported in the pathology of human meningiomas [12, 13]. A role for PLA₂-VI may, thus, be suggested in meningioma tumor growth. Together, these observations might suggest PLA₂-VI as a novel and interesting target for drug development for meningioma therapy. However, given the ubiquitous expression of PLA₂-VI and its role in glycerophospholipid metabolism, drug strategies targeting PLA₂-VI must exhibit selectivity to avoid undesired side effects.

In a second set of experiments we investigated if mRNAs derived from the nine sPLA₂ genes (i.e., PLA₂-IB, PLA₂-IIA, PLA₂-IID, PLA₂-IIE, PLA₂-IIF, PLA₂-III, PLA₂-V, PLA₂-X, PLA₂-XIIA, and PLA₂-XIIB) were detected in human meningiomas. PLA₂-IIE, PLA₂-IIF, and PLA₂-XIIB transcripts were not present at detectable levels in tumors while PLA₂-IID and PLA₂-X transcripts were detected in only a few number (4/26, 15%) of them. In contrast, PLA₂-IB, PLA₂-IIA, PLA₂-III, PLA₂-V and PLA₂-XIIA were detected in 88% (23/26), 88% (23/26), 77% (20/26), 65% (17/26), and 96% (25/26) of tumors, respectively (Figure 3). Mean C_T values are reported in Table 1. Results indicated the following rank of magnitude for sPLA₂ transcripts in meningiomas: PLA₂-XIIA > PLA₂-IIA > PLA₂-IB = PLA₂-V = PLA₂-III = PLA₂-IID > PLA₂-X. No difference ($P > .05$, Mann Whitney *U*-test) was found for sPLA₂ transcript amounts in relation to the tumor grade (Figure 3), nor the subtype of meningiomas, the presence of inflammatory infiltrated cells, of an associated edema,

FIGURE 3: Secreted PLA₂ transcripts in human meningiomas. Same legend as in Figure 2. (○) indicates patients with no detectable transcripts. No significant differences were documented between groups.

mitosis, brain invasion, vascularisation or necrosis (data not shown). PLA₂-XIIA, PLA₂-IIA, and PLA₂-IB might be implicated in meningioma growth. The physiologic roles of PLA₂-XIIA remain an open question. Whether PLA₂-XIIA exhibits a weak AA catalytic activity [14], a potential

role for this enzyme is suggested in membrane fusion or cell division [15]. PLA₂-XIIA was reported to inhibit bone morphogenetic protein (BMP) through the loss of activated Smad1/4 complexes [16]; a result of importance since BMP inhibits the tumorigenic potential of human glioblastomas

FIGURE 4: PAF-AH and PLA₂R transcripts in human meningiomas. Upper panel: plasma PAF-AH (PLA₂-VIIA) and intracellular PAF-AH (PLA₂-VIIB) in meningioma tumors. Fifteen-grade I and nine-grade II+III meningiomas were investigated. (○) indicates patients with no detectable transcript. No significant differences were documented between groups. Lower panel: PLA₂R transcripts in fifteen-grade I and nine-grade II+III meningiomas.

by triggering the Smad signaling cascade [17]. PLA₂-IB expression is mainly neuronal in human brain [18]. Apart from its lipolytic and pro-inflammatory activities, PLA₂-IB acts as receptor ligand to induce cell signaling and subsequent activation of cPLA₂, thus indirectly contributing to AA production [19]. However the role of PLA₂-IB as a ligand for the PLA₂R is still controversial. PLA₂-IIA elicits a mitogenic response and activates AA metabolism in astrocytoma cells [20], is critical for neuronal death via reactive oxygen species [21] and plays a role in cellular senescence [22]. Finally, studies have suggested the role of PLA₂-IIA, PLA₂-III, and PLA₂-V as potential prognostic markers in colorectal adenocarcinomas and prostate cancer [23, 24]. The data reported in Figure 3 suggest that levels of PLA₂-IIA, PLA₂-III, and PLA₂-V transcripts greatly varied between patients (see the Log scale). Would it be possible that their levels were related to patient outcomes in meningioma

tumors? Clearly investigation of a larger number of patients would be of interest to test this hypothesis.

In a third set of experiments we investigated PAF-AH enzymes that constitute another PLA₂ subfamily. As shown in Figure 4 (upper panel), PLA₂-VIIA (the plasma PAF-AH form) and PLA₂-VIIB (the intracellular PAF-AH form) were present in 100% (23/23) and 95% (22/23) meningioma tumors. Mean C_T values are reported in Table 1. No difference ($P > .05$, Mann Whitney U -test) was found for PAF-AH transcript amounts in relation to the tumor grade (Figure 4(a)), nor to other clinical data (data not shown). The present results confirm a previous study reporting PAF-AH enzymatic activity in meningioma homogenates [6]. The PAF-AH family exhibits unique substrate specificity toward PAF and oxidized phospholipids. Degradation of these bioactive phospholipids by PAF-AH may lead to the termination of inflammatory reaction. Its presence in human

TABLE 1: C_T values obtained during real-time PCR analysis.

	Mean $\pm$ SEM C_T (made on detectable samples)	Number of samples with a $C_T > 40$ (nondetectable samples)
18S	12.27 $\pm$ 0.18	0
PLA ₂ -IB	35.35 $\pm$ 0.29	3
PLA ₂ -IIA	33.69 $\pm$ 0.74	3
PLA ₂ -IID	34.91 $\pm$ 0.81	22
PLA ₂ -IIE	nd	26
PLA ₂ -IIF	nd	26
PLA ₂ -III	35.28 $\pm$ 0.50	6
PLA ₂ -IVA	32.3 $\pm$ 0.27	1
PLA ₂ -IVB	30.57 $\pm$ 0.33	0
PLA ₂ -IVC	33.9 $\pm$ 0.35	2
PLA ₂ -IVD	nd	26
PLA ₂ -V	35.81 $\pm$ 0.61	9
PLA ₂ -VI	30.78 $\pm$ 0.30	0
PLA ₂ -VIIA	33.19 $\pm$ 0.48	0
PLA ₂ -VIIB	30.25 $\pm$ 1.21	1
PLA ₂ -X	37.93 $\pm$ 0.26	22
PLA ₂ -XIIA	32.86 $\pm$ 0.33	1
PLA ₂ -XIIB	nd	26

Results are reported as mean $\pm$ SEM of 26 experiments excepted for PLA₂-VIIA and PLA₂-VIIB where 23 samples were analysed. nd: not detectable C_T .

meningioma is consistent with the presence of PAF in meningioma tumors [6, 25].

Finally in a fourth set of experiments we focused our attention on PLA₂R transcripts in human meningioma. As shown in Figure 4 (lower panel), PLA₂R transcripts were detected in 100% (23/23) meningioma tumors but without significant link with the tumor grade. The PLA₂R can act as a ligand for several sPLA₂ thus mediating a variety of biological responses (such as cell proliferation, cell migration, hormone release, lipid mediator production and cytokine production). In turn, PLA₂R can also play a negative role in sPLA₂ functions by downregulating their exaggerated reactions as PLA₂R is involved in the degradation/internalization of sPLA₂ [26]. Particularly, PLA₂R deficient mice exhibit resistance to endotoxic shock [27] and knockdown of the PLA₂R prevents the onset of replicative senescence and diminishes stress-induced senescence [28]. Finally PLA₂R was found to be upregulated in dermatofibrosarcoma [29].

In conclusion, numerous genes encoding multiples forms of cPLA₂, sPLA₂, and PAF-AH are expressed (at the mRNA level) in human meningiomas where they might act on tumor growth not only by acting on phospholipid remodeling but also by altering the local eicosanoid and/or cytokine networks. It is of evidence that immunohistochemistry would be of importance to confirm the relative expression of the different PLA₂ forms in human meningioma tumors. The discovery of specific receptors that bind sPLA₂ strongly indicate that these enzymes can exert various biological responses via binding to a receptor, in addition to their enzymatic activity. Of interest meningioma tumors expressed PLA₂R transcripts. Further studies are clearly needed to elucidate the contributions of sPLA₂, cPLA₂, iPLA₂, and PAF-AH in meningioma and to determine their possible relevance in the targeting of new therapeutic interventions.

Abbreviations

PLA₂: phospholipase A₂
sPLA₂: secreted phospholipase A₂
cPLA₂: cytosolic phospholipase A₂
iPLA₂: calcium independent phospholipase A₂
PCR: polymerase chain reaction.

Acknowledgments

This research was supported by Grants from “La Ligue Contre le Cancer” (comité de la Haute-Vienne) and “Lions Club de la Corrèze, Zone 33 district 103 Sud”. The founders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- [1] I. R. Whittle, C. Smith, P. Navoo, and D. Collie, “Meningiomas,” *The Lancet*, vol. 363, no. 9420, pp. 1535–1543, 2004.
- [2] N. Nathoo, G. H. Barnett, and M. Golubic, “The eicosanoid cascade: possible role in gliomas and meningiomas,” *Journal of Clinical Pathology*, vol. 57, no. 1, pp. 6–13, 2004.
- [3] I. Kudo and M. Murakami, “Phospholipase A₂ enzymes,” *Prostaglandins and Other Lipid Mediators*, vol. 68–69, pp. 3–58, 2002.
- [4] G. Lambeau and M. H. Gelb, “Biochemistry and physiology of mammalian secreted phospholipases A₂,” *Annual Review of Biochemistry*, vol. 77, pp. 495–520, 2008.
- [5] J. E. Burke and E. A. Dennis, “Phospholipase A₂ biochemistry,” *Cardiovascular Drugs and Therapy*, vol. 23, no. 1, pp. 49–59, 2009.
- [6] Y. Denizot, R. de Armas, F. Caire, I. Pommepuy, V. Truffinet, and F. Labrousse, “Platelet-activating factor and human

- meningiomas," *Neuropathology and Applied Neurobiology*, vol. 32, no. 6, pp. 674–678, 2006.
- [7] M. J. Petr, T. C. Oigitano, and R. D. Wurster, "PLA₂ activity regulates Ca²⁺ storage-dependent cellular proliferation," *Experimental Cell Research*, vol. 244, no. 1, pp. 310–318, 1998.
 - [8] D. N. Louis, H. Oghaki, O. D. Wiestler, and W. K. Cavenee, *WHO Classification of Tumours of the Central Nervous System*, IARC Press, Lyon, France, 2007.
 - [9] C. Vincent, R. Fiancette, M. Donnard, et al., "5-LOX, 12-LOX and 15-LOX in immature forms of human leukemic blasts," *Leukemia Research*, vol. 32, no. 11, pp. 1756–1762, 2008.
 - [10] R. Fiancette, C. Vincent, M. Donnard, et al., "Genes encoding multiple forms of phospholipase A₂ are expressed in immature forms of human leukemic blasts," *Leukemia*, vol. 23, no. 6, pp. 1196–1199, 2009.
 - [11] S. B. Hooks and B. S. Cummings, "Role of Ca²⁺-independent phospholipase A₂ in cell growth and signaling," *Biochemical Pharmacology*, vol. 76, no. 9, pp. 1059–1067, 2008.
 - [12] M. D. Johnson, M. O'Connell, F. Vito, and R. S. Bakos, "Increased STAT-3 and synchronous activation of Raf-1-MEK-1-MAPK, and phosphatidylinositol 3-kinase-Akt-mTOR pathways in atypical and anaplastic meningiomas," *Journal of Neuro-Oncology*, vol. 92, no. 2, pp. 129–136, 2009.
 - [13] Z. N. Chang, C.-L. Guo, I. Ahronowitz, A. O. Stemmer-Rachamimov, M. MacCollin, and F. P. Nunes, "A role for the p53 pathway in the pathology of meningiomas with NF2 loss," *Journal of Neuro-Oncology*, vol. 91, no. 3, pp. 265–270, 2009.
 - [14] M. H. Gelb, E. Valentin, F. Ghomashchi, M. Lazdunski, and G. Lambeau, "Cloning and recombinant expression of a structurally novel human secreted phospholipase A₂," *Journal of Biological Chemistry*, vol. 275, no. 51, pp. 39823–39826, 2000.
 - [15] M. Murakami, S. Masuda, S. Shimbara, et al., "Cellular arachidonate-releasing function of novel classes of secretory phospholipase A₂s (groups III and XII)," *Journal of Biological Chemistry*, vol. 278, no. 12, pp. 10657–10667, 2003.
 - [16] I. Munoz-Sanjuan and A. H. Brivanlou, "Induction of ectopic olfactory structures and bone morphogenetic protein inhibition by Rossy, a group XII secreted phospholipase A₂," *Molecular and Cellular Biology*, vol. 25, no. 9, pp. 3608–3619, 2005.
 - [17] S. G. M. Piccirillo, B. A. Reynolds, N. Zanetti, et al., "Bone morphogenetic proteins inhibit the tumorigenic potential of human brain tumour-initiating cells," *Nature*, vol. 444, no. 7120, pp. 761–765, 2006.
 - [18] M. Kolko, N. R. Christoffersen, H. Varoqui, and N. G. Bazan, "Expression and induction of secretory phospholipase A₂ group IB in brain," *Cellular and Molecular Neurobiology*, vol. 25, no. 7, pp. 1107–1122, 2005.
 - [19] E. Valentin and G. Lambeau, "Increasing molecular diversity of secreted phospholipases A₂ and their receptors and binding proteins," *Biochimica et Biophysica Acta*, vol. 1488, no. 1-2, pp. 59–70, 2000.
 - [20] L. Fuentes, M. Hernandez, M. L. Nieto, and M. Sanchez Crespo, "Biological effects of group IIA secreted phospholipase A₂," *FEBS Letters*, vol. 531, no. 1, pp. 7–11, 2002.
 - [21] G. H. Mathisen, I. H. Thorkildsen, and R. E. Paulsen, "Secretory PLA₂-IIA and ROS generation in peripheral mitochondria are critical for neuronal death," *Brain Research*, vol. 1153, no. 1, pp. 43–51, 2007.
 - [22] H. J. Kim, K. S. Kim, S. H. Kim, et al., "Induction of cellular senescence by secretory phospholipase A₂ in human dermal fibroblasts through an ROS-mediated p53 pathway," *Journals of Gerontology. Series A*, vol. 64, no. 3, pp. 351–362, 2009.
 - [23] T. Mirtti, V. J. O. Laine, H. Hiekkänen, et al., "Group IIA phospholipase A₂ as a prognostic marker in prostate cancer: relevance to clinicopathological variables and disease-specific mortality," *Acta Pathologica Microbiologica and Immunologica*, vol. 117, no. 3, pp. 151–161, 2009.
 - [24] C. M. Mounier, D. Wendum, E. Greenspan, J.-F. Fléjou, D. W. Rosenberg, and G. Lambeau, "Distinct expression pattern of the full set of secreted phospholipases A₂ in human colorectal adenocarcinomas: sPLA₂-III as a biomarker candidate," *British Journal of Cancer*, vol. 98, no. 3, pp. 587–595, 2008.
 - [25] Y. Hirashima, N. Hayashi, O. Fukuda, H. Ito, S. Endo, and A. Takaku, "Platelet-activating factor and edema surrounding meningiomas," *Journal of Neurosurgery*, vol. 88, no. 2, pp. 304–307, 1998.
 - [26] K. Hanasaki, "Mammalian phospholipase A₂: phospholipase A₂ receptor," *Biological and Pharmaceutical Bulletin*, vol. 27, no. 8, pp. 1165–1167, 2004.
 - [27] K. Hanasaki, Y. Yokota, J. Ishizaki, T. Itoh, and H. Arita, "Resistance to endotoxin shock in phospholipase A₂ receptor-deficient mice," *Journal of Biological Chemistry*, vol. 272, no. 52, pp. 32792–32797, 1997.
 - [28] A. Augert, C. Payré, Y. de Launoit, J. Gil, G. Lambeau, and D. Bernard, "The M-type receptor PLA₂R regulates senescence through the p53 pathway," *EMBO Reports*, vol. 10, no. 3, pp. 271–277, 2009.
 - [29] S. C. Linn, R. B. West, J. R. Pollack, et al., "Gene expression patterns and gene copy number changes in dermatofibrosarcoma protuberans," *American Journal of Pathology*, vol. 163, no. 6, pp. 2383–2395, 2003.