Review Article

Role of Exercise Training on Autonomic Changes and Inflammatory Profile Induced by Myocardial Infarction

Bruno Rodrigues,¹ Fabio S. Lira,² Fernanda M. Consolim-Colombo,³ Juraci A. Rocha,³ Erico C. Caperuto,¹ Kátia De Angelis,⁴ and Maria-Cláudia Irigoyen³

¹ Human Movement Laboratory, Sao Judas Tadeu University (USJT), Avenida Taquari 546, 03166-000 São Paulo, SP, Brazil

² Immunometabolism Research Group, Department of Physical Education, Universidade Estadual Paulista (UNESP), Rua Roberto Simonsen 305, 19060-900 Presidente Prudente, SP, Brazil

³ Hypertension Unit, Heart Institute (InCor), Medical School of University of São Paulo, Avenida Dr. Enéas de Carvalho Aguiar 44, 05403-900 São Paulo, SP, Brazil

⁴ Translational Physiology Laboratory, Universidade Nove de Julho (UNINOVE), Rua Vergueiro 235/249, Liberdade, 01504-001 São Paulo, SP, Brazil

Correspondence should be addressed to Bruno Rodrigues; prof.brodrigues@gmail.com

Received 24 January 2014; Revised 6 March 2014; Accepted 11 March 2014; Published 18 June 2014

Academic Editor: José Cesar Rosa

Copyright © 2014 Bruno Rodrigues et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The cardiovascular autonomic imbalance in patients after myocardial infarction (MI) provides a significant increase in mortality rate, and seems to precede metabolic, hormonal, and immunological changes. Moreover, the reduction in the parasympathetic function has been associated with inflammatory response in different pathological conditions. Over the years, most of the studies have indicated the exercise training (ET) as an important nonpharmacological tool in the management of autonomic dysfunction and reduction in inflammatory profile after a myocardial infarction. In this work, we reviewed the effects of ET on autonomic imbalance after MI, and its consequences, particularly, in the post-MI inflammatory profile. Clinical and experimental evidence regarding relationship between alterations in autonomic regulation and local or systemic inflammation response after MI were also discussed.

1. Introduction

Since the 50s, when cardiovascular diseases (CVDs) exceeded 50% as a cause of mortality worldwide, a detailed search for better understanding of the risk factors was initiated with the Framingham study. Cigarette smoking, hypertension, hypercholesterolemia, diabetes mellitus, physical inactivity, and obesity were identified as the main threats [1] and prevention strategies were initiated.

Despite all technological and scientific advances in the management and prevention of CVD, it remains the leading cause of morbidity and mortality in developed countries and it is fast becoming a major health challenge in developing countries, contributing significantly to high costs in public health [2]. Coronary artery disease (CAD) has a broad spectrum of clinical manifestations, in particular acute

coronary syndromes (ACS), that is, unstable angina and acute myocardial infarction (MI). MI is estimated to occur in the US every 44 seconds and about 49% of CVD deaths in the country are attributed to cardiac ischemic events [2].

Both during and after MI, neurohumoral changes occur in order to minimize the consequences of reduced ventricular function, which is caused by the obstruction of blood flow in the left ventricle (LV) of patients who had experienced an ischemic event. On the other hand, chronically, autonomic imbalance is a key element in the pathophysiology of heart failure (HF) after a MI [3–6]. Thus, strategies in order to detect, prevent, and attenuate the cardiovascular autonomic dysfunction, particularly associated with reduced vagal activity, have been seen as important interventions in the management of the changes triggered by MI.

FIGURE 1: Cardiovascular reflexes impairment and cardiac autonomic nervous system imbalance after myocardial infarction.

The metabolic, cardiovascular, autonomic, and anti-inflammatory benefits of a physically active life style have led many researchers to suggest exercise training (ET) as an important nonpharmacological tool in the prevention and treatment of CVD [7–11]. The effectiveness of ET as a great tool in the treatment of patients with established CAD (either with or without MI) has been widely reported in the literature [7, 12–16]. In this sense, the purpose of this paper was to review the effects of ET on autonomic imbalance and inflammatory profile after MI. Clinical and experimental evidence, or lack of them, regarding relationship between alterations on cardiac autonomic regulation and local or systemic inflammation response after MI will be discussed.

2. Myocardial Infarction and Autonomic Dysfunction

As illustrated in Figure 1, after MI, the reduction in cardiac output is associated with an imbalance of the autonomic nervous system in favor of increased sympathetic activity and reduced vagal activity and is usually accompanied by abnormalities in the cardiorespiratory reflex control, that is, impairment of baroreflex sensitivity and function, and increased activation of ergoreflex and chemoreflex [3, 5, 17, 18]. The sympathoinhibition by the arterial baroreflex is significantly suppressed, whereas sympathoexcitatory reflexes, including the cardiac sympathetic afferent reflex,

arterial chemoreceptor, and cardiopulmonary reflexes, are augmented [19]. Thus, the change in cardiovascular reflexes leads to a generalized activation of the sympathetic nervous system after MI in order to change the heart and peripheral hemodynamics. These changes are initially necessary; however, chronically, they are associated with reduced heart rate variability and increased blood pressure variability, which contributes with target organ damage, heart failure development, risk of arrhythmias, and sudden cardiac death [20].

In this sense, the reflex control of circulation commanded by arterial pressure receptors has been recognized as an important predictor of cardiovascular risk after cardiac event. The ATRAMI study (autonomic tone and reflexes after myocardial infarction) has provided clinical evidence of the prognostic value of baroreflex sensitivity and heart rate variability (HRV) in the mortality rate after MI, regardless of ejection fraction and ventricular arrhythmias [3]. Furthermore, in a study undertaken by Kleiger et al. [21], at the Multicenter Post Myocardial Infarction Program, it was found that the individuals who had lower HRV displayed a higher relative risk of mortality (five times) as compared with those who had higher HRV.

Additionally, experimental data have also pointed to a loss of autonomic function in animals after coronary artery occlusion. Indeed, our group has consistently demonstrated that after chronic MI the animals displayed baroreflex dysfunction and reduction in total HRV and in parasympathetic

modulation, as well as increased sympathetic modulation in relation to the noninfarcted animals [19, 22–28]. Furthermore, the changes of hormonal/signaling factors levels at specific sites (angiotensin II, nitric oxide, reactive oxygen species, arginine vasopressin, endothelin-1, atrial natriuretic peptide, prostaglandins, and aldosterone), as well as the increased concentration of proinflammatory cytokines in the central nervous system, may be potential candidate mechanisms underlying the increased sympathetic outflow [29–33].

The clinical prognosis of increased sympathetic activity after myocardial ischemia and HF is now well established [34], and, as such, beta adrenergic blockade has become a standard element in the therapy of these patients [35]. In contrast, the reduction in parasympathetic tonus, although already demonstrated more than 40 years [36], has received less attention. In this sense, more recently, experimental studies have demonstrated that vagal stimulation promoted an antifibrillatory effect and reduced mortality rate in animal models of HF [37, 38]. On the other hand, our group has recently tested the effectiveness of treatment with pyridostigmine bromide, an acetylcholinesterase inhibitor, for 7 days (0.14 mg/mL/day), on autonomic function in rats after MI. The MI treated group improved vagal tonus and decreased sympathetic tonus and MI area in relation to the MI placebo group [39].

Thus, interventions to detect, prevent, or attenuate cardiovascular autonomic dysfunction (particularly reduced vagal activity) have been welcomed as new and important strategies in the management of the MI-induced changes.

3. Autonomic Dysfunction and Inflammatory Response

Inflammatory processes combined with cytokine release are important steps in response to tissue injury and play an active role in cardiac remodeling and function after MI. Inflammatory cytokines, such as tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), interleukin-6 (IL-6), and interleukin-10 (IL-10), are released after acute ischemic injury and may regulate survival or apoptosis of myocytes, as well as triggering additional inflammatory cell response [40]. In this sense, Ammirati et al. [41] have observed 109 patients during an acute MI and found that those with additional elevations of IL-6 and IL-10 displayed a poor prognosis when compared to individuals with lower levels of these cytokines. Thus, the authors suggested that the degree of acute inflammation after ischemia may indicate a poor prognosis in MI patients.

Chronically, cytokines may mediate cardiac remodeling and repair, enabling the formation of collagen and matrix metalloproteinase, associated with integrin regulation, angiogenesis, and mobilization of progenitor cells [40]. After the initial increase of proinflammatory cytokines in the infarcted area, their levels usually decline toward baseline values one week after MI [42]. However, according to Ono et al. [42], if the size of MI is large, or if there are other stress factors in course, the gene expression of cytokines may remain significantly elevated in noninfarcted region following 20 weeks

of MI. Furthermore, these researchers showed that the cytokines levels, such as IL-1 β , are associated with ventricular diastolic diameter increase and collagen deposition in the infarcted area after 8 and 20 weeks of MI. Thus, acutely and/or chronically, the release of proinflammatory cytokines adversely affects the LV function, exerts a negative inotropic effect [43], induces abnormalities in cardiac metabolism, and promotes myocardial remodeling, leading to HF [44, 45]. Additionally, activation of the immune system promotes the development of endothelial dysfunction and skeletal muscle apoptosis in HF [46, 47].

The origin of immune activation after ischemia has been dealt with by various research studies but remains unclear. There are at least five hypotheses addressing the underlying mechanism of inflammatory response [48]: (1) the failure of the myocardium *per se* would be the main source of cytokine production [49]; (2) the circulatory decompensation would lead to increased intestinal translocation of bacterial endotoxin (lipopolysaccharide) to the systemic circulation, which in turn would activate circulating immune cells [50]; (3) the main source of proinflammatory mediators would be the body tissues exposed to hypoxia [51, 52]; (4) immune activation would be a consequence of increased [53] sympathetic stimulation; and (5) reduction in parasympathetic participation would work as the primary mediator of the inflammatory response activation [48].

Studies by Kevin J. Tracey group [54–62] have lent strength to the hypothesis of a direct relation between parasympathetic activation and immune system response. This group initially argued that the activation of the vagus nerve (electrically or by cholinergic agonists) would reduce inflammatory response in experimental models of sepsis, and subsequently they postulated the theory of “inflammatory reflex,” linking aspects related to neuroimmunomodulation. The reflex arc, mediated by the nervous system, is composed by efferent via, integrative areas of central nervous system, and efferent via. Briefly, Kevin J. Tracey’s group observed that inflammatory mediators (cytokines) produced in peripheral tissues may warn the central nervous system by a direct central action or by afferent stimulation of the vagus nerve. In this model, the integration of signals takes place in the central areas, triggering the activation of the parasympathetic efferent pathway, mediated mainly by the vagus nerve, whose neurotransmitter is acetylcholine. The cholinergic pathway innervates various components of the immune system (reticuloendothelial system), such as lymph nodes, liver, heart, spleen, and gastrointestinal tract. The activation of the vagus nerve leads to reduced production of cytokines, which in turn decreases the inflammatory response in models of septic and aseptic inflammation [55].

According to the “inflammatory reflex” theory, the vagus nerve seems to be the most important element in the efferent arm. In this sense, Blalock [63] has suggested that the immune system linked in such way to the central nervous system works as a “sixth sense,” being able to detect microbial invasion and other inflammatory substances and to retransmit this information to the brain, triggering responses that would interfere in the initial process. In fact, studies have viewed the parasympathetic hyperactivity, brought about by either drugs

or direct vagal stimulation, as a mechanism that reduces the release of cytokines and reactive oxygen species during an inflammatory process [56, 64].

In this context, Borovikova et al. [65] have also demonstrated that injection of endotoxin (lipopolysaccharide) in animals that underwent vagus nerve stimulation resulted in reduced systemic release of inflammatory cytokines and macrophages, without affecting the release of interleukin-10 (IL-10). However, the vagus nerve transection abolished this protection. Accordingly, in humans, significantly reduced HRV is associated with elevated levels of inflammatory cytokines (IL-6) and C-reactive protein (CRP) [66]. Lanza et al. [67] have shown that serum CRP levels were significantly associated with reduced HRV in patients with unstable angina. This association has also been observed in healthy individuals and patients with stable coronary artery disease and HF [68]. Thus, in accordance with “inflammatory reflex” theory, inflammatory products produced in ischemic ventricle activate afferent signals that are relayed to the nucleus tractus solitaries, and subsequent activation of vagus efferent activity would inhibit cytokine synthesis through the cholinergic anti-inflammatory pathway (Figure 2). However, as previously mentioned there is an imbalance in favor of increased sympathetic activity and reduced vagal activity after MI. Thus, it is possible to suggest that cardiovascular autonomic imbalance after ischemic event may blunt “inflammatory reflex” by reducing cholinergic anti-inflammatory pathway. However, despite the fact that some suggestions are pointed out on this direction in the literature [66–68], there is no strong evidence to support a cause-effect relationship.

In this sense, preventing and/or attenuating autonomic dysfunction triggered by MI would result in a less intense inflammatory response and, as such, cardiac and peripheral structure and function would be preserved.

4. Aerobic Exercise Training as Therapy

In recent years, our laboratory has investigated the changes induced by MI alone [25, 27] or associated with different risk factors, such as hypertension [69, 70], ovarian hormone deprivation [23], sinoaortic denervation [24], and diabetes [26–28, 71–73]. These studies have also tested different therapeutic approaches in the management of MI-induced changes. Among these therapeutic approaches, some deserve greater emphasis: therapy involving stem cells and mesenchymal bone marrow both *in situ* [70] and intravenously [69], gene therapy with vascular endothelium-derived growth factor (VEGF) [73], and ET [22, 23, 25, 27].

Moderate intensity aerobic ET is responsible for structural and hemodynamic adaptations in the cardiovascular system and promotes adjustments in autonomic nervous system. This can be exemplified by cardiac adaptations such as increase of stroke volume [74, 75], adjustments in diastolic and systolic functions [75], and positive changes in cavities diameter and ventricular mass [76, 77] as well as alterations in the rest heart rate (HR) [77].

Resting bradycardia is associated with a decrease in intrinsic HR as well as an altered autonomic balance, leading

FIGURE 2: Blunted “inflammatory reflex” by vagal efferent function reduction after myocardial infarction (adapted from Tracey [55]). Ach: acetylcholine; Nor: noradrenaline.

to parasympathetic dominance [78]. This is thought to be mediated, at least in part, by an increase in cardiac vagal tone [79]. The mechanisms by which exercise produces changes in autonomic control have yet to be fully understood; however, there is evidence of alterations in the central afferent and efferent pathways and in the effector organs (receptor function) [80, 81].

Aerobic ET is largely recommended for patients with cardiac disease, including those who have experienced an MI. A meta-analysis, based on 48 randomized controlled trials (8940 patients), showed that cardiac rehabilitation programs based on aerobic exercises reduced all-cause mortality by 20% and CVD mortality by 26% in patients after MI, angina pectoris, and/or CAD [14]. Additionally, in another meta-analysis based on 34 studies, the authors observed that aerobic ET (predominantly) after MI, even for a short period (1–3 months), was an effective therapeutic strategy, since the observed benefits extended far beyond the intervention period [16].

One of the most important adaptations to the ET, generally associated with reduced mortality rate, is the improvement in autonomic nervous system regulation. Classic research studies such as Billman et al. [12] and Hull Jr. et al. [13] have suggested that this intervention reduces mortality in individuals after an MI, particularly when associated with increased vagal component and decreased sympathetic activity. La Rovere et al. [82], using head-up tilt testing to evaluate heart rate variability (HRV) in MI patients, have observed that aerobically trained individuals presented a better autonomic response to orthostatic stress, with reduction in vagal modulation and an increase in sympathetic vasoconstrictor outflow due to initiation of the baroreceptors. Moreover, Iellamo et al. [83], in the first randomized controlled study in the area, have observed that aerobic ET results in marked enhancement of both baroreflex sensitivity and HRV in patients with coronary artery disease. These researchers have also suggested that the improvement in baroreflex sensitivity associated with aerobic ET is not limited to patients with prior MI but extends to coronary patients without MI, for whom risk-reducing strategies designed to avoid subsequent lethal events might be of paramount importance.

In la Rovere et al. [7] study, 95 consecutive male patients, survivors of a first uncomplicated MI, underwent four weeks of ET. The researchers found that aerobic ET was associated with increased survival together with an adequate modulation of the autonomic balance toward increase in vagal activity, as revealed by the increase in baroreflex sensitivity. In this context, several clinical studies have reported improvements in cardiovascular autonomic control, particularly in HRV, after ET among MI patients. Sandercock et al. [15] have observed that after eight-week cardiac rehabilitation program, MI participants had significant increases in HRV parameters as compared with those not participating in the training program. In an elegant study, Malfatto et al. [84] have reported improvement in HRV following 8 weeks of aerobic ET. After this period, participants were encouraged to continue exercising at home two to three times per week. After one year of training program participation, improvements in HRV index were still relevant. More recently, six months of aerobic (predominantly) ET significantly decreased muscle sympathetic nerve activity and the low-frequency component of systolic arterial pressure and increased the baroreflex sensitivity in MI patients. These changes were so marked that the differences between patients with MI and the normal control group were no longer observed after ET. These findings highlight the clinical importance of this nonpharmacological therapy based on ET in the long-term treatment of patients with MI [85].

In an experimental setting, our group has studied the role of aerobic ET in functional, biochemical, and molecular alterations, as well as in the mortality rate after MI in rats. Rondon et al. [22] have investigated the role of ET in MI-induced heart failure rats and demonstrated that aerobic ET increased peak oxygen uptake and the high frequency band of HRV, while reducing the low frequency band of HRV. In addition, these authors also observed that after ET protocol the heart failure animals presented improvement in baroreflex control of heart rate and renal sympathetic nerve activity, associated with increased aortic depressor nerve activity. However, we should bear in mind that, although these data suggest an association between increased aortic depressor nerve activity and improvement in baroreflex control, the cause-effect relationship remains unclear. Moreover, Jorge et al. [25] have demonstrated that early aerobic ET intervention (one week) after MI induced an improvement in LV systolic and diastolic functions. In the study, we have also observed a normalization of hemodynamic and regional blood flows and an improvement in cardiovascular autonomic function associated with increased baroreflex sensitivity. These benefits, in turn, resulted in an increase in functional capacity and a reduction in mortality rate in trained infarcted animals. Similar results were also demonstrated in streptozotocin-diabetic rats undergoing MI, for which aerobic ET promoted, in addition to the benefits observed in autonomic function, positive changes in the expression of proteins associated with intracellular calcium handling [86]. Studying female ovariectomized rats undergoing MI, Flores et al. [23] have demonstrated that eight weeks of aerobic ET was able to improve resting hemodynamic status and reflex control of the circulation (arterial and cardiopulmonary baroreflex),

possibly associated with the increase in vagal component observed in the study.

In line with la Rovere et al. [7] study, in which aerobic ET was associated with improvement in baroreflex sensitivity and increased survival rate in a 10-year followup, our group tested the hypothesis that autonomic ET benefits might remain for an extended period, even during detraining. Accordingly, in a study by Barboza et al. [27], MI animals underwent three months of aerobic ET (starting one week after MI) and one month of detraining. The authors demonstrated that one month of detraining did not alter the beneficial effects of ET on the MI area, LV morphometry and function, and baroreflex sensitivity, as well as overall survival rate in MI-detrained animals. These findings indicate not only that aerobic ET is an effective tool in the management of cardiovascular and autonomic MI derangements, but also that these positive changes were extended even beyond one month of detraining in rats.

In addition to the benefits of ET on autonomic function, it is now well established that cytokine production by exercise is different from that observed in response to severe infections or tissue injury, since exercises usually do not provide expressive changes in the classic proinflammatory cytokines, $\text{TNF-}\alpha$ and $\text{IL-1}\beta$ [87]. During exercise, IL-6 is the first cytokine present in the circulation, exponentially increasing when the exercise is in course, and fast declining after the exercise period [88–90]. In this sense, there is evidence in the literature that IL-6 may exert anti-inflammatory effects, since circulating IL-6 triggers an inhibitory effect on $\text{TNF-}\alpha$ and IL-1 production [91] and stimulates the production of IL-1 receptor antagonist and IL-10 [92]. IL-10 and IL-1 receptor antagonist production after exercise, in turn, contributes to the inhibition of the synthesis of a large spectrum of proinflammatory cytokines by different cells, particularly the ones of monocytic lineage [93].

Actually, the protective effects of ET on inflammatory markers have been widely discussed [94]. In this sense, Adamopoulos et al. [95] have demonstrated a reduction in inflammatory markers after 12 weeks of aerobic ET in patients with moderate to severe HF, suggesting a correlation between improvement in exercise tolerance and attenuation of inflammatory process. In an experimental model of HF, Batista Jr. et al. [96] have shown that aerobic ET in infarcted rats increased the ratio $\text{IL-10/tumor necrosis factor-}\alpha$ ($\text{TNF-}\alpha$) in the soleus muscle of animals, emphasizing the anti-inflammatory effect of exercise after an ischemic event.

Individuals with increased risk of CAD presented reduction in atherogenic cytokines, as interleukin-1 (IL-1), IL-6 , $\text{TNF-}\alpha$, and C reactive protein, as well as improving atheroprotective cytokines, as IL-10 , and transforming growth factor beta-1 after an aerobic ET program [97]. The reduction in plasma $\text{TNF-}\alpha$, IL-6 , and their receptors has been demonstrated in aerobically trained HF patients [98], suggesting an attenuation in the chronic inflammatory condition mediated by a regulation in peripheral inflammatory response [95, 98]. Similarly, the association between aerobic and resistance ET decreased the concentration of $\text{TNF-}\alpha$ receptors I and II in patients with HF [99].

Thus, aerobic ET seems to be an important strategy in the management MI-induced changes and HF, especially with regard to improving cardiovascular autonomic control and reducing chronic inflammatory response, consequently reducing cardiac work, decreasing risk of fatal arrhythmias, and increasing survival in affected individuals.

In order to understand the mechanisms associated with autonomic adjustments and anti-inflammatory role of exercise fully, it is necessary to do a more detailed search of the nature, intensity, and duration of ET after MI. The beneficial effects of aerobic ET are well known; however, autonomic and anti-inflammatory effects of resistance ET or high intensity aerobic ET are poorly defined and remain areas for future investigations. In this direction, our group recently demonstrated that resistance ET (moderate intensity) induced additional benefits in the low frequency band (+50%) and high frequency band of HRV (+45%), as well as in the low frequency band of systolic blood pressure variability (−46%) of trained infarcted rats compared to sedentary [100]. These findings suggest that resistance ET was effective in reducing cardiac and peripheral sympathetic modulation, as well as increasing cardiac parasympathetic modulation of infarcted rats.

5. Exercise Training on “Inflammatory Reflex” Response after Myocardial Infarction

Despite all data generated on the anti-inflammatory role of the exercise, the underlying mechanisms associated with improved inflammatory profile in patients after an ischemic event have yet to be fully understood. During the last decade, studies on the anti-inflammatory effects of exercise have pointed to three possible mechanisms [87, 101–106]: reduction in visceral fat mass; increased production and release of anti-inflammatory cytokines from contracting skeletal muscle (myokines); and reduced expression of toll-like receptors (TLRs) on monocytes and macrophages. Regarding the first mechanism, ET is able to reduce low-grade and systemic inflammation via a decrease in proinflammatory adipokine secretion, such as TNF, leptin, retinol-binding protein 4, lipocalin 2, IL-6, and IL-18, which is a direct result of lowering the amount of fat stored in abdominal and visceral depots [107–110].

Related to the second proposed mechanism, during exercise it seems that IL-6, produced by contractile skeletal muscle via a TNF-independent pathway, stimulates systemic appearance of anti-inflammatory cytokines, such as IL-1 receptor antagonist and IL-10, and inhibits the liberation of proinflammatory cytokine TNF- α [101, 111]. Additionally, IL-6 may be associated with increase of lipid turnover, stimulating lipolysis as well as fat oxidation [101]. In line with the third proposed mechanism about the anti-inflammatory effects of ET, it has been evidenced that TLRs may be involved in the link between a sedentary lifestyle and inflammation and disease [112]. Studies have observed that blood monocytes from trained individuals present decreased TLR4 expression, which is associated with decreased inflammatory

cytokine production, and reduced inflammatory response to endotoxin stimulation *in vitro* [103, 105, 113].

On the other hand, as previously mentioned, the “inflammatory reflex” proposed by Tracey [55] seems to have an important role in the development of the inflammatory process, since it is in line with the suggestion that the activation of the vagus nerve leads to a reduced production of cytokines by the reticuloendothelial system, such as lymph nodes, liver, heart, spleen, and gastrointestinal tract. ET, in turn, modulates cardiac autonomic control with reduction in sympathetic tonus and increases the role of the vagal tonus [7, 15, 85], thereby positively affecting the prognosis of MI patients. Thus, we suggest that in addition to the reduction in visceral fat mass, increased production anti-inflammatory myokines, and reduced expression of TLRs on monocytes and macrophages, the increase in vagal activation induced by ET may be an important mechanism that would explain, at least in part, the improvement in the inflammatory status of animals or patients after MI.

As evidenced in Figure 3, we proposed that chronic benefits of ET would be associated with significant improvements in baroreflex sensitivity, resulting in the increase in vagal activity and reduction in sympathetic activity (as observed to the heart and vessels [7, 12, 13, 15, 22, 23, 25, 27, 78, 82–86]) to important organs (reticulum endothelial system and other tissues), including brain, heart, liver, spleen, and gastrointestinal (GI) tract. This autonomic remodeling, in turn, would be directly associated with improved local and systemic inflammatory profile, reducing cardiovascular risk and enhancing the survival rate in patients who had suffered a MI. When studying infarcted rats undergoing 3 months of moderate-intensity aerobic ET, preliminary data from our group has noted that HRV parameters such as the high frequency band and root mean square of successive difference (RMSSD)—indexes of cardiac parasympathetic modulation—were negatively correlated with cardiac levels of TNF- α and IL-6 (unpublished data). Furthermore, the low frequency band of systolic blood pressure variability, an indicator of vascular sympathetic modulation, was positively associated with the TNF- α and IL-6 levels in the adipose tissue of these animals (unpublished data). Taken together, our preliminary findings suggest that ET can increase cardiac vagal modulation, as well as reducing sympathetic peripheral modulation, leading to a reduction in the inflammatory profile of infarcted rats.

Despite the vast literature raised in this review, many questions still remain about the benefits of ET after MI on cardiovascular autonomic function and inflammatory response, or if there is a direct relationship between them. At present, we do not know whether there is a cause-effect relationship between the improvement in vagal cholinergic activity promoted by ET and reduction in cardiac and systemic inflammatory status after a cardiac ischemic event. Or even, what is the relative importance of these mechanisms on cardiac morphology and function, as well as in the prognosis of infarcted patients.

FIGURE 3: Possible mechanisms associated with inflammatory profile reduction in patients after myocardial infarction undergoing exercise training.

Intuitively, we can expect that many anti-inflammatory mechanisms of exercise, known or unknown, are acting in combination in order to better respond to different stimulations applied. Furthermore, it is possible that the involved mechanisms and the magnitude of the benefits found in autonomic function and inflammatory profile

will depend on the mode, intensity and duration of exercise performed. However, the vast majority of the studies presented in this review used the moderate intensity aerobic ET. The absence of studies comparing different intensities, durations, and modalities of ET and its possible relationship with the anti-inflammatory cholinergic

pathway opens new perspectives for working in exercise cardiology.

6. Conclusions

There is consistent experimental and clinical evidence that the autonomic nervous system and inflammatory response play a key role in cardiac and peripheral dysfunctions after myocardial infarction. Exercise training has been associated with increased vagal tonus/modulation, as well as with reduction in sympathetic tonus/modulation and inflammatory profile in heart failure or after myocardial infarction. In recent years, a direct relationship between vagal activity and inflammatory status has been proposed. Here, we suggest that exercise training may improve autonomic function to different sites, leading to a reduction in the proinflammatory response observed after myocardial infarction. However, further experimental and clinical investigations are needed to test the role of exercise training in the modulation of this “inflammatory reflex” after myocardial infarction. The results of these future studies will improve the management of cardiovascular risk, quality of life, and survival of this population.

Conflict of Interests

The authors have no conflict of interests to declare.

Authors' Contribution

Kátia De Angelis and Maria-Cláudia Irigoyen contributed equally to this paper.

Acknowledgments

This work was funded by Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP-2013/14788-9). Bruno Rodrigues, Kátia De Angelis, and Maria-Cláudia Irigoyen received financial support from Conselho Nacional de Pesquisa e Desenvolvimento (CNPq-BPQ).

References

- [1] G. Thiene and C. Basso, “Myocardial infarction: a paradigm of success in modern medicine,” *Cardiovascular Pathology*, vol. 19, no. 1, pp. 1–5, 2010.
- [2] A. S. Go, D. Mozaffarian, V. L. Roger et al., “Heart disease and stroke statistics—2014 update: a report from the American Heart Association,” *Circulation*, vol. 129, no. 3, pp. e28–e292, 2014.
- [3] M. T. la Rovere, J. T. Bigger Jr., F. I. Marcus, A. Mortara, and P. J. Schwartz, “Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction,” *The Lancet*, vol. 351, no. 9101, pp. 478–484, 1998.
- [4] E. Braunwald and M. R. Bristow, “Congestive heart failure: fifty years of progress,” *Circulation*, vol. 102, no. 20, pp. 14–23, 2000.
- [5] C. E. Negrao and H. R. Middlekauff, “Adaptations in autonomic function during exercise training in heart failure,” *Heart Failure Reviews*, vol. 13, no. 1, pp. 51–60, 2008.
- [6] K. Thygesen, J. S. Alpert, A. S. Jaffe et al., “Third universal definition of myocardial infarction,” *Circulation*, vol. 126, no. 16, pp. 2020–2035, 2012.
- [7] M. T. la Rovere, C. Bersano, M. Gnemmi, G. Specchia, and P. J. Schwartz, “Exercise-induced increase in baroreflex sensitivity predicts improved prognosis after myocardial infarction,” *Circulation*, vol. 106, no. 8, pp. 945–949, 2002.
- [8] B. K. Pedersen and B. Saltin, “Evidence for prescribing exercise as therapy in chronic disease,” *Scandinavian Journal of Medicine and Science in Sports*, vol. 16, no. 1, pp. 3–63, 2006.
- [9] T. A. Lakka and D. E. Laaksonen, “Physical activity in prevention and treatment of the metabolic syndrome,” *Applied Physiology, Nutrition and Metabolism*, vol. 32, no. 1, pp. 76–88, 2007.
- [10] S. Ertek and A. Cicero, “Impact of physical activity on inflammation: effects on cardiovascular disease risk and other inflammatory conditions,” *Archives of Medical Science*, vol. 8, no. 5, pp. 794–804, 2012.
- [11] N. Mann and A. Rosenzweig, “Can exercise teach Us how to treat heart disease?” *Circulation*, vol. 126, no. 22, pp. 2625–2635, 2012.
- [12] G. E. Billman, P. J. Schwartz, and H. L. Stone, “The effects of daily exercise on susceptibility to sudden cardiac death,” *Circulation*, vol. 69, no. 6, pp. 1182–1189, 1984.
- [13] S. S. Hull Jr., E. Vanoli, P. B. Adamson, R. L. Verrier, R. D. Foreman, and P. J. Schwartz, “Exercise training confers anticipatory protection from sudden death during acute myocardial ischemia,” *Circulation*, vol. 89, no. 2, pp. 548–552, 1994.
- [14] R. S. Taylor, A. Brown, S. Ebrahim et al., “Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials,” *American Journal of Medicine*, vol. 116, no. 10, pp. 682–692, 2004.
- [15] G. R. H. Sandercock, R. Grocott-Mason, and D. A. Brodie, “Changes in short-term measures of heart rate variability after eight weeks of cardiac rehabilitation,” *Clinical Autonomic Research*, vol. 17, no. 1, pp. 39–45, 2007.
- [16] P. R. Lawler, K. B. Filion, and M. J. Eisenberg, “Efficacy of exercise-based cardiac rehabilitation post-myocardial infarction: a systematic review and meta-analysis of randomized controlled trials,” *American Heart Journal*, vol. 162, no. 4, pp. 571.e2–584.e2, 2011.
- [17] P. P. Ponikowski, T. P. Chua, D. P. Francis, A. Capucci, A. J. S. Coats, and M. F. Piepoli, “Muscle ergoreceptor overactivity reflects deterioration in clinical status and cardiorespiratory reflex control in chronic heart failure,” *Circulation*, vol. 104, no. 19, pp. 2324–2330, 2001.
- [18] P. Ponikowski, T. P. Chua, S. D. Anker et al., “Peripheral chemoreceptor hypersensitivity an ominous sign in patients with chronic heart failure,” *Circulation*, vol. 104, no. 5, pp. 544–549, 2001.
- [19] J. E. D. C. Lacerda, F. M. Consolim-Colombo, E. D. Moreira et al., “Influence of cardiopulmonary reflex on the sympathetic activity during myocardial infarction,” *Autonomic Neuroscience: Basic and Clinical*, vol. 133, no. 2, pp. 128–135, 2007.
- [20] J. S. Floras, “Sympathetic nervous system activation in human heart failure. Clinical implications of an updated model,” *Journal of the American College of Cardiology*, vol. 54, no. 5, pp. 375–385, 2009.
- [21] R. E. Kleiger, J. P. Miller, J. T. Bigger Jr., and A. J. Moss, “Decreased heart rate variability and its association with increased mortality after acute myocardial infarction,” *The American Journal of Cardiology*, vol. 59, no. 4, pp. 256–262, 1987.

- [22] E. Rondon, M. S. Brasileiro-Santos, E. D. Moreira et al., "Exercise training improves aortic depressor nerve sensitivity in rats with ischemia-induced heart failure," *American Journal of Physiology. Heart and Circulatory Physiology*, vol. 291, no. 6, pp. H2801–H2806, 2006.
- [23] L. J. Flores, D. Figueroa, I. C. Sanches et al., "Effects of exercise training on autonomic dysfunction management in an experimental model of menopause and myocardial infarction," *Menopause*, vol. 17, no. 4, pp. 712–717, 2010.
- [24] C. Mostarda, B. Rodrigues, M. Vane et al., "Autonomic impairment after myocardial infarction: role in cardiac remodelling and mortality," *Clinical and Experimental Pharmacology and Physiology*, vol. 37, no. 4, pp. 447–452, 2010.
- [25] L. Jorge, B. Rodrigues, K. T. Rosa et al., "Cardiac and peripheral adjustments induced by early exercise training intervention were associated with autonomic improvement in infarcted rats: role in functional capacity and mortality," *European Heart Journal*, vol. 32, no. 7, pp. 904–912, 2011.
- [26] B. Rodrigues, K. T. Rosa, A. Medeiros et al., "Hyperglycemia can delay left ventricular dysfunction but not autonomic damage after myocardial infarction in rodents," *Cardiovascular Diabetology*, vol. 10, article 26, 2011.
- [27] C. A. Barboza, L. Y. Rocha, C. T. Mostarda et al., "Ventricular and autonomic benefits of exercise training persist after detraining in infarcted rats," *European Journal of Applied Physiology*, vol. 113, no. 5, pp. 1137–1146, 2013.
- [28] B. Rodrigues, C. T. Mostarda, L. Jorge et al., "Impact of myocardial infarction on cardiac autonomic function in diabetic rats," *Journal of Diabetes and Its Complications*, vol. 27, no. 1, pp. 16–22, 2013.
- [29] J. Francis, Y. Chu, A. K. Johnson et al., "Acute myocardial infarction induces hypothalamic cytokine synthesis," *American Journal of Physiology. Heart and Circulatory Physiology*, vol. 286, no. 6, pp. H2264–H2271, 2004.
- [30] T. Kishi, Y. Hirooka, Y. Kimura, K. Ito, H. Shimokawa, and A. Takeshita, "Increased reactive oxygen species in rostral ventrolateral medulla contribute to neural mechanisms of hypertension in stroke-prone spontaneously hypertensive rats," *Circulation*, vol. 109, no. 19, pp. 2357–2362, 2004.
- [31] I. H. Zucker, H. D. Schultz, K. P. Patel, W. Wang, and L. Gao, "Regulation of central angiotensin type 1 receptors and sympathetic outflow in heart failure," *American Journal of Physiology. Heart and Circulatory Physiology*, vol. 297, no. 5, pp. H1557–H1566, 2009.
- [32] T. Kishi, Y. Hirooka, S. Konno, K. Ogawa, and K. Sunagawa, "Angiotensin II type 1 receptor-activated caspase-3 through ras/mitogen-activated protein kinase/extracellular signal-regulated kinase in the rostral ventrolateral medulla is involved in sympathoexcitation in stroke-prone spontaneously hypertensive rats," *Hypertension*, vol. 55, no. 2, pp. 291–297, 2010.
- [33] K. Ogawa, Y. Hirooka, T. Kishi, and K. Sunagawa, "Brain AT1 receptor activates the sympathetic nervous system through toll-like receptor 4 in mice with heart failure," *Journal of Cardiovascular Pharmacology*, vol. 58, no. 5, pp. 543–549, 2011.
- [34] J. S. Floras, "Sympathetic activation in human heart failure: diverse mechanisms, therapeutic opportunities," *Acta Physiologica Scandinavica*, vol. 177, no. 3, pp. 391–398, 2003.
- [35] M. Packer, M. R. Bristow, J. N. Cohn et al., "The effect of carvedilol on morbidity and mortality in patients with chronic heart failure," *New England Journal of Medicine*, vol. 334, no. 21, pp. 1349–1355, 1996.
- [36] D. L. Eckberg, M. Drabinsky, and E. Braunwald, "Defective cardiac parasympathetic control in patients with heart disease," *New England Journal of Medicine*, vol. 285, no. 16, pp. 877–883, 1971.
- [37] M. Li, C. Zheng, T. Sato, T. Kawada, M. Sugimachi, and K. Sunagawa, "Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats," *Circulation*, vol. 109, no. 1, pp. 120–124, 2004.
- [38] Y. Zhang, Z. B. Popović, S. Bibevski et al., "Chronic vagus nerve stimulation improves autonomic control and attenuates systemic inflammation and heart failure progression in a canine high-rate pacing model," *Circulation: Heart Failure*, vol. 2, no. 6, pp. 692–699, 2009.
- [39] R. N. de la Fuente, B. Rodrigues, I. C. Moraes-Silva et al., "Cholinergic stimulation with pyridostigmine improves autonomic function in infarcted rats," *Clinical and Experimental Pharmacology and Physiology*, vol. 40, no. 9, pp. 610–616, 2013.
- [40] M. Nian, P. Lee, N. Khaper, and P. Liu, "Inflammatory cytokines and postmyocardial infarction remodeling," *Circulation Research*, vol. 94, no. 12, pp. 1543–1553, 2004.
- [41] E. Ammirati, C. V. Cannistraci, N. A. Cristell et al., "Identification and predictive value of interleukin-6+ interleukin-10+ and interleukin-6-interleukin-10+ cytokine patterns in st-elevation acute myocardial infarction," *Circulation Research*, vol. 111, no. 10, pp. 1336–1348, 2012.
- [42] K. Ono, A. Matsumori, T. Shioi, Y. Furukawa, and S. Sasayama, "Cytokine gene expression after myocardial infarction in rat hearts: possible implication in left ventricular remodeling," *Circulation*, vol. 98, no. 2, pp. 149–156, 1998.
- [43] D. R. Meldrum, "Tumor necrosis factor in the heart," *American Journal of Physiology. Regulatory Integrative and Comparative Physiology*, vol. 274, no. 3, pp. R577–R595, 1998.
- [44] D. L. Mann, "Inflammatory mediators and the failing heart: past, present, and the foreseeable future," *Circulation Research*, vol. 91, no. 11, pp. 988–998, 2002.
- [45] A. Diwan, T. Tran, A. Misra, and D. L. Mann, "Inflammatory mediators and the failing heart: a translational approach," *Current Molecular Medicine*, vol. 3, no. 2, pp. 161–182, 2003.
- [46] S. D. Anker, P. P. Ponikowski, A. L. Clark et al., "Cytokines and neurohormones relating to body composition alterations in the wasting syndrome of chronic heart failure," *European Heart Journal*, vol. 20, no. 9, pp. 683–693, 1999.
- [47] R. Sharma, A. J. S. Coats, and S. D. Anker, "The role of inflammatory mediators in chronic heart failure: cytokines, nitric oxide, and endothelin-1," *International Journal of Cardiology*, vol. 72, no. 2, pp. 175–186, 2000.
- [48] E. A. Jankowska, P. Ponikowski, M. F. Piepoli, W. Banasiak, S. D. Anker, and P. A. Poole-Wilson, "Autonomic imbalance and immune activation in chronic heart failure—pathophysiological links," *Cardiovascular Research*, vol. 70, no. 3, pp. 434–445, 2006.
- [49] G. Torre-Amione, S. Kapadia, J. Lee et al., "Tumor necrosis factor- α and tumor necrosis factor receptors in the failing human heart," *Circulation*, vol. 93, no. 4, pp. 704–711, 1996.
- [50] J. Niebauer, H.-D. Volk, M. Kemp et al., "Endotoxin and immune activation in chronic heart failure: a prospective cohort study," *The Lancet*, vol. 353, no. 9167, pp. 1838–1842, 1999.
- [51] D. Hasper, M. Hummel, F. X. Kleber, I. Reindl, and H.-D. Volk, "Systemic inflammation in patients with heart failure," *European Heart Journal*, vol. 19, no. 5, pp. 761–765, 1998.

- [52] T. Bachetti and R. Ferrari, "The dynamic balance between heart function and immune activation," *European Heart Journal*, vol. 19, no. 5, pp. 681–682, 1998.
- [53] U. Müller-Werdan and K. Werdan, "Immune modulation by catecholamines—a potential mechanism of cytokine release in heart failure?" *Herz*, vol. 25, no. 3, pp. 271–273, 2000.
- [54] F. Rangwala, R. C. Drisdel, S. Rakhilin et al., "Neuronal α -bungarotoxin receptors differ structurally from other nicotinic acetylcholine receptors," *Journal of Neuroscience*, vol. 17, no. 21, pp. 8201–8212, 1997.
- [55] K. J. Tracey, "The inflammatory reflex," *Nature*, vol. 420, no. 6917, pp. 853–859, 2002.
- [56] H. Wang, H. Liao, M. Ochani et al., "Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis," *Nature Medicine*, vol. 10, no. 11, pp. 1216–1221, 2004.
- [57] V. A. Pavlov, M. Ochani, M. Gallowitsch-Puerta et al., "Central muscarinic cholinergic regulation of the systemic inflammatory response during endotoxemia," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 13, pp. 5219–5223, 2006.
- [58] K. J. Tracey, "Physiology and immunology of the cholinergic antiinflammatory pathway," *Journal of Clinical Investigation*, vol. 117, no. 2, pp. 289–296, 2007.
- [59] R. P. Sloan, H. McCreath, K. J. Tracey, S. Sidney, K. Liu, and T. Seeman, "RR interval variability is inversely related to inflammatory markers: the CARDIA study," *Molecular Medicine*, vol. 13, no. 3–4, pp. 178–184, 2007.
- [60] M. Rosas-Ballina and K. J. Tracey, "The neurology of the immune system: neural reflexes regulate immunity," *Neuron*, vol. 64, no. 1, pp. 28–32, 2009.
- [61] K. J. Tracey, "Reflex control of immunity," *Nature Reviews Immunology*, vol. 9, no. 6, pp. 418–428, 2009.
- [62] J. M. Huston and K. J. Tracey, "The pulse of inflammation: heart rate variability, the cholinergic anti-inflammatory pathway and implications for therapy," *Journal of Internal Medicine*, vol. 269, no. 1, pp. 45–53, 2011.
- [63] J. E. Blalock, "The immune system as the sixth sense," *Journal of Internal Medicine*, vol. 257, no. 2, pp. 126–138, 2005.
- [64] T. Tsutsumi, T. Ide, M. Yamato et al., "Modulation of the myocardial redox state by vagal nerve stimulation after experimental myocardial infarction," *Cardiovascular Research*, vol. 77, no. 4, pp. 713–721, 2008.
- [65] L. V. Borovikova, S. Ivanova, M. Zhang et al., "Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin," *Nature*, vol. 405, no. 6785, pp. 458–462, 2000.
- [66] D. Aronson, M. A. Mittleman, and A. J. Burger, "Interleukin-6 levels are inversely correlated with heart rate variability in patients with decompensated heart failure," *Journal of Cardiovascular Electrophysiology*, vol. 12, no. 3, pp. 294–300, 2001.
- [67] G. A. Lanza, G. A. Sgueglia, D. Cianflone et al., "Relation of heart rate variability to serum levels of C-reactive protein in patients with unstable angina pectoris," *American Journal of Cardiology*, vol. 97, no. 12, pp. 1702–1706, 2006.
- [68] M. H. Shishehbor, C. Alves, and V. Rajagopal, "Inflammation: implications for understanding the heart-brain connection," *Cleveland Clinic Journal of Medicine*, vol. 74, no. 1, pp. 37–41, 2007.
- [69] L. M. G. de Macedo Braga, K. Rosa, B. Rodrigues et al., "Systemic delivery of adult stem cells improves cardiac function in spontaneously hypertensive rats," *Clinical and Experimental Pharmacology and Physiology*, vol. 35, no. 2, pp. 113–119, 2008.
- [70] L. M. G. De Macedo Braga, S. Lacchini, B. D. Schaan et al., "In situ delivery of bone marrow cells and mesenchymal stem cells improves cardiovascular function in hypertensive rats submitted to myocardial infarction," *Journal of Biomedical Science*, vol. 15, no. 3, pp. 365–374, 2008.
- [71] C. Malfitano, T. C. Alba Loureiro, B. Rodrigues et al., "Hyperglycaemia protects the heart after myocardial infarction: aspects of programmed cell survival and cell death," *European Journal of Heart Failure*, vol. 12, no. 7, pp. 659–667, 2010.
- [72] B. Rodrigues, D. M. T. Figueroa, J. Fang et al., "Short-term diabetes attenuates left ventricular dysfunction and mortality rates after myocardial infarction in rodents," *Clinics*, vol. 66, no. 8, pp. 1437–1442, 2011.
- [73] B. Rodrigues, C. T. Mostarda, K. T. Rosa et al., "VEGF165 gene therapy improves left ventricular function and exercise capacity in diabetic rats after myocardial infarction: impact on mortality rate," *Journal of Diabetes & Metabolism*, vol. 4, article 006, 2011.
- [74] M. U. P. Brandao, M. Wajngarten, E. Rondon, M. C. P. Giorgi, F. Hironaka, and C. E. Negrão, "Left ventricular function during dynamic exercise in untrained and moderately trained subjects," *Journal of Applied Physiology*, vol. 75, no. 5, pp. 1989–1995, 1993.
- [75] A. D'Andrea, P. Caso, S. Severino et al., "Effects of different training protocols on left ventricular myocardial function in competitive athletes: a Doppler tissue imaging study," *Italian Heart Journal*, vol. 3, no. 1, pp. 34–40, 2002.
- [76] L. M. Shapiro and R. G. Smith, "Effect of training on left ventricular structure and function. An echocardiographic study," *British Heart Journal*, vol. 50, no. 6, pp. 534–539, 1983.
- [77] A. M. Dart, I. T. Meredith, and G. L. Jennings, "Effects of 4 weeks endurance training on cardiac left ventricular structure and function," *Clinical and Experimental Pharmacology and Physiology*, vol. 19, no. 11, pp. 777–783, 1992.
- [78] M. L. Smith, D. L. Hudson, H. M. Graitzer, and P. B. Raven, "Exercise training bradycardia: the role of autonomic balance," *Medicine and Science in Sports and Exercise*, vol. 21, no. 1, pp. 40–44, 1989.
- [79] K. de Angelis, R. B. Wichi, W. R. A. Jesus et al., "Exercise training changes autonomic cardiovascular balance in mice," *Journal of Applied Physiology*, vol. 96, no. 6, pp. 2174–2178, 2004.
- [80] P. C. Brum, G. J. Justo da Silva, E. Dias Moreira, F. Ida, C. E. Negrão, and E. M. Krieger, "Exercise training increases baroreceptor gain sensitivity in normal and hypertensive rats," *Hypertension*, vol. 36, no. 6, pp. 1018–1022, 2000.
- [81] F. Favret, K. K. Henderson, R. L. Clancy, J.-P. Richalet, and N. C. Gonzalez, "Exercise training alters the effect of chronic hypoxia on myocardial adrenergic and muscarinic receptor number," *Journal of Applied Physiology*, vol. 91, no. 3, pp. 1283–1288, 2001.
- [82] M. T. la Rovere, A. Mortara, G. Sandrone, and F. Lombardi, "Autonomic nervous system adaptations to short-term exercise training," *Chest*, vol. 101, no. 5, supplement, pp. 299S–303S, 1992.
- [83] F. Iellamo, J. M. Legramante, M. Massaro, G. Raimondi, and A. Galante, "Effects of a residential exercise training on baroreflex sensitivity and heart rate variability in patients with coronary artery disease: a randomized, controlled study," *Circulation*, vol. 102, no. 21, pp. 2588–2592, 2000.
- [84] G. Malfatto, M. Facchini, R. Bragato, G. Branzi, L. Sala, and G. Leonetti, "Short and long term effects of exercise training on the tonic autonomic modulation of heart rate variability after myocardial infarction," *European Heart Journal*, vol. 17, no. 4, pp. 532–538, 1996.
- [85] D. G. Martinez, J. C. Nicolau, R. L. Lage et al., "Effects of long-term exercise training on autonomic control in myocardial

- infarction patients," *Hypertension*, vol. 58, no. 6, pp. 1049–1056, 2011.
- [86] B. Rodrigues, L. Jorge, C. T. Mostarda et al., "Aerobic exercise training delays cardiac dysfunction and improves autonomic control of circulation in diabetic rats undergoing myocardial infarction," *Journal of Cardiac Failure*, vol. 18, no. 9, pp. 734–744, 2012.
- [87] N. P. Walsh, M. Gleeson, R. J. Shephard et al., "Position statement part one: immune function and exercise," *Exercise Immunology Review*, vol. 17, pp. 6–63, 2011.
- [88] B. K. Pedersen and L. Hoffman-Goetz, "Exercise and the immune system: regulation, integration, and adaptation," *Physiological Reviews*, vol. 80, no. 3, pp. 1055–1081, 2000.
- [89] K. Suzuki, S. Nakaji, M. Yamada, M. Totsuka, K. Sato, and K. Sugawara, "Systemic inflammatory response to exhaustive exercise. Cytokine kinetics," *Exercise Immunology Review*, vol. 8, pp. 6–48, 2002.
- [90] M. A. Febbraio and B. K. Pedersen, "Muscle-derived interleukin-6: mechanisms for activation and possible biological roles," *FASEB Journal*, vol. 16, no. 11, pp. 1335–1347, 2002.
- [91] R. Schindler, J. Mancilla, S. Endres, R. Ghorbani, S. C. Clark, and C. A. Dinarello, "Correlations and interactions in the production of interleukin-6 (IL-6), IL-1, and tumor necrosis factor (TNF) in human blood mononuclear cells: IL-6 suppresses IL-1 and TNF," *Blood*, vol. 75, no. 1, pp. 40–47, 1990.
- [92] A. Steensberg, C. P. Fischer, C. Keller, K. Møller, and B. K. Pedersen, "IL-6 enhances plasma IL-1ra, IL-10, and cortisol in humans," *American Journal of Physiology. Endocrinology and Metabolism*, vol. 285, no. 2, pp. E433–E437, 2003.
- [93] M. Pretolani, "Interleukin-10: an anti-inflammatory cytokine with therapeutic potential," *Clinical and Experimental Allergy*, vol. 29, no. 9, pp. 1164–1171, 1999.
- [94] A. M. W. Petersen and B. K. Pedersen, "The anti-inflammatory effect of exercise," *Journal of Applied Physiology*, vol. 98, no. 4, pp. 1154–1162, 2005.
- [95] S. Adamopoulos, J. Parissis, C. Kroupis et al., "Physical training reduces peripheral markers of inflammation in patients with chronic heart failure," *European Heart Journal*, vol. 22, no. 9, pp. 791–797, 2001.
- [96] M. L. Batista Jr., J. C. Rosa, R. D. Lopes et al., "Exercise training changes IL-10/TNF- α ratio in the skeletal muscle of post-MI rats," *Cytokine*, vol. 49, no. 1, pp. 102–108, 2010.
- [97] J. K. Smith, R. Dykes, J. E. Douglas, G. Krishnaswamy, and S. Berk, "Long-term exercise and atherogenic activity of blood mononuclear cells in persons at risk of developing ischemic heart disease," *Journal of the American Medical Association*, vol. 281, no. 18, pp. 1722–1727, 1999.
- [98] S. Adamopoulos, J. Parissis, D. Karatzas et al., "Physical training modulates proinflammatory cytokines and the soluble Fas/soluble Fas ligand system in patients with chronic heart failure," *Journal of the American College of Cardiology*, vol. 39, no. 4, pp. 653–663, 2002.
- [99] V. M. Conraads, P. Beckers, J. Bosmans et al., "Combined endurance/resistance training reduces plasma TNF- α receptor levels in patients with chronic heart failure and coronary artery disease," *European Heart Journal*, vol. 23, no. 23, pp. 1854–1860, 2002.
- [100] C. F. Grans, D. J. Feriani, M. E. V. Abssamra et al., "Resistance training after myocardial infarction in rats: role on cardiac and autonomic function," *Arquivos Brasileiros de Cardiologia*. In press.
- [101] C. Brandt and B. K. Pedersen, "The role of exercise-induced myokines in muscle homeostasis and the defense against chronic diseases," *Journal of Biomedicine and Biotechnology*, vol. 2010, Article ID 520258, 6 pages, 2010.
- [102] N. Mathur and B. K. Pedersen, "Exercise as a mean to control low-grade systemic inflammation," *Mediators of Inflammation*, vol. 2008, Article ID 109502, 6 pages, 2008.
- [103] M. G. Flynn and B. K. McFarlin, "Toll-like receptor 4: link to the anti-inflammatory effects of exercise?" *Exercise and Sport Sciences Reviews*, vol. 34, no. 4, pp. 176–181, 2006.
- [104] B. K. Pedersen and M. A. Febbraio, "Muscle as an endocrine organ: focus on muscle-derived interleukin-6," *Physiological Reviews*, vol. 88, no. 4, pp. 1379–1406, 2008.
- [105] M. Gleeson, B. McFarlin, and M. Flynn, "Exercise and toll-like receptors," *Exercise Immunology Review*, vol. 12, pp. 34–53, 2006.
- [106] N. Kawanishi, H. Yano, Y. Yokogawa, and K. Suzuki, "Exercise training inhibits inflammation in adipose tissue via both suppression of macrophage infiltration and acceleration of phenotypic switching from M1 to M2 macrophages in high-fat-diet-induced obese mice," *Exercise Immunology Review*, vol. 16, pp. 105–118, 2010.
- [107] J. S. Yudkin, "Inflammation, obesity, and the metabolic syndrome," *Hormone and Metabolic Research*, vol. 39, no. 10, pp. 707–709, 2007.
- [108] P. P. Mujumdar, P. J. Duerksen-Hughes, A. F. Firek, and D. A. Hessinger, "Long-term, progressive, aerobic training increases adiponectin in middle-aged, overweight, untrained males and females," *Scandinavian Journal of Clinical and Laboratory Investigation*, vol. 71, no. 2, pp. 101–107, 2011.
- [109] O. Ben Ounis, M. Elloumi, G. Lac et al., "Two-month effects of individualized exercise training with or without caloric restriction on plasma adipocytokine levels in obese female adolescents," *Annales d'Endocrinologie*, vol. 70, no. 4, pp. 235–241, 2009.
- [110] S. Lim, H. C. Sung, I.-K. Jeong et al., "Insulin-sensitizing effects of exercise on adiponectin and retinol-binding protein-4 concentrations in young and middle-aged women," *Journal of Clinical Endocrinology and Metabolism*, vol. 93, no. 6, pp. 2263–2268, 2008.
- [111] R. B. Nunes, J. P. Alves, L. P. Kessler, and P. dal Lago, "Aerobic exercise improves the inflammatory profile correlated with cardiac remodeling and function in chronic heart failure rats," *Clinics*, vol. 68, no. 6, pp. 876–882, 2013.
- [112] M. Gleeson, N. C. Bishop, D. J. Stensel, M. R. Lindley, S. S. Mastana, and M. A. Nimmo, "The anti-inflammatory effects of exercise: mechanisms and implications for the prevention and treatment of disease," *Nature Reviews Immunology*, vol. 11, no. 9, pp. 607–610, 2011.
- [113] G. I. Lancaster, Q. Khan, P. Drysdale et al., "The physiological regulation of toll-like receptor expression and function in humans," *Journal of Physiology*, vol. 563, no. 3, pp. 945–955, 2005.