

Research Article

HLA Haplotypes and Genotypes Frequencies in Brazilian Chronic Periodontitis Patients

Emília Ângela Sippert,¹ Cléverson de Oliveira e Silva,² Christiane Maria Ayo,¹ Silvia Barbosa Dutra Marques,³ Jeane Eliete Laguila Visentainer,⁴ and Ana Maria Sell⁴

¹Post Graduation Program of Biosciences Applied to Pharmacy, Biomedicine and Clinical Analysis Department, Maringá State University, 87020900 Maringá, PR, Brazil

²Dentistry Department, Maringá State University, 87020900 Maringá, PR, Brazil

³Hemocentro of Campinas, HLA Laboratory, University of Campinas, 13083878 Campinas, SP, Brazil

⁴Basic Health Sciences Department, Maringá State University, 87020900 Maringá, PR, Brazil

Correspondence should be addressed to Ana Maria Sell; anamsell@gmail.com

Received 9 March 2015; Revised 24 May 2015; Accepted 30 May 2015

Academic Editor: Paula Barrionuevo

Copyright © 2015 Emília Ângela Sippert et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Human leukocyte antigens (HLA) have a pivotal role in immune response and may be involved in antigen recognition of periodontal pathogens. However, the associations of HLA with chronic periodontitis (CP) have not been previously studied in the Brazilian population. In an attempt to clarify the issue of genetic predisposition to CP, we examined the distribution of HLA alleles, genotypes, and haplotypes in patients from Southern Brazil. One hundred and eight CP patients and 151 healthy and unrelated controls with age-, gender-, and ethnicity-matched were HLA investigated by polymerase chain reaction with sequence specific oligonucleotides. To exclude smoking as a predisposing factor, statistical analyses were performed in the total sample and in nonsmoking individuals. The significant results showed a positive association of the A*02/HLA-B*40 haplotype with CP (total samples: 4.2% versus 0%, $P_c = 0.03$; nonsmokers: 4.3% versus 0%, $P_c = 0.23$) and a lower frequency of HLA-B*15/HLA-DRB1*11 haplotype in CP compared to controls (total samples: 0.0% versus 4.3%, $P_c = 0.04$; nonsmokers: 0 versus 5.1%, $P_c = 1.0$). In conclusion, the HLA-A*02/B*40 haplotype may contribute to the development of CP, while HLA-B*15/DRB1*11 haplotype might indicate resistance to disease among Brazilians.

1. Introduction

Chronic periodontitis (CP) is a common complex disease of the oral cavity that is characterized by an inflammatory response to commensal and pathogenic oral bacteria [1, 2]. Due to bacterial infection, periodontal tissues become inflamed and are slowly destroyed by the action of the inflammatory process. If the disease is left untreated, teeth lose their ligamentous supporting structure to the alveolar bone, the alveolar bone is resorbed, and the teeth become mobile, finally resulting in teeth loss [1]. CP is considered the main cause of tooth loss among adults and is associated with severe quality of life impact [3].

The inflammatory response of the periodontal tissues to infection is influenced by environmental factors as well as by genetic factors [4, 5]. It is estimated that 50% of the expression

of periodontitis in CP could be attributed to genetic factors [5]. The observation that periodontitis is a complex disease entity with a multifactorial etiology has led to the search for risk factors that predispose to periodontitis in general as well as distinctive risk factors that might predispose to different clinical presentations of this group of diseases.

The human leukocyte antigens (HLA) play an important role in immune responsiveness and may be involved in antigen recognition of periodontal pathogens [6]. These cell-surface molecules have a key role in antigen presentation and activation of T cells. The polymorphisms of HLA can directly affect the binding capability of antigen peptides and thus affect the antigen-specific T-cell response [7]. Hence, these polymorphisms could represent an important susceptibility or resistance factor to periodontitis.

For many years, researchers have periodically screened populations of patients with different forms of periodontitis for associations with HLA antigens [8–14] and consistent results in relation to CP could not be obtained up to now. This study aimed to investigate differences in allelic group, genotype, and haplotype frequencies of HLA classes I and II in a sample of Brazilian patients with CP compared with a control group without CP.

2. Materials and Methods

2.1. Sample Selection. Between January and September 2012, a total of 259 individuals were selected from those who sought dental treatment at the dental clinics of Maringá State University (UEM) and Inga University (UNINGÁ) at Maringá, PR, Brazil (north/northwest region of the State of Parana, located in the southern region of Brazil, between 22°29'30"–26°42'59"S and 48°02'24"–54°37'38"W). Males and females, ethnically similar, aged over 34 years and with at least 20 teeth in the buccal cavity participated in this study. The criteria for exclusion were as follows: individuals with acute infections or diseases with known associations to HLA alleles such as diabetes, rheumatic diseases, systemic lupus erythematosus or narcolepsy, use of antibiotics during the last six months, and chronic usage of anti-inflammatory drugs or lactations and those who were pregnant.

After taking the patient's history, clinical periodontal examinations were conducted by two examiners. Clinical parameters of probing depth (PD) and clinical attachment level (CAL) were examined at six sites (mesiovestibular, vestibular, distovestibular, mesiolingual, lingual, and distolingual) of each tooth, as was bleeding on probing (BOP). After the periodontal examination, participants were categorized into two different groups: the CP group ($n = 108$) composed of individuals who had at least 5 sites in different teeth with $PD \geq 5$ mm, $CAL \geq 3$ mm, and more than 25% of BOP; and the control group ($n = 191$), formed by individuals who did not have sites with reduced CAL, displayed a PD of less than 4 mm, and exhibited less than 25% of BOP. Therefore, the control group was matched to the case group according to epidemiological characteristics such as ethnicity, gender, and geographical region. Information on the habit of smoking was obtained by interviewing the individual (anamnesis).

2.2. Ethics Information. All individuals who agreed to participate in this research were informed regarding the nature of the study and signed an informed consent form authorizing the use of their samples in the study, which was approved by the Ethics Committee for Human Research at Maringá State University (UEM-N°.719/2011, 02/12/2011).

2.3. Sample Collection and DNA Extraction. Blood samples (4 mL) were collected from the subjects in tubes containing anticoagulant (EDTA) and centrifuged at 210 g for 15 minutes, and the buffy-coat was conserved at -20°C until use. The genomic DNA was extracted using the salting-out method described by Miller et al. [15]. The concentration and quality of the DNA were analyzed by optical density in a Thermo Scientific Nanodrop 2000 apparatus (Wilmington, USA).

2.4. HLA Genotyping. HLA typing (HLA-A, HLA-B, HLA-C, DRB1, DQA1, and DQB1) was carried out using the polymerase chain reaction-sequence specific oligonucleotides technique (PCR-SSO; One Lambda, Canoga Park, CA, USA), low resolution to HLA-A, HLA-B, HLA-C, and DRB1, and high definition to HLA-DQA1 and DQB1. First, target DNA was PCR-amplified using group specific primers set, after the amplified product was biotinylated, which allowed later detection using R-phycoerythrin-conjugated streptavidin (SAPE), and hybridized with microspheres linked to specific conjugated fluorescent probes for HLA allele groups (One Lambda, Canoga Park, CA, USA). The fluorescent intensity varied based on reaction outcome and was expected to be 1000 or above for control positive probes. Reaction readings were carried out by flow cytometry using Luminex technology (One Lambda). Samples were analyzed through the HLA FUSION software (One Lambda Inc., San Diego, CA, USA).

2.5. Statistics. HLA specificity frequencies between the groups were compared using Fisher exact test. A two-sided P value (P) of <0.05 was considered statistically significant. Odds ratio (OR) values with a 95% confidence interval (95% CI) were also calculated to evaluate the risk of the individual developing the disease when having a particular HLA type. All the statistical analyses were performed using the R statistical environment [16]. However, to account for multiple comparisons, the observed P values were corrected (P_c) for the number of alleles when one locus was considered alone (Bonferroni correction). As the gametic phase of alleles from the different loci was not known and segregation analysis into families was not carried out, haplotypes and the calculation of their frequency were determined by the likelihood ratio test, through the Arlequin statistical software program [17]. Hardy-Weinberg equilibrium [18] was achieved by calculating the expected genotype frequencies and comparing them to the observed values, also using Arlequin statistical software program [17]. To exclude smoking as a predisposing factor statistical analyses were performed in the total sample (smokers, ex-smokers, and nonsmokers: patients versus control) as well as in nonsmoking patients versus nonsmoking controls. Only the haplotypes with frequencies above 1.0% were considered in this study.

3. Results

The study involved 259 individuals, 108 of whom were patients with CP, while 151 controls were healthy individuals. At the time of sample collection, the ages of the patients and controls ranged between 34 and 81 years (47.22% males and 52.78% females by patients and 35.10% males and 64.90% females by controls). Regarding the ethnic background, 59.26% of the patients with CP were reported to be Caucasians, 27.78% racially mixed, and 12.96% Afro-Brazilian. Concerning the control individuals, 69.54% were reported to be Caucasians, 23.84% racially mixed, and 6.62% Afro-Brazilian. No differences were observed between patients and controls according to gender, age, and ethnicity. Smoking was associated with CP and was more frequent in patients than in

TABLE 1: Characteristics of patients with chronic periodontitis and controls.

Category	Subcategory	Patients	Controls	Total
		(N = 108) n (%)	(N = 151) n (%)	(N = 259) n (%)
Gender	Female	57 (52.78)	98 (64.90)	157 (60.62)
	Male	51 (47.22)	53 (35.10)	102 (39.38)
Age	34–49 years	65 (60.18)	97 (64.24)	162 (62.55)
	50–65 years	35 (32.41)	50 (33.11)	85 (32.82)
	66–81 years	8 (7.41)	4 (2.65)	12 (4.63)
Ethnic background	Caucasian	64 (59.26)	105 (69.54)	169 (65.25)
	Afro-Brazilian	14 (12.96)	10 (6.62)	24 (9.27)
	Racially mixed	30 (27.78)	36 (23.84)	66 (25.48)
Smoking	Smoker (A)	23 (21.30)	17 (11.26)	40 (15.44)
	Nonsmoker (B)	46 (42.59)	108 (71.52)	154 (59.46)
	Ex-smoker (C)	39 (36.11)	26 (17.22)	65 (25.10)

B versus A: $P \leq 0.002$; OR = 0.32; 95% CI = 0.14–0.68.

B versus C: $P \leq 0.001$; OR = 0.29; 95% CI = 0.15–0.54.

A versus C: $P = 0.84$.

TABLE 2: HLA associations between chronic periodontitis patients and controls.

HLA	Total*			Nonsmokers**						
	Patients N = 216 n (%)	Controls N = 302 n (%)	P	P _c	OR (95% CI)	Patients N = 92 n (%)	Controls N = 216 n (%)	P	P _c	OR (95% CI)
A*02	49 (22.7)	86 (28.5)				17 (18.5)	66 (30.6)	0.035	0.63	0.52 (0.26–0.97)
A*32	3 (1.4)	15 (5.0)	0.030	0.45	0.27 (0.05–0.97)	1 (1.1)	15 (6.0)			
B*40	16 (7.4)	8 (2.6)	0.018	0.38	2.93 (1.16–8.07)	9 (9.8)	7 (3.2)	0.025	0.60	3.22 (1.03–10.55)
DRB1*08	17 (7.9)	11 (3.6)	0.048	0.58	2.26 (0.97–5.45)	8 (8.7)	8 (3.7)			
DQB1*06:01	0 (0)	7 (2.3)	0.045	0.63	0 (0–0.96)	0 (0.0)	5 (2.3)			
DQB1*06:09	5 (2.3)	0 (0.0)	0.012	0.17	undf (1.29–undf)	0 (0.0)	0 (0.0)			

undf = undefined.

*Total group (smokers, nonsmokers, and ex-smokers patients and controls).

**Nonsmoking group (patients and controls).

controls ($P = 0.043$; OR = 2.12; 95% CI = 1.02–4.51) as well as in ex-smokers compared to controls ($P = 0.001$, OR = 2.7; 95% IC = 1.46, 5.05) (Table 1).

In all groups, the distribution of HLA genotypes was confirmed to be in Hardy-Weinberg equilibrium ($P \geq 0.05$ in the patients and in the controls).

The most common HLA types were the following: HLA-A*01, *02, *03, and *24 and HLA-B*15, *35, *44, and *51, similar to other previous reports in the north/northwest region of the State of Parana [19]; this was an important control to demonstrate that the population was representative.

Table 2 presents the HLA classes I and II group allele frequencies distribution only for the significant differences between patients and controls. When comparing the HLA class I frequencies between CP and controls, HLA-A*32 had a lower frequency in patients with CP (smokers, nonsmokers,

and ex-smokers) and HLA-A*02 in nonsmokers group; HLA-B*40 had a higher frequency in CP total patients and also in nonsmokers patients group; no significant differences were found for HLA-C allelic groups. In relation to class II, HLA-DRB1*08 allelic group and DQB1*06:09 were overrepresented in patients compared to controls, and HLA-DQB1*06:01 was less frequent in patients compared with controls; however, neither of these associations was confirmed in the nonsmoker group.

Differences between CP and controls were found comparing the HLA genotypes frequencies for homo- and heterozygosis. HLA-C*03-C*06 genotype was increased in CP, whereas HLA-DQB1*03:01 homozygous genotype and DQB1*03:01/DQB1*03:02 were significantly less frequent in CP (Table 3). After Bonferroni correction, the associations lost the significance.

TABLE 3: HLA genotype frequency associations between chronic periodontitis patients and controls*.

HLA genotypes	Patients N = 108 n (%)	Controls N = 151 n (%)	P	P _c	OR (CI 95%)
C*03-C*06	4 (4.6)	0 (0)	0.012	0.56	undf (1.31–undf)
DQB1*03:01-DQB1*03:01	0 (0.0)	7 (4.6)	0.044	1.00	0 (0–0.95)
DQB1*03:01-DQB1*03:02	0 (0.0)	7 (4.6)	0.044	1.00	0 (0–0.95)

undf = undefined.

*Total group (smokers, nonsmokers, and ex-smoking patients and controls).

TABLE 4: HLA haplotype associations between chronic periodontitis patients and controls*.

HLA haplotypes	Patients N = 216 n (%)	Controls N = 302 n (%)	P	P _c	OR (95% CI)
A*02/B*40	9 (4.2)	0 (0.0)	<0.001	0.03	undf (2.83–undf)
A*03/B*51	5 (2.5)	0 (0.0)	0.012	0.36	undf (1.29–undf)
A*26/B*38	5 (2.3)	0 (0.0)	0.012	0.36	undf (1.29–undf)
B*40/C*03	12 (5.6)	5 (1.7)	0.022	0.62	3.49 (1.12–12.83)
B*18/C*12	4 (1.9)	0 (0.0)	0.03	0.93	undf (0.93–undf)
B*51/C*01	4 (1.9)	0 (0.0)	0.30	0.93	undf (0.93–undf)
B*15/DRB1*11	0 (0.0)	13 (4.3)	0.001	0.04	undf (0–0.45)
B*40/DRB1*07	6 (2.8)	0 (0.0)	0.005	0.19	undf (1.67–undf)
B*50/DRB1*04	4 (1.9)	0 (0.0)	0.030	1.14	undf (0.93–undf)
DQA1*01:02/DQB1*06:09	5 (2.3)	0 (0.0)	0.012	0.23	undf (1.30–undf)
DRB1*13/DQA1*01:02/DQB1*06:09	5 (2.3)	0 (0.0)	0.012	0.25	undf (1.29–undf)

undf = undefined.

*Total group (smokers, nonsmokers, and ex-smoking patients and controls).

Bold font: significant after Bonferroni correction.

Regarding haplotype frequencies (Tables 4 and 5) patients with CP had significantly higher frequencies for HLA-A*02/HLA-B*40 and a lower frequency for HLA-B*15/HLA-DRB1*11. Furthermore, patients with CP (only total samples) had higher haplotypes frequencies for B*18/C*12, B*51/C*01, B*50/DRB1*04, DQA1*01:02/DQB1*06:09, and DRB1*13/DQA1*01:02/DQB1*06:09 (Table 4); for non-smokers B*44/DRB1*07, B*35/DRB1*11, and B*14/DRB1*01 were more frequent and HLA-A*02/HLA-B*35 was less frequent in patients compared to controls (Table 5); A*03/B*51, A*26/B*38, B*40/C*03, and B*40/DRB1*07 (Tables 4 and 5) had higher frequencies for both groups, although for all of them the significance was lost after Bonferroni correction.

4. Discussion

While many studies have shown associations of HLA polymorphisms with aggressive periodontitis in different populations [8, 10–14, 20–24] there are few studies of HLA association with CP and the results are inconsistent [9–11, 20, 25–27]. To the best of our knowledge, this is the first study of HLA association with CP in the Brazilian population. Our results provide evidence that HLA classes I and II are associated with chronic periodontitis.

In general, discrepancies in results are observed in HLA case-control studies, and these discrepancies could be caused by differences in the choice of controls, disease diagnostic, racial background, and statistical analyses [28]. In this study, in order to avoid bias in the final results, the control population was selected after clinical examination with the same criteria used for patient exclusion, such as gestation and manifestations of infectious disease or known medical conditions that may contribute to development of CP. Patients with aggressive periodontitis were also excluded. Other confounding variables as gender, age, and ethnicity were considered. Gender and age ratio was similar between patients and controls. Reichert et al. [20] warned against the fact that gender could represent a confounding variable that should be considered in HLA and periodontitis studies; however, as HLA antigen expression has been reported to not vary as a function of gender [29, 30], this expression was probably not taken into consideration in many of the earlier studies on HLA association for periodontal diseases [10, 11, 13, 27, 31]. Age was not related to HLA expression but can be related to the disease, although severity and prevalence were mostly related to past disease history, social and behavior factors [32], and altered inflammatory responses [33].

Important to emphasize is that HLA varies according to population and ethnic group and that Brazil has an admixed

TABLE 5: HLA haplotype associations between nonsmokers, chronic periodontitis patients, and controls.

HLA haplotypes	Patients N = 92 n (%)	Controls N = 216 n (%)	P	P _c	OR (95% CI)
A*02/B*40	4 (4.3)	0 (0.0)	0.008	0.23	undf (1.58–undf)
A*02/B*35	0 (0.0)	11 (5.1)	0.038	1.00	0 (0–0.91)
A*03/B*51	3 (3.3)	0 (0.0)	0.026	0.75	undf (0.98–undf)
A*26/B*38	3 (3.3)	0 (0.0)	0.026	0.75	undf (0.98–undf)
B*40/C*03	7 (7.6)	4 (1.9)	0.019	0.51	4.34 (1.07–20.75)
B*44/B*07	4 (4.3)	1 (0.5)	0.029	0.78	9.69 (0.94–482.15)
B*15/DRB1*11	0 (0.0)	11 (5.1)	0.038	1.00	0 (0–0.91)
B*40/DRB1*07	4 (4.3)	1 (0.5)	0.029	0.81	9.69 (0.94–482.15)
B*35/DRB1*11	4 (4.3)	1 (0.5)	0.029	0.81	9.69 (0.94–482.15)
B*14/DRB1*01	4 (4.3)	0 (0.0)	0.008	0.22	undf (1.58–undf)

undf = undefined.

population. The population in Parana has been well defined regarding ethnicity and HLA distribution: in accordance with HLA phenotypic classification, the white population (majority) is predominantly of European origin (80.6%), with a smaller contribution of African (12.5%) and Amerindian (7.0%) genes [34]. Thus, regarding ethnic backgrounds, periodontitis patients and controls were similar, not representing a confounding variable. For this reason added to the high polymorphism of HLA, stratification by ethnicity was not realized.

In this study, as expected, smoking was associated with CP and having stopped smoking maintained susceptibility to disease. As smoking is a risk factor for the onset and progression of CP and the habit can obscure genetic risk factors [35] we analyzed nonsmoking patients and nonsmoking controls in addition to the total group (smokers, ex-smokers, and nonsmokers).

In the current study some trends to associations of HLA class I antigens with CP were found. HLA-A*32 was less frequent in total samples and HLA-A*02 was less frequent in nonsmoker patients representing possible protection to CP; however, significance was lost after Bonferroni correction. Added to these results, the HLA-A*02/B*35 haplotype was significantly less frequent in nonsmoker patients. Similar to our results, pioneering and recent investigations involving periodontitis and HLA have also found HLA*02 associated with protection to periodontitis [25, 27, 36, 37]. Thus, our results indicate a protective role of HLA-A*02. A2 could yield an efficient antimicrobial T-cell response reducing the disease [25].

Otherwise, HLA-B*40 was more frequent in both groups representing a risk effect against CP, although significance was also lost after Bonferroni correction. However, HLA-A*02/B*40 haplotype was significantly associated with susceptibility to CP. Added to this result, HLA-B*40/C*03 and HLA-B*40/DRB1*07 haplotypes tend to be associated with CP susceptibility. As HLA-A*02/B*40 was in linkage disequilibrium ($\Delta' > 0.70$, for all analyzed population: patients and controls) and HLA-A*02 has been associated as a protection factor, the susceptibility found for patients

with CP may be related to HLA-B*40. Individuals carrying B*40 (B60 and B61 antigens) had a three-fold higher risk of developing the disease. Contrarily, in another reporter, HLA-B*40 was associated with a lower clinical attachment loss in CP; however, this association was lost after Bonferroni correction [31].

Taking into account other haplotypes that were significantly associated in our population, the HLA-B*15-DRB1*11 combination must be highlighted conferring protection against CP in the total group of patients. HLA-B*15-DRB1*11 was in linkage disequilibrium ($\Delta' > 0.80$ for all analyzed population). HLA-B*15 could be associated with the disease: no associations between DRB1*11 and CP have been previously reported; however, in agreement with our results, Mauramo et al. [31] found that patients expressing HLA-B*15 had less clinical periodontal disease manifestation and better periodontal health over a long period of time. HLA-B*15 positive individuals might have a somewhat peculiar genetic response towards periodontal bacteria challenge contributing to CP development [31].

Another haplotype that lost the significance after Bonferroni correction in our study but must be highlighted was HLA-B*50/DRB1*04. This haplotype was a susceptibility factor to CP in our population and it is possible that the effect was associated with DRB1*04. According to previous reports, and in agreement with our results, HLA-DRB1*04 was associated with risk to CP [38, 39] and aggressive periodontitis [9, 12, 23, 30], as well as their alleles [8, 30]. DRB1*04:04 was considered a risk factor for bone loss [30, 40]. Contrarily, in a recent investigation, HLA-DRB1*04 or HLA-DRB1*04/DRB4*(DR53)/DQB1*03:02 haplotype had a decreased colonization risk of *Aggregatibacter actinomycetemcomitans* [9]. DRB1*04 was an immunogenetic susceptibility factor for type 1 diabetes [41] and for rheumatoid arthritis [30]; however, both diseases were included in our exclusion criteria.

According to other association studies between periodontitis and HLA, a meta-analysis focusing on Caucasian case-control studies conducted by Stein et al. [25] demonstrated no associations between HLA and CP, although for

aggressive periodontitis HLA-A*09 and B*15 appeared to represent susceptibility factors and HLA-A*02 and B*05 were potential protective factors. Other HLA-A, HLA-B, and HLA-C associations with periodontal disease previously described in the literature were as follows: susceptibility mediated by HLA-A*11, A*29, B*14, and C*08 and protection by HLA-A*03, HLA-A*31 and -A*30/A*31 genotype observed in German patients with CP [11] and HLA-B*57 as a protecting factor in German patients with CP or generalized aggressive periodontitis [9]; HLA-A*02 and HLA-B*05 associated with protection in the CP patients from USA [27, 36, 37]; HLA-A*09 positively associated with CP from France [26]; and HLA-B*51 related to fewer deep periodontal pockets in the CP in Swiss adults. HLA-DQA1*03 and HLA-DRB1*04 were higher and HLA-DQB1*06:03 might have protective effects against aggressive periodontitis in Iranian patients [8].

A possible role of HLA in the immune response and development of periodontal disease may be related to their ability to bind some processed peptides from bacteria antigens and expressing them on the surface of antigen presenting cells (peptide-HLA class II) or target cells (peptide-HLA class I) in order to present them to T cells (CD4 or CD8) [42]. The binding capacity of the bacteria peptide depends on HLA allotypic structure of their paratope. Failures in this link capacity can compromise the immune response and could be a risk of disease. The large individual capacity of immune response occurs due to the fact that HLA is highly polymorphic: each gene has multiple alleles and each individual has many expressed genes. These preliminary results indicate that HLA haplotypes might be involved in the susceptibility or risk for periodontal disease. However, further investigations of HLA haplotypes markers in relation to antigenic peptide-binding motifs are necessary in order to understand its relation in periodontitis.

The weaknesses of the case-control studies were related to the reproduction of results in function of ethnicity background of the individuals, sample size, diagnostic tools of disease, and HLA genotyping methods. Consequently, no strong association could be found, especially in CP. In addition, CP is a complex and multifactorial disease and several other genetic polymorphisms have been reported to be associated with bacterium response, inflammation, chronicity, and wide ranging systemic effects and disease development, such as *IL1*, *IL4*, *TNF*, *IL8*, and *IL10* cytokines genes, immunoglobulin G Fc receptor (FcγR), and TLR [35, 43–60]. Despite this fact, all information regarding genetic susceptibility to diseases is valuable and can be applied to therapy intervention or individual approach. For better results, we have tried to be very critical in the choice of the study population. CP inclusion criteria and CP and control exclusion criteria were defined in order to avoid confounding factors and were a representative of generally healthy and CP adults. As independent multiple comparisons were carried out and we considered that all genotypes examined had the same chance of being increased or decreased in CP, Bonferroni correction was applied for all significant *P* values.

The present study has potential limitations. We focused on the HLA allele group, not on alleles and epitopes analysis, and split antigens were not investigated. The major limitation

of this study is the relatively small sample size regarding the analysis of the nonsmoker group.

In these Brazilian CP patients, the HLA-A*02/B*40 haplotype was considered a risk factor for CP and HLA-B*15/DRB1*11 haplotype was considered a protective factor for disease. Susceptibility may possibly be associated with B*40 molecules and protection may be associated with B*15 and A*02. Further investigation relating antigenic peptide-binding motifs and their immunopathogenesis involving CP development should be investigated.

5. Conclusion

These results provide evidence that class I and II HLA polymorphisms are associated with chronic periodontitis. HLA-A*02/B*40 haplotype seems to represent susceptibility factors and HLA-B*15/DRB1*11 haplotype was potential protective factors against disease.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Emília Ângela Sippert, Christiane Maria Ayo, and Silvia Barbosa Dutra Marques carried out the molecular genetic studies, Cléverson de Oliveira e Silva participated in the clinical diagnostic of chronic periodontitis, Ana Maria Sell and Jeane Eliete Laguila Visentainer designed and coordinated the study, and Emília Ângela Sippert, Ana Maria Sell, and Jeane Eliete Laguila Visentainer wrote the paper. All authors read and approved the final paper.

Acknowledgments

The authors wish to thank all the volunteers who were part of this study. They thank Camila Rodrigues, Fabiano Cavalcanti de Melo, and Marco Antonio Braga for help with technical performance. They thank Roberto Zulli for the contribution in statistical analysis and Raquel Foglio for language review of this paper.

References

- [1] M. L. Laine, W. Crielaard, and B. G. Loos, "Genetic susceptibility to periodontitis," *Periodontology 2000*, vol. 58, no. 1, pp. 37–68, 2012.
- [2] A. B. Berezow and R. P. Darveau, "Microbial shift and periodontitis," *Periodontology 2000*, vol. 55, no. 1, pp. 36–47, 2011.
- [3] P. E. Petersen and H. Ogawa, "Strengthening the prevention of periodontal disease: the WHO approach," *Journal of Periodontology*, vol. 76, no. 12, pp. 2187–2193, 2005.
- [4] D. F. Kinane, M. Peterson, and P. G. Stathopoulou, "Environmental and other modifying factors of the periodontal diseases," *Periodontology 2000*, vol. 40, no. 1, pp. 107–119, 2006.
- [5] B. S. Michalowicz, S. R. Diehl, J. C. Gunsolley et al., "Evidence of a substantial genetic basis for risk of adult periodontitis," *Journal of Periodontology*, vol. 71, no. 11, pp. 1699–1707, 2000.

- [6] M. Baines and A. Ebringer, "HLA and disease," *Molecular Aspects of Medicine*, vol. 13, no. 4, pp. 263–378, 1992.
- [7] R. M. Zinkernagel and P. C. Doherty, "The discovery of MHC restriction," *Immunology Today*, vol. 18, no. 1, pp. 14–17, 1997.
- [8] M. M. Jazi, G. Solgi, H. S. Roosta et al., "HLA-DRB and HLA-DQA/HLA-DQB allele and haplotype frequencies in Iranian patients with aggressive periodontitis," *Journal of Periodontal Research*, vol. 48, no. 4, pp. 533–539, 2013.
- [9] S. Reichert, W. Altermann, J. M. Stein, H.-G. Schaller, H. K. G. MacHulla, and S. Schulz, "Individual composition of human leukocyte antigens and periodontopathogens in the background of periodontitis," *Journal of Periodontology*, vol. 84, no. 1, pp. 100–109, 2013.
- [10] J. Stein, S. Reichert, A. Gautsch, and H. K. G. Machulla, "Are there HLA combinations typical supporting for or making resistant against aggressive and/or chronic periodontitis?" *Journal of Periodontal Research*, vol. 38, no. 5, pp. 508–517, 2003.
- [11] H. K. G. Machulla, J. Stein, A. Gautsch, J. Langner, H.-G. Schaller, and S. Reichert, "HLA-A, B, Cw, DRB1, DRB3/4/5, DQB1 in German patients suffering from rapidly progressive periodontitis (RPP) and adult periodontitis (AP)," *Journal of Clinical Periodontology*, vol. 29, no. 6, pp. 573–579, 2002.
- [12] E. Firatli, A. Kantarci, I. Cebeci et al., "Association between HLA antigens and early onset periodontitis," *Journal of Clinical Periodontology*, vol. 23, no. 6, pp. 563–566, 1996.
- [13] L. Shapira, S. Eizenberg, M. N. Sela, A. Soskolne, and H. Brautbar, "HLA A9 and B15 are associated with the generalized form, but not the localized form, of early-onset periodontal diseases," *Journal of Periodontology*, vol. 65, no. 3, pp. 219–223, 1994.
- [14] A. Amer, G. Singh, C. Darke, and A. E. Dolby, "Association between HLA antigens and periodontal disease," *Tissue Antigens*, vol. 31, no. 2, pp. 53–58, 1988.
- [15] S. A. Miller, D. D. Dykes, and H. F. Polesky, "A simple salting out procedure for extracting DNA from human nucleated cells," *Nucleic Acids Research*, vol. 16, no. 3, article 1215, 1988.
- [16] R Core Team, *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, 2015, <http://www.r-project.org/>.
- [17] L. Excoffier, G. Laval, and S. Schneider, "Arlequin (version 3.0): an integrated software package for population genetics data analysis," *Evolutionary Bioinformatics Online*, vol. 1, pp. 47–50, 2005.
- [18] S. W. Guo and E. A. Thompson, "Performing the exact test of Hardy-Weinberg proportion for multiple alleles," *Biometrics*, vol. 48, no. 2, pp. 361–372, 1992.
- [19] D. S. A. Franceschi, L. T. Tsuneto, P. S. Mazini et al., "Class-I human leukocyte alleles in leprosy patients from southern Brazil," *Revista da Sociedade Brasileira de Medicina Tropical*, vol. 44, no. 5, pp. 616–620, 2011.
- [20] S. Reichert, J. Stein, A. Gautsch, H.-G. Schaller, and H. K. G. Machulla, "Gender differences in HLA phenotype frequencies found in German patients with generalized aggressive periodontitis and chronic periodontitis," *Oral Microbiology and Immunology*, vol. 17, no. 6, pp. 360–368, 2002.
- [21] H. Ohyama, S. Takashiba, K. Oyaizu et al., "HLA class II genotypes associated with early-onset periodontitis: DQB1 molecule primarily confers susceptibility to the disease," *Journal of Periodontology*, vol. 67, no. 9, pp. 888–894, 1996.
- [22] J. H. Moses, H. Tsichti, P. Donaldson, P. B. Smith, N. W. Johnson, and J. G. Bodmer, "HLA and susceptibility to juvenile periodontitis in Afro-Caribbeans," *Tissue Antigens*, vol. 43, no. 5, pp. 316–319, 1994.
- [23] J. Katz, J. Goultchin, R. Benoliel, and C. Brautbar, "Human leukocyte antigen (HLA) DR4. Positive association with rapidly progressing periodontitis," *Journal of Periodontology*, vol. 58, no. 9, pp. 607–610, 1987.
- [24] P. T. Klouda, S. R. Porter, C. Scully et al., "Association between HLA-A9 and rapidly progressive periodontitis," *Tissue Antigens*, vol. 28, no. 3, pp. 146–149, 1986.
- [25] J. M. Stein, H. K. G. MacHulla, R. Smeets, F. Lampert, and S. Reichert, "Human leukocyte antigen polymorphism in chronic and aggressive periodontitis among Caucasians: a meta-analysis," *Journal of Clinical Periodontology*, vol. 35, no. 3, pp. 183–192, 2008.
- [26] A. Balndin-Texier, M. Gueguin, R. Fauchet, M. Yardin, and G. Cathelineau, "The HLA-A9 antigen and chronic periodontitis," *Journal de Parodontologie*, vol. 5, no. 3, pp. 221–227, 1986.
- [27] D. Goteiner and M. J. Goldman, "Human lymphocyte antigen haplotype and resistance to periodontitis," *Journal of Periodontology*, vol. 55, no. 3, pp. 155–158, 1984.
- [28] A. G. Pacheco and M. O. Moraes, "Genetic polymorphisms of infectious diseases in case-control studies," *Disease Markers*, vol. 27, no. 3–4, pp. 173–186, 2009.
- [29] J. L. Tiwari and P. I. Terasaki, "HLA-DR and disease associations," *Progress in clinical and biological research*, vol. 58, pp. 151–163, 1981.
- [30] J. J. Bonfil, F. L. Dillier, P. Mercier et al., "A 'case control' study on the rôle of HLA DR4 in severe periodontitis and rapidly progressive periodontitis: identification of types and subtypes using molecular biology (PCR.SSO)," *Journal of Clinical Periodontology*, vol. 26, no. 2, pp. 77–84, 1999.
- [31] M. Mauramo, A. M. Ramseier, A. Buser et al., "Associations of HLA-A, -B and -DRB1 types with oral diseases in Swiss adults," *PLoS ONE*, vol. 9, no. 7, Article ID e103527, 2014.
- [32] P. R. Kamen, "Periodontal care," *Dental Clinics of North America*, vol. 41, no. 4, pp. 751–762, 1997.
- [33] O. A. Gonzalez, M. J. Novak, S. Kirakodu et al., "Comparative analysis of gingival tissue antigen presentation pathways in ageing and periodontitis," *Journal of Clinical Periodontology*, vol. 41, no. 4, pp. 327–339, 2014.
- [34] C. M. Probst, E. P. Bompeixe, N. F. Pereira et al., "HLA polymorphism and evaluation of European, African, and Amerindian contribution to the white and mulatto populations from Parana, Brazil," *Human Biology*, vol. 72, no. 4, pp. 597–617, 2000.
- [35] K. S. Kornman, A. Crane, H. Y. Wang et al., "The interleukin-1 genotype as a severity factor in adult periodontal disease," *Journal of Clinical Periodontology*, vol. 24, no. 1, pp. 72–77, 1997.
- [36] P. I. Terasaki, R. S. Kaslick, T. L. West, and A. I. Chasens, "Low HL A2 frequency and periodontitis," *Tissue Antigens*, vol. 5, no. 4, pp. 286–288, 1975.
- [37] R. S. Kaslick, T. L. West, A. I. Chasens, P. I. Terasaki, R. Lazzara, and S. Weinberg, "Association between HL-A2 antigen and various periodontal diseases in young adults," *Journal of Dental Research*, vol. 54, no. 2, p. 424, 1975.
- [38] C. S. Alley, R. A. Reinhardt, C. A. Maze et al., "HLA-D and T lymphocyte reactivity to specific periodontal pathogens in type 1 diabetic periodontitis," *Journal of Periodontology*, vol. 64, no. 10, pp. 974–979, 1993.

- [39] J. K. Dyer, M. A. Peck, R. A. Reinhardt et al., "HLA-D types and serum IgG responses to capnocytophaga in diabetes and periodontitis," *Journal of Dental Research*, vol. 76, no. 12, pp. 1825–1832, 1997.
- [40] C. E. Repeke, C. R. Cardoso, M. Claudino et al., "Non-inflammatory destructive periodontal disease: a clinical, microbiological, immunological and genetic investigation," *Journal of Applied Oral Science*, vol. 20, no. 1, pp. 113–121, 2012.
- [41] C. Alves, I. Meyer, N. Vieira, M. B. P. Toralles, and D. LeMaire, "Distribution and frequency of HLA alleles and haplotypes in Brazilians with type 1 diabetes mellitus," *Arquivos Brasileiros de Endocrinologia & Metabologia*, vol. 50, no. 3, pp. 436–444, 2006.
- [42] S. Nair, M. Faizuddin, and J. Dharmapalan, "Role of autoimmune responses in periodontal disease," *Autoimmune Diseases*, vol. 2014, Article ID 596824, 7 pages, 2014.
- [43] P. C. Trevilatto, A. P. de Souza Pardo, R. M. Scarel-Caminaga et al., "Association of IL1 gene polymorphisms with chronic periodontitis in Brazilians," *Archives of Oral Biology*, vol. 56, no. 1, pp. 54–62, 2011.
- [44] M. L. Laine, M. A. Farré, M. A. Garcíagonzález et al., "Polymorphisms of the interleukin-1 gene family, oral microbial pathogens, and smoking in adult periodontitis," *Journal of Dental Research*, vol. 80, no. 8, pp. 1695–1699, 2001.
- [45] S. B. Ferreira Jr., A. P. F. Trombone, C. E. Repeke et al., "An interleukin-1 β (IL-1 β) single-nucleotide polymorphism at position 3954 and red complex periodontopathogens independently and additively modulate the levels of IL-1 β in diseased periodontal tissues," *Infection and Immunity*, vol. 76, no. 8, pp. 3725–3734, 2008.
- [46] N. J. López, L. Jara, and C. Y. Valenzuela, "Association of interleukin-1 polymorphisms with periodontal disease," *Journal of Periodontology*, vol. 76, no. 2, pp. 234–243, 2005.
- [47] J. Craandijk, M. V. Van Krugten, C. L. Verweij, U. Van Der Velden, and B. G. Loos, "Tumor necrosis factor-alpha gene polymorphisms in relation to periodontitis," *Journal of Clinical Periodontology*, vol. 29, no. 1, pp. 28–34, 2002.
- [48] E. A. Gore, J. J. Sanders, J. P. Pandey, Y. Palesch, and G. M. P. Galbraith, "Interleukin-1beta+3953 allele 2: association with disease status in adult periodontitis," *Journal of Clinical Periodontology*, vol. 25, no. 10, pp. 781–785, 1998.
- [49] G. M. P. Galbraith, R. B. Steed, J. J. Sanders, and J. P. Pandey, "Tumor necrosis factor alpha production by oral leukocytes: influence of tumor necrosis factor genotype," *Journal of Periodontology*, vol. 69, no. 4, pp. 428–433, 1998.
- [50] L. I. Holla, A. Fassmann, P. Augustin, T. Halabala, V. Znojil, and J. Vanek, "The association of interleukin-4 haplotypes with chronic periodontitis in a Czech population," *Journal of Periodontology*, vol. 79, no. 10, pp. 1927–1933, 2008.
- [51] C. C. Pontes, J. R. Gonzales, A. B. Novaes Jr. et al., "Interleukin-4 gene polymorphism and its relation to periodontal disease in a Brazilian population of African heritage," *Journal of Dentistry*, vol. 32, no. 3, pp. 241–246, 2004.
- [52] T. Kobayashi, A. Murasawa, S. Ito et al., "Cytokine gene polymorphisms associated with rheumatoid arthritis and periodontitis in Japanese adults," *Journal of Periodontology*, vol. 80, no. 5, pp. 792–799, 2009.
- [53] Y. J. Kim, A. C. Viana, K. M. C. Curtis, S. R. P. Orrico, J. A. Cirelli, and R. M. Scarel-Caminaga, "Lack of association of a functional polymorphism in the interleukin 8 gene with susceptibility to periodontitis," *DNA and Cell Biology*, vol. 28, no. 4, pp. 185–190, 2009.
- [54] R. M. Scarel-Caminaga, P. C. Trevilatto, A. P. Souza, R. B. Brito, L. E. A. Camargo, and S. R. P. Line, "Interleukin 10 gene promoter polymorphisms are associated with chronic periodontitis," *Journal of Clinical Periodontology*, vol. 31, no. 6, pp. 443–448, 2004.
- [55] E. Angela Sippert, C. D. O. Silva, J. E. L. Visentainer, and A. M. Sell, "Association of duffy blood group gene polymorphisms with IL8 gene in chronic periodontitis," *PLoS ONE*, vol. 8, no. 12, Article ID e83286, 2013.
- [56] M. Claudino, A. P. F. Trombone, C. R. Cardoso et al., "The broad effects of the functional IL-10 promoter-592 polymorphism: modulation of IL-10, TIMP-3, and OPG expression and their association with periodontal disease outcome," *Journal of Leukocyte Biology*, vol. 84, no. 6, pp. 1565–1573, 2008.
- [57] P. R. Moreira, J. E. Costa, R. S. Gomez, K. J. Gollob, and W. O. Dutra, "TNFA and IL10 gene polymorphisms are not associated with periodontitis in Brazilians," *The Open Dentistry Journal*, vol. 3, no. 1, pp. 184–190, 2009.
- [58] T. Kobayashi, N. A. C. Westerdaal, A. Miyazaki et al., "Relevance of immunoglobulin G Fc receptor polymorphism to recurrence of adult periodontitis in Japanese patients," *Infection and Immunity*, vol. 65, no. 9, pp. 3556–3560, 1997.
- [59] A. P. Colombo, C. Eftimiadi, A. D. Haffajee, M. A. Cugini, and S. S. Socransky, "Serum IgG2 level, Gm(23) allotype and Fc gammaRIIa and Fc gammaRIIb receptors in refractory periodontal disease," *Journal of Clinical Periodontology*, vol. 25, no. 6, pp. 465–474, 1998.
- [60] Z. Heidari, H. Mahmoudzadeh-Sagheb, M. A. Rigi-Ladiz, M. Taheri, A. Moazenni-Roodi, and M. Hashemi, "Association of TGF- β 1 -509 C/T, 29 C/T and 788 C/T gene polymorphisms with chronic periodontitis: a case-control study," *Gene*, vol. 518, no. 2, pp. 330–334, 2013.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

