Research Article

Well-Posedness of the Boundary Value Problem for Parabolic Equations in Difference Analogues of Spaces of Smooth Functions

A. Ashyralyev

Received 15 June 2006; Accepted 27 December 2006

Recommended by F. E. Udwardia

The first and second orders of accuracy difference schemes for the approximate solutions of the nonlocal boundary value problem $v'(t) + Av(t) = f(t)$ ($0 \leq t \leq 1$), $v(0) = v(\lambda) + \mu$, $0 < \lambda \leq 1$, for differential equation in an arbitrary Banach space E with the strongly positive operator A are considered. The well-posedness of these difference schemes in difference analogues of spaces of smooth functions is established. In applications, the coercive stability estimates for the solutions of difference schemes for the approximate solutions of the nonlocal boundary value problem for parabolic equation are obtained.

Copyright © 2007 A. Ashyralyev. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction: difference schemes

It is known that (see, e.g., [1–5] and the references given therein) many applied problems in fluid mechanics and other areas of physics and mathematical biology were formulated into nonlocal mathematical models. However, such problems were not well investigated in general.

In [6], the well-posedness in the spaces of smooth functions of the nonlocal boundary value problem

$$v'(t) + Av(t) = f(t) \quad (0 \leq t \leq 1), \quad v(0) = v(\lambda) + \mu \quad (0 < \lambda \leq 1) \quad (1.1)$$

for differential equation in an arbitrary Banach space E with the strongly positive operator A was established. The importance of coercive (well-posedness) inequalities is well known [7, 8].

2 Mathematical Problems in Engineering

For the construction of difference schemes, we consider the uniform grid space

$$[0, 1]_\tau = \{t_k = k\tau, 0 \leq k \leq N, N\tau = 1\}. \quad (1.2)$$

Assume that $\tau \leq \lambda$. We consider the first order of accuracy implicit Rothe difference scheme

$$\begin{aligned} \frac{u_k - u_{k-1}}{\tau} + Au_k &= \varphi_k, \quad \varphi_k = f(t_k), t_k = k\tau, 1 \leq k \leq N, \\ u_0 &= u_{[\lambda/\tau]} + \mu, \end{aligned} \quad (1.3)$$

and the second order of accuracy implicit difference scheme

$$\begin{aligned} \frac{u_k - u_{k-1}}{\tau} + A\left(I + \frac{\tau A}{2}\right)u_k &= \left(I + \frac{\tau A}{2}\right)\varphi_k, \quad \varphi_k = f\left(t_k - \frac{\tau}{2}\right), t_k = k\tau, 1 \leq k \leq N, \\ u_0 &= \left(I - \left(\lambda - \left[\frac{\lambda}{\tau}\right]\tau\right)A\right)u_{[\lambda/\tau]} + \mu + \left(\lambda - \left[\frac{\lambda}{\tau}\right]\tau\right)\varphi_{[\lambda/\tau]}, \end{aligned} \quad (1.4)$$

approximately solving the boundary value problem (1.1).

Let $F_\tau(E)$ be the linear space of mesh functions $\varphi^\tau = \{\varphi_k\}_1^N$ with values in the Banach space E . Next on $F_\tau(E)$, we introduce the Banach spaces $C_\tau(E) = C([0, 1]_\tau, E)$, $C_\tau^{\beta, \gamma}(E) = C^{\beta, \gamma}([0, 1]_\tau, E)$ ($0 \leq \gamma \leq \beta < 1$) with the norms

$$\begin{aligned} \|\varphi^\tau\|_{C_\tau(E)} &= \max_{1 \leq k \leq N} \|\varphi_k\|_E, \\ \|\varphi^\tau\|_{C_\tau^{\beta, \gamma}(E)} &= \|\varphi^\tau\|_{C_\tau(E)} + \sup_{1 \leq k < k+r \leq N} \|\varphi_{k+r} - \varphi_k\|_E \frac{((k+r)\tau)^\gamma}{(r\tau)^\beta}. \end{aligned} \quad (1.5)$$

We introduce the fractional space $E_\alpha = E_\alpha(E, A)$ ($0 < \alpha < 1$), consisting of all $v \in E$ for which the following norm is finite:

$$\|v\|_{E_\alpha} = \sup_{\lambda > 0} \lambda^\alpha \|A(\lambda + A)^{-1}v\|_E. \quad (1.6)$$

The difference scheme (1.3) or (1.4) is said to be coercively stable (well-posed) in $F_\tau(E)$ if we have the coercive inequality

$$\left\| \{\tau^{-1}(u_k - u_{k-1})\}_1^N \right\|_{F_\tau(E)} \leq M[\|A\mu\|_{E'} + \|\varphi^\tau\|_{F_\tau(E)}], \quad E' \subset E, \quad (1.7)$$

where M is independent not only of φ^τ, μ but also of τ .

In [9, 10], the stability and coercive stability of the difference schemes (1.3) and (1.4) in $C_\tau^{\alpha, \alpha}(E)$ and $C_\tau(E_\alpha)$ ($0 < \alpha < 1$) spaces and almost coercive stability (with multiplier $\min\{\ln 1/\tau, 1 + |\ln \|A\|_{E \rightarrow E}|\}$) of the difference schemes (1.3) and (1.4) in $C_\tau(E)$ spaces are established.

In the present paper, the coercive stability of difference schemes (1.3) and (1.4) in $C_\tau^{\beta,\gamma}(E)$ ($0 \leq \gamma \leq \beta < 1$) and $C_\tau^{\beta,\gamma}(E_{\alpha-\beta})$ ($0 \leq \gamma \leq \beta \leq \alpha < 1$) spaces under the assumption that the operator $-A$ generates an analytic semigroup $\exp\{-tA\}$ ($t \geq 0$) with exponentially decreasing norm, when $t \rightarrow +\infty$,

$$\|\exp\{-tA\}\|_{E \rightarrow E} \leq M e^{-\delta t}, \quad \|A \exp\{-tA\}\|_{E \rightarrow E} \leq \frac{M}{t}, \quad t > 0, \delta, M > 0, \quad (1.8)$$

is established. In applications, this abstract result permits us to obtain the almost coercivity inequality and the coercive stability estimates for the solutions of difference schemes of the first and second orders of accuracy over time and of an arbitrary order of accuracy over space variables in the case of the nonlocal boundary value problem for the $2m$ -order multidimensional parabolic equation.

Finally, methods for numerical solutions of the evolution differential equations have been studied extensively by many researchers (see [8, 11–32] and the references therein).

2. Well-posedness of (1.3) and (1.4)

THEOREM 2.1. *Let τ be a sufficiently small number. Then the solutions of the difference schemes (1.3) and (1.4) in $C_\tau^{\beta,\gamma}(E)$ ($0 \leq \gamma \leq \beta$, $0 < \beta < 1$) obey the coercivity inequality*

$$\begin{aligned} & \left\| \{\tau^{-1}(u_k - u_{k-1})\}_1^N \right\|_{C_\tau^{\beta,\gamma}(E)} + \left\| \{\tau^{-1}(u_k - u_{k-1})\}_1^N \right\|_{C_\tau(E_1^{\beta,\gamma})} \\ & \leq \frac{M_1}{\beta(1-\beta)} \|\varphi^\tau\|_{C_\tau^{\beta,\gamma}(E)} + M_1 \|\mu + A^{-1}(\varphi_{[\lambda\tau]} - \varphi_1)\|_1^{\beta,\gamma}, \end{aligned} \quad (2.1)$$

where M_1 is independent not only of φ^τ , μ , β , γ , but also of τ .

Here, the space of traces $E_1^{\beta,\gamma} = E^{\beta,\gamma}(E)$ which consist of the elements $w \in E$ for which the norm

$$\begin{aligned} |w|_1^{\beta,\gamma} = \sup_{0 < \tau \leq \tau_0} & \left[\max_{1 \leq i \leq N} \|\tau^{-1}(I - R)R^{i-1}w\|_E \right. \\ & \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \|\tau^{-1}(I - R)(R^{i+r-1} - R^{i-1})w\|_E \right] \end{aligned} \quad (2.2)$$

is finite, where $R = (I + \tau A)^{-1}$ for (1.3) and $R = (I + \tau A + (\tau A)^2/2)^{-1}$ for (1.4).

Proof. Let us prove (2.1) for difference scheme (1.3). By [7, formula (0.2) in Chapter 2],

$$u_k = R^k u_0 + \sum_{j=1}^k R^{k-j+1} \varphi_j \tau, \quad k = 1, \dots, N, \quad (2.3)$$

4 Mathematical Problems in Engineering

for the solution of the first order of accuracy implicit difference scheme for the approximate solutions of Cauchy problem

$$u'(t) + Au(t) = f(t) \quad (0 \leq t \leq 1) \quad u(0) = u_0. \quad (2.4)$$

From this formula and the condition $u_0 = u_{[\lambda/\tau]} + \mu$, it follows that

$$u_0 = R^{[\lambda/\tau]}u_0 + \sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1}\varphi_j\tau + \mu. \quad (2.5)$$

Since the semigroup $\exp\{-tA\}$ obeys the exponential decay estimate (1.8), we have that

$$\|R^k\|_{E \rightarrow E} \leq M(1 + \delta\tau)^{-k}, \quad \|k\tau AR^k\|_{E \rightarrow E} \leq M, \quad k \geq 1. \quad (2.6)$$

From this estimate, it follows that the operator $I - R^{[\lambda/\tau]}$ has a bounded inverse $T_\tau = (I - R^{[\lambda/\tau]})^{-1}$ and

$$\|T_\tau\|_{E \rightarrow E} \leq M(\lambda, \delta). \quad (2.7)$$

Actually, we have that

$$\begin{aligned} T_\tau - (I - \exp\{-\lambda A\})^{-1} &= T_\tau(I - \exp\{-\lambda A\})^{-1}(R^{[\lambda/\tau]} - \exp\{-\lambda A\}), \\ R^{[\lambda/\tau]} - \exp\{-\lambda A\} &= \int_0^1 (I + s\tau A)^{-([\lambda/\tau]+1)} \left(\left(\lambda - \left\lfloor \frac{\lambda}{\tau} \right\rfloor \tau \right) A + s\tau\lambda A^2 \right) \exp\{-\lambda(1-s)A\} ds. \end{aligned} \quad (2.8)$$

Then, using the triangle inequality and the estimates

$$\|(I - \exp\{-\lambda A\})^{-1}\|_{E \rightarrow E} \leq M(\lambda, \delta), \quad (2.9)$$

$$\|R^{[\lambda/\tau]} - \exp\{-\lambda A\}\|_{E \rightarrow E} \leq M(\lambda, \delta)\tau, \quad (2.10)$$

we obtain estimate (2.7). The proof of (2.9) is based on the estimate (1.8) and it was proved in [33]. The proof of (2.10) is based on the estimates (1.8) and (2.6) and it was proved in [34].

So, we have the formula

$$\begin{aligned} u_k &= R^k u_0 + \sum_{j=1}^k R^{k-j+1}\varphi_j\tau, \quad k = 1, \dots, N, \\ u_0 &= T_\tau \left\{ \sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1}\varphi_j\tau + \mu \right\} \end{aligned} \quad (2.11)$$

for the solution of problem (1.3). By [7, Theorems 5.1 and 5.2 in Chapter 2],

$$\begin{aligned}
& \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_r^{\beta, \gamma}(E)} \\
& \leq M \left[\max_{1 \leq i \leq N} \left\| \tau^{-1}(I - R)R^{i-1}(u_0 - A^{-1}\varphi_1) \right\|_E \right. \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - R)(R^{i+r-1} - R^{i-1})(u_0 - A^{-1}\varphi_1) \right\|_E \\
& \quad \left. + \frac{M}{\beta(1-\beta)} \|\varphi^\tau\|_{C_r^{\beta, \gamma}(E)} \right]
\end{aligned} \tag{2.12}$$

for the solution of the first order of accuracy implicit difference scheme for the approximate solutions of Cauchy problem (2.4). The proof of estimate (2.1) for difference scheme (1.3) is based on the estimate (2.12) and the following estimate:

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1}(I - R)R^{i-1}(u_0 - A^{-1}\varphi_1) \right\|_E \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - R)(R^{i+r-1} - R^{i-1})(u_0 - A^{-1}\varphi_1) \right\|_E \\
& \leq M_1 \left[\|\mu + A^{-1}(\varphi_{[\lambda/\tau]} - \varphi_1)\|_1^{\beta, \gamma} + \frac{M}{\beta(1-\beta)} \|\varphi^\tau\|_{C_r^{\beta, \gamma}(E)} \right]
\end{aligned} \tag{2.13}$$

for the solution of problem (1.3). Using formula (2.11) and estimate (2.7), we obtain

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1}(I - R)R^{i-1}(u_0 - A^{-1}\varphi_1) \right\|_E \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - R)(R^{i+r-1} - R^{i-1})(u_0 - A^{-1}\varphi_1) \right\|_E \\
& \leq M(\lambda, \delta) \left[\max_{1 \leq i \leq N} \left\| \tau^{-1}(I - R)R^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1}(\varphi_j - \varphi_{[\lambda/\tau]})\tau \right. \right. \right. \\
& \quad \left. \left. \left. + \mu - (I - R^{[\lambda/\tau]})A^{-1}(\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \right. \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - R)(R^{i+r-1} - R^{i-1}) \right. \\
& \quad \left. \left. \times \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1}(\varphi_j - \varphi_{[\lambda/\tau]})\tau + \mu - (I - R^{[\lambda/\tau]})A^{-1}(\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \right].
\end{aligned} \tag{2.14}$$

6 Mathematical Problems in Engineering

By [7, Theorem 5.2 in Chapter 2],

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^y \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \\
& \leq \sup_{0 < \tau \leq \tau_0} \left[\max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \right. \\
& \quad \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^y \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \right] \\
& = \left\| \mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]}) \right\|_1^{\beta, y}.
\end{aligned} \tag{2.15}$$

Similar to estimate (2.12) and using estimates (2.6), we can show that

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + R^{[\lambda/\tau]} A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^y \\
& \times \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + R^{[\lambda/\tau]} A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \\
& \leq \frac{M}{\beta(1-\beta)} \|\varphi^\tau\|_{C_\tau^{\beta, y}(E)}.
\end{aligned} \tag{2.16}$$

From this estimate and (2.15), estimate (2.13) follows. Now let us consider the difference scheme (1.4). In a similar manner with the difference scheme (1.3), we can obtain the formula

$$\begin{aligned}
u_k &= D^k u_0 + \sum_{j=1}^k \left(I + \frac{\tau}{2} A \right) D^{k-j+1} \varphi_j \tau, \quad k = 1, \dots, N, \\
u_0 &= T_\tau \left\{ \left(I - \left(\lambda - \left[\frac{\lambda}{\tau} \right] \tau \right) A \right) \sum_{j=1}^{[\lambda/\tau]-1} \left(I + \frac{\tau}{2} A \right) D^{[\lambda/\tau]-j+1} \varphi_j \tau \right. \\
& \quad \left. + \mu + D \left(\left(1 + \frac{\lambda}{\tau} - \left[\frac{\lambda}{\tau} \right] \right) I + \frac{\tau}{2} A \right) \tau \varphi_{[\lambda/\tau]} \right\}
\end{aligned} \tag{2.17}$$

for the solution of problem (1.4). Here

$$T_\tau = \left(I - \left(I - \left(\lambda - \left[\frac{\lambda}{\tau} \right] \tau \right) A \right) D^{[\lambda/\tau]} \right)^{-1}, \quad D = \left(I + \tau A + \frac{(\tau A)^2}{2} \right)^{-1}. \tag{2.18}$$

By [7, Theorem 5.1 in Chapter 3],

$$\begin{aligned}
& \left\| \{ \tau^{-1}(u_k - u_{k-1}) \}_1^N \right\|_{C_r^{\beta, \gamma}(E)} \\
& \leq M \left[\max_{1 \leq i \leq N} \left\| A \left(I + \frac{\tau}{2} A \right) D^i (u_0 - A^{-1} \varphi_1) \right\|_E \right. \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| A \left(I + \frac{\tau}{2} A \right) (D^{i+r} - D^i) (u_0 - A^{-1} \varphi_1) \right\|_E \\
& \quad \left. + \frac{M}{\beta(1-\beta)} \|\varphi^\tau\|_{C_r^{\beta, \gamma}(E)} \right]
\end{aligned} \tag{2.19}$$

for the solution of the second order of accuracy implicit difference scheme for the approximate solutions of Cauchy problem (2.4). The proof of estimate (2.1) for difference scheme (1.4) is based on the estimate (2.19) and the following estimate:

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) D^{i-1} (u_0 - A^{-1} \varphi_1) \right\|_E \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) (u_0 - A^{-1} \varphi_1) \right\|_E \\
& \leq M_1 \left[\left\| \mu + A^{-1} (\varphi_{[\lambda/\tau]} - \varphi_1) \right\|_1^{\beta, \gamma} + \frac{M}{\beta(1-\beta)} \|\varphi^\tau\|_{C_r^{\beta, \gamma}(E)} \right]
\end{aligned} \tag{2.20}$$

for the solution of problem (1.4). We have that

$$\|T_\tau\|_{E \rightarrow E} \leq M(\lambda, \delta). \tag{2.21}$$

Actually, we can write

$$\begin{aligned}
T_\tau - (I - \exp\{-\lambda A\})^{-1} &= T_\tau (I - \exp\{-\lambda A\})^{-1} \\
&\times \left(\left(I - \left(\lambda - \left\lfloor \frac{\lambda}{\tau} \right\rfloor \tau \right) A \right) D^{[\lambda/\tau]} - \exp\{-\lambda A\} \right).
\end{aligned} \tag{2.22}$$

Then, using the triangle inequality and estimates (2.9),

$$\left\| \left(\lambda - \left\lfloor \frac{\lambda}{\tau} \right\rfloor \tau \right) A D^{[\lambda/\tau]} \right\|_{E \rightarrow E} \leq M(\lambda, \delta) \tau, \tag{2.23}$$

$$\|D^{[\lambda/\tau]} - \exp\{-\lambda A\}\|_{E \rightarrow E} \leq M(\lambda, \delta) \tau, \tag{2.24}$$

we obtain estimate (2.7). Estimate (2.23) follows from

$$\|D^k\|_{E \rightarrow E} \leq M, \quad \|k\tau A D^k\|_{E \rightarrow E} \leq M, \quad k \geq 1. \tag{2.25}$$

The proof of (2.24) is based on estimates (1.8) and (2.25) and it was proved in [34].

8 Mathematical Problems in Engineering

Using formula (2.17) and estimate (2.21), we obtain

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1}(I - D)D^{i-1}(u_0 - A^{-1}\varphi_1) \right\|_E \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - D)(D^{i+r-1} - D^{i-1})(u_0 - A^{-1}\varphi_1) \right\|_E \\
& \leq M(\lambda) \left[\max_{1 \leq i \leq N} \left\| \tau^{-1}(I - D)D^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1}(\varphi_j - \varphi_{[\lambda/\tau]})\tau \right. \right. \right. \\
& \quad \left. \left. \left. + \mu - (I - D^{[\lambda/\tau]})A^{-1}(\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \right. \\
& \quad \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - D)(D^{i+r-1} - D^{i-1}) \right. \right. \\
& \quad \left. \left. \times \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1}(\varphi_j - \varphi_{[\lambda/\tau]})\tau + \mu - (I - D^{[\lambda/\tau]})A^{-1}(\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \right]. \tag{2.26}
\end{aligned}$$

By [7, Theorem 5.2 in Chapter 3],

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1}(I - D)D^{i-1}(\mu + A^{-1}(-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - D)(D^{i+r-1} - D^{i-1})(\mu + A^{-1}(-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \\
& \leq \sup_{0 < \tau \leq \tau_0} \left[\max_{1 \leq i \leq N} \left\| \tau^{-1}(I - D)D^{i-1}(\mu + A^{-1}(-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \right. \\
& \quad \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \left\| \tau^{-1}(I - D)(D^{i+r-1} - D^{i-1})(\mu + A^{-1}(-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \right] \\
& = \left\| \mu + A^{-1}(-\varphi_1 + \varphi_{[\lambda/\tau]}) \right\|_1^{\beta, \gamma}. \tag{2.27}
\end{aligned}$$

Similar to estimate (2.19) and using estimates (2.25), we can show that

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1}(I - D)D^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1}(\varphi_j - \varphi_{[\lambda/\tau]})\tau + D^{[\lambda/\tau]}A^{-1}(\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta}((i+r)\tau)^\gamma \\
& \times \left\| \tau^{-1}(I - D)(D^{i+r-1} - D^{i-1}) \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1}(\varphi_j - \varphi_{[\lambda/\tau]})\tau \right. \right. \\
& \quad \left. \left. + D^{[\lambda/\tau]}A^{-1}(\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_E \leq \frac{M}{\beta(1 - \beta)} \|\varphi^\tau\|_{C_\tau^{\beta, \gamma}(E)}. \tag{2.28}
\end{aligned}$$

From these estimates, estimate (2.20) follows. $\square$

Remark 2.2. The parameter γ can be chosen freely in $[0, \beta)$, which increases the number of spaces $C_\tau^{\beta, \gamma}(E)$ ($0 \leq \gamma \leq \beta$, $0 < \beta < 1$) of grid functions in which difference schemes (1.3) and (1.4) are well-posed.

THEOREM 2.3. *Let τ be a sufficiently small number. Then the solutions of the difference schemes (1.3) and (1.4) in $C_\tau^{\beta, \gamma}(E_{\alpha-\beta})$ ($0 \leq \gamma \leq \beta \leq \alpha < 1$) satisfy the following coercivity inequalities:*

$$\begin{aligned} & \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \\ & \leq \frac{M_1}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} + M_1 |\mu + A^{-1}(\varphi_{[\lambda\tau]} - \varphi_1)|_{1+\alpha-\beta}^{\beta, \gamma}, \quad E_{\alpha-\beta}^{\beta, \gamma} = E^{\beta, \gamma}(E_{\alpha-\beta}), \end{aligned} \quad (2.29)$$

where M_1 is independent not only of φ^τ , φ , α , β , γ , but also of τ .

Proof. Let us prove (2.29) for difference scheme (1.3). By [7, Theorem 5.3 in Chapter 2],

$$\begin{aligned} & \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \\ & \leq M \left[\max_{1 \leq i \leq N} \|\tau^{-1}(I - R)R^{i-1}(u_0 - A^{-1}\varphi_1)\|_{E_{\alpha-\beta}} \right. \\ & \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \|\tau^{-1}(I - R)(R^{i+r-1} - R^{i-1})(u_0 - A^{-1}\varphi_1)\|_{E_{\alpha-\beta}} \\ & \quad \left. + \frac{M}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \right] \end{aligned} \quad (2.30)$$

for the solution of the first order of accuracy implicit difference scheme for the approximate solutions of Cauchy problem (2.4). The proof of estimate (2.29) for difference scheme (1.3) is based on estimate (2.30) and the following estimate:

$$\begin{aligned} & \max_{1 \leq i \leq N} \|\tau^{-1}(I - R)R^{i-1}(u_0 - A^{-1}\varphi_1)\|_{E_{\alpha-\beta}} \\ & \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \|\tau^{-1}(I - R)(R^{i+r-1} - R^{i-1})(u_0 - A^{-1}\varphi_1)\|_{E_{\alpha-\beta}} \\ & \leq M_1 \left[|\mu + A^{-1}(\varphi_{[\lambda\tau]} - \varphi_1)|_{1+\alpha-\beta}^{\beta, \gamma} + \frac{M}{\alpha(1-\alpha)} \|\varphi^\tau\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \right] \end{aligned} \quad (2.31)$$

for the solution of problem (1.3). Using formula (2.11) and estimate (2.7), we obtain

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& \leq M(\lambda) \left[\max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau \right. \right. \right. \\
& \quad \left. \left. \left. + \mu - (I - R^{[\lambda/\tau]}) A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \right. \\
& \quad \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) \right. \right. \\
& \quad \left. \left. \times \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + \mu - (I - R^{[\lambda/\tau]}) A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \right]. \tag{2.32}
\end{aligned}$$

By [7, Theorem 5.2 in Chapter 2],

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \\
& \leq \sup_{0 < \tau \leq \tau_0} \left[\max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \right. \\
& \quad \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \right] \\
& = \left| \mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]}) \right|_{1+\alpha-\beta}^{\beta, \gamma}. \tag{2.33}
\end{aligned}$$

Similar to estimate (2.30) and using estimates (2.6), we can show that

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - R) R^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + R^{[\lambda/\tau]} A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - R) (R^{i+r-1} - R^{i-1}) \right. \\
& \quad \left. \times \left(\sum_{j=1}^{[\lambda/\tau]} R^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + R^{[\lambda/\tau]} A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \\
& \leq \frac{M}{\alpha(1-\alpha)} \left\| \varphi^\tau \right\|_{C_r^{\beta, \gamma}(E_{\alpha-\beta})}. \tag{2.34}
\end{aligned}$$

From this estimate and (2.33), estimate (2.31) follows. Now let us consider the difference scheme (1.4). By [7, Theorem 5.3 in Chapter 3],

$$\begin{aligned}
& \left\| \{ \tau^{-1}(u_k - u_{k-1}) \}_1^N \right\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \\
& \leq M \left[\max_{1 \leq i \leq N} \left\| A \left(I + \frac{\tau}{2} A \right) D^i (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \right. \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| A \left(I + \frac{\tau}{2} A \right) (D^{i+r} - D^i) (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& \quad \left. + \frac{M}{\alpha(1-\alpha)} \left\| \varphi^\tau \right\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \right]
\end{aligned} \tag{2.35}$$

for the solution of the second order of accuracy implicit difference scheme for the approximate solutions of Cauchy problem (2.4). The proof of estimate (2.29) for difference scheme (1.4) is based on estimate (2.35) and the following estimate:

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) D^{i-1} (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& \leq M_1 \left[\left\| \mu + A^{-1} (\varphi_{[\lambda/\tau]} - \varphi_1) \right\|_{1+\alpha-\beta}^{\beta, \gamma} + \frac{M}{\beta(1-\beta)} \left\| \varphi^\tau \right\|_{C_\tau^{\beta, \gamma}(E_{\alpha-\beta})} \right]
\end{aligned} \tag{2.36}$$

for the solution of problem (1.4). Using formula (2.17) and estimate (2.21), we obtain

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) D^{i-1} (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) (u_0 - A^{-1} \varphi_1) \right\|_{E_{\alpha-\beta}} \\
& \leq M(\lambda) \left[\max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) D^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau \right. \right. \right. \\
& \quad \left. \left. + \mu - (I - D^{[\lambda/\tau]}) A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \\
& \quad + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) \right. \\
& \quad \left. \times \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + \mu - (I - D^{[\lambda/\tau]}) A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \right].
\end{aligned} \tag{2.37}$$

By [7, Theorem 5.3 in Chapter 3],

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) d^{i-1} (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_E \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \\
& \leq \sup_{0 < \tau \leq \tau_0} \left[\max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) D^{i-1} (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \right. \\
& \quad \left. + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) (\mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]})) \right\|_{E_{\alpha-\beta}} \right] \\
& = \left\| \mu + A^{-1} (-\varphi_1 + \varphi_{[\lambda/\tau]}) \right\|_{1+\alpha-\beta}^{\beta, \gamma}.
\end{aligned} \tag{2.38}$$

Similar to estimate (2.35) and using estimates (2.25), we can show that

$$\begin{aligned}
& \max_{1 \leq i \leq N} \left\| \tau^{-1} (I - D) D^{i-1} \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + D^{[\lambda/\tau]} A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \\
& + \sup_{1 \leq i < i+r \leq N} (r\tau)^{-\beta} ((i+r)\tau)^\gamma \left\| \tau^{-1} (I - D) (D^{i+r-1} - D^{i-1}) \right. \\
& \quad \left. \times \left(\sum_{j=1}^{[\lambda/\tau]} D^{[\lambda/\tau]-j+1} (\varphi_j - \varphi_{[\lambda/\tau]}) \tau + D^{[\lambda/\tau]} A^{-1} (\varphi_1 - \varphi_{[\lambda/\tau]}) \right) \right\|_{E_{\alpha-\beta}} \\
& \leq \frac{M}{\alpha(1-\alpha)} \left\| \varphi^\tau \right\|_{C_r^{\beta, \gamma}(E_{\alpha-\beta})}.
\end{aligned} \tag{2.39}$$

From these estimates, estimate (2.36) follows. $\square$

Remark 2.4. The spaces $C_r^{\beta, \gamma}(E_{\alpha-\beta})$ of grid functions, in which coercive solvability has been established, depend on the parameters α , β , and γ . However, the constants in the coercive inequalities depend only on α . Hence, we can choose the parameters β and γ freely, which increases the number of spaces of grid functions in which difference schemes (1.3) and (1.4) are well-posed.

Remark 2.5. Using the coercive stability estimates of Theorems 2.1–2.3 and by passing to the limit for $\tau \rightarrow 0$, one can recover theorems on coercive solvability of the nonlocal-boundary value problem (1.1) [6].

3. Applications

We consider the boundary value problem on the range $\{0 \leq t \leq 1, x \in \mathbb{R}^n\}$ for $2m$ -order multidimensional parabolic equation

$$\begin{aligned} \frac{\partial v(t, x)}{\partial t} + \sum_{|\tau|=2m} a_\tau(x) \frac{\partial^{|\tau|} v(t, x)}{\partial x_1^{\tau_1} \cdots \partial x_n^{\tau_n}} + \delta v(t, x) &= f(t, x), \quad 0 \leq t \leq 1, \\ v(0, x) &= v(\lambda, x) + \mu(x), \quad 0 < \lambda \leq 1, x \in \mathbb{R}^n, \quad |\tau| = \tau_1 + \cdots + \tau_n, \end{aligned} \quad (3.1)$$

where $a_r(x)$, $f(t, x)$ and $\mu(x)$ are given sufficiently smooth functions and $\delta > 0$ is a sufficiently large positive constant.

Now, the abstract theorems given above are applied in the investigation of difference schemes for approximate solution of (3.1). The discretization of problem (3.1) is carried out in two steps. Let us define the grid space $\mathbb{R}_h^n$ ($0 < h \leq h_0$) as the set of all points of the Euclidean space $\mathbb{R}^n$ whose coordinates are given by

$$x_k = s_k h, \quad s_k = 0, \pm 1, \pm 2, \dots, \quad k = 1, \dots, n. \quad (3.2)$$

In the first step, let us give the difference operator A_h^x by the formula

$$A_h^x u^h = \sum_{2m \leq |r| \leq S} b_r^x D_h^r u^h + \delta u^h. \quad (3.3)$$

The coefficients are chosen in such a way that the operator A_h^x approximates in a specified way the operator

$$\sum_{|r|=2m} a_r(x) \frac{\partial^{|\tau|}}{\partial x_1^{\tau_1} \cdots \partial x_n^{\tau_n}} + \delta. \quad (3.4)$$

We will assume that for $|\xi_k h| \leq \pi$ and fixed x , the symbol $A^x(\xi h, h)$ of the operator $A_h^x - \delta$ satisfies the inequalities

$$(-1)^m A^x(\xi h, h) \geq M_1 |\xi|^{2m}, \quad |\arg A^x(\xi h, h)| \leq \phi < \phi_0 \leq \frac{\pi}{2}. \quad (3.5)$$

With the help of A_h^x , we arrive at the nonlocal boundary value problem

$$\begin{aligned} \frac{dv^h(t, x)}{dt} + A_h^x v^h(t, x) &= f^h(t, x), \quad 0 \leq t \leq 1, \\ v^h(0, x) &= v^h(\lambda, x) + \mu^h(x), \quad x \in \mathbb{R}_h^n, \end{aligned} \quad (3.6)$$

for an infinite system of ordinary differential equations.

In the second step, we replace problem (3.6) by the difference schemes

$$\begin{aligned}
\frac{u_k^h(x) - u_{k-1}^h(x)}{\tau} + A_h^x u_k^h(x) &= \varphi_k^h(x), \quad \varphi_k^h(x) = f^h(t_k, x), \quad t_k = k\tau, \\
1 \leq k \leq N, \quad u_0^h(x) &= u_{[\lambda/\tau]}^h(x) + \mu^h(x), \quad x \in \mathbb{R}_h^n, \\
\frac{u_k^h(x) - u_{k-1}^h(x)}{\tau} + A_h^x \left(I + \frac{\tau A_h^x}{2} \right) u_k^h(x) &= \left(I + \frac{\tau A_h^x}{2} \right) \varphi_k^h(x), \\
\varphi_k^h(x) &= f^h\left(t_k - \frac{\tau}{2}, x\right), \quad t_k = k\tau, \quad 1 \leq k \leq N, \\
u_0^h(x) &= \left(I - \left(\lambda - \left\lfloor \frac{\lambda}{\tau} \right\rfloor \tau \right) A_h^x \right) u_{[\lambda/\tau]}^h(x) + \left(\lambda - \left\lfloor \frac{\lambda}{\tau} \right\rfloor \tau \right) \varphi_{[\lambda/\tau]}^h(x), \quad x \in \mathbb{R}_h^n.
\end{aligned} \tag{3.7}$$

Let us give a number of corollaries of the abstract theorems given above. To formulate our result, we need to introduce the spaces $C_h = C(\mathbb{R}_h^n)$ and $C_h^\beta = C^\beta(\mathbb{R}_h^n)$ of all bounded grid functions $u^h(x)$ defined on $\mathbb{R}_h^n$, equipped with the norms

$$\begin{aligned}
\|u^h\|_{C_h} &= \sup_{x \in \mathbb{R}_h^n} |u^h(x)|, \\
\|u^h\|_{C_h^\beta} &= \sup_{x \in \mathbb{R}_h^n} |u^h(x)| + \sup_{x, y \in \mathbb{R}_h^n} \frac{|u^h(x) - u^h(x+y)|}{|y|^\beta}.
\end{aligned} \tag{3.8}$$

THEOREM 3.1. *The solutions of the difference schemes (3.7) satisfy the coercivity estimates:*

$$\begin{aligned}
&\left\| \left\{ \tau^{-1} (u_k^h - u_{k-1}^h) \right\}_1^{N-1} \right\|_{C_\tau^{\beta, \gamma}(C_h^\gamma)} \\
&\leq M(\alpha, \nu) \left[\sum_{|r|=2m} \|D_h^r u^h\|_{C_h^{\gamma+2m\alpha}} + \|\varphi^{\tau, h}\|_{C_\tau^{\beta, \gamma}(C_h^\gamma)} \right], \quad 0 < 2m\alpha + \nu < 1, \quad \nu > 0,
\end{aligned} \tag{3.9}$$

where $M(\alpha, \nu)$ does not depend on $\varphi^{\tau, h}$, μ^h , h , and τ .

The proof of this theorem is based on the abstract theorems (Theorems 2.1 and 2.3) and the positivity of the operator A_h^x in C_h [35] and on the coercivity inequality for an elliptic operator A_h^x in C_h^β [7] and on the following theorem.

THEOREM 3.2 (see [7]). *For any $0 < \beta < 1/2m$, the norms in the spaces $E_\beta(C_h, A_h^x)$ and $C_h^{2m\beta}$ are equivalent uniformly in h .*

Acknowledgment

The author would like to thank Professor Pavel Sobolevskii (Jerusalem, Israel) for his helpful suggestions for the improvement of this paper.

References

- [1] M. Dehghan, "On the numerical solution of the diffusion equation with a nonlocal boundary condition," *Mathematical Problems in Engineering*, vol. 2003, no. 2, pp. 81–92, 2003.
- [2] J. R. Cannon, S. Pérez Esteva, and J. van der Hoek, "A Galerkin procedure for the diffusion equation subject to the specification of mass," *SIAM Journal on Numerical Analysis*, vol. 24, no. 3, pp. 499–515, 1987.
- [3] N. Gordeziani, P. Natalini, and P. E. Ricci, "Finite-difference methods for solution of nonlocal boundary value problems," *Computers & Mathematics with Applications*, vol. 50, no. 8-9, pp. 1333–1344, 2005.
- [4] R. Dautray and J.-L. Lions, *Analyse mathématique et calcul numérique pour les sciences et les techniques. Volume 1—11*, Masson, Paris, France, 1988.
- [5] A. Ashyralyev and Y. Ozdemir, "Stability of difference schemes for hyperbolic-parabolic equations," *Computers & Mathematics with Applications*, vol. 50, no. 8-9, pp. 1443–1476, 2005.
- [6] A. Ashyralyev, A. Hanalyev, and P. E. Sobolevskii, "Coercive solvability of the nonlocal boundary value problem for parabolic differential equations," *Abstract and Applied Analysis*, vol. 6, no. 1, pp. 53–61, 2001.
- [7] A. Ashyralyev and P. E. Sobolevskii, *Well-Posedness of Parabolic Difference Equations*, vol. 69 of *Operator Theory: Advances and Applications*, Birkhäuser, Basel, Switzerland, 1994.
- [8] R. Rautmann, " $H^{2,r}$ -convergent approximation schemes to the Navier-Stokes equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 30, no. 4, pp. 1915–1926, 1997.
- [9] A. Ashyralyev and I. Karatay, "On the second order of accuracy difference schemes of the nonlocal boundary value problem for parabolic equations," *Functional Differential Equations*, vol. 10, no. 1-2, pp. 45–63, 2003.
- [10] A. Ashyralyev, I. Karatay, and P. E. Sobolevskii, "On well-posedness of the nonlocal boundary value problem for parabolic difference equations," *Discrete Dynamics in Nature and Society*, vol. 2004, no. 2, pp. 273–286, 2004.
- [11] A. Ashyralyev and P. E. Sobolevskii, "The theory of interpolation of linear operators and the stability of difference schemes," *Doklady Akademii Nauk SSSR*, vol. 275, no. 6, pp. 1289–1291, 1984 (Russian).
- [12] A. Ashyralyev, "Nonlocal boundary-value problems for abstract parabolic equations: well-posedness in Bochner spaces," *Journal of Evolution Equations*, vol. 6, no. 1, pp. 1–28, 2006.
- [13] A. Ashyralyev and P. E. Sobolevskii, "Difference schemes of a high order of accuracy for parabolic equations with variable coefficients," *Doklady Akademii Nauk Ukrainsoi SSR. Seriya A. Fiziko-Matematicheskie i Tekhnicheskie Nauki*, no. 6, pp. 3–7, 1988 (Russian).
- [14] A. Ashyralyev and P. E. Sobolevskii, "Well-posed solvability of the Cauchy problem for difference equations of parabolic type," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 24, no. 2, pp. 257–264, 1995.
- [15] A. Ashyralyev and P. E. Sobolevskii, *New Difference Schemes for Partial Differential Equations*, vol. 148 of *Operator Theory: Advances and Applications*, Birkhäuser, Basel, Switzerland, 2004.
- [16] D. Guidetti, B. Karasözen, and S. Piskarev, "Approximation of abstract differential equations," *Journal of Mathematical Sciences*, vol. 122, no. 2, pp. 3013–3054, 2004.
- [17] A. Ashyralyev, S. Piskarev, and L. Weis, "On well-posedness of difference schemes for abstract parabolic equations in $L^p([0, T]; E)$ spaces," *Numerical Functional Analysis and Optimization*, vol. 23, no. 7-8, pp. 669–693, 2002.
- [18] M. Crouzeix, S. Larsson, S. Piskarëv, and V. Thomée, "The stability of rational approximations of analytic semigroups," *BIT. Numerical Mathematics*, vol. 33, no. 1, pp. 74–84, 1993.
- [19] I. P. Gavrilyuk and V. L. Makarov, "Exponentially convergent parallel discretization methods for the first order evolution equations," *Computational Methods in Applied Mathematics*, vol. 1, no. 4, pp. 333–355, 2001.

- [20] I. P. Gavrilyuk and V. L. Makarov, "Algorithms without accuracy saturation for evolution equations in Hilbert and Banach spaces," *Mathematics of Computation*, vol. 74, no. 250, pp. 555–583, 2005.
- [21] I. P. Gavrilyuk and V. L. Makarov, "Exponentially convergent algorithms for the operator exponential with applications to inhomogeneous problems in Banach spaces," *SIAM Journal on Numerical Analysis*, vol. 43, no. 5, pp. 2144–2171, 2005.
- [22] D. Gordeziani, H. Meladze, and G. Avalishvili, "On one class of nonlocal in time problems for first-order evolution equations," *Zhurnal Obchyslyuval'noi ta Prykladnoi Matematyky*, vol. 88, no. 1, pp. 66–78, 2003.
- [23] D. G. Gordeziani and G. A. Avalishvili, "Time-nonlocal problems for Schrödinger-type equations—I: problems in abstract spaces," *Differential Equations*, vol. 41, no. 5, pp. 703–711, 2005.
- [24] R. P. Agarwal, M. Bohner, and V. B. Shakhmurov, "Maximal regular boundary value problems in Banach-valued weighted space," *Boundary Value Problems*, no. 1, pp. 9–42, 2005.
- [25] V. B. Shakhmurov, "Coercive boundary value problems for regular degenerate differential-operator equations," *Journal of Mathematical Analysis and Applications*, vol. 292, no. 2, pp. 605–620, 2004.
- [26] J. I. Ramos, "Linearly-implicit, approximate factorization, exponential methods for multi-dimensional reaction-diffusion equations," *Applied Mathematics and Computation*, vol. 174, no. 2, pp. 1609–1633, 2006.
- [27] X.-Z. Liu, X. Cui, and J.-G. Sun, "FDM for multi-dimensional nonlinear coupled system of parabolic and hyperbolic equations," *Journal of Computational and Applied Mathematics*, vol. 186, no. 2, pp. 432–449, 2006.
- [28] A. V. Gulín and V. A. Morozova, "On the stability of a nonlocal difference boundary value problem," *Differential Equations*, vol. 39, no. 7, pp. 962–967, 2003 (Russian).
- [29] A. V. Gulín, N. I. Ionkin, and V. A. Morozova, "On the stability of a nonlocal two-dimensional difference problem," *Differential Equations*, vol. 37, no. 7, pp. 970–978, 2001 (Russian).
- [30] A. Lunardi, *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, vol. 16 of *Progress in Nonlinear Differential Equations and Their Applications*, Birkhäuser, Basel, Switzerland, 1995.
- [31] Y. G. Wang and M. Oberguggenberger, "Nonlinear parabolic equations with regularized derivatives," *Journal of Mathematical Analysis and Applications*, vol. 233, no. 2, pp. 644–658, 1999.
- [32] W.-J. Beyn and B. M. Garay, "Estimates of variable stepsize Runge-Kutta methods for sectorial evolution equations with nonsmooth data," *Applied Numerical Mathematics*, vol. 41, no. 3, pp. 369–400, 2002.
- [33] A. Ashyralyev, H. Akca, and L. Bizevski, "On a semilinear evolution nonlocal Cauchy problem," in *Some Problems of Applied Mathematics*, pp. 29–44, Fatih University, Istanbul, Turkey, 2000.
- [34] A. Ashyralyev, "An error estimation of the solution of the purely implicit difference schemes for parabolic equations with nonsmooth data," in *Applied Mathematics and Computer Software*, pp. 26–28, Moscow, Russia, 1985.
- [35] Yu. A. Smirnitskii and P. E. Sobolevskii, "Positivity of multidimensional difference operators in the C-norm," *Uspekhi Matematicheskikh Nauk*, vol. 36, no. 4, pp. 202–203, 1981 (Russian).

A. Ashyralyev: Department of Mathematics, Faculty of Arts and Science, Fatih University,
34900 Istanbul, Turkey
Email address: aashyr@fatih.edu.tr