Research Article

Dissipation Effects in the Ratchetlike Fermi Acceleration

Cesar Manchein and Marcus W. Beims

Departamento de Física, Universidade Federal do Paraná, 81531-980 Curitiba, PR, Brazil

Correspondence should be addressed to Cesar Manchein, cmanchein@fisica.ufpr.br

Received 23 April 2009; Accepted 29 June 2009

Recommended by Edson Denis Leonel

Ac-driven asymmetric pulses can be used to control the Fermi acceleration between three different motions, **A**: the *accelerated* mode, **D**: the *decelerated* mode, and **H**: the *hyperaccelerated* mode. In this paper we show that dissipation strongly affects the particles velocity, reducing the possibility for an accurate control of the dynamics. The saturation time, where the mean velocity starts to be constant due to dissipation, decays with a power law $\sim \gamma^{-\beta}$, where γ is the dissipation parameter and β is close to 1. The value of the saturated mean velocity also decays with a power law with exponent $\beta \sim 0.6$ for case **H**, and $\beta \sim 0.3$ for case **A**. In case **D** this velocity is almost constant for small dissipations.

Copyright © 2009 C. Manchein and M. W. Beims. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fermi acceleration is a topic which got attention in various areas of physics, ranging from nonlinear physics [1–8], atom optics [9–11], plasma physics [12, 13] to astrophysics [14–16]. After the first model proposed by Fermi [17], essentially two different versions became common in the literature. In the first one, the Fermi-Ulam (FU) model, a bouncing particle moves between a fixed surface and a parallel oscillating surface [18]. In this case the regular islands in the phase space prevent the Fermi acceleration. A simplified version of this model was proposed to improve simulations [1], called the static wall model. It essentially ignores the displacement of the moving wall but keeps the information for the momentum transfer as the wall was oscillating. The dynamics of the static model was studied in different aspects [1, 3–5] and for different models [6], and the relevant result for the purpose of the present work is that invariant curves in the phase space, found for higher velocities, prevent the particle to increase its kinetic energy without bounds. Recently the hopping wall approximation was proposed [7, 8] which takes into account the effect of the wall displacement and allows the analytical estimation of the particle mean velocity. Compared to the simplified static model, the particle acceleration is enhanced. The second kind of Fermi accelerated model was proposed in 1977 by Pustyl'nikov [19], who considered a particle on

a periodically oscillating horizontal surface in the presence of a gravitational field. Different from the FU model, for some initial conditions and control parameters, the particle energy can grow indefinitely.

In this paper we analyze the effect of dissipation in the simplified FU model when an ac-driven asymmetric pulse controls the Fermi acceleration (deceleration) [20]. Small dissipation is inevitable in real systems and its influence on the dynamics of conservative systems is of great interest [21–23] since elliptic periodic orbits become small sinks and attractors start to exist [24, 25]. In the context of Fermi models dissipation effects have been inserted in two ways: frictional force [26] and inelastic collisions at the walls [27–29]. Here we use the second approach and the pulse is a *deformed* sawtooth driving law for the moving wall. This Ratchetlike pulse differs from the ac-driven asymmetric pulses (*symmetric* sawtooth) used for the Fermi acceleration in the early work of Lieberman and Lichtenberg [1] and proposed recently to control the motion of magnetic flux quanta [30] and to analyse the relative efficiency of mechanism leading to increased acceleration in the hopping wall approximation [8]. In the simplified Fermi model [1] the particle is free to move between the elastic impacts with the walls. Consider that the moving wall oscillates between two extrema with amplitude v_0 . The gravitational force is considered zero. The system is described by a two-dimensional map $M_{1(2)}(V_n, \phi_n) = (V_{n+1}, \phi_{n+1})$ which gives, respectively, the velocity of the particle, and the phase of the moving wall, immediately after the particle suffers a collision with the wall. Considering dimensionless variables the dissipative FU map with the *deformed* sawtooth wall is written as

$$M_1 : \begin{cases} V_{n+1} = \left| (1 - \gamma)V_n + \frac{v_0}{\eta_1}(\phi_n - \eta_1) \right|, \\ \phi_{n+1} = \phi_n + \mu \frac{(\eta_1 + \eta_2)}{V_{n+1}} \mod(\eta_1 + \eta_2), \end{cases} \quad (1)$$

for $\phi_n < \eta_1$, and

$$M_2 : \begin{cases} V_{n+1} = \left| (1 - \gamma)V_n + \frac{v_0}{\eta_2}(\phi_n - \eta_1) \right|, \\ \phi_{n+1} = \phi_n + \mu \frac{(\eta_1 + \eta_2)}{V_{n+1}} \mod(\eta_1 + \eta_2), \end{cases} \quad (2)$$

for $\phi_n \geq \eta_1$, where n is the iteration number and μ is the maximum distance between the walls. For $\gamma \neq 0$ we have dissipation effects. Since in this simplified model the displacement of the moving wall is ignored, the modulus function is used to avoid errors due to successive collisions which may occur in the original model. In other words, if after a collision with the wall the particle continues to have a negative velocity (a successive collision will occur in the original model), the particle moves beyond the wall. The modulus for the velocity injects the particle back and fixes the problem.

The time asymmetry of the oscillating wall in (1) and (2) is controlled by varying the parameters (η_1, η_2) . The *deformed* sawtooth (Ratchetlike) is obtained when $\eta_1 \neq \eta_2$. Figures 1(a)–1(c) show the time behaviour of the oscillating wall (the pulse) for different values of the asymmetry $\Delta\eta = \eta_2 - \eta_1$. The deformed sawtooth pulse is obtained when $\Delta\eta \neq 0.0$. Such pulses can be easily obtained from pulse generators. For the case of no dissipation three different

motions were obtained and controlled [20]. They are **A**: for $\Delta\eta = 0$ (symmetric sawtooth) the *accelerated* case (see Figure 2(a)); **D**: for $\Delta\eta < 0$ (asymmetric sawtooth) the *decelerated* case (see Figure 3(a)); **H**: for $\Delta\eta > 0$ (asymmetric sawtooth) the *hyperaccelerated* case (see Figure 4(a)). This classification, different from [31], is based on how fast the average velocity grows or decreases. At next we discuss separately the effects of dissipation in each case. To do this we show the corresponding phase space dynamics and determine the mean particles velocity at a given time n from

$$\langle V \rangle(n) = \frac{1}{n+1} \sum_{i=0}^n \frac{1}{\xi} \sum_{j=1}^{\xi} V_{n,j}, \quad (3)$$

where the index i refers to the i th iteration of the sample j , and ξ is the number of initial conditions. We iterate the map (1) or (2) for times $n = 1 \times 10^8$ and 3000 initial conditions in the interval $0 < \phi \leq \eta_1 + \eta_2$ and $0 \leq V \leq 10^{-3}$.

First we consider the accelerated mode which is similar to the simplified model studied by Lichtenberg and Lieberman [1, 2], where the harmonic force was considered. This case is shown in Figure 2. As the particle velocity increases, regular islands are observed and $\langle V \rangle$ increases slowly until ~ 10 (see Figures 2(a) and 2(c)). The regular islands prevent the particle velocity to increase very fast. We mention that all initial conditions start inside the chaotic region at low velocities. The growth rate of $\langle V \rangle$ depends on the number of regular islands inside the phase space. For very small dissipations ($\gamma = 10^{-6}$) the regular islands are transformed into sinks, as can be observed by comparing Figures 2(a) and 2(b) (observe the dark regions), and attract the chaotic trajectories which pass nearby. As a consequence the mean velocity cannot increase as before (compare Figures 2(c) and 2(d)). This effect increases for higher values of the dissipation parameter, as can be observed in Figures 2(e)–2(f). Therefore in case **A** the dissipation decreases the mean velocity (to a constant plateau) for any values of γ when compared to the nondissipative case. There are two additional information we can get from Figures 2(d)–2(f), the saturation time (t_s), where the mean velocity starts to be constant in time, and the saturated mean velocity (v_s). Plotting both quantities as a function of γ in the interval $10^{-6} \leq \gamma \leq 10^{-1}$ we obtain the decay $t_c \sim \gamma^{-1.0}$ and $v_c \sim \gamma^{-0.3}$.

The second case to discuss is the dissipation in the *hyperaccelerated* case **H** (see Figure 3(a)). For a very small asymmetry $\Delta\eta = 0.01$ the phase space is totally filled and no regular islands are observed (see Figure 3(a)). As shown in [20], the Lyapunov exponents approach zero for this case. The corresponding $\langle V \rangle$ increases very fast until $\langle V \rangle \sim 3 \times 10^4$ (see Figure 3(c)), showing that the accelerated mode is enhanced when compared to case **A**. By adding small dissipation ($\gamma = 10^{-6}$) into the system we observe in Figure 3(b) that again many sinks appear. In this case however, since no regular islands were observed for $\gamma = 0$, we are not able to relate the sinks to regular islands. On the other hand, we can say that the dissipation is able to find attracting points which were not visible in the conservative limit. Also here small dissipation prevents the hyperacceleration of particles. The mean particles velocity also decreases (to a constant value) due to the small sinks which appear in the phase space shown in Figure 3(b). For higher values of the dissipation parameter the maximal mean velocity decreases. Observe Figures 3(d)–3(f) for $\gamma = 0.0, 10^{-6}, 10^{-5}, 10^{-4}, 10^{-1}$. For the hyperaccelerated case we found that the saturated quantities decay like $t_c \sim \gamma^{-1.1}$ and $v_c \sim \gamma^{-0.6}$.

Figure 1: (a)–(c) The shape of the pulses used in the simulations. The deformed sawtooth effect is obtained when $\Delta\eta = \eta_2 - \eta_1 \neq 0.0$. For (a) $\Delta\eta = 0$ we have the symmetric *accelerated* case **A**, (b) $\Delta\eta < 0$ we have the deformed *decelerated* case **D**, and (c) $\Delta\eta > 0$ the deformed *hyperaccelerated* case **H**.

Now we discuss the decelerated case **D** ($\Delta\eta = -0.01$). Here many regular islands appear in phase space (see Figure 4(a)) which again prevent the acceleration to increase without bounds, as in case **A**. The regular islands are very large so that the maximal instantaneous velocity is around ~ 2.5 , while for case **A** it was ~ 70.0 . The corresponding $\langle V \rangle$ remains here almost constant (see Figure 4(c)). As in the other cases, when small dissipation is added ($\gamma = 10^{-6}$) into the system, regular islands are transformed into sinks. However here the mean velocity *increases* (see Figure 4(d)). This increasing in the mean velocity is easy to explain. Since the first (from below) four islands from the conservative system are very large (see Figure 4(a)), almost all initial conditions cannot reach velocities higher than ~ 1.1 . It is a kind of upper partial barrier for the velocities. However, when small dissipation is added, these four islands are transformed into sinks so that the chaotic trajectories can penetrate the islands and the upper barrier does not exist anymore. This makes the mean velocity to increase because the positions of the sinks are higher (in velocity) than those of the upper

Figure 2: Evolution of 500 chaotic orbits on the phase space $V \times \phi$ for the parameters $\mu = 10$, $v_0 = 0.2$ for case **A** ($\Delta\eta = 0.0$, $\eta_1 = \eta_2 = 1.00$). For (a) $\gamma = 0.0$ and (b) $\gamma = 10^{-6}$. The mean values of the velocity, calculated over 3000 trajectories, are shown in (c)–(f) and are related to the cases $\gamma = 0.0, 10^{-6}, 10^{-5}, 10^{-4}, 10^{-1}$.

Figure 3: Evolution of 500 chaotic orbits on the phase space $V \times \phi$ for the parameters $\mu = 10$, $v_0 = 0.2$ for case **H** ($\Delta\eta = 0.01$, $\eta_1 = 1.00$, $\eta_2 = 1.01$). For (a) $\gamma = 0.0$ and (b) $\gamma = 10^{-6}$. The mean values of the velocity, calculated over 3000 trajectories are shown in (c)–(f) and are related to the cases $\gamma = 0.0, 10^{-6}, 10^{-5}, 10^{-4}, 10^{-1}$.

Figure 4: Evolution of 500 chaotic orbits on the phase space $V \times \phi$ for the parameters $\mu = 10$, $v_0 = 0.2$ for case **D** ($\Delta\eta = -0.01$, $\eta_1 = 1.01$, $\eta_2 = 1.00$). For (a) $\gamma = 0.0$ and (b) $\gamma = 10^{-6}$. The mean values of the velocity, calculated over 3000 trajectories, are shown in (c)–(f) and are related to the cases $\gamma = 0.0, 10^{-6}, 10^{-5}, 10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}$.

barrier. Such increase in the mean velocity when dissipation is present was also observed in [25]. For the decelerated case we found for the saturated time $t_c \sim \gamma^{-1.2}$ while v_c is constant in the interval $10^{-6} \leq \gamma \leq 10^{-2}$ and decreases for $\gamma = 0.1$.

To conclude, for a long time the Fermi acceleration has been studied in different models and applications [1–19, 30], but just recently [20] a deformed sawtooth (Ratchetlike) pulse was proposed to control the Fermi de(acceleration). With changing the asymmetry parameter from the Ratchetlike pulse it was possible to get Fermi hyperacceleration and deceleration. By switching the pulse between hyperacceleration and deceleration modes an accurate control of the particles velocity was achieved. Beside the remarkable control of velocities obtained in the dissipation free problem, we observe here that dissipation effects transform regular islands into sinks, change the mean velocity and reduce the ability to control accurately the particle dynamics. The saturation time decays with a power law $t_c \sim \gamma^{-\beta}$, where β is close to 1.0 (± 0.01) for all considered cases. On the other hand, the saturated mean velocity also obeys the power law decay, but with exponents $\beta \sim 0.6$ for case **H**, and $\beta \sim 0.3$ for case **A**. In the decelerated case the saturated velocity is almost constant for small dissipations. Our results motivate further analysis related to the control of Fermi de(acceleration) for larger Ratchetlike asymmetries of the ac-driven pulse in the presence of dissipation [32] and in the hopping wall approximation. In the later case we expect to obtain a better efficiency to control the particles velocity since the energy gain per collision is not underestimated as in the present model. In addition, it would be interesting to analyse the implementation of the Ratchetlike ac-driven pulse applied to the walls of quasi-one-dimensional billiards which are coupled to thermal baths at different temperatures. In such cases it is desirable to optimize and achieve the control of directed heat conduction [33], which could leave to an important increase of the thermal efficiency in physical devices such as rectifiers and thermal transistors [34].

Acknowledgment

The authors thank CNPq and FINEP (under project CTINFRA-1) for financial support.

References

- [1] M. Lieberman and A. J. Lichtenberg, "Stochastic and adiabatic behavior of particles accelerated by periodic forces," *Physical Review A*, vol. 5, no. 4, pp. 1852–1866, 1972.
- [2] A. J. Lichtenberg and M. A. Lieberman, *Regular and Chaotic Dynamics*, Applied Mathematical Sciences, Springer, New York, NY, USA, 2nd edition, 1992.
- [3] E. D. Leonel, J. K. L. da Silva, and S. O. Kamphorst, "On the dynamical properties of a Fermi accelerator model," *Physica A*, vol. 331, no. 3–4, pp. 435–447, 2004.
- [4] E. D. Leonel, P. V. E. McClintock, and J. K. L. da Silva, "Fermi-Ulam Accelerator Model under Scaling Analysis," *Physical Review Letters*, vol. 93, Article ID 014101, 2004.
- [5] E. D. Leonel and P. V. E. McClintock, "A crisis in the dissipative Fermi accelerator model," *Journal of Physics A*, vol. 38, no. 23, pp. 425–430, 2005.
- [6] R. E. de Carvalho, F. C. Souza, and E. D. Leonel, "Fermi acceleration on the annular billiard," *Physical Review E*, vol. 73, no. 6, Article ID 066229, 10 pages, 2006.
- [7] A. K. Karlis, P. K. Papachristou, F. K. Diakonou, V. Constantoudis, and P. Schmelcher, "Hyperacceleration in a stochastic Fermi-Ulam model," *Physical Review Letters*, vol. 97, no. 19, Article ID 194102, 4 pages, 2006.
- [8] A. K. Karlis, P. K. Papachristou, F. K. Diakonou, V. Constantoudis, and P. Schmelcher, "Fermi acceleration in the randomized driven Lorentz gas and the Fermi-Ulam model," *Physical Review E*, vol. 76, no. 1, Article ID 016214, 17 pages, 2007.
- [9] A. Steane, P. Szriftgiser, P. Desbiolles, and J. Dalibard, "Phase modulation of atomic de Broglie waves," *Physical Review Letters*, vol. 74, no. 25, pp. 4972–4975, 1995.

- [10] F. Saif, I. Bialynicki-Birula, M. Fortunato, and W. P. Schleich, "Fermi accelerator in atom optics," *Physical Review A*, vol. 58, pp. 4779–4783, 1998.
- [11] F. Saif, "Classical and quantum chaos in atom optics," *Physics Reports*, vol. 419, no. 6, pp. 207–258, 2005.
- [12] A. V. Milovanov and L. M. Zeleny, "'Strange' Fermi processes and power-law nonthermal tails from a self-consistent fractional kinetic equation," *Physical Review E*, vol. 64, no. 5, Article ID 052101, 4 pages, 2001.
- [13] G. Michalek, M. Ostrowski, and R. Schlickeiser, "Cosmic-Ray Momentum Diffusion in Magnetosonic versus Alfvénic Turbulent Field," *Solar Physics*, vol. 184, p. 339, 2001, <http://www.springerlink.com>.
- [14] A. Veltri and V. Carbone, "Radiative intermittent events during Fermi's stochastic acceleration," *Physical Review Letters*, vol. 92, no. 14, Article ID 143901, 4 pages, 2004.
- [15] K. Kobayakawa, Y. S. Honda, and T. Tamura, "Acceleration by oblique shocks at supernova remnants and cosmic ray spectra around the knee region," *Physical Review D*, vol. 66, no. 8, Article ID 083004, 11 pages, 2002.
- [16] M. A. Malkov, "Ion leakage from quasiparallel collisionless shocks: implications for injection and shock dissipation," *Physical Review E*, vol. 58, no. 4, pp. 4911–4928, 1998.
- [17] E. Fermi, "On the origin of the Cosmic Radiation," *Physical Review*, vol. 75, p. 1169, 1949.
- [18] S. M. Ulam, "On some statistical properties of dynamical systems," in *Proceedings of the 4th Berkeley Symposium on Mathematical Statistics and Probability*, vol. 3, pp. 315–320, University of California Press, Berkeley, Calif, USA, 1961.
- [19] L. D. Pustyl'nikov, "Stable and oscillating motions in nonautonomous dynamical systems. II," *Trudy Moskovskogo Matematicheskogo Obščestva*, vol. 34, pp. 3–103, 1977, English translation in *Transactions of the Moscow Mathematical Society*, vol. 2, p. 1, 1978.
- [20] C. Manchein and M. W. Beims, "Ratchetlike pulse controlling the Fermi deceleration and hyperacceleration," *International Journal of Computational Science and Engineering*, vol. 1, p. 99, 2009.
- [21] Y.-C. Lai and C. Grebogi, "Complexity in Hamiltonian-driven dissipative chaotic dynamical systems," *Physical Review E*, vol. 54, p. 4667, 1996.
- [22] P. C. Rech, M. W. Beims, and J. A. C. Gallas, "Basin size evolution between dissipative and conservative limits," *Physical Review E*, vol. 71, no. 1, Article ID 17207, 4 pages, 2005.
- [23] C. Manchein, J. Rosa, and M. W. Beims, "Chaotic motion at the emergence of the time averaged energy decay," *Physica D*, vol. 238, no. 16, pp. 1688–1694, 2009.
- [24] U. Feudel, C. Grebogi, B. R. Hunt, and J. A. Yorke, "Map with more than 100 coexisting low-period periodic attractors," *Physical Review E*, vol. 54, no. 1, pp. 71–81, 1996.
- [25] R. E. de Carvalho and C. V. Abud, "Dissipation as a mechanism of energy gain," *Physical Review E*, vol. 77, Article ID 036204, 2008.
- [26] D. F. Tavares, "A simplified Fermi accelerator model under quadratic frictional force," *Brazilian Journal of Physics*, vol. 38, no. 1, p. 58, 2008.
- [27] E. D. Leonel and R. E. de Carvalho, "A family of crisis in a dissipative Fermi accelerator model," *Physics Letters A*, vol. 364, no. 6, pp. 475–479, 2007.
- [28] A. L. P. Livorati, D. G. Ladeira, and E. D. Leonel, "Scaling investigation of Fermi acceleration on a dissipative bouncer model," *Physical Review E*, vol. 78, no. 5, Article ID 056205, 12 pages, 2008.
- [29] D. G. Ladeira and E. D. Leonel, "Dynamical properties of a dissipative hybrid Fermi-Ulam-bouncer model," *Chaos*, vol. 17, no. 1, Article ID 013119, 7 pages, 2007.
- [30] D. Cole, S. Bending, S. Savellév, A. Grigorenko, T. Tamegai, and F. Nori, "Ratchet without spatial asymmetry for controlling the motion of magnetic flux quanta using time-asymmetric drives," *Nature Materials*, vol. 5, pp. 305–311, 2006.
- [31] A. Loskutov, "Dynamical chaos: systems of classical mechanics," *Physics-Uspokhi*, vol. 50, no. 9, pp. 939–964, 2007.
- [32] C. Manchein and M. W. Beims, "Controlling the Fermi acceleration by strong dissipation," in preparation.
- [33] R. Klages, *Microscopic Chaos, Fractals and Transport in Nonequilibrium Statistical Mechanics*, vol. 24 of *Advanced Series in Nonlinear Dynamics*, World Scientific, River Edge, NJ, USA, 2007.
- [34] G. Casati, C. Mejía-Monasterio, and T. Prosen, "Increasing thermoelectric efficiency: a dynamical systems approach," *Physical Review Letters*, vol. 101, no. 1, Article ID 016601, 4 pages, 2008.