Research Article

Exact Solutions for the Generalized BBM Equation with Variable Coefficients

Cesar A. Gómez¹ and Alvaro H. Salas^{2,3}

¹ Department of Mathematics, Universidad Nacional de Colombia, Bogotá, Colombia

² Department of Mathematics and Statistics, Universidad Nacional de Colombia, Manizales, Cll 45, Cra 30, Colombia

³ Department of Mathematics, Universidad de Caldas, Manizales, Colombia

Correspondence should be addressed to Alvaro H. Salas, asalash2002@yahoo.com

Received 23 November 2009; Accepted 21 January 2010

Academic Editor: Jihuan He

Copyright © 2010 C. A. Gómez and A. H. Salas. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The variational iteration algorithm combined with the exp-function method is suggested to solve the generalized Benjamin-Bona-Mahony equation (BBM) with variable coefficients. Periodic and soliton solutions are formally derived in a general form. Some particular cases are considered.

1. Introduction

The BBM equation

$$u_t + uu_x + u_x - \mu u_{xxt} = 0, \quad (1.1)$$

which describes approximately the unidirectional propagation of long waves in certain nonlinear dispersive systems, has been proposed by Benjamin et al. in 1972 [1] as a more satisfactory model than the KdV equation [2]

$$u_t + uu_x + u_{xxx} = 0. \quad (1.2)$$

It is easy to see that (1.1) can be derived from the equal width EW-equation [3]:

$$u_t + uu_x - \mu u_{xxt} = 0, \quad (1.3)$$

by means of the change of variable $u = u + 1$, that is, by replacing u with $u + 1$. This last equation is considered as an equally valid and accurate model for the same wave phenomena

simulated by (1.1) and (1.2). On the other hand, some researches analyzed the generalized KdV equation with variable coefficients

$$u_t + \sigma(t)u^p u_x + \mu(t)u_{xxx} = 0, \quad (1.4)$$

because this model has important applications in several fields of science [4–7].

Motivated by these facts, we will consider here the generalized EW-equation with variable coefficients

$$u_t + \sigma(t)u^p u_x - \mu(t)u_{xxt} = 0. \quad (1.5)$$

Using the solutions of (1.5) we obtain exact solutions to the generalized BBM equation

$$u_t + \sigma(t)(u + 1)^p u_x - \mu(t)u_{xxt} = 0, \quad (1.6)$$

of order $p > 0$.

2. Exact Solutions to Generalized BBM Equation

2.1. The Variational Iteration Method

Consider the following nonlinear equation:

$$Lu(x, t) + Nu(x, t) = g(x, t), \quad (2.1)$$

where L and N are linear and nonlinear operators, respectively, and $g(x, t)$ is an inhomogeneous term. According to the variational iteration method (VIM) [8–14], a functional correction to (2.1) is given by

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \theta(\tau) (Lu_n(x, \tau) + N\tilde{u}_n(x, \tau) - g(x, \tau)) d\tau, \quad (2.2)$$

where $\theta(\tau)$ is a general Lagrange's multiplier, which can be identified via the variational theory; the subscript $n \geq 0$ denotes the n th order approximation and $\tilde{u}$ is a restricted variation which means $\delta\tilde{u} = 0$. In this method, we first determine the Lagrange multiplier $\theta(\tau)$ that will be identified optimally via integration by parts. The successive approximation u_{n+1} of the solution u will be readily obtained upon using the determined Lagrangian multiplier and any selective function u_0 . One of the advantages of the VIM, is the free choice of the initial solution $u_0(x, t)$. If we consider a special form to u_0 with arbitrary parameters, using the relations

$$u_n(x, t) = u_{n+1}(x, t), \quad \frac{\partial^k}{\partial t^k} u_n(x, t) = \frac{\partial^k}{\partial t^k} u_{n+1}(x, t), \quad (2.3)$$

we can obtain a set of algebraic equations in the unknowns given by the parameters that appear in u_0 . Solving this system, we have exact solutions to (2.1). To solve (1.5), we construct the following functional equation

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \theta(\tau)(Lu_n(x, \tau) + N\tilde{u}_n(x, \tau))d\tau, \quad (2.4)$$

where

$$\begin{aligned} Lu_n(x, \tau) &= (u_n)_\tau(x, \tau), \\ N\tilde{u}_n(x, \tau) &= \sigma(\tau)(\tilde{u} + 1)^p \tilde{u}_x(x, \tau) - \mu(\tau)\tilde{u}_{xx\tau}(x, \tau). \end{aligned} \quad (2.5)$$

Taking in (2.4) variation with respect to the independent variable u_n , and noticing that $\delta N\tilde{u}_n = 0$ we have

$$\begin{aligned} \delta u_{n+1}(x, t) &= \delta u_n(x, t) + \delta \int_0^t \theta(\tau)(Lu_n(x, \tau) + N\tilde{u}_n(x, \tau))d\tau \\ &= \delta u_n(x, t) + \theta(t)\delta u_n(x, t) - \int_0^t \theta'(\tau)\delta u_n(x, \tau)d\tau = 0. \end{aligned} \quad (2.6)$$

This yields the stationary conditions

$$\begin{aligned} 1 + \theta(t) &= 0, \\ \theta'(t) &= 0. \end{aligned} \quad (2.7)$$

Therefore,

$$\theta(t) = -1. \quad (2.8)$$

Substituting this value into (2.4) we obtain the formula

$$u_{n+1}(x, t) = u_n(x, t) - \int_0^t (Lu_n(x, \tau) + N\tilde{u}_n(x, \tau))d\tau. \quad (2.9)$$

Using the wave transformation

$$\xi = x + \lambda t + \xi_0, \quad (2.10)$$

setting

$$\frac{\partial}{\partial t} u_1(\xi) = \frac{\partial}{\partial t} u_0(\xi), \quad (2.11)$$

and performing one integration, (2.9) reduces to

$$\lambda u_0(\xi) + \frac{\sigma(t)}{p+1} u_0^{p+1}(\xi) - \lambda \mu(t) u_0''(\xi) = 0, \quad (2.12)$$

where for sake of simplicity we set the constant of integration equal to zero. With the change of variable

$$u_0(\xi) = v^{2/p}(\xi), \quad (2.13)$$

equation (2.12) converts to

$$\lambda v^2(\xi) - \frac{2\mu(t)(2-p)}{p^2} \lambda (v')^2 - \frac{2\mu(t)}{p} \lambda v(\xi) v''(\xi) + \frac{\sigma(t)}{p+1} v(\xi)^4 = 0. \quad (2.14)$$

Observe that if $v(\xi)$ is a solution to (2.14), then $-v(\xi)$ is also a solution to this equation.

2.2. The Exp-Function Method

Recently, He and Wu [15] have introduced the Exp-function method to solve nonlinear differential equations. In particular, the Exp-function method is an effective method for solving nonlinear equations with high nonlinearity. The method has been used in a satisfactory way by other authors to solve a great variety of nonlinear wave equations [15–21]. The Exp-function method is very simple and straightforward, and can be briefly revised as follows: Given the nonlinear partial differential equation

$$F(u, u_x, u_t, u_{xx}, u_{xt}, u_{tt}, \dots) = 0, \quad (2.15)$$

it is transformed to ordinary differential equation

$$F(u, u', u'', u''', u_{xt}, \dots) = 0, \quad (2.16)$$

by mean of wave transformation $\xi = x + \lambda t + \xi_0$. Solutions to (2.16) can then be found using the expression

$$u(\xi) = \frac{\sum_{n=-c}^d a_n \exp(n\xi)}{\sum_{n=-p}^q b_n \exp(n\xi)}, \quad (2.17)$$

where c, d, p , and q are positive integers which are unknown to be determined later, a_n and b_n are unknown constants.

After balancing, we substitute (2.17) into (2.16) to obtain an algebraic systems in the variable $\zeta = \exp(n\xi)$. Solving the algebraic system we can obtain exact solutions to (2.16) and reversing, solutions to (2.15) in the original variables.

3. Solutions to (2.14) by the Exp-Function Method

Using the Exp-function method, we suppose that solutions to (2.14) can be expressed in the form

$$v(\xi) = \frac{\sum_{n=-1}^1 a_n \exp(nr\xi)}{\sum_{m=-1}^1 b_m \exp(mr\xi)} = \frac{a_{-1} \exp(-r\xi) + a_0 + a_1 \exp(r\xi)}{b_{-1} \exp(-r\xi) + b_0 + b_1 \exp(r\xi)}. \quad (3.1)$$

We obtain following solutions to (2.14):

$$\begin{aligned} v_1 &= \pm \frac{2\lambda k(p+1)(p+2)}{2(p+1)(p+2)\lambda \exp\left(\left(p/2\sqrt{\mu(t)}\right)\xi\right) - k^2\sigma(t) \exp\left(-\left(p/2\sqrt{\mu(t)}\right)\xi\right)}, & \lambda &= \lambda(t), \\ v_2 &= \pm \frac{2\lambda k(p+1)(p+2)}{\sigma(t)k^2 \exp\left(\left(p/2\sqrt{\mu(t)}\right)\xi\right) - 2\lambda(p+1)(p+2) \exp\left(-\left(p/2\sqrt{\mu(t)}\right)\xi\right)}, & \lambda &= \lambda(t). \end{aligned} \quad (3.2)$$

Some special solutions are obtained if

$$\lambda = \lambda(t) = \pm \frac{k^2}{2(p^2 + 3p + 2)} \sigma(t). \quad (3.3)$$

This choice gives solutions

$$v_3 = \pm \frac{k}{2} \operatorname{csch}\left(\frac{p}{2\sqrt{\mu(t)}}\xi\right), \quad \lambda = \frac{k^2}{2(p^2 + 3p + 2)} \sigma(t), \quad (3.4)$$

$$v_4 = \frac{k}{2} \operatorname{sech}\left(\frac{p}{2\sqrt{\mu(t)}}\xi\right), \quad \lambda = -\frac{k^2}{2(p^2 + 3p + 2)} \sigma(t), \quad (3.5)$$

$$v_5 = \pm \frac{k}{2} \operatorname{csc}\left(\frac{p}{2\sqrt{\mu(t)}}\xi\right), \quad \lambda = -\frac{k^2}{2(p^2 + 3p + 2)} \sigma(t). \quad (3.6)$$

Solution (3.6) follows from (3.4) with the identifications $\mu(t) \rightarrow -\mu(t)$ and $k \rightarrow -k\sqrt{-1}$.

$$v_6 = -\frac{k}{2} \sec\left(\frac{p}{2\sqrt{\mu(t)}}\xi\right), \quad \lambda = -\frac{k^2}{2(p^2 + 3p + 2)}\sigma(t). \quad (3.7)$$

Solution (3.7) follows from (3.4) with the identifications $\mu(t) \rightarrow -\mu(t)$ and $k \rightarrow -k$.

4. Particular Cases

4.1. Case 1: Solutions to (2.14) When $p = 2$

Equation (2.14) takes the form

$$\lambda v^2(\xi) - \lambda \mu(t) v(\xi) v''(\xi) + \frac{1}{3} \sigma(t) v(\xi)^4 = 0. \quad (4.1)$$

From (3.2) with $p = 2$:

$$\begin{aligned} v_7 &= \pm \frac{24\lambda k}{24\lambda \exp\left(\left(1/\sqrt{\mu(t)}\right)\xi\right) - k^2 \sigma(t) \exp\left(-\left(1/\sqrt{\mu(t)}\right)\xi\right)}, \\ v_8 &= \pm \frac{24\lambda k}{k^2 \sigma(t) \exp\left(\left(1/\sqrt{\mu(t)}\right)\xi\right) - 24\lambda \exp\left(-\left(1/\sqrt{\mu(t)}\right)\xi\right)}. \end{aligned} \quad (4.2)$$

From (3.3)–(3.7) with $p = 2$:

$$\begin{aligned} v_9 &= \pm \frac{k}{2} \operatorname{csch}\left(\frac{1}{\sqrt{\mu(t)}}\xi\right), & \lambda &= \frac{k^2}{24}\sigma(t), \\ v_{10} &= \pm \frac{k}{2} \operatorname{sech}\left(\frac{1}{\sqrt{\mu(t)}}\xi\right), & \lambda &= -\frac{k^2}{24}\sigma(t), \\ v_{11} &= \pm \frac{k}{2} \csc\left(\frac{1}{\sqrt{-\mu(t)}}\xi\right), & \lambda &= -\frac{k^2}{24}\sigma(t), \\ v_{12} &= \pm \frac{k}{2} \sec\left(\frac{1}{\sqrt{\mu(t)}}\xi\right), & \lambda &= -\frac{k^2}{24}\sigma(t). \end{aligned} \quad (4.3)$$

Other exact solutions are:

$$\begin{aligned}
 v_{13} &= \pm \frac{(3a^2 \exp(2\sqrt{-2/\mu(t)}\xi) + 2\sqrt{55}a \exp(\sqrt{-2/\mu(t)}\xi) - 22)k}{3a^2 \exp(2\sqrt{-2/\mu(t)}\xi) + 22a \exp(\sqrt{-2/\mu(t)}\xi) + 22}, & \lambda &= -\frac{1}{3}k^2\sigma(t), \\
 v_{14} &= \pm \frac{(3a^2 \pm 2\sqrt{55}a \exp(\sqrt{-2/\mu(t)}\xi) - 22 \exp(2\sqrt{-2/\mu(t)}\xi))k}{3a^2 + 22a \exp(\sqrt{-2/\mu(t)}\xi) + 22 \exp(2\sqrt{-2/\mu(t)}\xi)}, & \lambda &= -\frac{1}{3}k^2\sigma(t), \\
 v_{15} &= \pm k \left(1 - \frac{44(8 + \sqrt{55})}{3a(11 + \sqrt{55}) \exp(\sqrt{-2/\mu(t)}\xi) + 22(8 + \sqrt{55})} \right), & \lambda &= -\frac{1}{3}k^2\sigma(t), \\
 v_{16} &= \pm \frac{k(a \pm \sinh(\sqrt{-2/\mu(t)}\xi))}{\sqrt{a^2 + 1} \pm \cosh(\sqrt{-2/\mu(t)}\xi)}, & \lambda &= -\frac{1}{3}k^2\sigma(t), \\
 v_{17} &= \pm \frac{k(a \pm \cosh(\sqrt{-2/\mu(t)}\xi))}{\sqrt{a^2 + 1} \pm \sinh(\sqrt{-2/\mu(t)}\xi)}, & \lambda &= -\frac{1}{3}k^2\sigma(t), \\
 v_{18} &= \pm \frac{k \cos(\sqrt{2/\mu(t)}\xi)}{1 \pm \sin(\sqrt{2/\mu(t)}\xi)}, & \lambda &= \frac{k^2}{3}\sigma(t).
 \end{aligned} \tag{4.4}$$

4.2. Case 2: Solutions to (2.14) When $p = 4$

Equation (2.14) takes the form

$$\lambda v^2(\xi) + \mu(t)\lambda(v')^2 - \frac{\lambda}{2}\mu(t)v(\xi)v''(\xi) + \frac{1}{5}\sigma(t)v(\xi)^4 = 0. \tag{4.5}$$

From (3.2) with $p = 4$:

$$\begin{aligned}
 v_{19} &= \pm \frac{60\lambda k}{60\lambda \exp\left(\left(2/\sqrt{\mu(t)}\right)\xi\right) - k^2\sigma(t) \exp\left(-\left(2/\sqrt{\mu(t)}\right)\xi\right)}, & \lambda &= \lambda(t), \\
 v_{20} &= \pm \frac{60\lambda k}{\sigma(t)k^2 \exp\left(\left(2/\sqrt{\mu(t)}\right)\xi\right) - 60\lambda \exp\left(-\left(2/\sqrt{\mu(t)}\right)\xi\right)}, & \lambda &= \lambda(t).
 \end{aligned} \tag{4.6}$$

From (3.3)–(3.7) with $p = 4$:

$$\begin{aligned}
 v_{21} &= \pm \frac{k}{2} \operatorname{csch} \left(\frac{2}{\sqrt{\mu(t)}} \xi \right), & \lambda &= \frac{k^2}{60} \sigma(t), \\
 v_{22} &= \pm \frac{k}{2} \operatorname{sech} \left(\frac{2}{\sqrt{\mu(t)}} \xi \right), & \lambda &= -\frac{k^2}{60} \sigma(t), \\
 v_{23} &= \pm \frac{k}{2} \operatorname{csc} \left(\frac{2}{\sqrt{-\mu(t)}} \xi \right), & \lambda &= -\frac{k^2}{60} \sigma(t), \\
 v_{24} &= \pm \frac{k}{2} \operatorname{sec} \left(\frac{2}{\sqrt{-\mu(t)}} \xi \right), & \lambda &= -\frac{k^2}{60} \sigma(t).
 \end{aligned} \tag{4.7}$$

Other exact solutions are:

$$\begin{aligned}
 v_{25} &= \pm \frac{k \left(a \exp \left(\left(2/\sqrt{-\mu(t)} \right) \xi \right) - 4 \right)^2}{a^2 \exp \left(\left(4/\sqrt{-\mu(t)} \right) \xi \right) + 16a \exp \left(\left(2/\sqrt{-\mu(t)} \right) \xi \right) + 16}, & \lambda &= -\frac{1}{5} k^2 \sigma(t), \\
 v_{26} &= \pm \frac{k \left(4 \exp \left(\left(2/\sqrt{-\mu(t)} \right) \xi \right) - a \right)^2}{a^2 + 16a \exp \left(\left(2/\sqrt{-\mu(t)} \right) \xi \right) + 16 \exp \left(\left(4/\sqrt{-\mu(t)} \right) \xi \right)}, & \lambda &= -\frac{1}{5} k^2 \sigma(t), \\
 v_{27} &= \pm 2k \left(1 - \frac{3}{2 \pm \cos \left(\left(2/\sqrt{\mu(t)} \right) \xi \right)} \right), & \lambda &= -\frac{4}{5} k^2 \sigma(t), \\
 v_{28} &= \pm 2k \left(1 - \frac{3}{2 \pm \sin \left(\left(2/\sqrt{\mu(t)} \right) \xi \right)} \right), & \lambda &= -\frac{4}{5} k^2 \sigma(t).
 \end{aligned} \tag{4.8}$$

It is clear that using (2.13) we obtain solutions to (1.5). Finally, observe that if $u_0(x, t)$ is a solution of (1.5), then the solutions $u(x, t)$ to the generalized BBM equation (1.6) are obtained as follows:

$$u(x, t) = u_0(x, t) - 1. \tag{4.9}$$

5. Conclusions

We have considered the generalized EW-equation with variable coefficients and the generalized BBM-equation with variable coefficients. We obtained analytic solutions by using the variational iteration method combined with the exp-function method. With the aid of *Mathematica* we have derived a lot of different types of solutions for these two models. Combined formal soliton-like solutions as well as kink solutions have been formally derived.

The results obtained show that the technique used here can be considered as a powerful method to analyze other types of nonlinear wave equations.

According to [22], there are alternative iteration algorithms, which might be useful for future work. Furthermore, various modifications of the exp-function method have been appeared in open literature, for example, the double exp-function method [23, 24].

Other methods for solving nonlinear differential equations may be found in [25–35].

We think that the results presented in this paper are new in the literature.

References

- [1] T. B. Benjamin, J. L. Bona, and J. J. Mahony, "Model equations for long waves in nonlinear dispersive systems," *Philosophical Transactions of the Royal Society of London. Series A*, vol. 272, no. 1220, pp. 47–78, 1972.
- [2] D. J. Korteweg and G. de Vries, "On the change of long waves advancing in a rectangular canal and a new type of long stationary wave," *Philosophical Magazine*, vol. 39, pp. 422–443, 1835.
- [3] P. J. Morrison, J. D. Meiss, and J. R. Cary, "Scattering of regularized-long-wave solitary waves," *Physica D*, vol. 11, no. 3, pp. 324–336, 1984.
- [4] R. M. Miura, "Korteweg-de Vries equation and generalizations. I. A remarkable explicit nonlinear transformation," *Journal of Mathematical Physics*, vol. 9, pp. 1202–1204, 1968.
- [5] N. Nirmala, M. J. Vedan, and B. V. Baby, "Auto-Bäcklund transformation, Lax pairs, and Painlevé property of a variable coefficient Korteweg-de Vries equation. I," *Journal of Mathematical Physics*, vol. 27, no. 11, pp. 2640–2643, 1986.
- [6] Z. Liu and C. Yang, "The application of bifurcation method to a higher-order KdV equation," *Journal of Mathematical Analysis and Applications*, vol. 275, no. 1, pp. 1–12, 2002.
- [7] Y. Zhang, S. Lai, J. Yin, and Y. Wu, "The application of the auxiliary equation technique to a generalized mKdV equation with variable coefficients," *Journal of Computational and Applied Mathematics*, vol. 223, no. 1, pp. 75–85, 2009.
- [8] J.-H. He, "Variational iteration method for autonomous ordinary differential systems," *Applied Mathematics and Computation*, vol. 114, no. 2-3, pp. 115–123, 2000.
- [9] J.-H. He, "Homotopy perturbation method: a new nonlinear analytical technique," *Applied Mathematics and Computation*, vol. 135, no. 1, pp. 73–79, 2003.
- [10] J.-H. He, "The homotopy perturbation method nonlinear oscillators with discontinuities," *Applied Mathematics and Computation*, vol. 151, no. 1, pp. 287–292, 2004.
- [11] A.-M. Wazwaz, "The variational iteration method for rational solutions for KdV, $K(2,2)$, Burgers, and cubic Boussinesq equations," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 18–23, 2007.
- [12] E. Yusufoglu and A. Bekir, "The variational iteration method for solitary patterns solutions of gBBM equation," *Physics Letters. A*, vol. 367, no. 6, pp. 461–464, 2007.
- [13] J.-M. Zhu, Z.-M. Lu, and Y.-L. Liu, "Doubly periodic wave solutions of Jaulent-Miodek equations using variational iteration method combined with Jacobian-function method," *Communications in Theoretical Physics*, vol. 49, no. 6, pp. 1403–1406, 2008.
- [14] C. A. Gómez and A. H. Salas, "The variational iteration method combined with improved generalized tanh-coth method applied to Sawada-Kotera equation," *Applied Mathematics and Computation*, 2009. In press.
- [15] J.-H. He and X.-H. Wu, "Exp-function method for nonlinear wave equations," *Chaos, Solitons and Fractals*, vol. 30, no. 3, pp. 700–708, 2006.
- [16] S. Zhang, "Exp-function method exactly solving the KdV equation with forcing term," *Applied Mathematics and Computation*, vol. 197, no. 1, pp. 128–134, 2008.
- [17] J.-H. He and L.-N. Zhang, "Generalized solitary solution and compacton-like solution of the Jaulent-Miodek equations using the Exp-function method," *Physics Letters. A*, vol. 372, no. 7, pp. 1044–1047, 2008.
- [18] S. Zhang, "Application of Exp-function method to a KdV equation with variable coefficients," *Physics Letters. A*, vol. 365, no. 5-6, pp. 448–453, 2007.
- [19] A. H. Salas, "Exact solutions for the general fifth KdV equation by the exp function method," *Applied Mathematics and Computation*, vol. 205, no. 1, pp. 291–297, 2008.

- [20] A. H. Salas, C. A. Gómez, and J. E. Castillo Hernández, "New abundant solutions for the Burgers equation," *Computers & Mathematics with Applications*, vol. 58, no. 3, pp. 514–520, 2009.
- [21] S. Zhang, "Exp-function method: solitary, periodic and rational wave solutions of nonlinear evolution equations," *Nonlinear Science Letters A*, vol. 1, pp. 143–146, 2010.
- [22] J.-H. He, G.-C. Wu, and F. Austin, "The variational iteration method which should be followed," *Nonlinear Science Letters A*, vol. 1.1, pp. 1–30, 2010.
- [23] Z.-D. Dai, C.-J. Wang, S.-Q. Lin, D.-L. Li, and G. Mu, "The three-wave method for nonlinear evolution equations," *Nonlinear Science Letters A*, vol. 1, no. 1, pp. 77–82, 2010.
- [24] H.-M. Fu and Z.-D. Dai, "Double Exp-function method and application," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 10, no. 7, pp. 927–933, 2009.
- [25] A. H. Salas, "Some solutions for a type of generalized Sawada-Kotera equation," *Applied Mathematics and Computation*, vol. 196, no. 2, pp. 812–817, 2008.
- [26] A. H. Salas and C. A. Gómez, "Computing exact solutions for some fifth KdV equations with forcing term," *Applied Mathematics and Computation*, vol. 204, no. 1, pp. 257–260, 2008.
- [27] A. H. Salas, C. A. Gómez, and J. G. Escobar, "Exact solutions for the general fifth order KdV equation by the extended tanh method," *Journal of Mathematical Sciences: Advances and Applications*, vol. 1, no. 2, pp. 305–310, 2008.
- [28] A. H. Salas and C. A. Gómez, "A practical approach to solve coupled systems of nonlinear PDE's," *Journal of Mathematical Sciences: Advances and Applications*, vol. 3, no. 1, pp. 101–107, 2009.
- [29] A. H. Salas and C. A. Gómez, "El software Mathematica en la búsqueda de soluciones exactas de ecuaciones diferenciales no lineales en derivadas parciales mediante el uso de la ecuación de Riccati," in *Memorias del Primer Seminario Internacional de Tecnologías en Educación Matemática*, vol. 1, pp. 379–387, Universidad Pedagógica Nacional, Santafé de Bogotá, Colombia, 2005.
- [30] A. H. Salas, H. J. Castillo, and J. G. Escobar, "About the seventh-order Kaup-Kupershmidt equation and its solutions," September 2008, <http://arxiv.org>, arXiv:0809.2865.
- [31] A. H. Salas and J. G. Escobar, "New solutions for the modified generalized Degasperis-Procesi equation," September 2008, <http://arxiv.org/abs/0809.2864>.
- [32] A. H. Salas, "Two standard methods for solving the Ito equation," May 2008, <http://arxiv.org/abs/0805.3362>.
- [33] A. H. Salas, "Some exact solutions for the Caudrey-Dodd-Gibbon equation," May 2008, <http://arxiv.org/abs/0805.3362>.
- [34] A. H. Salas S and C. A. Gómez S, "Exact solutions for a third-order KdV equation with variable coefficients and forcing term," *Mathematical Problems in Engineering*, vol. 2009, Article ID 737928, 13 pages, 2009.
- [35] C. A. Gómez S and A. H. Salas, "The Cole-Hopf transformation and improved tanh-coth method applied to new integrable system (KdV6)," *Applied Mathematics and Computation*, vol. 204, no. 2, pp. 957–962, 2008.