Research Article

A Sextuple Product Identity with Applications

Jun-Ming Zhu^{1,2}

¹ Department of Mathematics, East China Normal University, Shanghai 200241, China

² Department of Mathematics, Luoyang Normal University, Luoyang, Henan 471022, China

Correspondence should be addressed to Jun-Ming Zhu, junming_zhu@163.com

Received 21 February 2011; Revised 20 March 2011; Accepted 24 March 2011

Academic Editor: Ming Li

Copyright © 2011 Jun-Ming Zhu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We get a new proof of a sextuple product identity depending on the Laurent expansion of an analytic function in an annulus. Many identities, including an identity for $(q; q)_{\infty}^4$, are obtained from this sextuple product identity.

1. Introduction

For convenience, we let $|q| < 1$ throughout the paper. We employ the standard notation

$$(a; q)_{\infty} = \prod_{n=0}^{\infty} (1 - aq^n), \quad (a, b, \dots, c; q)_{\infty} = (a; q)_{\infty} (b; q)_{\infty} \cdots (c; q)_{\infty}. \quad (1.1)$$

Series product has been an interesting topic. The Jacobi triple product is one of the most famous series-product identity. We announce it in the following (see, e.g., [1, page 35, Entry 19] or [2, Equation (2.1)]):

$$\left(q, z, \frac{q}{z}; q\right)_{\infty} = \sum_{n=-\infty}^{\infty} (-1)^n q^{(1/2)n(n-1)} z^n, \quad z \neq 0. \quad (1.2)$$

It is well known that an analytic function has a unique Laurent expansion in an annulus. Bailey [3] used this property to prove the quintuple product identity. By this approach, Cooper [4, 5] and Kongsiriwong and Liu [2] proved many types of the Macdonald identities and some other series-product identities. In this paper, we use this method to deal with a sextuple product identity.

In Section 2, we present the sextuple product identity ((2.1) below) and its proof. Our identity is equivalent to [2, Equation (8.16)] by Kongsiriwong and Liu, which is the simplification of [2, Equation (6.13)]. Kongsiriwong and Liu got [2, Equation (8.16)] from a more general identity. In this section, we give it a direct proof.

In Section 3, we get many identities from this sextuple product identity.

To simplify notation, we often write $\sum_n$ for $\sum_{n=-\infty}^{\infty}$ in the following when no confusion occurs.

2. A New Proof of the Sextuple Product Identity

The starting point of our investigation in this section is the identity in the following theorem.

Theorem 2.1. *For any complex number z with $z \neq 0$, one has*

$$\begin{aligned} \left(q, z, \frac{q}{z}; q\right)_{\infty} \left(q^3, z^3, \frac{q^3}{z^3}; q^3\right)_{\infty} &= \left(q^{12}, -q^6, -q^6; q^{12}\right)_{\infty} \sum_n q^{2n^2-2n} z^{4n} \\ &\quad + 2 \left(q^{12}, -q^{12}, -q^{12}; q^{12}\right)_{\infty} \sum_n q^{2n^2+1} z^{4n+2} \\ &\quad - \left(q^3, -q^3, -q^3; q^3\right)_{\infty} \sum_n q^{(1/2)(n^2-n)} z^{2n+1}. \end{aligned} \quad (2.1)$$

Before the proof of Theorem 2.1, we need some preparations. The two identities in the following lemma are from [6]. We write them in this version.

Lemma 2.2. *One has*

$$\begin{aligned} \left(q^8, q^3, q^5; q^8\right)_{\infty} \left(q^{24}, q^9, q^{15}; q^{24}\right)_{\infty} &+ q^2 \left(q^8, q, q^7; q^8\right)_{\infty} \left(q^{24}, q^3, q^{21}; q^{24}\right)_{\infty} \\ &= \left(q^2, -q^2, -q^2; q^2\right)_{\infty} \left(q^6, q^3, q^3; q^6\right)_{\infty}, \end{aligned} \quad (2.2)$$

$$\begin{aligned} \left(q^8, q, q^7; q^8\right)_{\infty} \left(q^{24}, q^9, q^{15}; q^{24}\right)_{\infty} &- q \left(q^8, q^3, q^5; q^8\right)_{\infty} \left(q^{24}, q^3, q^{21}; q^{24}\right)_{\infty} \\ &= \left(q^2, q, q; q^2\right)_{\infty} \left(q^6, -q^6, -q^6; q^6\right)_{\infty}. \end{aligned} \quad (2.3)$$

Proof. For (2.2), see [6, Equation (3.18)]. Equation (2.3) is from [6, Equation (3.21)]. Its proof is similar to that of [6, Equation (3.18)]. $\square$

The lemma above is used to prove the following two identities.

Lemma 2.3. *One has*

$$\begin{aligned} & (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty + (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty \\ &= 2(q^4, -q^4, -q^4; q^4)_\infty (q^{12}, -q^6, -q^6; q^{12})_\infty, \end{aligned} \quad (2.4)$$

$$\begin{aligned} & (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty - (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty \\ &= 2q(q^4, -q^2, -q^2; q^4)_\infty (q^{12}, -q^{12}, -q^{12}; q^{12})_\infty. \end{aligned} \quad (2.5)$$

Proof. By (1.2), we have

$$\begin{aligned} (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty &= \frac{1}{4} (q, -1, -q; q)_\infty (q^3, -1, -q^3; q^3)_\infty \\ &= \frac{1}{4} \sum_m q^{(1/2)(m^2-m)} \sum_n q^{(3/2)(n^2-n)} = \sum_m q^{2m^2+m} \sum_n q^{6n^2+3n} \\ &= \sum_m q^{8m^2+2m} \sum_n q^{24n^2+6n} + q^4 \sum_m q^{8m^2+6m} \sum_n q^{24n^2+18n} \\ &\quad + q^3 \sum_m q^{8m^2+2m} \sum_n q^{24n^2+18n} + q \sum_m q^{8m^2+6m} \sum_n q^{24n^2+6n}, \end{aligned} \quad (2.6)$$

$$\begin{aligned} (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty &= \frac{1}{2} (q, i, -iq; q)_\infty (q^3, -i, iq^3; q^3)_\infty \\ &= \frac{1}{2} \sum_m (-1)^m q^{(1/2)(m^2-m)} i^m \sum_n (-1)^n q^{(3/2)(n^2-n)} i^{3n} \\ &= \sum_m (-1)^m q^{2m^2+m} \sum_n (-1)^n q^{6n^2+3n} \\ &= \sum_m q^{8m^2+2m} \sum_n q^{24n^2+6n} + q^4 \sum_m q^{8m^2+6m} \sum_n q^{24n^2+18n} \\ &\quad - q^3 \sum_m q^{8m^2+2m} \sum_n q^{24n^2+18n} - q \sum_m q^{8m^2+6m} \sum_n q^{24n^2+6n}. \end{aligned} \quad (2.7)$$

Adding (2.6) and (2.7), we have

$$\begin{aligned} & (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty + (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty \\ &= 2 \sum_m q^{8m^2+2m} \sum_n q^{24n^2+6n} + 2q^4 \sum_m q^{8m^2+6m} \sum_n q^{24n^2+18n} \\ &= 2(q^{16}, -q^6, -q^{10}; q^{16})_\infty (q^{48}, -q^{18}, -q^{30}; q^{48})_\infty \\ &\quad + 2q^4(q^{16}, -q^2, -q^{14}; q^{16})_\infty (q^{48}, -q^6, -q^{42}; q^{48})_\infty. \end{aligned} \quad (2.8)$$

By (2.2), we have (2.4).

Subtracting (2.7) from (2.6), we obtain

$$\begin{aligned}
 & (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty - (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty \\
 &= 2q^3 \sum_m q^{8m^2+2m} \sum_n q^{24n^2+18n} + 2q \sum_m q^{8m^2+6m} \sum_n q^{24n^2+6n} \\
 &= 2q^3 (q^{16}, -q^6, -q^{10}; q^{16})_\infty (q^{48}, -q^6, -q^{42}; q^{48})_\infty \\
 &\quad + 2q (q^{16}, -q^2, -q^{14}; q^{16})_\infty (q^{48}, -q^{18}, -q^{32}; q^{48})_\infty.
 \end{aligned} \tag{2.9}$$

Replacing q in (2.3) by $-q^2$ and, then, applying the resulting identity to the above equation, we get (2.5). This completes the proof. $\square$

Proof of Theorem 2.1. Set

$$f(z, q) = \left(q, z, \frac{q}{z}; q \right)_\infty \left(q^3, z^3, \frac{q^3}{z^3}; q^3 \right)_\infty. \tag{2.10}$$

Then f is an analytic function of z in the annulus $0 < |z| < \infty$. Put

$$f(z, q) = \sum_n a_n(q) z^n, \quad 0 < |z| < \infty. \tag{2.11}$$

By (2.10), we can easily verify

$$f(z, q) = z^4 f(zq, q). \tag{2.12}$$

Combining (2.11) and (2.12) gives

$$\sum_m a_m(q) z^m = \sum_m q^{m-4} a_{m-4}(q) z^m. \tag{2.13}$$

Equate the coefficients of z^m on both sides to get

$$a_m(q) = q^{m-4} a_{m-4}(q). \tag{2.14}$$

Using the above relation, we obtain

$$\begin{aligned}
 a_{4m-1}(q) &= q^{2m^2-3m} a_{-1}(q), & a_{4m}(q) &= q^{2m^2-2m} a_0(q), \\
 a_{4m+1}(q) &= q^{2m^2-m} a_1(q), & a_{4m+2}(q) &= q^{2m^2} a_2(q).
 \end{aligned} \tag{2.15}$$

Substituting the above four identities into (2.11), we have

$$\begin{aligned} f(z, q) = & a_{-1}(q) \sum_m q^{2m^2-3m} z^{4m-1} + a_0(q) \sum_m q^{2m^2-2m} z^{4m} \\ & + a_1(q) \sum_m q^{2m^2-m} z^{4m+1} + a_2(q) \sum_m q^{2m^2} z^{4m+2}. \end{aligned} \quad (2.16)$$

By (2.10), we also have

$$f(z, q) = f\left(\frac{q}{z}, q\right). \quad (2.17)$$

This gives

$$\sum_m a_m(q) z^m = \sum_m q^{-m} a_{-m}(q) z^m. \quad (2.18)$$

Then we have

$$a_m(q) = q^{-m} a_{-m}(q). \quad (2.19)$$

Set $m = 1$ to get

$$a_1(q) = q^{-1} a_{-1}(q). \quad (2.20)$$

By this relation, (2.16) reduces to

$$\begin{aligned} f(z, q) = & a_0(q) \sum_m q^{2m^2-2m} z^{4m} + a_1(q) \sum_m q^{(1/2)(m^2-m)} z^{2m+1} \\ & + a_2(q) \sum_m q^{2m^2} z^{4m+2}. \end{aligned} \quad (2.21)$$

Now, it remains to determine $a_0(q)$, $a_1(q)$, and $a_2(q)$.

Putting $z = 1$ in (2.21) gives

$$0 = a_0(q) \sum_m q^{2m^2-2m} + a_1(q) \sum_m q^{(1/2)(m^2-m)} + a_2(q) \sum_m q^{2m^2}. \quad (2.22)$$

Set $z = -1$ in (2.21) to get

$$\begin{aligned} & 4(q, -q, -q; q)_\infty \left(q^3, -q^3, -q^3; q^3 \right)_\infty \\ & = a_0(q) \sum_m q^{2m^2-2m} - a_1(q) \sum_m q^{(1/2)(m^2-m)} + a_2(q) \sum_m q^{2m^2}. \end{aligned} \quad (2.23)$$

Taking $z = i$ in (2.21) and noting that $\sum_m (-1)^m q^{(1/2)(m^2-m)} = 0$, we have

$$(q, i, -iq; q)_\infty (q^3, -i, iq^3; q^3)_\infty = a_0(q) \sum_m q^{2m^2-2m} - a_2(q) \sum_m q^{2m^2}. \quad (2.24)$$

Subtracting (2.23) from (2.22) and noting that $\sum_m q^{(1/2)(m^2-m)} = 2(q, -q, -q; q)_\infty$, we obtain

$$a_1(q) = -\left(q^3, -q^3, -q^3; q^3\right)_\infty. \quad (2.25)$$

Add (2.22) and (2.23) to get

$$2(q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty = a_0(q) \sum_m q^{2m^2-2m} + a_2(q) \sum_m q^{2m^2}. \quad (2.26)$$

Adding (2.24) and (2.26) and, then, using (1.2) in the resulting equation, we obtain

$$\begin{aligned} (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty + (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty \\ = 2a_0(q) (q^4, -q^4, -q^4; q^4)_\infty. \end{aligned} \quad (2.27)$$

By (2.4), we have

$$a_0(q) = (q^{12}, -q^6, -q^6; q^{12})_\infty. \quad (2.28)$$

Similarly, subtracting (2.24) from (2.26) and, then using (1.2), we have

$$\begin{aligned} (q, -q, -q; q)_\infty (q^3, -q^3, -q^3; q^3)_\infty - (q, iq, -iq; q)_\infty (q^3, -iq^3, iq^3; q^3)_\infty \\ = a_2(q) (q^4, -q^2, -q^2; q^4)_\infty. \end{aligned} \quad (2.29)$$

Applying (2.5) to this equation gives

$$a_2(q) = 2q (q^{12}, -q^{12}, -q^{12}; q^{12})_\infty, \quad (2.30)$$

which completes the proof. $\square$

3. Some Applications

In this section, we deduce many modular identities from Theorem 2.1.

Corollary 3.1. *One has*

$$\begin{aligned}
 3(q; q)_\infty^3 (q^3; q^3)_\infty^3 &= (q^{12}, -q^6, -q^6; q^{12})_\infty \sum_n 2n(4n-1)q^{2n^2-2n} \\
 &\quad + 2(q^{12}, -q^{12}, -q^{12}; q^{12})_\infty \sum_n (2n+1)(4n+1)q^{2n^2+1} \\
 &\quad - (q^3, -q^3, -q^3; q^3)_\infty \sum_n n(2n+1)q^{1/2(n^2-n)}.
 \end{aligned} \tag{3.1}$$

Proof. Dividing both sides of (2.1) by $(1-z)^2$, letting $z \rightarrow 1$, and then using L'Hospital's rule twice on the right-hand side gives (3.1). $\square$

Corollary 3.2. *One has*

$$\begin{aligned}
 &(q^{24}, -q^{12}, -q^{12}; q^{24})_\infty (q^8, -q^4, -q^4; q^8)_\infty + 4q^4 (q^{24}, -q^{24}, -q^{24}; q^{24})_\infty (q^8, -q^8, -q^8; q^8)_\infty \\
 &\quad + 2q (q^6, -q^6, -q^6; q^6)_\infty (q^2, -q^2, -q^2; q^2)_\infty \\
 &= (q^2, -q, -q; q^2)_\infty (q^6, -q^3, -q^3; q^6)_\infty,
 \end{aligned} \tag{3.2}$$

$$\begin{aligned}
 &(q^{36}, -q^{18}, -q^{18}; q^{36})_\infty (q^{12}, -q^4, -q^4; q^{12})_\infty + 2q^5 (q^{36}, -q^{36}, -q^{36}; q^{36})_\infty (q^{12}, -q^2, -q^{10}; q^{12})_\infty \\
 &\quad - q (q^9, -q^9, -q^9; q^9)_\infty (q^3, -q, -q^2; q^3)_\infty \\
 &= (q; q)_\infty (q^3; q^3)_\infty,
 \end{aligned} \tag{3.3}$$

$$\begin{aligned}
 &(q^{12}, -q^6, -q^6; q^{12})_\infty (q^4, -q^4, -q^4; q^4)_\infty - q (q^{12}, -q^{12}, -q^{12}; q^{12})_\infty (q^4, -q^2, -q^2; q^4)_\infty \\
 &= \frac{(q^2; q^2)_\infty (q^6; q^6)_\infty}{(-q; q^2)_\infty (-q^3; q^6)_\infty},
 \end{aligned} \tag{3.4}$$

$$\begin{aligned}
& \left(q^{60}, -q^{30}, -q^{30}; q^{60} \right)_{\infty} \left(q^{20}, -q^8, -q^{12}; q^{20} \right)_{\infty} + 2q^9 \left(q^{60}, -q^{60}, -q^{60}; q^{60} \right)_{\infty} \left(q^{20}, -q^2, -q^{18}; q^{20} \right)_{\infty} \\
& - q^2 \left(q^{15}, -q^{15}, -q^{15}; q^{15} \right)_{\infty} \left(q^5, -q, -q^4; q^5 \right)_{\infty} \\
& = \frac{(q; q)_{\infty}}{(q, q^4; q^5)_{\infty}} \frac{(q^5; q^5)_{\infty}}{(q^3, q^{12}; q^{15})_{\infty}}, \tag{3.5}
\end{aligned}$$

$$\begin{aligned}
& \left(q^{60}, -q^{30}, -q^{30}; q^{60} \right)_{\infty} \left(q^{20}, -q^4, -q^{16}; q^{20} \right)_{\infty} + 2q^7 \left(q^{60}, -q^{60}, -q^{60}; q^{60} \right)_{\infty} \left(q^{20}, -q^6, -q^{14}; q^{20} \right)_{\infty} \\
& - q \left(q^{15}, -q^{15}, -q^{15}; q^{15} \right)_{\infty} \left(q^5, -q^2, -q^3; q^5 \right)_{\infty} \\
& = \frac{(q; q)_{\infty}}{(q^2, q^3; q^5)_{\infty}} \frac{(q^5; q^5)_{\infty}}{(q^6, q^9; q^{15})_{\infty}}. \tag{3.6}
\end{aligned}$$

Proof. Replace q in (2.1) by q^2 and, then, z by $-q$. Using (1.2) in the resulting identity gives (3.2).

Replace q in (2.1) by q^3 and, then, z by q . Using (1.2) in the resulting identity gives (3.3).

Replace q in (2.1) by q^4 and, then, z by q . Using (1.2) and the fact that $(q^4, -q, -q^3; q^4)_{\infty} = (q, -q, -q; q)_{\infty}$ in the resulting identity, we obtain

$$\begin{aligned}
& \left(\left(q^{48}, -q^{24}, -q^{24}; q^{48} \right)_{\infty} + 2q^6 \left(q^{48}, -q^{48}, -q^{48}; q^{48} \right)_{\infty} \right) \left(q^4, -q^4, -q^4; q^4 \right)_{\infty} \\
& - q \left(q^{12}, -q^{12}, -q^{12}; q^{12} \right)_{\infty} \left(q^4, -q^2, -q^2; q^4 \right)_{\infty} = \frac{(q^2; q^2)_{\infty}}{(-q; q^2)_{\infty}} \frac{(q^6; q^6)_{\infty}}{(-q^3; q^6)_{\infty}}. \tag{3.7}
\end{aligned}$$

By (1.2), we have

$$\begin{aligned}
\left(q^{12}, -q^6, -q^6; q^{12} \right)_{\infty} &= \sum_n q^{6n^2} = \sum_n q^{6(2n)^2} + \sum_n q^{6(2n+1)^2} \\
&= \left(q^{48}, -q^{24}, -q^{24}; q^{48} \right)_{\infty} + 2q^6 \left(q^{48}, -q^{48}, -q^{48}; q^{48} \right)_{\infty}. \tag{3.8}
\end{aligned}$$

Combining (3.7) and (3.8) gives (3.4).

Replace q in (2.1) by q^5 and, then, z by q^2 . Using (1.2) in the resulting identity gives (3.5).

Replace q in (2.1) by q^5 and, then, z by q . Using (1.2) in the resulting identity gives (3.6). $\square$

Obviously, using the same method above, we can get more identities from (2.1).

Now, we deduce a formula for $(q; q)_{\infty}^4$ from (2.1).

Corollary 3.3. *One has*

$$\begin{aligned} (q; q)_\infty^4 &= 2 \sum_m q^{2m^2} \sum_n 2n q^{6n^2+2n} + 2q \sum_m q^{2m^2+2m} \sum_n (2n-1) q^{6n^2-4n} \\ &\quad + \sum_m q^{2m^2+m} \sum_n (2n+1) q^{(1/2)(3n^2+n)}. \end{aligned} \quad (3.9)$$

Proof. Denote the left-hand side of (2.1) by $f(z)$ and the right-hand side of (2.1) by $g(z)$. Let z_0 be a zero point of $f(z)$. Because (2.1) holds in $0 < |z| < \infty$, z_0 is also a zero point of $g(z)$. If $az_0 = 1$, we have

$$\lim_{z \rightarrow z_0} \frac{f(z)}{1-az} = \lim_{z \rightarrow z_0} \frac{g(z)}{1-az}. \quad (3.10)$$

Setting $z_0 = a = 1$ in (3.10) and by L'Hospital's rule on the right-hand side, we have

$$\begin{aligned} 0 &= \left(q^3, -q^3, -q^3; q^3 \right)_\infty \sum_n (2n+1) q^{(1/2)(n^2-n)} \\ &\quad - 2 \left(q^{12}, -q^{12}, -q^{12}; q^{12} \right)_\infty \sum_n (4n+2) q^{2n^2+1} \\ &\quad - \left(q^{12}, -q^6, -q^6; q^{12} \right)_\infty \sum_n 4n q^{2n^2-2n}. \end{aligned} \quad (3.11)$$

Let $\omega = e^{(2/3)\pi i}$. Putting $z_0 = \omega$ and $a = \omega^2$ in (3.10) and noting $\omega^{3n} = 1$ for any integer n , we have

$$\begin{aligned} 3(1-\omega) \left(q^3; q^3 \right)_\infty^4 &= \left(q^3, -q^3, -q^3; q^3 \right)_\infty \sum_n (2n+1) q^{(1/2)(n^2-n)} \omega^{2(n-1)} \\ &\quad - 2 \left(q^{12}, -q^{12}, -q^{12}; q^{12} \right)_\infty \sum_n (4n+2) q^{2n^2+1} \omega^{n-1} \\ &\quad - \left(q^{12}, -q^6, -q^6; q^{12} \right)_\infty \sum_n 4n q^{2n^2-2n} \omega^n. \end{aligned} \quad (3.12)$$

Taking $z_0 = \omega^2$ and $a = \omega$ in (3.10), we obtain

$$\begin{aligned} 3(1-\omega^2) \left(q^3; q^3 \right)_\infty^4 &= \left(q^3, -q^3, -q^3; q^3 \right)_\infty \sum_n (2n+1) q^{(1/2)(n^2-n)} \omega^{n-1} \\ &\quad - 2 \left(q^{12}, -q^{12}, -q^{12}; q^{12} \right)_\infty \sum_n (4n+2) q^{2n^2+1} \omega^{2(n-1)} \\ &\quad - \left(q^{12}, -q^6, -q^6; q^{12} \right)_\infty \sum_n 4n q^{2n^2-2n} \omega^{2n}. \end{aligned} \quad (3.13)$$

Adding the above three identities together gives

$$\begin{aligned}
9(q^3; q^3)_\infty^4 &= (q^3, -q^3, -q^3; q^3)_\infty \sum_n (2n+1) q^{(1/2)(n^2-n)} (1 + \omega^{n-1} + \omega^{2(n-1)}) \\
&\quad - 2(q^{12}, -q^{12}, -q^{12}; q^{12})_\infty \sum_n (4n+2) q^{2n^2+1} (1 + \omega^{n-1} + \omega^{2(n-1)}) \\
&\quad - (q^{12}, -q^6, -q^6; q^{12})_\infty \sum_n 4n q^{2n^2-2n} (1 + \omega^n + \omega^{2n}).
\end{aligned} \tag{3.14}$$

Using the fact

$$1 + \omega^n + \omega^{2n} = \begin{cases} 3, & n \equiv 0 \pmod{3}, \\ 0, & n \not\equiv 0 \pmod{3} \end{cases} \tag{3.15}$$

in the above identity and, then, replacing q^3 by q , we get

$$\begin{aligned}
(q; q)_\infty^4 &= (q, -q, -q; q)_\infty \sum_n (2n+1) q^{(1/2)(3n^2+n)} \\
&\quad - 4q(q^4, -q^4, -q^4; q^4)_\infty \sum_n (2n+1) q^{6n^2+4n} \\
&\quad - 2(q^4, -q^2, -q^2; q^4)_\infty \sum_n 2n q^{6n^2-2n}.
\end{aligned} \tag{3.16}$$

Replacing n in the last two sums on the right-hand side of the above identity by $-n$ and, then, applying (1.2) to the resulting equation, we get Corollary 3.3. $\square$

4. Conclusion

Besides the Jacobi triple product (1.2), well-known series-product identities are known as the quintuple product identity, the Winquist identity, and so forth. The formula (2.1) is also such an identity. Recently, we also obtain some other identities of this kind, including the simplifications of the formulae [2, Equations (6.12) and (6.14)], with a different method. These identities are widely used in number theory, combinatorics, and many other fields. literature on this topic abounds. In (2.1), if we replace z by e^{2iz} , then the right-hand side of (2.1) turns into fourier series. For recent papers on the applications of fourier analysis, we refer the readers to [7–9].

Acknowledgment

This research is supported by the Shanghai Natural Science Foundation (Grant no. 10ZR1409100), the National Science Foundation of China (Grant no. 10771093), and the Natural Science Foundation of Education Department of Henan Province of China (Grant no. 2007110025).

References

- [1] B. C. Berndt, *Ramanujan's Notebooks. Part III*, Springer, New York, NY, USA, 1991.
- [2] S. Kongsiriwong and Z.-G. Liu, "Uniform proofs of q -series-product identities," *Results in Mathematics*, vol. 44, no. 3-4, pp. 312–339, 2003.
- [3] W. N. Bailey, "On the simplification of some identities of the Rogers-Ramanujan type," *Proceedings of the London Mathematical Society. Third Series*, vol. 1, no. 3, pp. 217–221, 1951.
- [4] S. Cooper, "A new proof of the Macdonald identities for A_{n-1} ," *Journal of the Australian Mathematical Society. Series A*, vol. 62, no. 3, pp. 345–360, 1997.
- [5] S. Cooper, "The Macdonald identities for G_2 and some extensions," *New Zealand Journal of Mathematics*, vol. 26, no. 2, pp. 161–182, 1997.
- [6] S.-L. Chen and S.-S. Huang, "New modular relations for the Göllnitz-Gordon functions," *Journal of Number Theory*, vol. 93, no. 1, pp. 58–75, 2002.
- [7] E. G. Bakhoun and C. Toma, "Mathematical transform of traveling-wave equations and phase aspects of quantum interaction," *Mathematical Problems in Engineering*, vol. 2010, Article ID 695208, 15 pages, 2010.
- [8] C. Cattani, "Shannon wavelets for the solution of integrodifferential equations," *Mathematical Problems in Engineering*, vol. 2010, Article ID 408418, 22 pages, 2010.
- [9] M. Li, S. C. Lim, and S. Chen, "Exact solution of impulse response to a class of fractional oscillators and its stability," *Mathematical Problems in Engineering*, vol. 2011, Article ID 657839, 9 pages, 2011.