Research Article

Existence and Iteration of Monotone Positive Solution of BVP for an Elastic Beam Equation

Jian-Ping Sun and Xian-Qiang Wang

Department of Applied Mathematics, Lanzhou University of Technology, Lanzhou, Gansu 730050, China

Correspondence should be addressed to Jian-Ping Sun, jpsun@lut.cn

Received 5 October 2010; Accepted 28 January 2011

Academic Editor: Jitao Sun

Copyright © 2011 J.-P. Sun and X.-Q. Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the existence of monotone positive solution of boundary value problem for an elastic beam equation. By applying iterative techniques, we not only obtain the existence of monotone positive solution but also establish iterative scheme for approximating the solution. It is worth mentioning that the iterative scheme starts off with zero function, which is very useful and feasible for computational purpose. An example is also included to illustrate the main results.

1. Introduction

It is well-known that beam is one of the basic structures in architecture. The deformations of an elastic beam in equilibrium state can be described by the following equation of deflection curve:

$$\frac{d^2}{dx^2} \left(EI_z \frac{d^2 u}{dx^2} \right) = q(x), \quad (1.1)$$

where E is Yang's modulus constant, I_z is moment of inertia with respect to z axes, and $q(x)$ is loading at x . If the loading of beam considered is in relation to deflection and rate of change of deflection, we need to study more general equation

$$u^{(4)}(x) = f(x, u(x), u'(x)). \quad (1.2)$$

According to different forms of supporting, various boundary value problems (BVPs) should be considered.

Owing to its importance in engineering, physics, and material mechanics, BVPs for elastic beam equations have attracted much attention from many authors see, for example, [1–15] and the references therein. However, almost all of the papers we mentioned focused their attention on the existence of solutions or positive solutions. In the existing literature, there are few papers concerned with the computational methods of solutions or positive solutions. It is worth mentioning that Zhang [16] obtained the existence of positive solutions and established iterative schemes for approximating the solutions for an elastic beam equation with a corner. The main tools used were monotone iterative techniques. For monotone iterative methods, one can refer [17–19] and the references therein.

Motivated greatly by the above-mentioned excellent works, in this paper we investigate the existence and iteration of monotone positive solution for the following elastic beam equation BVP

$$\begin{aligned} u^{(4)}(t) &= f(t, u(t), u'(t)), \quad t \in (0, 1), \\ u(0) &= u'(1) = u''(0) = u'''(1) = 0. \end{aligned} \quad (1.3)$$

In material mechanics, the equation in (1.3) describes the deflection or deformation of an elastic beam under a certain force. The boundary conditions in (1.3) mean that the elastic beam is simply fixed at the end $t = 0$ and fastened with a sliding clamp at the end $t = 1$. By applying iterative techniques, we not only obtain the existence of monotone positive solution but also establish iterative scheme for approximating the solution. It is worth mentioning that the iterative scheme starts off with zero function, which is very useful and feasible for computational purpose. An example is also included to illustrate the main results.

Throughout this paper, we always assume that the following condition is satisfied:

(A) $f \in C([0, 1] \times [0, +\infty) \times [0, +\infty), [0, +\infty))$.

2. Preliminary

In order to obtain the main results of this paper, we first present several fundamental lemmas in this section.

Lemma 2.1. *Let $y \in C[0, 1]$. Then, the BVP*

$$\begin{aligned} u^{(4)}(t) &= y(t), \quad t \in (0, 1), \\ u(0) &= u'(1) = u''(0) = u'''(1) = 0, \end{aligned} \quad (2.1)$$

has a unique solution

$$u(t) = \int_0^1 G(t, s) y(s) ds, \quad t \in [0, 1], \quad (2.2)$$

where

$$G(t, s) = \frac{1}{6} \begin{cases} s(6t - 3t^2 - s^2), & 0 \leq s \leq t \leq 1, \\ t(6s - 3s^2 - t^2), & 0 \leq t \leq s \leq 1. \end{cases} \quad (2.3)$$

Lemma 2.2. For any $(t, s) \in [0, 1] \times [0, 1]$, we have

$$0 \leq \frac{\partial G(t, s)}{\partial t} \leq s(1 - t), \quad \frac{1}{3}t^2s \leq G(t, s) \leq \frac{1}{2}(2t - t^2)s. \quad (2.4)$$

Proof. For any fixed $s \in [0, 1]$, it is easy to know that

$$\frac{\partial G(t, s)}{\partial t} = \begin{cases} s(1 - t), & 0 \leq s \leq t \leq 1, \\ s(1 - t) - \frac{1}{2}(s - t)^2, & 0 \leq t \leq s \leq 1, \end{cases} \quad (2.5)$$

which shows that

$$0 \leq \frac{\partial G(t, s)}{\partial t} \leq s(1 - t), \quad \text{for } (t, s) \in [0, 1] \times [0, 1], \quad (2.6)$$

and so,

$$G(t, s) = \int_0^t \frac{\partial G(\tau, s)}{\partial \tau} d\tau \leq \int_0^t s(1 - \tau) d\tau = \frac{1}{2}(2t - t^2)s, \quad \text{for } (t, s) \in [0, 1] \times [0, 1]. \quad (2.7)$$

On the other hand, it follows from the expression of $G(t, s)$ that

$$\begin{aligned} G(t, s) &= \frac{1}{6}s(6t - 3t^2 - s^2) \geq \frac{1}{6}s(6t - 4t^2) \geq \frac{1}{3}t^2s, \quad 0 \leq s \leq t \leq 1, \\ G(t, s) &= \frac{1}{6}t(6s - 3s^2 - t^2) \geq \frac{1}{6}t(6s - 4s^2) \geq \frac{1}{3}t^2s, \quad 0 \leq t \leq s \leq 1. \end{aligned} \quad (2.8)$$

□

Let $E = C^1[0, 1]$ be equipped with the norm $\|u\| = \max\{\|u\|_\infty, \|u'\|_\infty\}$. Then, E is a Banach space. Denote

$$K = \left\{ u \in E : u(t) \geq \frac{2}{3}t^2\|u\|_\infty, \quad u'(t) \geq 0, \quad \text{for } t \in [0, 1] \right\}. \quad (2.9)$$

Then, it is easy to verify that K is a cone in E . Note that this induces an order relation $\lesssim$ in E by defining $u \lesssim v$ if and only if $v - u \in K$. Now, we define an operator T on K by

$$(Tu)(t) = \int_0^1 G(t, s)f(s, u(s), u'(s))ds, \quad t \in [0, 1]. \quad (2.10)$$

Obviously, fixed points of T are monotone and nonnegative solutions of the BVP (1.3).

Lemma 2.3. $T : K \rightarrow K$ is completely continuous.

Proof. First, we prove $T(K) \subset K$. Suppose that $u \in K$. In view of Lemma 2.2, on the one hand,

$$\begin{aligned} 0 &\leq (Tu)(t) \\ &= \int_0^1 G(t, s) f(s, u(s), u'(s)) ds \\ &\leq \frac{1}{2} (2t - t^2) \int_0^1 s f(s, u(s), u'(s)) ds \\ &\leq \frac{1}{2} \int_0^1 s f(s, u(s), u'(s)) ds, \quad t \in [0, 1], \end{aligned} \tag{2.11}$$

which shows that

$$\|Tu\|_\infty \leq \frac{1}{2} \int_0^1 s f(s, u(s), u'(s)) ds. \tag{2.12}$$

On the other hand,

$$\begin{aligned} (Tu)(t) &= \int_0^1 G(t, s) f(s, u(s), u'(s)) ds \\ &\geq \frac{1}{3} t^2 \int_0^1 s f(s, u(s), u'(s)) ds, \quad t \in [0, 1], \end{aligned} \tag{2.13}$$

which together with (2.12) implies that

$$(Tu)(t) \geq \frac{2}{3} t^2 \|Tu\|_\infty, \quad t \in [0, 1]. \tag{2.14}$$

Again, by Lemma 2.2, we have

$$(Tu)'(t) = \int_0^1 \frac{\partial G(t, s)}{\partial t} f(s, u(s), u'(s)) ds \geq 0, \quad t \in [0, 1]. \tag{2.15}$$

So, it follows from (2.14), (2.15) and the definition of K that $T(K) \subset K$.

Next, we show that T is a compact operator. Let $D \subset K$ be a bounded set. Then, there exists $M_1 > 0$ such that $\|u\| \leq M_1$ for any $u \in D$. For any $\{y_k\}_{k=1}^\infty \subset T(D)$, there exist $\{x_k\}_{k=1}^\infty \subset D$ such that $y_k = Tx_k$. Denote

$$M_2 = \sup \{ f(t, x, y) : (t, x, y) \in [0, 1] \times [0, M_1] \times [0, M_1] \}. \tag{2.16}$$

Then, for any positive integer k , it follows from Lemma 2.2 that

$$\begin{aligned}
 |y_k(t)| &= |(Tx_k)(t)| \\
 &= \left| \int_0^1 G(t,s) f(s, x_k(s), x'_k(s)) ds \right| \\
 &\leq \frac{1}{2} (2t - t^2) \int_0^1 s f(s, x_k(s), x'_k(s)) ds \\
 &\leq \frac{M_2}{4}, \quad t \in [0, 1],
 \end{aligned} \tag{2.17}$$

which indicates that $\{y_k\}_{k=1}^\infty$ is uniformly bounded. Similarly, we have

$$\begin{aligned}
 |y'_k(t)| &= |(Tx_k)'(t)| \\
 &= \left| \int_0^1 \frac{\partial G(t,s)}{\partial t} f(s, x_k(s), x'_k(s)) ds \right| \\
 &\leq (1-t) \int_0^1 s f(s, x_k(s), x'_k(s)) ds \\
 &\leq \frac{M_2}{2}, \quad t \in [0, 1].
 \end{aligned} \tag{2.18}$$

This shows that $\{y'_k\}_{k=1}^\infty$ is uniformly bounded, which implies that $\{y_k\}_{k=1}^\infty$ is equicontinuous. By Arzela-Ascoli theorem, we know that $\{y_k\}_{k=1}^\infty$ has a convergent subsequence in $C[0, 1]$. Without loss of generality, we may assume that $\{y_k\}_{k=1}^\infty$ converges in $C[0, 1]$.

On the other hand, for any $\epsilon > 0$, by the uniform continuity of $\partial G(t, s)/\partial t$, we know that there exists a $\delta > 0$ such that for any $t_1, t_2 \in [0, 1]$ with $|t_1 - t_2| < \delta$,

$$\left| \frac{\partial G(t_1, s)}{\partial t} - \frac{\partial G(t_2, s)}{\partial t} \right| < \frac{\epsilon}{M_2 + 1}, \quad s \in [0, 1]. \tag{2.19}$$

So, for any positive integer k , $t_1, t_2 \in [0, 1]$ with $|t_1 - t_2| < \delta$, we get

$$\begin{aligned}
 |y'_k(t_1) - y'_k(t_2)| &= |(Tx_k)'(t_1) - (Tx_k)'(t_2)| \\
 &= \int_0^1 \left| \frac{\partial G(t_1, s)}{\partial t} - \frac{\partial G(t_2, s)}{\partial t} \right| f(s, x_k(s), x'_k(s)) ds \\
 &< \frac{M_2 \epsilon}{M_2 + 1} \\
 &< \epsilon,
 \end{aligned} \tag{2.20}$$

which shows that $\{y'_k\}_{k=1}^\infty$ is equicontinuous. Again, it follows from Arzela-Ascoli theorem that $\{y'_k\}_{k=1}^\infty$ has a convergent subsequence in $C[0, 1]$. Therefore, $\{y_k\}_{k=1}^\infty$ has a convergent subsequence in K .

Finally, we prove that T is continuous. Suppose that $u_m, u \in K$ and $\|u_m - u\| \rightarrow 0$ ($m \rightarrow \infty$). Then, there exists $M_3 > 0$ such that for any positive integer m , $\|u_m\| \leq M_3$. Denote

$$M_4 = \sup\{f(t, x, y) : (t, x, y) \in [0, 1] \times [0, M_3] \times [0, M_3]\}. \quad (2.21)$$

Then, for any positive integer m and $t \in [0, 1]$, by Lemma 2.2, we have

$$\begin{aligned} G(t, s)f(s, u_m(s), u'_m(s)) &\leq \frac{M_4}{2}(2t - t^2)s \leq \frac{M_4}{2}s, \quad s \in [0, 1], \\ \frac{\partial G(t, s)}{\partial t}f(s, u_m(s), u'_m(s)) &\leq M_4(1 - t)s \leq M_4s, \quad s \in [0, 1]. \end{aligned} \quad (2.22)$$

So, it follows from Lebesgue dominated convergence theorem that

$$\begin{aligned} \lim_{m \rightarrow \infty} (Tu_m)(t) &= \lim_{m \rightarrow \infty} \int_0^1 G(t, s)f(s, u_m(s), u'_m(s))ds \\ &= \int_0^1 G(t, s)f(s, u(s), u'(s))ds \\ &= (Tu)(t), \quad t \in [0, 1], \\ \lim_{m \rightarrow \infty} (Tu_m)'(t) &= \lim_{m \rightarrow \infty} \int_0^1 \frac{\partial G(t, s)}{\partial t}f(s, u_m(s), u'_m(s))ds \\ &= \int_0^1 \frac{\partial G(t, s)}{\partial t}f(s, u(s), u'(s))ds \\ &= (Tu)'(t), \quad t \in [0, 1], \end{aligned} \quad (2.23)$$

which implies that T is continuous. Therefore, $T : K \rightarrow K$ is completely continuous. $\square$

3. Main Results

Theorem 3.1. Assume that $f(t, 0, 0) \neq 0$ for $t \in (0, 1)$, and there exists a constant $a > 0$ such that

$$f(t, u_1, v_1) \leq f(t, u_2, v_2) \leq 2a, \quad 0 \leq t \leq 1, \quad 0 \leq u_1 \leq u_2 \leq a, \quad 0 \leq v_1 \leq v_2 \leq a. \quad (3.1)$$

If we construct a iterative sequence $v_{n+1} = Tv_n$, $n = 0, 1, 2, \dots$, where $v_0(t) \equiv 0$ for $t \in [0, 1]$, then $\{v_n\}_{n=1}^\infty$ converges to v^* in $C^1[0, 1]$, which is a monotone positive solution of the BVP (1.3) and satisfy

$$0 < v^*(t) \leq a \quad \text{for } t \in (0, 1], \quad 0 \leq (v^*)'(t) \leq a \quad \text{for } t \in [0, 1]. \quad (3.2)$$

Proof. Let $K_a = \{u \in K : \|u\| \leq a\}$. We assert that $T : K_a \rightarrow K_a$. In fact, if $u \in K_a$, then

$$0 \leq u(s) \leq \max_{0 \leq s \leq 1} u(s) \leq \|u\| \leq a, \quad 0 \leq u'(s) \leq \max_{0 \leq s \leq 1} u'(s) \leq \|u\| \leq a, \quad \text{for } s \in [0, 1], \quad (3.3)$$

which together with the condition (3.1) and Lemma 2.2 implies that

$$\begin{aligned} 0 \leq (Tu)(t) &= \int_0^1 G(t, s) f(s, u(s), u'(s)) ds < a, \quad t \in [0, 1], \\ 0 \leq (Tu)'(t) &= \int_0^1 \frac{\partial G(t, s)}{\partial t} f(s, u(s), u'(s)) ds \leq a, \quad t \in [0, 1]. \end{aligned} \quad (3.4)$$

Hence, we have shown that $T : K_a \rightarrow K_a$.

Now, we assert that $\{v_n\}_{n=1}^\infty$ converges to v^* in $C^1[0, 1]$, which is a monotone positive solution of the BVP (1.3) and satisfies

$$0 < v^*(t) \leq a \quad \text{for } t \in (0, 1], \quad 0 \leq (v^*)'(t) \leq a \quad \text{for } t \in [0, 1]. \quad (3.5)$$

Indeed, in view of $v_0 \in K_a$ and $T : K_a \rightarrow K_a$, we have $v_n \in K_a$, $n = 1, 2, \dots$. Since the set $\{v_n\}_{n=0}^\infty$ is bounded and T is completely continuous, we know that the set $\{v_n\}_{n=1}^\infty$ is relatively compact.

In what follows, we prove that $\{v_n\}_{n=0}^\infty$ is monotone by induction. First, by Lemma 2.2, we have

$$\begin{aligned} &\frac{1}{3}t^2 \int_0^1 s f(s, 0, 0) ds \\ &\leq v_1(t) - v_0(t) = (Tv_0)(t) = \int_0^1 G(t, s) f(s, 0, 0) ds \leq \frac{1}{2} \int_0^1 s f(s, 0, 0) ds, \quad t \in [0, 1], \end{aligned} \quad (3.6)$$

which implies that

$$v_1(t) - v_0(t) \geq \frac{2}{3}t^2 \|v_1 - v_0\|_\infty, \quad t \in [0, 1]. \quad (3.7)$$

At the same time, it is obvious that

$$v_1'(t) - v_0'(t) = v_1'(t) = (Tv_0)'(t) = \int_0^1 \frac{\partial G(t, s)}{\partial t} f(s, 0, 0) ds \geq 0, \quad t \in [0, 1]. \quad (3.8)$$

It follows from (3.7) and (3.8) that $v_1 - v_0 \in K$, which shows that $v_0 \lesssim v_1$. Next, we assume that $v_{k-1} \lesssim v_k$. Then, in view of Lemma 2.2 and (3.1), we have

$$\begin{aligned}
 v_{k+1}(t) - v_k(t) &= \int_0^1 G(t, s) [f(s, v_k(s), v'_k(s)) - f(s, v_{k-1}(s), v'_{k-1}(s))] ds \\
 &\leq \frac{1}{2} \int_0^1 s [f(s, v_k(s), v'_k(s)) - f(s, v_{k-1}(s), v'_{k-1}(s))] ds, \quad t \in [0, 1], \\
 v_{k+1}(t) - v_k(t) &= \int_0^1 G(t, s) [f(s, v_k(s), v'_k(s)) - f(s, v_{k-1}(s), v'_{k-1}(s))] ds \\
 &\geq \frac{1}{3} t^2 \int_0^1 s [f(s, v_k(s), v'_k(s)) - f(s, v_{k-1}(s), v'_{k-1}(s))] ds, \quad t \in [0, 1],
 \end{aligned} \tag{3.9}$$

which implies that

$$v_{k+1}(t) - v_k(t) \geq \frac{2}{3} t^2 \|v_{k+1} - v_k\|_\infty, \quad t \in [0, 1]. \tag{3.10}$$

At the same time, by Lemma 2.2 and (3.1), we also have

$$v'_{k+1}(t) - v'_k(t) = \int_0^1 \frac{\partial G(t, s)}{\partial t} [f(s, v_k(s), v'_k(s)) - f(s, v_{k-1}(s), v'_{k-1}(s))] ds \geq 0, \quad t \in [0, 1]. \tag{3.11}$$

It follows from (3.10) and (3.11) that $v_{k+1} - v_k \in K$, which indicates that $v_k \lesssim v_{k+1}$. Thus, we have shown that $v_n \lesssim v_{n+1}$, $n = 0, 1, 2, \dots$

Since $\{v_n\}_{n=1}^\infty$ is relatively compact and monotone, there exists a $v^* \in K_a$ such that $\|v_n - v^*\| \rightarrow 0$ ($n \rightarrow \infty$), which together with the continuity of T and the fact that $v_{n+1} = Tv_n$ implies that $v^* = Tv^*$. Moreover, in view of $f(t, 0, 0) \neq 0$ for $t \in (0, 1)$, we know that the zero function is not a solution of the BVP (1.3). Thus, $\|v^*\|_\infty > 0$. So, it follows from $v^* \in K_a$ that

$$0 < v^*(t) \leq a \quad \text{for } t \in (0, 1], \quad 0 \leq (v^*)'(t) \leq a \quad \text{for } t \in [0, 1]. \tag{3.12}$$

□

4. An Example

Example 4.1. Consider the BVP

$$\begin{aligned}
 u^{(4)}(t) &= \frac{1}{2} t u(t) + \frac{1}{8} (u'(t))^2 + 1, \quad t \in (0, 1), \\
 u(0) &= u'(1) = u''(0) = u'''(1) = 0.
 \end{aligned} \tag{4.1}$$

If we let $f(t, u, v) = (1/2)tu + (1/8)v^2 + 1$ for $(t, u, v) \in [0, 1] \times [0, +\infty) \times [0, +\infty)$, then all the hypotheses of Theorem 3.1 are fulfilled with $a = 1$. It follows from Theorem 3.1 that the BVP (4.1) has a monotone positive solution v^* satisfying

$$0 < v^*(t) \leq 1 \quad \text{for } t \in (0, 1], \quad 0 \leq (v^*)'(t) \leq 1 \quad \text{for } t \in [0, 1]. \quad (4.2)$$

Moreover, the iterative scheme is

$$\begin{aligned} v_0(t) &\equiv 0, \quad t \in [0, 1], \\ v_{n+1}(t) &= \frac{1}{6} \int_0^t s(6t - 3t^2 - s^2) \left(\frac{1}{2} s v_n(s) + \frac{1}{8} (v'_n(s))^2 + 1 \right) ds \\ &\quad + \frac{1}{6} \int_t^1 t(6s - 3s^2 - t^2) \left(\frac{1}{2} s v_n(s) + \frac{1}{8} (v'_n(s))^2 + 1 \right) ds, \quad t \in [0, 1], \quad n = 0, 1, 2, \dots \end{aligned} \quad (4.3)$$

The first, second, third, and fourth terms of the scheme are as follows:

$$\begin{aligned} v_0(t) &\equiv 0, \\ v_1(t) &= \frac{1}{3}t - \frac{1}{6}t^3 + \frac{1}{24}t^4, \\ v_2(t) &= \frac{73}{216}t + \frac{5}{144}t^2 - \frac{79}{432}t^3 + \frac{1}{64}t^4 + \frac{25}{1152}t^5 - \frac{23}{8640}t^6 - \frac{1}{192}t^7 + \frac{47}{11520}t^8 - \frac{11}{8640}t^9 + \frac{1}{6912}t^{10}, \\ v_3(t) &= \frac{378577}{1119744}t + \frac{6935}{186624}t^2 - \frac{404629}{2239488}t^3 + \frac{60361}{8957952}t^4 + \frac{71689}{2985984}t^5 + \frac{110633}{14929920}t^6 - \frac{22181}{1492992}t^7 \\ &\quad + \frac{373759}{119439360}t^8 + \frac{2467627}{477757440}t^9 - \frac{2449943}{716636160}t^{10} - \frac{800341}{1592524800}t^{11} + \frac{16256149}{11466178560}t^{12} \\ &\quad - \frac{3556357}{7166361600}t^{13} - \frac{655267}{4777574400}t^{14} + \frac{70313}{477757440}t^{15} - \frac{25471}{2388787200}t^{16} - \frac{363911}{9555148800}t^{17} \\ &\quad + \frac{7781}{298598400}t^{18} - \frac{25973}{2866544640}t^{19} + \frac{8917}{4777574400}t^{20} - \frac{31}{143327232}t^{21} + \frac{25}{2293235712}t^{22}. \end{aligned} \quad (4.4)$$

Acknowledgment

This work is supported by the National Natural Science Foundation of China (10801068).

References

- [1] A. R. Aftabizadeh, "Existence and uniqueness theorems for fourth-order boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 116, no. 2, pp. 415–426, 1986.
- [2] R. P. Agarwal, "On fourth order boundary value problems arising in beam analysis," *Differential and Integral Equations*, vol. 2, no. 1, pp. 91–110, 1989.

- [3] Z. Bai, "The upper and lower solution method for some fourth-order boundary value problems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 67, no. 6, pp. 1704–1709, 2007.
- [4] Z. Bai and H. Wang, "On positive solutions of some nonlinear fourth-order beam equations," *Journal of Mathematical Analysis and Applications*, vol. 270, no. 2, pp. 357–368, 2002.
- [5] J. Ehme, P. W. Eloe, and J. Henderson, "Upper and lower solution methods for fully nonlinear boundary value problems," *Journal of Differential Equations*, vol. 180, no. 1, pp. 51–64, 2002.
- [6] J. R. Graef and B. Yang, "Positive solutions of a nonlinear fourth order boundary value problem," *Communications on Applied Nonlinear Analysis*, vol. 14, no. 1, pp. 61–73, 2007.
- [7] C. P. Gupta, "Existence and uniqueness theorems for the bending of an elastic beam equation," *Applicable Analysis*, vol. 26, no. 4, pp. 289–304, 1988.
- [8] Y. Li, "Positive solutions of fourth-order boundary value problems with two parameters," *Journal of Mathematical Analysis and Applications*, vol. 281, no. 2, pp. 477–484, 2003.
- [9] B. Liu, "Positive solutions of fourth-order two point boundary value problems," *Applied Mathematics and Computation*, vol. 148, no. 2, pp. 407–420, 2004.
- [10] J. Liu and W. Xu, "Positive solutions for some beam equation boundary value problems," *Boundary Value Problems*, vol. 2009, Article ID 393259, 9 pages, 2009.
- [11] R. Ma, "Positive solutions of fourth-order two-point boundary value problems," *Annals of Differential Equations*, vol. 15, no. 3, pp. 305–313, 1999.
- [12] R. Ma and H. Wang, "On the existence of positive solutions of fourth-order ordinary differential equations," *Applicable Analysis*, vol. 59, no. 1–4, pp. 225–231, 1995.
- [13] Q. Yao, "Positive solutions for eigenvalue problems of fourth-order elastic beam equations," *Applied Mathematics Letters*, vol. 17, no. 2, pp. 237–243, 2004.
- [14] Q. Yao, "Existence and multiplicity of positive solutions to a singular elastic beam equation rigidly fixed at both ends," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 69, no. 8, pp. 2683–2694, 2008.
- [15] Q. Yao, "Successively iterative technique of a classical elastic beam equation with Carathéodory nonlinearity," *Acta Applicandae Mathematicae*, vol. 108, no. 2, pp. 385–394, 2009.
- [16] X. Zhang, "Existence and iteration of monotone positive solutions for an elastic beam equation with a corner," *Nonlinear Analysis. Real World Applications*, vol. 10, no. 4, pp. 2097–2103, 2009.
- [17] B. Ahmad and Juan J. Nieto, "The monotone iterative technique for three-point second-order integrodifferential boundary value problems with p -Laplacian," *Boundary Value Problems*, vol. 2007, Article ID 57481, 9 pages, 2007.
- [18] Q. Yao, "Monotone iterative technique and positive solutions of Lidstone boundary value problems," *Applied Mathematics and Computation*, vol. 138, no. 1, pp. 1–9, 2003.
- [19] Q. Yao, "Monotonically iterative method of nonlinear cantilever beam equations," *Applied Mathematics and Computation*, vol. 205, no. 1, pp. 432–437, 2008.