Research Article

On the Polynomial Basis of $GF(2^n)$ Having a Small Number of Trace-One Elements

Jiantao Wang,¹ Dong Zheng,² and Zheng Huang¹

¹School of Information Security Engineering, Shanghai Jiao Tong University, Shanghai 200240, China

²National Engineering Laboratory for Wireless Security, Xi'an University of Posts and Telecommunications, Xi'an 710121, China

Correspondence should be addressed to Dong Zheng; dzhengcrypto@hotmail.com

Received 15 March 2016; Revised 9 August 2016; Accepted 20 October 2016

Academic Editor: Fazal M. Mahomed

Copyright © 2016 Jiantao Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the Galois fields $GF(2^n)$, a polynomial basis with a small number of trace-one elements is desirable for its convenience in computing. To find new irreducible polynomials $g(x)$ over $GF(2)$ with this property, we research into the auxiliary polynomial $f(x) = (x+1)g(x)$ with roots $\{1, \alpha_1, \alpha_2, \dots, \alpha_n\}$, such that the symmetric polynomials $s_k = 1 + \alpha_1^k + \alpha_2^k + \dots + \alpha_n^k$ are relative to the symmetric polynomials of $g(x)$. We introduce a new class of polynomials with the number “1” occupying most of the values in its s_k . This indicates that the number “0” occupies most of the values of the traces of the elements $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$. This new class of polynomial gives us an indirect way to find irreducible polynomials having a small number of trace-one elements in their polynomial bases.

1. Introduction

Let $GF(2^n)$ denote the finite field with 2^n elements. Let α be a root of an irreducible polynomial $g(x)$ of degree n over $GF(2)$. The set $\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$ forms a basis of $GF(2^n)$, called a polynomial basis. An element $\beta \in GF(2^n)$ can be represented as

$$\beta = a_0 + a_1\alpha + a_2\alpha^2 + \dots + a_{n-1}\alpha^{n-1}. \quad (1)$$

The trace of β is defined as

$$\text{tr}(\beta) = \beta + \beta^2 + \dots + \beta^{2^{n-1}}, \quad (2)$$

which is the linear combination of the traces of elements of the polynomial basis; that is,

$$\begin{aligned} \text{tr}(\beta) &= a_0 \text{tr}(1) + a_1 \text{tr}(\alpha) + a_2 \text{tr}(\alpha^2) + \dots \\ &\quad + a_{n-1} \text{tr}(\alpha^{n-1}). \end{aligned} \quad (3)$$

If the number of basis elements α^k for which $\text{tr}(\alpha^k) = 1$ is small, the computation of $\text{tr}(\beta)$ in software and the implementation in hardware are faster. It is benefit for halving a point on an elliptic curve over $GF(2^n)$ [1, 2] or generating pseudo random sequences using elliptic curves [3]. Thus it

is of interest to find irreducible polynomials having a small number of trace-one elements in their polynomial bases. As the value of $\text{tr}(\alpha^k)$ equals the symmetric polynomial $\sum_{t=0}^{n-1} (\alpha^{2^t})^k$, the authors in [4] got the conditions for trinomials and pentanomials to have this property using the Newton function for symmetric polynomials [5, 6].

We presented an indirect strategy that is different from the above one, which is inspired by extending the work of [7] in [8]. We still use the Newton function to investigate the polynomial $f(x)$ with the following two properties. First, the polynomial $f(x)$ should have the factorization $f(x) = (x+1)g(x)$ with $g(x)$ irreducible. Second, let $\beta_0 = 1, \beta_1, \beta_2, \dots, \beta_n$ be the roots of $f(x)$, the number of different values of the symmetric polynomials, $s_k = \sum_{t=0}^n \beta_t^k$ equaling one should be as many as possible. With these two properties, we can identify $g(x)$ as having less trace-one elements in its polynomial basis. A class of polynomials $f(x)$ suiting our demands is introduced, and we showed their existence by numerical experiments.

The paper is organized as follows. In Section 2 we give some basic definitions which are needed for our result. The new class of polynomials is introduced in Section 3. Section 4 draws the conclusion of our work.

2. Preliminaries

Let $g(x)$ be a polynomial over $\text{GF}(2)$ of degree n whose roots are $\alpha_1, \alpha_2, \dots, \alpha_n$. The elementary symmetric polynomials $a_1, a_2, \dots, a_n$ are the coefficients of $g(x)$:

$$\begin{aligned} g(x) &= (x - \alpha_1)(x - \alpha_2) \cdots (x - \alpha_n) \\ &= x^n - a_1 x^{n-1} + a_2 x^{n-2} - \cdots + (-1)^{n-1} a_{n-1} x \\ &\quad + (-1)^n a_n. \end{aligned} \quad (4)$$

Another particular kind of symmetric polynomial is defined as

$$s_k = s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = \alpha_1^k + \alpha_2^k + \cdots + \alpha_n^k, \quad (5)$$

where k is an integer.

The Newton function is a relation between s_k and the coefficients of the polynomial; it is defined as follows [5, p.12]:

$$\begin{aligned} s_k - s_{k-1}a_1 + \cdots + (-1)^i s_{k-i}a_i + \cdots + (-1)^{k-1} s_1 a_{k-i} \\ + (-1)^k k a_k = 0, \quad a_j = 0 \text{ for } j > n. \end{aligned} \quad (6)$$

It is widely known that if $g(x)$ is irreducible, then the traces of α^k equal the value of s_k .

Let $f(x) = (x+1)g(x)$; the roots of $f(x)$ are $\beta_0 = 1, \beta_1 = \alpha_1, \beta_2 = \alpha_2, \dots, \beta_n = \alpha_n$. We have

$$\begin{aligned} s_k(\beta_0, \beta_1, \beta_2, \dots, \beta_n) &= \beta_0^k + \beta_1^k + \beta_2^k + \cdots + \beta_n^k \\ &= 1 + s_k(\alpha_1, \alpha_2, \dots, \alpha_n) \\ &= 1 + \text{tr}(\alpha^k). \end{aligned} \quad (7)$$

So when more ones appeared in the set of the values of $s_k(\beta_0, \beta_1, \dots, \beta_n)$, more zeros appeared in the traces of elements of the polynomial basis of $g(x)$.

3. The Three Piece Polynomials

We first introduce a class of polynomials $f(x)$ with roots $\alpha_1, \alpha_2, \dots, \alpha_n$ over $\text{GF}(2)$ fitting the conditions that the value set of $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$ has plenty of ones and then search for the polynomials with the decomposition $f(x) = (x+1)g(x)$ where $g(x)$ is irreducible.

3.1. The Three Piece Polynomials

Definition 1. Let

$$f(x) = a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \cdots + a_{n-1} x + a_n \quad (8)$$

be a polynomial over $\text{GF}(2)$ with nonzero constant term. If $(a_0, a_1, \dots, a_n) = (\text{ones}(n_1), \text{zeros}(n_2), \text{ones}(n_3))$ where $\text{ones}(k)$ and $\text{zeros}(k)$ mean k consecutive bits of ones and zeros, respectively, we call $f(x)$ three-piece polynomial over $\text{GF}(2)$ and denote it by its coefficients as $(n_1, n_2, n_3)_{\text{TP}}$.

Lemma 2. Let $f(x) = a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \cdots + a_{n-1} x + a_n$ be a three-piece polynomial with coefficients $(n_1, n_2, n_3)_{\text{TP}}$ over $\text{GF}(2)$. Let $\alpha_1, \alpha_2, \dots, \alpha_n$ be the roots of $f(x)$; if $n_1 \geq 2$, then

$$s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = 1, \quad 0 < k < n_1. \quad (9)$$

Proof. If $n_1 \geq 2$, then $a_0 = a_1 = \cdots = a_{n_1-1} = 1$, and the Newton formula tells us that $s_1 = a_1 = 1$ and $s_2 = s_1 a_1 + 2a_2 = 1$. Suppose $s_k = 1$ is true for $1 \leq k \leq n_1 - 2$; then $a_{k+1} = 1$ and

$$\begin{aligned} s_k &= a_1 s_{k-1} + a_2 s_{k-2} + \cdots + a_{k-1} s_1 + k a_k \\ &= (k-1) \cdot 1 + k = 1. \end{aligned} \quad (10)$$

We have

$$\begin{aligned} s_{k+1} &= a_1 s_k + a_2 s_{k-1} + \cdots + a_k s_1 + (k+1) a_{k+1} \\ &= k \cdot 1 + k + 1 = 1. \end{aligned} \quad (11)$$

□

We mainly divide our theorems into two parts according to the parity of n_1 . First, let us consider the condition that n_1 is even.

Theorem 3. Let $f(x)$ be denoted as above. If both n_1 and n_2 are even, then the values of functions $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, $0 \leq k \leq n-1$ all equal one.

Proof. From Lemma 2, we know that $s_k = 1$ for $1 \leq k \leq n_1 - 1$. We have

$$\begin{aligned} s_{n_1} &= a_1 s_{n_1-1} + a_2 s_{n_1-2} + \cdots + a_{n_1-1} s_1 + n_1 a_{n_1} \\ &= (n_1 - 1) \cdot 1 + 0 = 1, \end{aligned} \quad (12)$$

as n_1 is even and $a_{n_1} = 0$. Suppose $s_{n_1+k} = 1$ is true for $0 \leq k \leq n_2 - 2$; then

$$\begin{aligned} s_{n_1+k+1} &= a_1 s_{n_1+k} + a_2 s_{n_1+k-1} + \cdots + a_{n_1-1} s_{k+2} \\ &\quad + a_{n_1} s_{k+1} + \cdots + a_{n_1+k} s_1 \\ &\quad + (n_1 + k + 1) a_{n_1+k+1} = (n_1 - 1) \cdot 1 + 0 \\ &= 1, \end{aligned} \quad (13)$$

as $a_{n_1+k+1} = 0$. Here we come to the next even number $n_1 + n_2$ with $a_{n_1+n_2} = 1$,

$$\begin{aligned} s_{n_1+n_2} &= a_1 s_{n_1+n_2-1} + a_2 s_{n_1+n_2-2} + \cdots + a_{n_1+n_2-1} s_1 \\ &\quad + (n_1 + n_2) a_{n_1+n_2} = (n_1 - 1) \cdot 1 + 0 = 1. \end{aligned} \quad (14)$$

Suppose $s_{n_1+n_2+k} = 1$ is true for $0 \leq k \leq n_3 - 2$; then

$$\begin{aligned} s_{n_1+n_2+k+1} &= a_1 s_{n_1+n_2+k} + a_2 s_{n_1+n_2+k-1} + \cdots \\ &\quad + a_{n_1-1} s_{n_2+k+2} + a_{n_1} s_{n_2+k+1} + \cdots \\ &\quad + a_{n_1+n_2+k} s_1 \\ &\quad + (n_1 + n_2 + k + 1) a_{n_1+n_2+k+1} \\ &= (n_1 - 1) \cdot 1 + (k + 1) \cdot 1 \\ &\quad + (n_1 + n_2 + k + 1) = 1, \end{aligned} \quad (15)$$

as $a_{n_1+n_2} = a_{n_1+n_2+1} = \cdots = a_{n_1+n_2+n_3-1} = 1$. □

In the following proofs, we still focus on the changes of the value in the summations of the symmetric polynomial s_k . To simplify narration, we will divide the summations in s_k into four parts $\{D_1, D_2, D_3, D_4\}$ according to the subscript of a_i . That is, the summations $a_i s_{k-i}$ with $1 \leq i < n_1$ are contained in D_1 , the summation $a_i s_{k-i}$ with $n_1 \leq i < n_1 + n_2$ comprises D_2 , the summations $a_i s_{k-i}$ in D_3 are with $n_1 + n_2 \leq i < n$, and the ending part $D_4 = k a_k$.

Theorem 4. Let $f(x)$ be denoted as above. If n_1 is even, n_2 is odd, and $n_1 + n_2 > n_3$, then the values of functions $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, $0 \leq k \leq n-1$, are

$$s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = \begin{cases} 1, & \text{if } 1 \leq k \leq n_1 + n_2 - 1, \\ 0, & \text{if } k \geq n_1 + n_2, \quad k - n_1 - n_2 \equiv 0 \pmod{n_1}, \\ 1, & \text{if } k \geq n_1 + n_2, \quad k - n_1 - n_2 > 0 \pmod{n_1}. \end{cases} \quad (16)$$

Proof. The first case is contained in Theorem 3.

For the remaining cases, we declare they are periodic. We first show one period in the following. That is, for $n_1 + n_2 \leq k \leq n_1 + n_2 + n_1$ we have

$$\begin{aligned} s_k &= \sum_{i \leq n_1-1} a_i s_{k-i} + \sum_{n_1=i}^{n_1+n_2-1} a_i s_{k-i} + \sum_{i=n_1+n_2}^{k-1} a_i s_{k-i} + k a_k \\ &= \sum_{i \leq n_1-1} s_{k-i} + \sum_{i=n_1}^{n_1+n_2-1} 0 \cdot s_{k-i} + \sum_{i=n_1+n_2}^{k-1} s_{k-i} + k \\ &= D_1 + D_2 + D_3 + D_4. \end{aligned} \quad (17)$$

By the assumption $n_1 + n_2 > n_3$, we get $s_{k-i} = 1$ for $n_1 + n_2 \leq i \leq k-1 < n_1 + n_2 + n_1$ and $D_2 + D_3 + D_4 = 0 + k - n_1 - n_2 + k = 1$; these simplify the above equation to $s_k = D_1 + 1$.

For the value of $s_{n_1+n_2+1}$, as $s_{n_1+n_2}$ appears in the D_1 of $s_{n_1+n_2+1}$ and its value is " 0^3 " we have $s_{n_1+n_2+1} = 1$.

We go on our proof by induction. Suppose $s_t = 1$ holds for all t satisfying $n_1 + n_2 < t < k < n_1 + n_2 + n_1$; only the summation with $s_{n_1+n_2}$ equals zero in D_1 of s_k , so $s_k = n_1 - 2 + 1 = 1$.

For $s_{n_1+n_2+n_1}$, as $s_{n_1+n_2}$ goes into the summation in D_2 , the value of the summation D_1 turns back to $n_1 - 1$ and we have $s_{n_1+n_2+n_1} = 0$. Now, we finished computing the value of s_k in one period in our declaration, which is $n_1 + n_2 < k \leq n_1 + n_2 + n_1$.

The condition $n_1 + n_2 > n_3$ guarantees the subsequent computations can still use $s_k = D_1 + 1$ for (17), as $s_{k-i} = 1$ and $D_2 + D_3 + D_4 = 0 + k - n_1 - n_2 + k = 1$ still holds. So the above method will work for the next periods, and our declaration holds. $\square$

Next we will deal with the situation that n_1 is odd, which is more complicated. It has been proved that $s_k = 1$ for $k \leq n_1 - 1$ in Lemma 2.

Lemma 5. Let $f(x)$ be denoted as above. If n_1 is odd, then the values of functions $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, $n_1 \leq k \leq n_1 + n_2 - 1$, are

$$s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = \begin{cases} 0, & k - n_1 \equiv 0 \pmod{n_1}, \\ 1, & k - n_1 > 0 \pmod{n_1}. \end{cases} \quad (18)$$

Proof. We follow the idea in the proof of Theorem 4. For $n_1 \leq k \leq n_1 + n_2 - 1$, let

$$\begin{aligned} s_k &= \sum_{i \leq n_1-1} a_i s_{k-i} + \sum_{i=n_1}^{k-1} a_i s_{k-i} + k a_k \\ &= \sum_{i \leq n_1-1} s_{k-i} + \sum_{i=n_1}^{k-1} 0 \cdot s_{k-i} + k \cdot 0 = D_1. \end{aligned} \quad (19)$$

We have $s_{n_1} = n_1 - 1 = 0$ and it fixes the value of D_1 in s_t for $n_1 < t \leq n_1 + n_1 - 1$ to be $n_1 - 2$ which makes $s_t = 1$. If $n_1 < n_2$ then $s_{n_1+n_1} = n_1 - 1 = 0$ and the next period starts; otherwise the equation holds within one period. $\square$

In the following, we will deal with s_k with $k \geq n_1 + n_2$; remember that n_1 is odd. We define a comparative sequence here to simplify the narration. Let b_i be the coefficient of a polynomial $m(x)$ with

$$b_i = \begin{cases} a_i, & 1 \leq i < n_1 + n_2, \\ 0, & n_1 + n_2 \leq i, \end{cases} \quad (20)$$

and then s_k of $m(x)$, which is denoted by t_k , follows the period shown in Lemma 5. That is,

$$t_k = D_1 = \begin{cases} 0, & k - n_1 \equiv 0 \pmod{n_1}, \\ 1, & k - n_1 > 0 \pmod{n_1}. \end{cases} \quad (21)$$

Theorem 6. Let $f(x)$ and $m(x)$ be denoted as above. If n_1 is odd, n_2 is even, and $n_1 + n_2 > n_3$, then the values of functions $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, $n_1 + n_2 \leq k \leq n_1 + n_2 + n_3 - 1$ have two cases.

$$s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = \begin{cases} 1 - t_k, & k - n_1 - n_2 \equiv 0 \pmod{2n_1}, \\ t_k, & k - n_1 - n_2 > 0 \pmod{2n_1}. \end{cases} \quad (22)$$

Proof. For $n_1 + n_2 \leq k \leq n_1 + n_2 + n_1$, we have

$$\begin{aligned} s_k &= \sum_{i \leq n_1-1} a_i s_{k-i} + \sum_{i=n_1}^{n_1+n_2-1} a_i s_{k-i} + \sum_{i=n_1+n_2}^{k-1} a_i s_{k-i} + k a_k \\ &= D_1 + D_2 + D_3 + D_4 = D_1 + (k - n_1 - n_2) + k \\ &= D_1 + 1. \end{aligned} \quad (23)$$

For $n_1 + n_2 + n_1 \leq k < n_1 + n_2 + 2n_1$, we have $D_3 = k - 1 - n_1 - n_2$ as $s_{n_1} = 0$ enters the summations of D_3 , so $s_k = D_1$. By induction, we have

$$s_k = \begin{cases} D_1 + 1, & 0 \leq k - n_1 - n_2 \pmod{2n_1} < n_1, \\ D_1, & n_1 \leq k - n_1 - n_2 \pmod{2n_1} < 2n_1. \end{cases} \quad (24)$$

The value of s_k for $k - n_1 - n_2 \equiv 0 \pmod{2n_1}$ is opposite to the corresponding t_k .

We have $s_k = t_k$ for $0 < k - n_1 - n_2 \pmod{2n_1} < n_1$ as the value of $s_{n_1+n_2}$ is changed in the D_1 of s_k .

As all the summations in s_k and t_k are the same for $n_1 \leq k - n_1 - n_2 \pmod{2n_1} < 2n_1$, we get $s_k = t_k$ and our claim follows. $\square$

Using the same method, we can also prove the following theorem.

Theorem 7. Let $f(x)$ and $g(x)$ be denoted as above. If n_1 is odd, n_2 is odd, and $n_1 + n_2 > n_3$, then the values of functions $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, $n_1 + n_2 \leq k \leq n_1 + n_2 + n_3 - 1$ have two cases.

$$s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = \begin{cases} 1 - t_k, & k - 2n_1 - n_2 \equiv 0 \pmod{2n_1}, \\ t_k, & k - 2n_1 - n_2 > 0 \pmod{2n_1}, \end{cases} \quad (25)$$

That is, the changed value appears at the position k where $k - n_1 - n_2 \equiv n_1 \pmod{2n_1}$.

TABLE 1: Examples of three piece polynomials $(6, 12, 15)_{\text{TP}}$, $(6, 13, 14)_{\text{TP}}$, $(5, 12, 15)_{\text{TP}}$, and $(5, 13, 14)_{\text{TP}}$ with their coefficients a_i and the value of symmetric polynomials s_k .

$(a_0, a_1, \dots, a_n)$	Polynomial
11111100000000000001111111111111	$(6, 12, 15)_{\text{TP}}$
11111100000000000000111111111111	$(6, 13, 14)_{\text{TP}}$
11111000000000000001111111111111	$(5, 12, 15)_{\text{TP}}$
11111000000000000000111111111111	$(5, 13, 14)_{\text{TP}}$
$(s_1, s_2, \dots, s_{n-1})$	
11111111111111111111111111111111	$(6, 12, 15)_{\text{TP}}$
11111111111111111111011111011110	$(6, 13, 14)_{\text{TP}}$
111101111011110111110111010110	$(5, 12, 15)_{\text{TP}}$
11110111101111011111011011110	$(5, 13, 14)_{\text{TP}}$

Corollary 8. Let $f(x)$ and $g(x)$ be denoted as above. If $n_1 = 1$ and $n_1 + n_2 > n_3$, then the values of function $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, $1 \leq k \leq n_1 + n_2 + n_3 - 1$ are

$$s_k(\alpha_1, \alpha_2, \dots, \alpha_n) = \begin{cases} 0, & \text{if } 1 \leq k \leq n_1 + n_2 - 1, \\ n_2 + 1 \pmod{2}, & \text{if } k \geq n_1 + n_2, k \equiv n_2 + 1 \pmod{2}, \\ n_2 \pmod{2}, & \text{if } k \geq n_1 + n_2, k \equiv n_2 \pmod{2}. \end{cases} \quad (26)$$

Proof. As all the coefficients a_k , $1 \leq k \leq n_1 + n_2 - 1$ equal zero, the first case is straightforward. For $n_1 + n_2 \leq k \leq n_1 + n_2 + n_3 - 1$, we have $s_k = D_1 + D_2 + D_3 + D_4 = D_4 = k$ and the corollary follows. $\square$

We have discussed the four cases concerning the three-piece polynomials over $\text{GF}(2)$. The condition $n_1 + n_2 > n_3$ guarantees that the above cases cover all the possible kinds of three-piece polynomials. The number “1” does occupy a great proportion of the values of $s_k(\alpha_1, \alpha_2, \dots, \alpha_n)$, as the examples shown in Table 1.

3.2. Experiment Results. Next, we search for the three-piece polynomial $f(x)$ whose decomposition is $f(x) = (x + 1)g(x)$ with $g(x)$ irreducible. As there exists a trinomial or pentanomial basis for $\text{GF}(2^n)$ having at most four elements of trace one for each $n \in [2, 10000]$ [4], the polynomial $g(x)$ with less number of trace-one elements that appeared in its polynomial basis is acceptable.

With the formula $\text{tr}(1) \equiv n \pmod{2}$, we choose the number of trace-one elements to be at most four, and the result showed that the three-piece polynomials fitting these requirements exist for a great number of n , and we guess they exist for $n \geq 40$. The proof for this conjecture is difficult as it remains open to prove the existence of irreducible polynomials when almost all coefficients of them are prescribed [9].

If we want to get a polynomial basis with less than two trace-one elements from a three-piece polynomial, then almost every odd degree will upset us. These two results are illustrated in Tables 2 and 3.

For example, we take $n = 88$ in Table 2 to see how it works. The three-piece polynomial is $(6, 75, 8)_{\text{TP}}$ representing

TABLE 2: Three piece polynomials $f(x) = (x + 1)g(x)$ with $f(x)$ irreducible. The number of trace-one elements in the polynomial basis formed by $g(x)$ is no more than four.

Degree	$(n_1, n_2, n_3)_{\text{TP}}/\text{none}$
20	(7, 5, 9)
21	(7, 12, 3)
22	(2, 17, 4)
23	(5, 10, 9)
24	(7, 9, 9)
25	(7, 16, 3)
26	(9, 7, 11)
27	(9, 6, 13)
28	(12, 15, 2)
29	None
30	(2, 25, 4)
31	(11, 18, 3)
32	(2, 23, 8)
33	(13, 4, 17)
34	(8, 17, 10)
35	(9, 22, 5)
36	(2, 31, 4)
37	(7, 28, 3)
38	None
39	None
40	(4, 31, 6)
41	(13, 20, 9)
42	(4, 25, 14)
43	(9, 18, 17)
44	(6, 19, 20)
45	(13, 12, 21)
46	(4, 29, 14)
47	(13, 14, 21)
48	(6, 27, 16)
49	(15, 16, 19)
50	(12, 25, 14)
51	(15, 18, 19)
52	(4, 35, 14)
53	(13, 16, 25)
54	(8, 21, 26)
55	(17, 30, 9)
56	(15, 25, 17)
57	(17, 12, 29)
58	(6, 45, 8)
59	(13, 26, 21)
60	(4, 43, 14)
61	(15, 40, 7)
62	(4, 53, 6)
63	(19, 34, 11)
64	(8, 47, 10)
65	(19, 40, 7)
66	(6, 45, 16)

TABLE 2: Continued.

Degree	$(n_1, n_2, n_3)_{\text{TP}}/\text{none}$
67	(19, 26, 23)
68	(12, 31, 26)
69	(19, 36, 15)
70	(12, 45, 14)
71	(29, 34, 9)
72	(2, 63, 8)
73	(19, 32, 23)
74	(14, 29, 32)
75	(17, 34, 25)
76	(16, 31, 30)
77	(21, 20, 37)
78	(8, 57, 14)
79	(21, 42, 17)
80	(6, 67, 8)
81	(21, 48, 13)
82	(12, 57, 14)
83	(23, 22, 39)
84	(14, 39, 12)
85	(21, 40, 25)
86	(4, 73, 10)
87	(25, 54, 9)
88	(6, 75, 8)
89	(33, 52, 5)
90	(14, 61, 16)
91	(23, 26, 43)
92	(10, 59, 24)
93	(23, 68, 3)
94	(2, 89, 4)
95	(17, 34, 45)
96	(16, 63, 18)
97	(25, 40, 33)
98	(2, 89, 8)
99	(25, 66, 9)
100	(18, 51, 32)
101	(31, 64, 7)

TABLE 3: Three piece polynomials $f(x) = (x + 1)g(x)$ with $g(x)$ irreducible. The number of trace-one elements in the polynomial basis formed by $g(x)$ is no more than two.

Degree	$(n_1, n_2, n_3)_{\text{TP}}/\text{none}$
20	None
21	None
22	(2, 17, 4)
23	None
24	(8, 7, 10)
25	None
26	None
27	(13, 6, 9)

TABLE 3: Continued.

Degree	$(n_1, n_2, n_3)_{\text{TP}}/\text{none}$
28	(12, 15, 2)
29	None
30	(2, 25, 4)
31	None
32	(8, 23, 2)
33	None
34	(8, 17, 10)
35	None
36	(2, 31, 4)
37	None
38	None
39	None
40	(4, 31, 6)
41	None
42	(14, 25, 4)
43	(13, 26, 5)
44	(10, 23, 12)
45	None
46	(12, 21, 14)
47	None
48	(8, 31, 10)
49	None
50	(12, 25, 14)
51	None
52	(12, 39, 2)
53	None
54	(18, 17, 20)
55	None
56	None
57	None
58	(6, 45, 8)
59	None
60	(14, 43, 4)
61	None
62	(4, 53, 6)
63	None
64	(8, 47, 10)
65	None
66	(16, 45, 6)
67	(33, 6, 29)
68	(20, 47, 2)
69	None
70	(12, 45, 14)
71	None
72	(8, 63, 2)
73	None
74	(30, 29, 16)
75	(37, 18, 21)

TABLE 3: Continued.

Degree	$(n_1, n_2, n_3)_{TP}/\text{none}$
76	(30, 31, 16)
77	None
78	(14, 57, 8)
79	None
80	(6, 67, 8)
81	None
82	(12, 57, 14)
83	None
84	(18, 63, 4)
85	None
86	(10, 73, 4)
87	None
88	(6, 75, 8)
89	None
90	(14, 61, 16)
91	None
92	(12, 79, 2)
93	None
94	(2, 89, 4)
95	(29, 58, 9)
96	(16, 63, 18)
97	None
98	(8, 89, 2)
99	None
100	(32, 51, 18)
101	None

the polynomial $f(x) = x^{88} + x^{87} + x^{86} + x^{85} + x^{84} + x^{83} + x^7 + x^6 + x^5 + x^4 + x^3 + x^2 + x + 1$. The value set of s_k of $f(x)$ is

$$\left(\underbrace{1 \dots 1}_{80 \text{ ones}} 0111110 \right). \quad (27)$$

We decompose $f(x)$ as $f(x) = (x+1)g(x) = (x+1)(x^{87} + x^{85} + x^{83} + x^6 + x^4 + x^2 + 1)$ with $g(x)$ irreducible, and the value set of s_k of $g(x)$ is

$$\left(\underbrace{0 \dots 0}_{80 \text{ zeros}} 100000 \right). \quad (28)$$

Combining with the fact that $\text{tr}(1) = 87 \pmod{2} \equiv 1$, we know the polynomial basis of $g(x)$ has two trace-one elements.

4. Conclusion

We investigated the three-piece polynomials to get the polynomial basis with small number of trace-one elements. The regular pattern shown by the value of s_k of the three-piece polynomial is the reason to choose them. Our numerical experiments showed that the three-piece polynomial relative

to polynomial basis with small number of trace-one elements exists for many degrees, but the proof of the existence remains unsolved.

Competing Interests

The authors declare that they have no competing interests.

Acknowledgments

This work is supported by the National Natural Science Foundation of China under Grant nos. 61272037 and 61472472 and “New Generation of Broadband Wireless Communications Network” Ministry of Industry and Information Technology major projects (Project no. 2013ZX03002004).

References

- [1] K. Fong, D. Hankerson, J. López, and A. Menezes, “Field inversion and point halving revisited,” *IEEE Transactions on Computers*, vol. 53, no. 8, pp. 1047–1059, 2004.
- [2] E. Knudsen, “Elliptic scalar multiplication using point halving,” in *Advances in Cryptology—ASIACRYPT ’99*, vol. 1716 of *Lecture Notes in Computer Science*, pp. 135–149, Springer, 1999.
- [3] G. Gong, T. A. Berson, and D. R. Stinson, “Elliptic curve pseudorandom sequence generators,” in *Selected Areas in Cryptography*, vol. 1758 of *Lecture Notes in Computer Science*, pp. 34–48, Springer, Berlin, Germany, 2000.
- [4] O. Ahmadi and A. Menezes, “On the number of trace-one elements in polynomial bases for $\mathbb{F}_2^n$,” *Designs, Codes and Cryptography*, vol. 37, no. 3, pp. 493–507, 2005.
- [5] H. M. Edwards, *Galois Theory*, Springer, Berlin, Germany, 1997.
- [6] R. Lidl and H. Niederreiter, *Introduction to Finite Fields and Their Applications*, Cambridge University Press, 1986.
- [7] R. G. Swan, “Factorization of polynomials over finite fields,” *Pacific Journal of Mathematics*, vol. 12, pp. 1099–1106, 1962.
- [8] A. W. Hales and D. W. Newhart, “Swan’s theorem for binary tetranomials,” *Finite Fields and Their Applications*, vol. 12, no. 2, pp. 301–311, 2006.
- [9] D. Panario, “Open problems for polynomials over finite fields and applications,” in *Open Problems in Mathematics and Computational Science*, pp. 111–126, Springer, 2014.

Hindawi

Submit your manuscripts at
<http://www.hindawi.com>