

Review Article

Epigenetic Modulation on Tau Phosphorylation in Alzheimer's Disease

Chao-Chao Yu ¹, Tao Jiang,¹ Ao-Fei Yang ^{2,3}, Yan-Jun Du,¹ Miao Wu,^{2,3}
and Li-Hong Kong ¹

¹Hubei University of Chinese Medicine, Wuhan, Hubei, China

²Hubei Provincial Hospital of TCM, Wuhan, Hubei, China

³Hubei Province Academy of Traditional Chinese Medicine, Wuhan, Hubei, China

Correspondence should be addressed to Li-Hong Kong; xiyu1618@sina.com

Received 11 February 2019; Accepted 28 March 2019; Published 10 April 2019

Academic Editor: Nicoletta Berardi

Copyright © 2019 Chao-Chao Yu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tau hyperphosphorylation is a typical pathological change in Alzheimer's disease (AD) and is involved in the early onset and progression of AD. Epigenetic modification refers to heritable alterations in gene expression that are not caused by direct changes in the DNA sequence of the gene. Epigenetic modifications, such as noncoding RNA regulation, DNA methylation, and histone modification, can directly or indirectly affect the regulation of tau phosphorylation, thereby participating in AD development and progression. This review summarizes the current research progress on the mechanisms of epigenetic modification associated with tau phosphorylation.

1. Introduction

Alzheimer's disease (AD), also known as senile dementia, is a common neurodegenerative disorder among the elderly. Mild memory impairment is the primary first sign of AD. As the disease progresses, cognitive functions, such as comprehension, intelligence, emotion, and language proficiency, along with self-care abilities gradually decline in AD patients. Notably, the onset of AD is closely associated with aging [1]. In fact, AD has become one of the major health-threatening disorders among the elderly, having the 6th highest mortality rate in the United States (US) and with a rapidly rising prevalence rate of 1 million new cases per year. It is estimated that there will be 132 million AD patients by the year 2050. As the global population continues to age, AD has become one of the top medical and social concerns worldwide [2]. The pathogenesis of AD is very complex and involves β -amyloid protein metabolism disorder and deposition, neurofibrillary tangle (NFT) formation due to abnormal or excessive tau phosphorylation, cholinergic neuron damage, neuroinflammation, abnormal epigenetic modification, intestinal microbiota dysbiosis, abnormal glucose and lipid metabolism, and

oxidative stress [3]. The interrelationships between these mechanisms thus create a complex pathogenic network.

Epigenetic modifications refer to heritable alterations in gene expression by means of DNA regulation, RNA methylation, histone modification, and noncoding RNA, which are not caused by changes in the DNA sequence of the gene. Epigenetic modifications can act as a medium between the external environment and the genome. Importantly, environmental changes and stress responses can induce intracellular epigenetic modifications leading to gene transcription or gene silencing [4]. Therefore, understanding the regulatory mechanisms underlying epigenetic modification will provide new strategies for the prevention and treatment of AD. There is currently no clear correlation between early onset AD (5-10% of total AD) or late onset AD and gene mutations [5, 6]. Although the β -amyloid cascade hypothesis emphasizes the dominance of senile β -amyloid plaques in the pathogenesis of AD [7], many of the β -amyloid-targeting drugs developed in recent years have demonstrated poor efficacy and safety in the treatment of AD patients [8]; therefore, the β -amyloid theory has been increasingly questioned. Previous studies have demonstrated that the severity and




FIGURE 1: Isoforms of tau protein. The six isoforms of tau are by alternative splicing of exons 2, 3, and 10.

progression of AD are closely associated with the number of NFTs formed and less with the number of senile β -amyloid plaques developed [9, 10]. Importantly, paired helical filaments (PHFs), which are formed via tau protein aggregation, are a major component of NFTs [11]. Thus, the ability to target tau protein signaling has become a major goal of drug research and development for AD. In addition, epigenetic modulation on tau phosphorylation is now the primary focus of targeted drug development. Here, we will provide a review on the epigenetic modifications of tau phosphorylation identified in AD.

2. Structure and Biological Function of Microtubule-Associated Tau Proteins

Tau is a microtubule-associated protein (MAP) encoded by the 16 exon-containing *MAPT* gene on chromosome 17 (17q21.31). Tau proteins are rich in neurons in the frontal, temporal, hippocampal, and entorhinal regions of the brain. Intraneuronal tau is predominantly localized in the axons, and it is also present in somatodendritic compartments in much lower levels and contributes to synapse physiology [12, 13]. Markedly, tau proteins bind with a higher affinity to axons than to the cell body or dendrites of a neuron. Depending on the presence or absence of 1 or 2 amino acid insertions in the N-terminus (0N, 1N, and 2N) of the protein and the insertion of either 3 or 4 repeated amino acid sequences in the microtubule-binding domain (3R and 4R) of the protein, a healthy adult human brain can express up to six tau isomers, a result of selective splicing of the *MAPT* gene at exons 2, 3, and 10. The ratio of 4R/3R tau is normally close to 1 : 1 [14, 15]. However, an imbalance in this ratio can lead to neurodegenerative diseases, such as dementia and AD [16, 17]. These tau isomers are comprised of 352, 381, 383, 410, 412, and 411 amino acids and present a molecular

weight of approximately 37 to 46 kilodaltons (kDa) [18, 19]. Genetic studies have revealed a relationship between the structural changes in tau and the development of disease pathology. Over 50 mutations in the *MAPT* gene have been identified to date [15], which have been shown to cause abnormal 4R-tau elevation and excessive tau aggregation via interference with the tau protein structure or exon 10 splicing. H1 and H2 are the two haplotypes of *MAPT*. The H1 haplotype is closely associated with an increased risk of late onset AD [20] and Parkinson's disease [21], whereas the H2 haplotype is associated with a reduced risk of late onset AD. The tau protein (Figure 1) is primarily comprised of a N-terminus projection domain, a proline-rich domain, a C-terminus microtubule-binding domain, and a tail domain. The main biological functions of the tau protein include the promotion of microtubule formation, as well as the assembly and promotion of microtubule stability in the cytoskeleton to ensure normal axoplasmic transport and synaptic plasticity [14, 22]. Tau interacts with a large number of partners, thereby acting as the center in cellular protein-protein interaction networks [19]. Interactions between tau and microtubules are mediated by the microtubule-binding repeats and are highly dynamic. The binding of tau to microtubules occurs via 3 or 4 imperfect 18-amino acid repeats (R1–R4) which are located in the microtubule-binding region with a single repeat as the basic microtubule interacting unit [23]. The microtubule-binding region is also involved in the binding of tau to actin filaments and is required for tau aggregation [24]. Several other tau interaction partners have been identified in addition to microtubules. These include membrane-associated proteins such as annexin A2, which contributes to tau's axonal localisation [25]; src-family non-receptor tyrosine kinases such as Fyn [26], which may be associated with mediating amyloid-beta toxicity at the post-synapse [27]; and protein phosphatase 2A, which serves as

the primary tau phosphatase [28]. Evidence indicates that tau oligomers rather than higher aggregates represent the toxic species [29].

The tau protein contains approximately 77 serine/threonine phosphorylation sites. In addition to phosphorylation, other posttranslational modifications on tau including acetylation, methylation, ubiquitination, small ubiquitin-like modifier (SUMO) modification, nitration, glycosylation, truncation, and splicing [14] have been reported and may contribute differentially to physiological functions of tau and disease [30]. In particular, the phosphorylation of tau is the main posttranslational modification event [31]. It should be noted that tau phosphorylation not only just causes damage to neurons but also exerts protective effects. Evidence indicated that tau phosphorylation at the Ser396 and Ser404 sites can render cells antiapoptotic by stabilizing beta-catenin [32]. And tau hyperphosphorylation at the Thr205, Thr231, Ser262, and Ser396 sites can attenuate the endoplasmic reticulum stress- and death-associated protein kinase-induced apoptosis [33, 34]. In addition, tau phosphorylation may also play an important role in adult hippocampal neurogenesis [35]. But when tau is hyperphosphorylated, its affinity for microtubules can be reduced [14]. Hyperphosphorylated tau has been found in the brain of AD patients, with the level of phosphorylation being 3 to 4 times more than that observed in normal individuals [31, 36, 37]. Consequently, tau hyperphosphorylation is currently recognized as an early pathology in AD pathogenesis [38]. The dynamic imbalance between tau phosphorylation and dephosphorylation is mainly caused by abnormal tau protein kinase and protein phosphatase activities. These kinases include glycogen synthase kinase- (GSK-) 3β , cyclin AMP- (cAMP-) dependent protein kinase A (PKA), mitogen-activated protein kinase (MAPK), protein kinase (PKC), calmodulin kinase II (CaMK II), microtubule affinity regulating kinase (MARK), and protein phosphatase type 2A (PP2A) [31, 39]. Notably, aberrant tau phosphorylation can lead to microtubule collapse, axon degeneration, and axoplasmic transport disorders, which can subsequently affect neurotransmitter synthesis, transport, release, and uptake, thereby resulting in neurodegeneration [40]. Therefore, the modulation of protein kinases and protein phosphatase activities during tau phosphorylation is currently a major direction of anti-AD drug research and development [41–43].

3. Regulation of Tau Phosphorylation by Epigenetic Modification

3.1. Noncoding RNA and Tau Phosphorylation. Noncoding RNAs (ncRNAs), including microRNAs (miRNAs), long ncRNAs (lncRNAs), and circular RNAs (circRNAs), are types of non-protein-coding transcription factors that regulate cell function via the regulation of gene expression [44]. There is increasing evidence demonstrating that abnormal ncRNA expression in the brain can affect AD development and progression through multiple molecular pathways [45]. In particular, 20- to 24-nucleotide-long miRNAs are endogenous ncRNAs that have been well studied and are known to play a role in AD pathogenesis. These miRNAs are widely

found in the central nervous system (CNS) and play an important regulatory role in neural development, differentiation, and maturation. Furthermore, the miRNA-mediated regulation of target genes is considered a type of post-transcriptional regulation. miRNAs can interact with the 3' untranslated region (3'-UTR) of the target gene messenger RNAs (mRNAs) via complementary base pairing and induce the degradation or transcriptional suppression of the target mRNA, thereby affecting gene expression. An increasing number of clinical and laboratory studies have now shown that miRNAs play an important regulatory function in the expression of AD-associated genes, including amyloid precursor protein (*APP*), β -site APP cleaving enzyme 1 (*BACE1*), *GSK-3 β* , and Sirtuin 1 (*SIRT1*) [46]. In addition, miRNAs circulating in the peripheral blood and cerebrospinal fluid (CSF) are also considered potential early diagnostic markers [47] and drug targets [45] for AD. Previous studies have found that many miRNAs, including miR-124, miR-9, miR-132, and miR-137, can alter the 4R/3R tau ratio in neurons by modulating the splicing process of the *MAPT* gene [45].

To this end, a study by Santa-Maria et al. showed that miRNA-219 is downregulated in the brain of AD patients, and subsequent cellular experiments showed that miRNA-219 binds directly to the 3'-UTR of the tau mRNA and represses tau synthesis [48]. Furthermore, miRNA-132 has also been found to be significantly downregulated in AD [49]. miRNA-132 is involved in tau metabolism, as miRNA-132 inhibition can increase amyloid-beta peptide ($A\beta$) deposition [50] and tau hyperphosphorylation, whereas miRNA-132 upregulation can promote ITPKB and p-ERK1/2 expression, thereby inhibiting tau hyperphosphorylation [51]. It was previously found that the upregulation of miRNA-132 reduced total, phosphorylated, acetylated, and cleaved tau protein levels through the regulation of tau acetyltransferase EP300, *GSK-3 β* , RNA binding fox-1 homolog 1 (*Rbfox1*), calpain 2, and caspases 3/7 protein levels [52]. These changes in turn promote axon extension and bifurcation, enhance synapse plasticity, and prevent neuronal loss. The inhibition of miRNA-132/miRNA-212 can also promote tau protein overexpression, hyperphosphorylation, and aggregation, resulting in cognitive dysfunction [53]. Therefore, given the multitarget properties of miRNA-132, its regulation may be a new potential prevention and treatment strategy for AD [49]. Aside from miRNA-132, many other miRNAs also participate in and influence tau metabolism. For example, the downregulation of brain-derived neurotrophic factors (BDNFs), which are important for the regulation of synapse plasticity, as well as neural growth and differentiation, is closely associated with anxiety and progressive memory loss in AD patients [54]. Evidence indicated that the downregulation of BDNFs is accompanied by the upregulation of miRNA-322 in a mouse model of AD, and further research revealed that miRNA-322 is involved in the phosphorylation of tau proteins via targeted regulation of *BDNF* gene expression and the activity of the TrkB receptor [55]. Binding of BDNF to the TrkB receptor can activate several downstream intracellular signaling cascades including the phosphatidylinositol 3-kinase- (PI3K-) Akt pathway

and the Ras-mitogen-activated protein kinase (MAPK) pathway that affect tau phosphorylation [54, 56], while the effects of miRNA-322 on downstream signaling pathways associated with tau phosphorylation such as the PI3K/Akt/GSK-3 β or MAPK/ERK1/2 pathway remained elusive. Ubiquitin carboxy-terminal hydrolase L1 (UCHL1) is a target of miRNA-922, and inhibition of UCHL1 expression by miRNA-922 promotes tau hyperphosphorylation [57]. Moreover, *in vitro* and *in vivo* experiments demonstrated that miRNA-146a inhibits the expression of the rho-associated, coiled-coil-containing protein kinase 1 (ROCK1) gene and then suppresses tau hyperphosphorylation via ROCK1 regulation through the protein phosphatase and tensin homolog (PTEN) [58]. Evidence has shown that the binding of UCHL1 to PTEN is important for PTEN phosphorylation which promotes tau dephosphorylation [59–61]. In addition, miRNA-12-3p can regulate the expression of the *Caveolin-1* gene and modulate the Caveolin-1-PI3K/AKT/GSK-3 β signaling pathway to inhibit tau hyperphosphorylation and neuronal apoptosis [62]. Wang et al. reported that the retinoic acid receptor alpha (RARA) is a target gene of miRNA-138 and miRNA-138 can modulate RARA/GSK-3 β to promote tau hyperphosphorylation [63]. Sun et al. showed that the knockout of miRNA-195 activates Cdk5/p25 signals and promotes the phosphorylation of tau at Ser202, Thr205, Ser262, Thr231, and Ser422 residues. A subsequent study by the authors demonstrated that miRNA-195 can bind to the 3'UTR of the Cdk5r1 mRNA to downregulate the protein expression of p35 and miRNA-195 upregulation in turn suppresses p25 activity, thereby inhibiting tau hyperphosphorylation [64]. miRNA-125b can directly inhibit the expression of Bcl-w to indirectly enhance the activities of tau phosphorylation-associated kinases including Cdk5, p35, and p44/42-MAPK, thus promoting tau hyperphosphorylation [65]. In addition, *in vitro* experiments have demonstrated that members of the miRNA-15 family, such as miRNA-15, miRNA-16, miRNA-195, miRNA-497 [66], and miRNA-26a [67], directly target other tau phosphorylation-associated genes (e.g., *ERK1* and *GSK-3 β*) to participate in the development of AD. miRNA-98 is involved in the regulation of tau phosphorylation and β -amyloid synthesis via the regulation of the insulin-like growth factor-1 (IGF-1) expression [68]. IGF-1 plays a major role in regulating tau phosphorylation in the aging brain [69], and insulin- or IGF-1-activated PI3K/Akt/GSK-3 β signaling may be involved in several tauopathies [70, 71]. Evidence showed that the inhibition of tyrosine-protein phosphatase nonreceptor type 1 (PTPN1) can suppress A β -induced tau phosphorylation by targeting Akt and GSK-3 β [72], and PTPN1 was a direct target of miR-124 as validated by the luciferase reporter assay [73]. Rebuilding the miR-124/PTPN1 pathway by suppression of miR-124 or overexpression of PTPN1 restored synaptic dysfunction and memory loss in AD [73]. In addition, Kim et al. identified that the death-associated protein kinase 1 (DAPK1) overexpression increased tau protein stability and phosphorylation at multiple AD-related sites including Ser262, Ser396, and Thr231 [74]. DAPK1 was also a direct target of miR-26a, and miR-26a/DAPK1 signaling cascades were associated with cellular

pathologies in neurodegenerative disorders such as Parkinson's disease [75].

Furthermore, Xiong et al. reported that miRNA-218 can modulate GSK-3 β and phosphatase 2A activities by regulating the expression of the protein tyrosine phosphatase alpha (PTP α) [76]. This in turn affects the homeostasis between phosphorylated and dephosphorylated tau proteins. These aforementioned miRNAs were all involved in regulating several signaling pathways which play a significant role in tau phosphorylation.

In addition to those reported miRNAs associated with tau phosphorylation-related signaling pathways, there are also other miRNAs not targeting these pathways but eventually promoting tau hyperphosphorylation. A whole-genome expression analysis indicated that methyl-CpG-binding protein-2 (MeCP2) was a key regulator of tauopathy [77]; a further study confirmed the direct regulation of MeCP2 by miR-132, and the miR-132/MeCP2/dynamin 1 pathway participated in hTau-induced neuronal endocytosis deficiency [78]. In addition, the activation of N-methyl-D-aspartate (NMDA) receptor NR2A, which can also be regulated by miR-125b [79], can decrease tau phosphorylation via the PKC/GSK-3 β pathway [80]. The protooncogene tyrosine-protein kinase Fyn is a nonreceptor tyrosine kinase primarily expressed in the axons of neurons, which is involved in the regulation of nervous system development and in neuroinflammation, as well as synapse function and neural plasticity [81]. Previous studies have shown that the interaction between tau and Fyn impairs the stability of receptor complexes in the postsynaptic density (PSD) structure and plays an important role in AD pathogenesis [82]. Liu et al. reported that *Fyn* is a target gene of miRNA-106b. *Fyn* overexpression can promote tau phosphorylation at the Tyr18 site, and miRNA-106b upregulation can inhibit *Fyn*-induced Tyr18 phosphorylation [83].

Moreover, tau acetylation promotes not only tau autophosphorylation but also abnormal tau aggregation. The acetyltransferase p300 [84] and deacetylase SIRT1 [85] are involved in the regulation of tau acetylation. Reduced SIRT1 levels in the brain of AD patients lead to tau hyperacetylation and consequently tau hyperphosphorylation. Numerous studies have now found that the *SIRT1* gene is directly targeted by miRNA-9, miRNA-212, miRNA-181c, and miRNA-132 [86, 87]. In summary, miRNAs can directly or indirectly (Table 1) affect the expression and activity of several tau phosphorylation-associated proteins and signaling pathways and then modulate tau phosphorylation. Therefore, regulation of these miRNAs may serve as a potential strategy for the development of effective anti-AD therapeutics.

3.2. DNA Methylation and Tau Phosphorylation. DNA methylation is a common form of epigenetic modification. This process occurs when the cytosine-guanine dinucleotide (CpG) is modified into 5-methylcytosine by the addition of a methyl group, donated by the S-adenosylmethionine (SAM), on the C5 of CpG in the presence of DNA methyltransferases (i.e., DNMT1, DNMT3A, DNMT3B, and DNMT3L) [90]. The synthesis of SAM is closely associated

TABLE 1: MicroRNAs (miRNAs) associated with tau phosphorylation in Alzheimer's disease.

Dysregulated miRNA(s)	Level in AD	Target site(s)	Reference
miRNA-132	Downregulated	ITPKB, MeCP2, GSK-3 β , and SIRT1	[51, 52, 78, 87, 88]
miRNA-322	Downregulated	BDNF	[55]
miRNA-922	Downregulated	UCHL1	[57]
miRNA-146a	Upregulated	ROCK1	[58]
miRNA-124-3p	Downregulated	Caveolin-1	[62]
miRNA-138	Upregulated	RARA	[63]
miRNA-195	Downregulated	Cdk5r1	[64]
miRNA-125b	Upregulated	Bcl-w, DUSP6, PPP1CA, NMDA, and GSK-3 β	[65, 79, 89]
miRNA-15	Downregulated	ERK1	[66]
miRNA-98	Upregulated	IGF-1	[68]
miRNA-124	Upregulated	PTPN1	[73]
miRNA-26a	Downregulated	DAPK1	[75]
miRNA-106b	Downregulated	Fyn	[83]
miRNA-218	Upregulated	PTP α	[76]
miRNA-212	Downregulated	SIRT1	[86, 87]

ITPKB: inositol 1,4,5-trisphosphate 3-kinase B; MeCP2: methyl-CpG-binding protein-2; GSK-3 β : glycogen synthase kinase-3 β ; SIRT1: Sirtuin 1; BDNF: brain-derived neurotrophic factor; UCHL1: ubiquitin carboxy-terminal hydrolase L1; ROCK1: rho-associated, coiled-coil-containing protein kinase 1; RARA: retinoic acid receptor alpha; DUSP6: dual-specificity phosphatase 6; PPP1CA: protein phosphatase 1 catalytic subunit alpha isoform; NMDA: N-methyl-D-aspartate; Bcl-w: Bcl-2-like protein 2; ERK1: extracellular-regulated kinase; IGF-1: insulin-like growth factor 1; PTPN1: tyrosine-protein phosphatase nonreceptor type 1; DAPK1: death-associated protein kinase 1; PTP α : protein tyrosine phosphatase α .

with vitamin B and folic acid, with CpG being the primary site of DNA methylation. CpG sites are densely found in certain regions of the genome and can be up to 200 base pairs (bp) in length. The CpG island (CpGI) is a region of the genome in which CpG sites comprise 60% to 70% of the sequence and the CpG observed/expected ratio (ObsCpG/ExpCpG) has been shown to be less than (<) 0.6. CpGIs are primarily found upstream of the promoter region/transcription initiation site [91]. The methylation of CpGIs in the promoter region can repress the transcription of target genes and is therefore considered a form of transcriptional regulation [92]. Numerous studies have demonstrated that DNA methylation plays an important role in the aberrant expression of AD-associated genes. Further, immunohistochemical (IHC) analysis of pathological brain tissue sections from deceased AD patients has revealed decreased DNA methylation levels in the prefrontal cortex [93], entorhinal cortex [94], and hippocampus [95]. Moreover, reduced DNA methylation can promote the activation of astrocytes and microglia and proinflammatory cytokine secretion during aging, which consequently results in the vicious cycle of a number of pathological processes [96].

Current clinical and basic research studies have confirmed the presence of abnormal methylation levels in the promoter regions of tau phosphorylation-related genes. For instance, analysis of DNA methylation in the promoter region of the *GSK-3 β* gene from the prefrontal cortex tissue of deceased AD patients indicated that the *GSK-3 β* promoter region is methylated at low levels during early AD development. The mRNA of *GSK-3 β* is upregulated during this period, but the protein expression levels of *GSK-3 β* remain unchanged [97]. Some studies have found that vitamin B deficiency can lead to low levels of cytosine methylation in

the *GSK-3 β* promoter region and hence the *GSK-3 β* overexpression [98]. Using chromatin immunoprecipitation (ChIP) and bisulfite sequencing technologies, Li et al. found that the promoter region of *Cdk5* has a low level of cytosine methylation in the hippocampal CA1 region of a rat model with *A β ₁₋₄₂*-induced memory deficiency. The upregulation of *Cdk5* expression leads to tau hyperphosphorylation and suppressed long-term synaptic potentiation, resulting in spatial learning and memory impairment in this rat model [99]. In addition, it was previously reported that AD patients have increased methylation in the promoter region of the *dual-specificity phosphatase 22 (DUSP22)* gene and downregulated *DUSP22* expression, which in turn inhibited PKA-mediated tau phosphorylation and cAMP response element-binding protein (CREB) activation [100] and affected synapse plasticity and long-term memory formation [101]. Besides, DNA demethylation regulated by ten-eleven translocation proteins (Tet1-3) that oxidize 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC) [102] could also affect tau phosphorylation. Several studies have proved that Tet1 activity functions in active DNA demethylation and gene regulation during learning and memory [103–105]. It is known that BDNF is a key component in the maintenance of synaptic plasticity and synaptogenesis in the hippocampus [56] and is closely related to tau hyperphosphorylation [54, 55, 106]. Ambigapathy et al. reported that Tet1 and ERK1/2 were critical partners regulating BDNF chromatin status and promoter accessibility [107]. It is reasonably assumed that BDNF DNA demethylation regulated by Tet1 could influence the tau phosphorylation levels. These studies demonstrate that targeted regulation of methylation levels of tau phosphorylation-related genes is a potential treatment strategy for AD.

3.3. Histone Modifications and Tau Phosphorylation. A nucleosome is the basic unit of DNA packaging, which consists of a segment of DNA wound around histone proteins. H1/H5, H2A, H2B, H3, and H4 are the primary histone proteins important for maintaining the chromosome configuration in DNA material. H2A, H2B, H3, and H4 are core histone proteins, whereas H1/H5 are linker histones. Histones can be modified by acetylation, methylation, phosphorylation, ubiquitination, SUMO modification, and glycosylation. These modifications can affect gene transcription by modulating the spatial conformation of chromatin. For example, acetylation and methylation of lysine (K) and arginine (R) in the N-terminus of histones can neutralize the positive charges on these residues, leading to a reduced affinity between the DNA and histones and loosened chromatin structure (euchromatin) that are favorable for the binding of transcription factors to DNA and hence gene transcription. In contrast, deacetylation of histones tightens the spatial conformation of chromatin (heterochromatin) and suppresses gene transcription [108]. Enzymes that are mainly involved in histone acetylation include histone acetylase (HAT) and histone deacetylase (HDAC). Several studies have recently shown that histone modification plays a role in AD development and progression. It was reported that AD patients and AD mouse models have elevated levels of HDAC2 in the brain [109], and inhibition or knockout of HDAC2 can significantly improve cognitive dysfunction [110]. Furthermore, both AD patients and the 3xTg and APP_{SDI} mouse model have an elevated level of Lys12 acetylation on histone H4 (H4K12) as seen in brain tissue samples [111]. Notably, treatment with an HDAC inhibitor can induce hippocampal axonal regeneration, increase axon number, and improve learning and memory in CK-p25 mice [112]. The HDAC inhibitor phenylbutyrate can reduce tau hyperphosphorylation, increase axon density, and improve the spatial learning and memory impairment seen in Tg2576 AD mice [113]. In addition, a study by Li et al. showed that increased histone H3 acetylation can lead to tau hyperphosphorylation and impaired synaptic plasticity by promoting Cdk5 transcription and expression [99]. Hippocampal HDAC2 overexpression in 3xTg-AD mice resulted in the deacetylation of the hepatocyte nuclear factor 4 alpha (HNF-4 α), which allows HNF-4 α to bind to the promoter of miRNA-101b and repress miRNA-101b expression. This subsequently upregulates AMPK expression and tau hyperphosphorylation, resulting in a reduced density and abnormal morphology of dendrites and consequently cognitive dysfunction in AD mice [114]. Aside from the regulatory role of histone acetylation in tau phosphorylation, Mastroeni et al. also found that an increased Lys4 methylation of histone H3 (H3K4me3) may be involved in tau pathology as an early event in AD pathology. Analysis of brain tissue sections from deceased AD patients revealed that the level of H3K4me3 in the cytoplasm of neurons is elevated as the Braak staging increased [115]. HDAC6 is another important epigenetic component of the etiopathogenesis of AD, and its specific role in AD has been extensively discussed in previous reviews [116, 117]. HDAC6 significantly increased during AD progression [118]. Recent evidence showed that the

inhibition of HDAC6 can reverse tau phosphorylation and restore microtubule stability, leading to the normalization of synaptosomal mitochondrial function and synaptic integrity [119, 120]. This evidence indicates that HDAC6 inhibitors may be a promising avenue for therapeutic intervention in AD and other neurodegenerative diseases. However, how HDAC6 impacts genes or signaling cascades related to tau phosphorylation is less reported which warrants further investigation.

4. Summary and Future Directions

AD is a pathologically complex neurodegenerative disease, and elucidating the underlying molecular mechanisms of related epigenetic modifications has provided new insights into the understanding of AD pathogenesis, creating new strategies for AD prevention and treatment. As previously mentioned, tau hyperphosphorylation is a key early event in AD pathology, and its development and progression is closely associated with aberrant epigenetic modifications such as miRNAs, DNA methylation, and histone modification. However, whether aberrant epigenetic modifications are the cause or consequence of AD development is still unclear. A review of the mechanisms by which epigenetic modification participates and regulates tau phosphorylation shows that ncRNAs, DNA methylation, and histone modification can directly or indirectly affect the expression and activity of tau-related kinase genes, forming a complex epigenetic regulatory network (Figure 2). Yet, it is still unclear which type of epigenetic modification plays a dominant role in tau hyperphosphorylation, as well as in NFT formation and deposition. This particular point is especially important for the development of specific antitau hyperphosphorylation agents. Therefore, the exact mechanisms by which epigenetic modification participates in tau phosphorylation will need to be further investigated.

Since the different phosphorylation sites of the tau protein can yield different effects which may be protective for neurons instead of causing damage, the potential effects of selected phosphorylation sites modified by epigenetic mechanisms of tau protein kinases and protein phosphatases should be considered. Also, to which degree the epigenetic modulation on tau phosphorylation will be protective should be investigated. So far, no existing evidence mentioned above has taken these issues into account which are important and need to be studied further. Importantly, the occurrence of the altered expression of distinct miRNAs, aberrant DNA methylation, and histone modification involved in tau hyperphosphorylation could be used as new and promising biomarkers for AD in the future. This approach could provide a scientific foundation for the preclinical use of drugs.

There is increasing evidence pointing to the brain stem nucleus as a possible initial site of AD pathology and dissemination. In particular, the raphe nuclei and locus coeruleus may be early regions of NFT formation and aggregation [121–124]. A magnetic resonance imaging (MRI) study revealed that AD patients have altered brain stem volume and morphology [125]. Furthermore, symptoms of cognitive dysfunction in early AD, such as depression and abnormal




FIGURE 2: Epigenetic modulation on tau phosphorylation and possible impacts on synapses and neurons. Dysregulated epigenetic modification on genes associated with the tau phosphorylation process could lead to NFT aggregation which can then cause neuroinflammation, axonal transport dysfunction, and mitochondrial and synaptic plasticity injuries.

emotion, in addition to changes in breathing and electrocardiogram (ECG) readings, are closely associated with the secretion of 5-hydroxytryptamine (5-HT) and norepinephrine by the locus coeruleus and raphe nuclei [126–128]. The locus coeruleus is the primary region of norepinephrine synthesis, and noradrenergic neurons in the nuclei project to various regions of the brain in a long-range and extensive manner. Braak et al. showed that NFT formation occurs earlier in the locus coeruleus than in any other brain regions and may be present without any significant clinical symptoms [129]. Neurons in the raphe nuclei are long-range projecting neurons that secrete 5-HT. NFT deposition in the raphe nuclei may explain the abnormal emotional symptoms, such as depression and irritability, that develop during early AD [130] and has also been shown to be associated with the progressive exacerbation of clinical symptoms [131, 132]. Many studies have demonstrated that hyperphosphorylation-mediated tau deposition occurs earlier in the locus coeruleus and raphe nuclei than in other regions of the brain [133] and is often accompanied by neuronal apoptosis [134]. Therefore, early targeted regulation of tau hyperphosphorylation, inhibition of NFT deposition, and formation in the locus coeruleus and raphe nuclei are especially important for the prevention of AD. As key regulators of early AD pathology, epigenetic modifications are also involved in tau hyperphosphorylation in the brainstem nucleus. Andres-Benito et al. found that the neurons in the locus coeruleus in an aging brain presented low katanin-interacting protein gene (*KIAA0566*) methylation levels along with downregulated mRNA and NFT deposition [135]. Researchers have compared the association between the NFT pathology and related

miRNA levels in the locus coeruleus, entorhinal cortex, hippocampal CA1 region, and dentate gyrus between Braak stages I and II and stages III and IV and found that miRNA-27a-3p, miRNA-124-3p, and miRNA-143-3p levels in the locus coeruleus, but not any in other regions of the brain, are already elevated in Braak stages I and II and are significantly increased in stages III and IV. Only miRNA-143-3p is elevated in the entorhinal cortex, with all the other miRNA levels remaining unchanged in the hippocampal CA1 region [136]. These findings demonstrate that abnormal epigenetic modification in the locus coeruleus is likely to be involved in the development and progression of tau hyperphosphorylation during early AD pathology. However, the mechanisms by which these epigenetic modifications regulate NFT aggregation in the raphe nuclei are still elusive and will need to be further investigated. Understanding the epigenetic modification mechanisms underlying these AD-related pathologies in associated brain nuclei may provide new insights into the elucidation of AD pathogenesis and the development of actionable drug targets.

However, for now, clinical epigenetics would not be useful as a potential therapeutic strategy to ameliorate AD because epigenetic-based therapy may affect numerous targets due to the lack of locus specificity. DNA methyltransferase inhibitors (DNMTi) and the histone deacetylase inhibitor (HDACi) class are classified as broad reprogrammers because of their large-scale effects on genomic sites [137]. ncRNA-oriented drugs may also yield potential non-specific off-target effects [138]. Despite these challenges, a range of epigenetic biomarkers for AD diagnosis are in development [139–141]. Integrating all epigenetic aspects and

considering epigenetic factors as highly dynamic and interactive players with cellular metabolism by adopting multiomics technologies such as epigenomics, transcriptomics, metabolomics, and proteomics could help in the discovery of novel diagnostic biomarkers or potential drugs for AD.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

Authors' Contributions

Chao-Chao Yu and Tao Jiang contributed equally to this manuscript. Ao-Fei Yang and Miao Wu illustrated the figures and Li-Hong Kong and Yan-Jun Du helped revise the manuscript. The authors approved the final manuscript.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Nos. 81373741, 81473786, and 81873380) and the "Acupuncture for Learning and Memory Disorders Related Diseases"-Preponderant Discipline Team Project funded by the Hubei University of Chinese Medicine.

References




- [1] R. Nativio, G. Donahue, A. Berson et al., "Dysregulation of the epigenetic landscape of normal aging in Alzheimer's disease," *Nature Neuroscience*, vol. 21, no. 4, pp. 497–505, 2018.
- [2] M. Prince, A. Wimo, M. Guerchet, G. C. Ali, Y. T. Wu, and M. Prina, "World Alzheimer report 2015. The global impact of dementia. An analysis of prevalence, incidence, cost and trends," <https://www.alz.co.uk/research/world-report-2015>.
- [3] F. T. Hane, B. Y. Lee, and Z. Leonenko, "Recent progress in Alzheimer's disease research, part 1: pathology," *Journal of Alzheimer's Disease*, vol. 57, no. 1, pp. 1–28, 2017.
- [4] A. Brunet and S. L. Berger, "Epigenetics of aging and aging-related disease," *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, vol. 69, Supplement 1, pp. S17–S20, 2014.
- [5] Alzheimer's Association, "2016 Alzheimer's disease facts and figures," *Alzheimer's & Dementia*, vol. 12, no. 4, pp. 459–509, 2016.
- [6] M. Robinson, B. Y. Lee, and F. T. Hanes, "Recent progress in Alzheimer's disease research, part 2: genetics and epidemiology," *Journal of Alzheimer's Disease*, vol. 61, no. 1, p. 459, 2018.
- [7] D. J. Selkoe and J. Hardy, "The amyloid hypothesis of Alzheimer's disease at 25 years," *EMBO Molecular Medicine*, vol. 8, no. 6, pp. 595–608, 2016.
- [8] D. Mehta, R. Jackson, G. Paul, J. Shi, and M. Sabbagh, "Why do trials for Alzheimer's disease drugs keep failing? A discontinued drug perspective for 2010-2015," *Expert Opinion on Investigational Drugs*, vol. 26, no. 6, pp. 735–739, 2017.
- [9] M. Marquí, M. Siao Tick Chong, A. Antón-Fernández et al., "[F-18]-AV-1451 binding correlates with postmortem neurofibrillary tangle Braak staging," *Acta Neuropathologica*, vol. 134, no. 4, pp. 619–628, 2017.
- [10] P. V. Arriagada, J. H. Growdon, E. T. Hedley-Whyte, and B. T. Hyman, "Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease," *Neurology*, vol. 42, no. 3, pp. 631–639, 1992.
- [11] J. Wang, J. T. Yu, M. S. Tan, T. Jiang, and L. Tan, "Epigenetic mechanisms in Alzheimer's disease: implications for pathogenesis and therapy," *Ageing Research Reviews*, vol. 12, no. 4, pp. 1024–1041, 2013.
- [12] L. M. Ittner and J. Gotz, "Amyloid- β and tau—a toxic *pas de deux* in Alzheimer's disease," *Nature Reviews Neuroscience*, vol. 12, no. 2, pp. 67–72, 2011.
- [13] A. Ittner and L. M. Ittner, "Dendritic tau in Alzheimer's disease," *Neuron*, vol. 99, no. 1, pp. 13–27, 2018.
- [14] T. Guo, W. Noble, and D. P. Hanger, "Roles of tau protein in health and disease," *Acta Neuropathologica*, vol. 133, no. 5, pp. 665–704, 2017.
- [15] M. G. Spillantini and M. Goedert, "Tau pathology and neurodegeneration," *The Lancet Neurology*, vol. 12, no. 6, pp. 609–622, 2013.
- [16] I. D'Souza and G. D. Schellenberg, "Regulation of tau isoform expression and dementia," *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, vol. 1739, no. 2-3, pp. 104–115, 2005.
- [17] T. M. Caffrey, C. Joachim, S. Paracchini, M. M. Esiri, and R. Wade-Martins, "Haplotype-specific expression of exon 10 at the human *MAPT* locus," *Human Molecular Genetics*, vol. 15, no. 24, pp. 3529–3537, 2006.
- [18] M. L. Caillet-Boudin, L. Buee, N. Sergeant, and B. Lefebvre, "Regulation of human *MAPT* gene expression," *Molecular Neurodegeneration*, vol. 10, no. 1, p. 28, 2015.
- [19] S. S. Khan and G. S. Bloom, "Tau: the center of a signaling nexus in Alzheimer's disease," *Frontiers in Neuroscience*, vol. 10, p. 31, 2016.
- [20] M. Allen, M. Kachadoorian, Z. Quicksall et al., "Association of *MAPT* haplotypes with Alzheimer's disease risk and *MAPT* brain gene expression levels," *Alzheimer's Research & Therapy*, vol. 6, no. 4, p. 39, 2014.
- [21] J. Simón-Sánchez, C. Schulte, J. M. Bras et al., "Genome-wide association study reveals genetic risk underlying Parkinson's disease," *Nature Genetics*, vol. 41, no. 12, pp. 1308–1312, 2009.
- [22] L. Dehmelt and S. Halpain, "The MAP2/Tau family of microtubule-associated proteins," *Genome Biology*, vol. 6, no. 1, p. 204, 2005.
- [23] Y. I. Pachima, L. Y. Zhou, P. Lei, and I. Gozes, "Microtubule-tau interaction as a therapeutic target for Alzheimer's disease," *Journal of Molecular Neuroscience*, vol. 58, no. 2, pp. 145–152, 2016.
- [24] H. Ksiazek-Reding and S. H. Yen, "Structural stability of paired helical filaments requires microtubule-binding domains of tau: a model for self-association," *Neuron*, vol. 6, no. 5, pp. 717–728, 1991.
- [25] A. Gauthier-Kemper, C. Weissmann, N. Golovyashkina et al., "The frontotemporal dementia mutation R406W blocks tau's interaction with the membrane in an annexin A2-dependent manner," *Journal of Cell Biology*, vol. 192, no. 4, pp. 647–661, 2011.
- [26] G. Lee, S. T. Newman, D. L. Gard, H. Band, and G. Panchamoorthy, "Tau interacts with src-family non-receptor tyrosine kinases," *Journal of Cell Science*, vol. 111, no. 21, pp. 3167–3177, 1998.

- [27] L. M. Ittner, Y. D. Ke, F. Delerue et al., "Dendritic function of tau mediates amyloid- β toxicity in Alzheimer's disease mouse models," *Cell*, vol. 142, no. 3, pp. 387–397, 2010.
- [28] J. M. Sontag and E. Sontag, "Protein phosphatase 2A dysfunction in Alzheimer's disease," *Frontiers in Molecular Neuroscience*, vol. 7, p. 16, 2014.
- [29] T. L. Spires-Jones, K. J. Kopeikina, R. M. Koffie, A. de Calignon, and B. T. Hyman, "Are tangles as toxic as they look?," *Journal of Molecular Neuroscience*, vol. 45, no. 3, pp. 438–444, 2011.
- [30] E. M. Mandelkow and E. Mandelkow, "Biochemistry and cell biology of tau protein in neurofibrillary degeneration," *Cold Spring Harbor Perspectives in Medicine*, vol. 2, no. 7, 2012.
- [31] J. Oliveira, M. Costa, M. S. C. de Almeida, O. A. B. da Cruz e Silva, and A. G. Henriques, "Protein phosphorylation is a key mechanism in Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 58, no. 4, pp. 953–978, 2017.
- [32] H. L. Li, H. H. Wang, S. J. Liu et al., "Phosphorylation of tau antagonizes apoptosis by stabilizing β -catenin, a mechanism involved in Alzheimer's neurodegeneration," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 9, pp. 3591–3596, 2007.
- [33] X. A. Liu, J. Song, Q. Jiang, Q. Wang, Q. Tian, and J. Z. Wang, "Expression of the hyperphosphorylated tau attenuates ER stress-induced apoptosis with upregulation of unfolded protein response," *Apoptosis*, vol. 17, no. 10, pp. 1039–1049, 2012.
- [34] D. X. Duan, G. S. Chai, Z. F. Ni et al., "Phosphorylation of tau by death-associated protein kinase 1 antagonizes the kinase-induced cell apoptosis," *Journal of Alzheimer's Disease*, vol. 37, no. 4, pp. 795–808, 2013.
- [35] X. P. Hong, C. X. Peng, W. Wei et al., "Essential role of tau phosphorylation in adult hippocampal neurogenesis," *Hippocampus*, vol. 20, no. 12, pp. 1339–1349, 2010.
- [36] E. Kopke, Y. C. Tung, S. Shaikh, A. C. Alonso, K. Iqbal, and I. Grundke-Iqbal, "Microtubule-associated protein tau. Abnormal phosphorylation of a non-paired helical filament pool in Alzheimer disease," *The Journal of Biological Chemistry*, vol. 268, no. 32, pp. 24374–24384, 1993.
- [37] I. Grundke-Iqbal, K. Iqbal, Y. C. Tung, M. Quinlan, H. M. Wisniewski, and L. I. Binder, "Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 83, no. 13, pp. 4913–4917, 1986.
- [38] H. Braak and E. Braak, "Neuropathological stageing of Alzheimer-related changes," *Acta Neuropathologica*, vol. 82, no. 4, pp. 239–259, 1991.
- [39] R. H. Ma, Y. Zhang, X. Y. Hong, J. F. Zhang, J. Z. Wang, and G. P. Liu, "Role of microtubule-associated protein tau phosphorylation in Alzheimer's disease," *Journal of Huazhong University of Science and Technology [Medical Sciences]*, vol. 37, no. 3, pp. 307–312, 2017.
- [40] C. Li and J. Gotz, "Tau-based therapies in neurodegeneration: opportunities and challenges," *Nature Reviews Drug Discovery*, vol. 16, no. 12, pp. 863–883, 2017.
- [41] S. Y. Hung and W. M. Fu, "Drug candidates in clinical trials for Alzheimer's disease," *Journal of Biomedical Science*, vol. 24, no. 1, p. 47, 2017.
- [42] Y. Yang and J. Z. Wang, "Nature of tau-associated neurodegeneration and the molecular mechanisms," *Journal of Alzheimer's Disease*, vol. 62, no. 3, pp. 1305–1317, 2018.
- [43] Y. Huang and L. Mucke, "Alzheimer mechanisms and therapeutic strategies," *Cell*, vol. 148, no. 6, pp. 1204–1222, 2012.
- [44] K. V. Morris and J. S. Mattick, "The rise of regulatory RNA," *Nature Reviews Genetics*, vol. 15, no. 6, pp. 423–437, 2014.
- [45] A. Di Meco and D. Pratico, "MicroRNAs as therapeutic targets for Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 53, no. 2, pp. 367–372, 2016.
- [46] J. Zhao, D. Yue, Y. Zhou et al., "The role of microRNAs in A β deposition and tau phosphorylation in Alzheimer's disease," *Frontiers in Neurology*, vol. 8, p. 342, 2017.
- [47] T. Kiko, K. Nakagawa, T. Tsuduki, K. Furukawa, H. Arai, and T. Miyazawa, "MicroRNAs in plasma and cerebrospinal fluid as potential markers for Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 39, no. 2, pp. 253–259, 2014.
- [48] I. Santa-Maria, M. E. Alaniz, N. Renwick et al., "Dysregulation of microRNA-219 promotes neurodegeneration through post-transcriptional regulation of tau," *The Journal of Clinical Investigation*, vol. 125, no. 2, pp. 681–686, 2015.
- [49] E. Salta and B. De Strooper, "MicroRNA-132: a key noncoding RNA operating in the cellular phase of Alzheimer's disease," *The FASEB Journal*, vol. 31, no. 2, pp. 424–433, 2017.
- [50] J. Hernandez-Rapp, S. Rainone, C. Goupil et al., "MicroRNA-132/212 deficiency enhances A β production and senile plaque deposition in Alzheimer's disease triple transgenic mice," *Scientific Reports*, vol. 6, no. 1, article 30953, 2016.
- [51] E. Salta, A. Sierksma, E. Vanden Eynden, and B. De Strooper, "miR-132 loss de-represses ITPKB and aggravates amyloid and TAU pathology in Alzheimer's brain," *EMBO Molecular Medicine*, vol. 8, no. 9, pp. 1005–1018, 2016.
- [52] R. El Fatimy, S. Li, Z. Chen et al., "MicroRNA-132 provides neuroprotection for tauopathies via multiple signaling pathways," *Acta Neuropathologica*, vol. 136, no. 4, pp. 537–555, 2018.
- [53] P. Y. Smith, J. Hernandez-Rapp, F. Jolivet et al., "miR-132/212 deficiency impairs tau metabolism and promotes pathological aggregation in vivo," *Human Molecular Genetics*, vol. 24, no. 23, pp. 6721–6735, 2015.
- [54] H. Tanila, "The role of BDNF in Alzheimer's disease," *Neurobiology of Disease*, vol. 97, Part B, pp. 114–118, 2017.
- [55] J. Zhang, Z. Liu, Y. Pei, W. Yang, C. Xie, and S. Long, "MicroRNA-322 cluster promotes tau phosphorylation via targeting brain-derived neurotrophic factor," *Neurochemical Research*, vol. 43, no. 3, pp. 736–744, 2018.
- [56] J. H. Song, J. T. Yu, and L. Tan, "Brain-derived neurotrophic factor in Alzheimer's disease: risk, mechanisms, and therapy," *Molecular Neurobiology*, vol. 52, no. 3, pp. 1477–1493, 2015.
- [57] Z. B. Zhao, L. Wu, R. Xiong et al., "MicroRNA-922 promotes tau phosphorylation by downregulating ubiquitin carboxy-terminal hydrolase L1 (UCHL1) expression in the pathogenesis of Alzheimer's disease," *Neuroscience*, vol. 275, pp. 232–237, 2014.
- [58] G. Wang, Y. Huang, L. L. Wang et al., "MicroRNA-146a suppresses ROCK1 allowing hyperphosphorylation of tau in Alzheimer's disease," *Scientific Reports*, vol. 6, no. 1, article 26697, 2016.
- [59] N. Nayeem, F. Kerr, H. Naumann, J. Linehan, S. Lovestone, and S. Brandner, "Hyperphosphorylation of tau and neurofilaments and activation of CDK5 and ERK1/2 in PTEN-deficient cerebella," *Molecular and Cellular Neurosciences*, vol. 34, no. 3, pp. 400–408, 2007.

- [60] X. Zhang, Y. W. Zhang, S. Liu et al., "Tumor suppressor PTEN affects tau phosphorylation: deficiency in the phosphatase activity of PTEN increases aggregation of an FTDP-17 mutant tau," *Molecular Neurodegeneration*, vol. 1, no. 1, p. 7, 2006.
- [61] S. Vemula, J. Shi, P. Hanneman, L. Wei, and R. Kapur, "ROCK1 functions as a suppressor of inflammatory cell migration by regulating PTEN phosphorylation and stability," *Blood*, vol. 115, no. 9, pp. 1785–1796, 2010.
- [62] Q. Kang, Y. Xiang, D. Li et al., "MiR-124-3p attenuates hyperphosphorylation of Tau protein-induced apoptosis via caveolin-1-PI3K/Akt/GSK3 β pathway in N2a/APP695swe cells," *Oncotarget*, vol. 8, no. 15, pp. 24314–24326, 2017.
- [63] X. Wang, L. Tan, Y. Lu et al., "MicroRNA-138 promotes tau phosphorylation by targeting retinoic acid receptor alpha," *FEBS Letters*, vol. 589, no. 6, pp. 726–729, 2015.
- [64] L. H. Sun, T. Ban, C. D. Liu et al., "Activation of Cdk5/p25 and tau phosphorylation following chronic brain hypoperfusion in rats involves microRNA-195 down-regulation," *Journal of Neurochemistry*, vol. 134, no. 6, pp. 1139–1151, 2015.
- [65] J. Banzhaf-Strathmann, E. Benito, S. May et al., "MicroRNA-125b induces tau hyperphosphorylation and cognitive deficits in Alzheimer's disease," *The EMBO Journal*, vol. 33, no. 15, pp. 1667–1680, 2014.
- [66] S. S. Hébert, A. S. Papadopoulou, P. Smith et al., "Genetic ablation of *Dicer* in adult forebrain neurons results in abnormal tau hyperphosphorylation and neurodegeneration," *Human Molecular Genetics*, vol. 19, no. 20, pp. 3959–3969, 2010.
- [67] J. S. Mohamed, M. A. Lopez, and A. M. Boriak, "Mechanical stretch up-regulates microRNA-26a and induces human airway smooth muscle hypertrophy by suppressing glycogen synthase kinase-3 β ," *Journal of Biological Chemistry*, vol. 285, no. 38, pp. 29336–29347, 2010.
- [68] Y. K. Hu, X. Wang, L. Li, Y. H. Du, H. T. Ye, and C. Y. Li, "MicroRNA-98 induces an Alzheimer's disease-like disturbance by targeting insulin-like growth factor 1," *Neuroscience Bulletin*, vol. 29, no. 6, pp. 745–751, 2013.
- [69] C. M. Cheng, V. Tseng, J. Wang, D. Wang, L. Matyakhina, and C. A. Bondy, "Tau is hyperphosphorylated in the insulin-like growth factor-I null brain," *Endocrinology*, vol. 146, no. 12, pp. 5086–5091, 2005.
- [70] L. Yang, H. Wang, L. Liu, and A. Xie, "The role of insulin/IGF-1/PI3K/Akt/GSK3 β signaling in Parkinson's disease dementia," *Frontiers in Neuroscience*, vol. 12, p. 73, 2018.
- [71] G. Bedse, F. Di Domenico, G. Serviddio, and T. Cassano, "Aberrant insulin signaling in Alzheimer's disease: current knowledge," *Frontiers in Neuroscience*, vol. 9, p. 204, 2015.
- [72] T. Kanno, A. Tsuchiya, A. Tanaka, and T. Nishizaki, "Combination of PKC ϵ activation and PTP1B inhibition effectively suppresses A β -induced GSK-3 β activation and tau phosphorylation," *Molecular Neurobiology*, vol. 53, no. 7, pp. 4787–4797, 2016.
- [73] X. Wang, D. Liu, H. Z. Huang et al., "A novel microRNA-124/PTPN1 signal pathway mediates synaptic and memory deficits in Alzheimer's disease," *Biological Psychiatry*, vol. 83, no. 5, pp. 395–405, 2018.
- [74] B. M. Kim, M. H. You, C. H. Chen et al., "Death-associated protein kinase 1 has a critical role in aberrant tau protein regulation and function," *Cell Death & Disease*, vol. 5, no. 5, article e1237, 2014.
- [75] Y. Su, M. F. Deng, W. Xiong et al., "MicroRNA-26a/death-associated protein kinase 1 signaling induces synucleinopathy and dopaminergic neuron degeneration in Parkinson's disease," *Biological Psychiatry*, 2018.
- [76] Y. S. Xiong, F. F. Liu, D. Liu et al., "Opposite effects of two estrogen receptors on tau phosphorylation through disparate effects on the miR-218/PTPA pathway," *Aging Cell*, vol. 14, no. 5, pp. 867–877, 2015.
- [77] N. M. Maphis, S. Jiang, J. Binder et al., "Whole genome expression analysis in a mouse model of tauopathy identifies MECP2 as a possible regulator of tau pathology," *Frontiers in Molecular Neuroscience*, vol. 10, p. 69, 2017.
- [78] A. J. Xie, T. Y. Hou, W. Xiong et al., "Tau overexpression impairs neuronal endocytosis by decreasing the GTPase dynamin 1 through the miR-132/MeCP2 pathway," *Aging Cell*, no. article e12929, 2019.
- [79] H. Tang, M. Ma, Y. Wu et al., "Activation of MT2 receptor ameliorates dendritic abnormalities in Alzheimer's disease via C/EBP α /miR-125b pathway," *Aging Cell*, vol. 18, no. 2, article e12902, 2019.
- [80] A. De Montigny, I. Elhiri, J. Allyson, M. Cyr, and G. Massicotte, "NMDA reduces Tau phosphorylation in rat hippocampal slices by targeting NR2A receptors, GSK3 β , and PKC activities," *Neural Plasticity*, vol. 2013, Article ID 261593, 10 pages, 2013.
- [81] H. B. Nygaard, "Targeting Fyn kinase in Alzheimer's disease," *Biological Psychiatry*, vol. 83, no. 4, pp. 369–376, 2018.
- [82] H. B. Nygaard, C. H. van Dyck, and S. M. Strittmatter, "Fyn kinase inhibition as a novel therapy for Alzheimer's disease," *Alzheimer's Research & Therapy*, vol. 6, no. 1, p. 8, 2014.
- [83] W. Liu, J. Zhao, and G. Lu, "miR-106b inhibits tau phosphorylation at Tyr18 by targeting Fyn in a model of Alzheimer's disease," *Biochemical and Biophysical Research Communications*, vol. 478, no. 2, pp. 852–857, 2016.
- [84] S. W. Min, X. Chen, T. E. Tracy et al., "Critical role of acetylation in tau-mediated neurodegeneration and cognitive deficits," *Nature Medicine*, vol. 21, no. 10, pp. 1154–1162, 2015.
- [85] L. Rizzi and M. Roriz-Cruz, "Sirtuin 1 and Alzheimer's disease: an up-to-date review," *Neuropeptides*, vol. 71, pp. 54–60, 2018.
- [86] R. B. Weinberg, E. J. Mufson, and S. E. Counts, "Evidence for a neuroprotective microRNA pathway in amnesic mild cognitive impairment," *Frontiers in Neuroscience*, vol. 9, p. 430, 2015.
- [87] A. Hadar, E. Milanesi, M. Walczak et al., "SIRT1, miR-132 and miR-212 link human longevity to Alzheimer's disease," *Scientific Reports*, vol. 8, no. 1, p. 8465, 2018.
- [88] Z. Tan, Y. Chen, W. Xie, X. Liu, Y. Zhu, and Y. Zhu, "Nimodipine attenuates tau phosphorylation at Ser396 via miR-132/GSK-3 β pathway in chronic cerebral hypoperfusion rats," *European Journal of Pharmacology*, vol. 819, pp. 1–8, 2018.
- [89] L. Zhang, H. Dong, Y. Si et al., "miR-125b promotes tau phosphorylation by targeting the neural cell adhesion molecule in neuropathological progression," *Neurobiology of Aging*, vol. 73, pp. 41–49, 2019.
- [90] Z. Siegfried and I. Simon, "DNA methylation and gene expression," *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*, vol. 2, no. 3, pp. 362–371, 2010.
- [91] M. Gardiner-Garden and M. Frommer, "CpG islands in vertebrate genomes," *Journal of Molecular Biology*, vol. 196, no. 2, pp. 261–282, 1987.

- [92] K. Jabbari and G. Bernardi, "Cytosine methylation and CpG, TpG (CpA) and TpA frequencies," *Gene*, vol. 333, pp. 143–149, 2004.
- [93] D. Mastroeni, A. McKee, A. Grover, J. Rogers, and P. D. Coleman, "Epigenetic differences in cortical neurons from a pair of monozygotic twins discordant for Alzheimer's disease," *PLoS One*, vol. 4, no. 8, article e6617, 2009.
- [94] D. Mastroeni, A. Grover, E. Delvaux, C. Whiteside, P. D. Coleman, and J. Rogers, "Epigenetic changes in Alzheimer's disease: decrements in DNA methylation," *Neurobiology of Aging*, vol. 31, no. 12, pp. 2025–2037, 2010.
- [95] L. Chouliaras, D. Mastroeni, E. Delvaux et al., "Consistent decrease in global DNA methylation and hydroxymethylation in the hippocampus of Alzheimer's disease patients," *Neurobiology of Aging*, vol. 34, no. 9, pp. 2091–2099, 2013.
- [96] V. L. Keleshian, H. R. Modi, S. I. Rapoport, and J. S. Rao, "Retracted: aging is associated with altered inflammatory, arachidonic acid cascade, and synaptic markers, influenced by epigenetic modifications, in the human frontal cortex," *Journal of Neurochemistry*, vol. 125, no. 1, pp. 63–73, 2013.
- [97] V. Nicolia, V. Ciraci, R. A. Cavallaro, I. Ferrer, S. Scarpa, and A. Fusco, "GSK3 β 5'-flanking DNA methylation and expression in Alzheimer's disease patients," *Current Alzheimer Research*, vol. 14, no. 7, pp. 753–759, 2017.
- [98] V. Nicolia, A. Fusco, R. A. Cavallaro, A. Di Luzio, and S. Scarpa, "B vitamin deficiency promotes tau phosphorylation through regulation of GSK3 β and PP2A," *Journal of Alzheimer's Disease*, vol. 19, no. 3, pp. 895–907, 2010.
- [99] L. Li, C. Zhang, X. Zi, Q. Tu, and K. Guo, "Epigenetic modulation of Cdk5 contributes to memory deficiency induced by amyloid fibrils," *Experimental Brain Research*, vol. 233, no. 1, pp. 165–173, 2015.
- [100] J. V. Sanchez-Mut, E. Aso, H. Heyn et al., "Promoter hypermethylation of the phosphatase DUSP22 mediates PKA-dependent TAU phosphorylation and CREB activation in Alzheimer's disease," *Hippocampus*, vol. 24, no. 4, pp. 363–368, 2014.
- [101] B. E. Lonze, A. Riccio, S. Cohen, and D. D. Ginty, "Apoptosis, axonal growth defects, and degeneration of peripheral neurons in mice lacking CREB," *Neuron*, vol. 34, no. 3, pp. 371–385, 2002.
- [102] J. U. Guo, Y. Su, C. Zhong, G. L. Ming, and H. Song, "Hydroxylation of 5-methylcytosine by TET1 promotes active DNA demethylation in the adult brain," *Cell*, vol. 145, no. 3, pp. 423–434, 2011.
- [103] G. A. Kaas, C. Zhong, D. E. Eason et al., "TET1 controls CNS 5-methylcytosine hydroxylation, active DNA demethylation, gene transcription, and memory formation," *Neuron*, vol. 79, no. 6, pp. 1086–1093, 2013.
- [104] M. A. Bradley-Whitman and M. A. Lovell, "Epigenetic changes in the progression of Alzheimer's disease," *Mechanisms of Ageing and Development*, vol. 134, no. 10, pp. 486–495, 2013.
- [105] A. Rudenko, M. M. Dawlaty, J. Seo et al., "Tet1 is critical for neuronal activity-regulated gene expression and memory extinction," *Neuron*, vol. 79, no. 6, pp. 1109–1122, 2013.
- [106] A. H. Xu, Y. Yang, Y. X. Sun, and C. D. Zhang, "Exogenous brain-derived neurotrophic factor attenuates cognitive impairment induced by okadaic acid in a rat model of Alzheimer's disease," *Neural Regeneration Research*, vol. 13, no. 12, pp. 2173–2181, 2018.
- [107] G. Ambigapathy, Z. Zheng, and J. Keifer, "Regulation of *BDNF* chromatin status and promoter accessibility in a neural correlate of associative learning," *Epigenetics*, vol. 10, no. 10, pp. 981–993, 2015.
- [108] J. Penney and L. H. Tsai, "Histone deacetylases in memory and cognition," *Science Signaling*, vol. 7, no. 355, article re12, 2014.
- [109] M. Gonzalez-Zuñiga, P. S. Contreras, L. D. Estrada et al., "c-Abl stabilizes HDAC2 levels by tyrosine phosphorylation repressing neuronal gene expression in Alzheimer's disease," *Molecular Cell*, vol. 56, no. 1, pp. 163–173, 2014.
- [110] J. Graff, B. T. Woldemichael, D. Berchtold, G. Dewarrat, and I. M. Mansuy, "Dynamic histone marks in the hippocampus and cortex facilitate memory consolidation," *Nature Communications*, vol. 3, no. 1, p. 991, 2012.
- [111] B. Plagg, D. Ehrlich, K. Kniewallner, J. Marksteiner, and C. Humpel, "Increased acetylation of histone H4 at lysine 12 (H4K12) in monocytes of transgenic Alzheimer's mice and in human patients," *Current Alzheimer Research*, vol. 12, no. 8, pp. 752–760, 2015.
- [112] A. Fischer, F. Sananbenesi, X. Wang, M. Dobbin, and L. H. Tsai, "Recovery of learning and memory is associated with chromatin remodelling," *Nature*, vol. 447, no. 7141, pp. 178–182, 2007.
- [113] A. Ricobaraza, M. Cuadrado-Tejedor, A. Perez-Mediavilla, D. Frechilla, J. Del Rio, and A. Garcia-Osta, "Phenylbutyrate ameliorates cognitive deficit and reduces tau pathology in an Alzheimer's disease mouse model," *Neuropsychopharmacology*, vol. 34, no. 7, pp. 1721–1732, 2009.
- [114] D. Liu, H. Tang, X. Y. Li et al., "Targeting the HDAC2/HNF-4A/miR-101b/AMPK pathway rescues tauopathy and dendritic abnormalities in Alzheimer's disease," *Molecular Therapy*, vol. 25, no. 3, pp. 752–764, 2017.
- [115] D. Mastroeni, E. Delvaux, J. Nolz et al., "Aberrant intracellular localization of H3k4me3 demonstrates an early epigenetic phenomenon in Alzheimer's disease," *Neurobiology of Aging*, vol. 36, no. 12, pp. 3121–3129, 2015.
- [116] L. Zhang, S. Sheng, and C. Qin, "The role of HDAC6 in Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 33, no. 2, pp. 283–295, 2013.
- [117] C. Simoes-Pires, V. Zwick, A. Nurisso, E. Schenker, P. A. Carrupt, and M. Cuendet, "HDAC6 as a target for neurodegenerative diseases: what makes it different from the other HDACs?," *Molecular Neurodegeneration*, vol. 8, no. 1, p. 7, 2013.
- [118] L. Mahady, M. Nadeem, M. Malek-Ahmadi, K. Chen, S. E. Perez, and E. J. Mufson, "Frontal cortex epigenetic dysregulation during the progression of Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 62, no. 1, pp. 115–131, 2018.
- [119] J. Ma, X. Huo, M. B. Jarpe, A. Kavelaars, and C. J. Heijnen, "Pharmacological inhibition of HDAC6 reverses cognitive impairment and tau pathology as a result of cisplatin treatment," *Acta Neuropathologica Communications*, vol. 6, no. 1, p. 103, 2018.
- [120] H.-Y. Lee, S.-J. Fan, F.-I. Huang et al., "5-Aroylindoles act as selective histone deacetylase 6 inhibitors ameliorating Alzheimer's disease phenotypes," *Journal of Medicinal Chemistry*, vol. 61, no. 16, pp. 7087–7102, 2018.
- [121] A. Elobeid, H. Soininen, and I. Alafuzoff, "Hyperphosphorylated tau in young and middle-aged subjects," *Acta Neuropathologica*, vol. 123, no. 1, pp. 97–104, 2012.

- [122] H. Braak and K. Del Tredici, "The preclinical phase of the pathological process underlying sporadic Alzheimer's disease," *Brain*, vol. 138, no. 10, pp. 2814–2833, 2015.
- [123] P. Theofilas, S. Dunlop, H. Heinsen, and L. T. Grinberg, "Turning on the light within: subcortical nuclei of the isodentritic core and their role in Alzheimer's disease pathogenesis," *Journal of Alzheimer's Disease*, vol. 46, no. 1, pp. 17–34, 2015.
- [124] C. M. Dengler-Crish, M. A. Smith, and G. N. Wilson, "Early evidence of low bone density and decreased serotonergic synthesis in the dorsal raphe of a tauopathy model of Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 55, no. 4, pp. 1605–1619, 2017.
- [125] J. H. Lee, J. Ryan, C. Andreescu, H. Aizenstein, and H. K. Lim, "Brainstem morphological changes in Alzheimer's disease," *NeuroReport*, vol. 26, no. 7, pp. 411–415, 2015.
- [126] J. K. Holth, T. K. Patel, and D. M. Holtzman, "Sleep in Alzheimer's disease—beyond amyloid," *Neurobiology of Sleep and Circadian Rhythms*, vol. 2, pp. 4–14, 2017.
- [127] G. Simic, G. Stanic, M. Mladinov, N. Jovanov-Milosevic, I. Kostovic, and P. R. Hof, "Does Alzheimer's disease begin in the brainstem?," *Neuropathology and Applied Neurobiology*, vol. 35, no. 6, pp. 532–554, 2009.
- [128] X. Guo, M. Waern, K. Sjögren et al., "Midlife respiratory function and incidence of Alzheimer's disease: a 29-year longitudinal study in women," *Neurobiology of Aging*, vol. 28, no. 3, pp. 343–350, 2007.
- [129] H. Braak, D. R. Thal, E. Ghebremedhin, and K. Del Tredici, "Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years," *Journal of Neuropathology & Experimental Neurology*, vol. 70, no. 11, pp. 960–969, 2011.
- [130] U. Rub, K. Del Tredici, C. Schultz, D. R. Thal, E. Braak, and H. Braak, "The evolution of Alzheimer's disease-related cytoskeletal pathology in the human raphe nuclei," *Neuropathology and Applied Neurobiology*, vol. 26, no. 6, pp. 553–567, 2000.
- [131] J. J. Rodriguez, H. N. Noristani, and A. Verkhatsky, "The serotonergic system in ageing and Alzheimer's disease," *Progress in Neurobiology*, vol. 99, no. 1, pp. 15–41, 2012.
- [132] M. K. P. Lai, S. W. Y. Tsang, P. T. Francis et al., "Postmortem serotonergic correlates of cognitive decline in Alzheimer's disease," *Neuroreport*, vol. 13, no. 9, pp. 1175–1178, 2002.
- [133] A. J. Ehrenberg, A. K. Nguy, P. Theofilas et al., "Quantifying the accretion of hyperphosphorylated tau in the locus coeruleus and dorsal raphe nucleus: the pathological building blocks of early Alzheimer's disease," *Neuropathology and Applied Neurobiology*, vol. 43, no. 5, pp. 393–408, 2017.
- [134] P. Theofilas, A. J. Ehrenberg, A. Nguy et al., "Probing the correlation of neuronal loss, neurofibrillary tangles, and cell death markers across the Alzheimer's disease Braak stages: a quantitative study in humans," *Neurobiology of Aging*, vol. 61, pp. 1–12, 2018.
- [135] P. Andres-Benito, R. Delgado-Morales, and I. Ferrer, "Altered regulation of KIAA0566, and katanin signaling expression in the locus coeruleus with neurofibrillary tangle pathology," *Frontiers in Cellular Neuroscience*, vol. 12, p. 131, 2018.
- [136] F. Llorens, K. Thüne, P. Andrés-Benito et al., "MicroRNA expression in the locus coeruleus, entorhinal cortex, and hippocampus at early and middle stages of Braak neurofibrillary tangle pathology," *Journal of Molecular Neuroscience*, vol. 63, no. 2, pp. 206–215, 2017.
- [137] P. A. Jones, J.-P. J. Issa, and S. Baylin, "Targeting the cancer epigenome for therapy," *Nature Reviews Genetics*, vol. 17, no. 10, pp. 630–641, 2016.
- [138] M. Esteller, "Non-coding RNAs in human disease," *Nature Reviews Genetics*, vol. 12, no. 12, pp. 861–874, 2011.
- [139] P. D. Fransquet and J. Ryan, "Micro RNA as a potential blood-based epigenetic biomarker for Alzheimer's disease," *Clinical Biochemistry*, vol. 58, pp. 5–14, 2018.
- [140] S. Nagaraj, K. M. Zoltowska, K. Laskowska-Kaszub, and U. Wojda, "MicroRNA diagnostic panel for Alzheimer's disease and epigenetic trade-off between neurodegeneration and cancer," *Ageing Research Reviews*, vol. 49, pp. 125–143, 2019.
- [141] A. L. Baird, S. Westwood, and S. Lovestone, "Blood-based proteomic biomarkers of Alzheimer's disease pathology," *Frontiers in Neurology*, vol. 6, p. 236, 2015.



Hindawi

Submit your manuscripts at
www.hindawi.com

