Research Article

Pomegranate-Derived Polyphenols Reduce Reactive Oxygen Species Production via SIRT3-Mediated SOD2 Activation

Chong Zhao,¹ Takenori Sakaguchi,² Kosuke Fujita,¹ Hideyuki Ito,³ Norihisa Nishida,⁴ Akifumi Nagatomo,⁴ Yukimasa Tanaka-Azuma,⁴ and Yoshinori Katakura^{1,2,5}

¹Graduate School of Systems Life Sciences, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

²Graduate School of Bioscience & Biotechnology, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

³Faculty of Health and Welfare Science, Okayama Prefectural University, 111 Kuboki, Soja, Okayama 719-1197, Japan

⁴Morishita Jintan Co., Ltd., 2-11-1 Tsudayamate, Hirakata, Osaka 573-0128, Japan

⁵Faculty of Agriculture, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

Correspondence should be addressed to Yoshinori Katakura; katakura@grt.kyushu-u.ac.jp

Received 12 May 2016; Revised 12 August 2016; Accepted 21 September 2016

Academic Editor: Gabriele Saretzki

Copyright © 2016 Chong Zhao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pomegranate-derived polyphenols are expected to prevent life-style related diseases. In this study, we evaluated the ability of 8 pomegranate-derived polyphenols, along with other polyphenols, to augment *SIRT3*, a mammalian *SIR2* homolog localized in mitochondria. We established a system for screening foods/food ingredients that augment the *SIRT3* promoter in Caco-2 cells and identified 3 *SIRT3*-augmenting pomegranate-derived polyphenols (eucalbanin B, pomegranin A, and eucarpanin T₁). Among them, pomegranin A activated superoxide dismutase 2 (SOD2) through *SIRT3*-mediated deacetylation, thereby reducing intracellular reactive oxygen species. The other *SIRT3*-augmenting polyphenols tested also activated SOD2, suggesting antioxidant activity. Our findings clarify the underlying mechanisms involved in the antioxidant activity of pomegranin A.

1. Introduction

Calorie restriction (CR) is the only experimental manipulation that is known to reduce the incidence of age-related disorders such as diabetes, cancer, and cardiovascular diseases in mammals [1]. Sirtuins are NAD⁺-dependent deacetylases that have been found to mediate the effects of CR and regulate lifespan in lower organisms [2]. In mammals, there are seven sirtuins (*SIRT1*–*SIRT7*) localized in diverse positions within cells [3]. *SIRT3*, a mammalian *SIR2* homolog localized in the mitochondria, regulates several aspects of mitochondrial function, including metabolism, energy homeostasis, and modulation of the response to oxidative stress [4]. Mitochondrial free radical theory of ageing suggests that accumulated oxidative stress caused by reactive oxygen species (ROS) is a major factor in determining lifespan [5]. Recently, increasing attention has been paid to the anti-aging effects of *SIRT3*, including a reduction in oxidative stress by targeting superoxide dismutase 2 (SOD2) enzyme [6] and preventing

age-related hearing loss by activating mitochondrial isocitrate dehydrogenase 2 (IDH2) [7]. Consequently, *SIRT3*-activating compounds are promising candidates for suppressing oxidative stress and reducing age-related disorders.

The pomegranate, *Punica granatum* L., is known to contain many kinds of ellagitannins and anthocyanins [8, 9]. A large body of *in vitro* and *in vivo* evidence indicates that pomegranate peel extracts possess antioxidant, anticarcinogenic, and anti-inflammatory properties that may be useful in the treatment and prevention of cancer, cardiovascular disease, diabetes, dental conditions, bacterial infections, and ultraviolet-induced skin damage [10–20]. Furthermore, *in vivo* experiments have shown that pomegranate juice exhibits an inhibitory effect toward age-related diseases such as Alzheimer's disease [21] and Parkinson's disease [22]. In the present study, we focused on the functionalities of pomegranate-derived polyphenols through augmentation of *SIRT3*. Pomegranate-derived polyphenols are difficult to be absorbed in intestine; thus in the present study, we

TABLE 1: Polyphenols.

Classification	Name	Origin
Polyphenol	Punicalin	Pomegranate
	Punicalagin	Pomegranate
	Oenothien B	Pomegranate
	Eucalbanin B	Pomegranate
	Eucarpanin T ₁	Pomegranate
	Pomegranin A	Pomegranate
	Urolithin A	Ellagitannin metabolite
	Chlorogenic acid	Coffee
	Resveratrol	Grape
Anthocyanin	Tellimagrandin II	Rose
	Cyanidin 3-glucoside chloride	Pomegranate
	Delphinidin 3-glucoside chloride	Pomegranate
Flavonoid	Tiliroside	Rose hip
	Saponarin	Green barley
	Isovitexin	Green barley
	Fisetin	Green barley
	Kaempferol	Tea
Terpenoid	Cafestol	Coffee beans
	Kahweol	Coffee beans
Lignan	Sesamin	Sesame
	Sesamolin	Sesame
	Episesamin	Sesame
	Sesamol	Sesame

investigated the effects of these polyphenols on intestinal cells by using human colorectal cancer cell line, Caco-2, as a model cell [23].

2. Materials and Methods

2.1. Cell Culture and Reagents. The Caco-2 human colorectal cancer cell line (Riken Bioresource Center, Tsukuba, Japan) was cultured in Dulbecco's modified Eagle's medium (DMEM; Nissui, Tokyo, Japan) supplemented with 10% heat-inactivated fetal bovine serum (Life Technologies, Gaithersburg, MD, USA) at 37°C in an atmosphere containing 5% CO₂.

2.2. Polyphenols. Polyphenols used in this study are listed in Table 1. The structure of ellagitannins is described elsewhere [9].

2.3. Screening System for Foods Augmenting the SIRT3 Promoter. We amplified the human SIRT3 promoter (−653 to −1) by PCR using human genomic DNA as a template, which was then cloned into the promoter-less Enhanced Green Fluorescent Protein- (EGFP-) expressing vector (pEGFP-C3, Takara, Shiga, Japan) and designated pSIRT3p-EGFP. Caco-2 cells transduced with pSIRT3p-EGFP were used to evaluate human SIRT3 promoter activity. Changes in the EGFP

fluorescence derived from pSIRT3p-EGFP were monitored by an IN Cell Analyzer 1000 (GE Healthcare, Amersham Place, UK).

2.4. Quantitative Reverse Transcription Polymerase Chain Reaction. RNA was extracted using the High Pure RNA Isolation Kit (Roche, Tokyo, Japan), and cDNA was prepared as previously described [24]. Quantitative reverse transcription polymerase chain reaction (qRT-PCR) was performed using Thunderbird SYBR qPCR Mix (Toyobo, Osaka, Japan) and a Thermal Cycler Dice Real Time System TP-800 instrument (Takara, Shiga, Japan) as previously described. Samples were analyzed in triplicate, and SIRT3 expression levels were normalized to the corresponding ACTB (β-actin) levels. The PCR primer sequences used were as follows: SIRT3 forward primer 5'-CTGTACAGCAACCTCCAGCA-3' and reverse primer 5'-CTCCTTGGCCAAAGTGAAAA-3'; ACTB forward primer 5'-TGGCACCCAGCACAATGA-A-3' and reverse primer 5'-CTAAGTCATAGTCCGCCT-AGAAGCA-3'.

2.5. Assays for Antioxidant Defense System. SOD2 activity was measured with the SOD Assay Kit-WST (Dojindo, Kumamoto, Japan), following the manufacturer's instructions. The amount of reduced glutathione (GSH) and oxidized glutathione (GSSG) was measured with the GSSG/GSH Quantification Kit (Dojindo), following the manufacturer's instructions.

2.6. Immunoprecipitation. SOD2 cDNA fragments derived from pBI-EGFP-MnSOD (#16612; Addgene, Cambridge, MA, USA) were cloned into the FLAG-pcDNA3 vector and designated as pcDNA3-Flag-SOD2. Caco-2 cells were transduced with pcDNA3-Flag-SOD2 and treated with polyphenols (10 μM) for 48 h. Proteins were extracted by NP-40 lysis buffer (0.5% NP-40, 5 mM ethylenediaminetetraacetic acid (EDTA), 2 mM Na₃VO₄, 10 mM Tris-Cl pH 7.6, 150 mM NaCl, 5 μg/mL aprotinin, and 1 mM phenylmethylsulfonyl fluoride (PMSF)). SOD2 was immunoprecipitated with anti-Flag antibodies (M2; Sigma, St. Louis, MO, USA) and then immobilized with PureProteome Protein G Magnetic Beads (Merck Millipore, Billerica, MA, USA) overnight at 4°C. Immunoprecipitates were eluted with sample buffer (1 M Tris-HCl pH 6.8, 10% SDS, and 10% 2 ME).

2.7. Immunoblotting. Cell lysates were resolved by electrophoresis using 12% SDS-PAGE and transferred to a Hybond P membrane (GE Healthcare). The membrane was probed with an anti-Flag M2 antibody, anti-acetylated-lysine antibody (Cell Signaling Technology, Danvers, MA, USA), or anti-β-actin antibody (G043; Applied Biological Materials Inc., Richmond, BC, Canada). Horseradish peroxidase-labeled anti-rabbit IgG antibody (GE Healthcare) and anti-mouse IgG antibody (GE Healthcare) were used as secondary antibodies. The proteins were detected using ImmunoStar LD (Wako Pure Chemical, Osaka, Japan) and visualized with a LAS-1000 Lumino Image analyzer (Fuji Film, Tokyo, Japan).

2.8. Retrovirus Production and Transduction. Viral supernatants were produced after transfecting 293 T cells with pGag-pol, pVSV-G, and individual expression vectors (pSUPER-puro, pSUPER-puro-shSIRT1, and pSUPER-puro-shSIRT3) using the HilyMax reagent (Dojindo) as previously described [25]. The cells were cultured at 37°C in DMEM supplemented with 10% FBS for 24 h. Medium was replaced with fresh DMEM supplemented with 2% FBS and incubated for an additional 24 h. Viral supernatant was collected and supplemented with 10 mg/mL polybrene (Merck Millipore). The target cells were infected with this viral supernatant for 24 h at 37°C. After infection, the cells were selected with 3 µg/mL puromycin (Enzo Life Sciences, Farmingdale, NY) for 3 days.

2.9. Short Hairpin RNA. The oligonucleotides containing siRNA-expressing sequences targeting human *SIRT1* and *SIRT3* were annealed (shSIRT1 top: 5'-GATCCCCGCAAGCCTTTCTGAATCTATTTCTGAAGAGATAGATTTCAGAAAGGCTTTGCTTTTTA-3', shSIRT1 bottom: AGCTTAAAAAGCAAGCCTTTCTGAATCTATCTCTTCGAAATAGATTTCAGAAAGGCTTTGCGGG-3'; shSIRT3 top: 5'-GATCCCCCAACGTCACCTACTACTTTTTCGAAGAGAAAGTAGTGAGTGACGTTGGGTTTTA-3', shSIRT3 bottom: 5'-AGCTTAAAAACCAACGTCACCTACTACTTTCTCTTCGAAAAAGTAGTGAGTGACGTTGGGGGG-3') and cloned into the pSUPER.retro.puro (OligoEngine, Seattle, WA, USA).

2.10. Evaluation of ROS Generation. Production of intracellular ROS in Caco-2 cells was measured using BES-H₂O₂-Ac (Wako Pure Chemical) according to the manufacturer's protocol. Caco-2 cells were treated with 10 µM pomegranin A for 48 h or 10 mM N-acetyl cysteine (NAC) for 2 h before analyzing ROS production. The medium was removed and cells were washed twice with 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer (pH 7.4). The cells were incubated with 5 µM BES-H₂O₂-Ac and Hoechst 33342 (Dojin) in HEPES (pH 7.4) at 37°C for 30 min, followed by detection with an IN Cell Analyzer 1000 (GE Healthcare).

2.11. Statistical Analysis. All experiments were performed at least 3 times, and the corresponding data are shown. The results are presented as mean ± standard deviation. Statistical significance was determined by one-way ANOVA with Tukey's *post hoc* test. Statistical significance was defined as $P < 0.05$ (* $P < 0.05$; ** $P < 0.01$).

3. Results

3.1. Identification of Polyphenols That Augment SIRT3 Transcription in Caco-2 Cells. Previously, we established a system for screening foods/food ingredients that augment the *SIRT1* promoter in Caco-2 cells and identified *Lactobacillus brevis* T2102 as *SIRT1*-activating lactic acid bacterium [26]. Here, we established a similar system for screening foods/food ingredients that augment the *SIRT3* promoter in Caco-2 cells, where pSIRT3p-EGFP was transduced. We tested 11 polyphenols, 5 flavonoids, 2 terpenoids, and 4 lignans in this assay. As shown in Figure 1(a), several samples including

tellimagrandin II, fisetin, oenothien B, eucalbanin B, resveratrol, eucarpanin T₁, pomegranin A, and kaempferol strongly augmented *SIRT3* promoter activity in Caco-2 cells. In addition, we tested whether these polyphenols augment endogenous *SIRT3* expression in Caco-2 cells by qRT-PCR. Dose-dependent change in polyphenol-induced *SIRT3* augmentation showed that treatment with 10 µM polyphenols is most effective and that polyphenols, including eucalbanin B, pomegranin A, eucarpanin T₁, and fisetin, increased endogenous *SIRT3* expression in Caco-2 cells to a level comparable to that observed in the positive control samples treated with kaempferol and resveratrol (Figure 1(b)).

3.2. Effects of SIRT3-Augmenting Polyphenols on SOD2 Activity. Qiu et al. reported that in 293 T cells *SIRT3* activates SOD2, a major mitochondrial antioxidant enzyme [6]; therefore, we examined whether *SIRT3*-augmenting polyphenols activate SOD2 activity in Caco-2 cells. Our results show that all polyphenols tested activate SOD2 activity in Caco-2 cells (Figure 2(a)), and foods/food ingredients that activate SOD2 activity can be screened by our system, which targets the *SIRT3* promoter. Among these polyphenols, eucalbanin B and pomegranin A are polyphenols that have not been reported as a sirtuin activator and reproducibly activate SOD2 as well as augmenting *SIRT3* expression in independent experiment; thus we used these polyphenols for further experiments.

Qiu et al. reported that *SIRT3* deacetylates and activates SOD2 [6]; therefore, we wanted to clarify the molecular mechanisms for activation of SOD2 by *SIRT3*-augmenting polyphenols. We tested whether *SIRT3*-augmenting polyphenols deacetylate SOD2 in Caco-2 cells, with a focus on the activity of eucalbanin B and pomegranin A, which are derived from pomegranate [9]. Caco-2 cells were transfected with Flag-SOD2 and treated with the test polyphenols; SOD2 was then immunoprecipitated with anti-Flag antibodies. The acetylation status of immunoprecipitated SOD2 was evaluated by immunoblotting using anti-acetylated-lysine antibodies (Figure 2(b)). Results show that eucalbanin B and pomegranin A significantly attenuated acetylation of SOD2 (Figures 2(b) and 2(c)), suggesting that these *SIRT3*-augmenting polyphenols activate SOD2 by deacetylation.

3.3. Pomegranin A Activation of SOD2 Is Dependent on SIRT3. To further clarify the involvement of *SIRT3* in the activation of SOD2 by eucalbanin B and pomegranin A, we examined their effects on SOD2 activity in *SIRT1*- or *SIRT3*-silenced Caco-2 cells (Figure 3). Results show that eucalbanin B and pomegranin A did not attenuate SOD2 activation in *SIRT1*-silenced Caco-2 cells. Furthermore, activation of SOD2 by pomegranin A, but not by eucalbanin B, was significantly reduced in *SIRT3*-silenced Caco-2 cells, suggesting that pomegranin A activates SOD2 in a *SIRT3*-dependent manner. However, molecular mechanisms for eucalbanin B-induced activation of SOD2 remain unclear.

3.4. Pomegranin A Reduced ROS Levels in Caco-2 Cells. As described above, pomegranin A activates SOD2 in Caco-2 cells, suggesting that it has antioxidant properties. Therefore, we investigated the effects of pomegranin A on intracellular

FIGURE 1: Identification of polyphenols that augment *SIRT3* transcription in Caco-2 cells. Polyphenols (10 μM) and their corresponding amount of DMSO were added to Caco-2 cells transduced with *SIRT3p*-EGFP and changes in EGFP fluorescence were monitored by an IN Cell Analyzer 1000 (a). Dose-dependent effects of polyphenols on the expression of endogenous *SIRT3* in Caco-2 cells were assessed by qRT-PCR. Control Caco-2 cells were treated with the same volume of DMSO used for treating with 1, 10, and 50 μM of polyphenols. (b) The results are expressed as mean ± standard deviation. Statistical significance between the respective concentration and its corresponding in the control was determined by one-way ANOVA with Tukey's *post hoc* test. Statistical significance was defined as $P < 0.05$ (* $P < 0.05$; ** $P < 0.01$). C3g, cyanidin 3-glucoside chloride; D3g, delphinidin 3-glucoside chloride; CA, chlorogenic acid.

ROS levels using an IN Cell Analyzer 1000 that allowed collection of fluorescent images derived from BES-H₂O₂. We evaluated intracellular ROS levels by analyzing cell size (x -axis) as well as the percentage of BES-H₂O₂-positive/negative cells (y -axis). The scatter plot shows that the relative number of cells with high ROS levels decreased upon treatment with pomegranin A and NAC. Specifically, the relative number of cells with high ROS levels decreased from 20% (control) to 14% and 4.5% by the treatment with pomegranin A and NAC, respectively (Figure 4(a)). Furthermore, the reduction in ROS levels observed in cells treated with pomegranin A was attenuated in *SIRT3*-silenced Caco-2 cells (Figure 4(b)). These results indicate that pomegranin A-associated decreases in intracellular ROS levels are dependent on *SIRT3* expression.

3.5. Effects of *SIRT3*-Augmenting Polyphenols on Glutathione Antioxidant System. Someya et al. reported that *SIRT3* increased ratio of reduced-to-oxidized glutathione through direct deacetylation of mitochondrial isocitrate dehydrogenase 2 [7]; then we examined whether *SIRT3*-augmenting polyphenols activate glutathione antioxidant system as well as SOD2 in Caco-2 cells. Results show that 3 polyphenols among *SIRT3*-augmenting polyphenols increased the ratio of GSH/GSSG, although all polyphenols activate SOD2 activity (Figure 5).

4. Discussion

Antioxidant activity and *in vivo* functionalities of pomegranate extract such as repair and maintenance of hepatic and intestinal function and attenuation of inflammation have

been extensively studied [13–20]. Ellagitannins, anthocyanidins, and related polyphenols are thought to be efficient antioxidants extracted from pomegranate (*Punica granatum* L.). Previous studies have shown that three major anthocyanidins (delphinidin, cyanidin, and pelargonidin) contribute to the antioxidant activity of pomegranate fruit [27]. Recently, studies have shown that new ellagitannin oligomers isolated from pomegranate exhibit potent inhibitory effects on the formation of advanced glycation end-products (AGEs), which have been associated with life-style related diseases including Alzheimer's disease [9]. Therefore, a better understanding of the mechanisms underlying the antioxidant activity of pomegranate-derived polyphenols is critical for their further development as anti-ageing chemopreventive agents. For this purpose, we report here that pomegranin A has potent antioxidant properties that are dependent on *SIRT3*-mediated SOD2 activation in Caco-2 cells.

Several food-derived bioactive polyphenols have been reported to attenuate cellular oxidative stress, and thus it has been suggested that they have the potential to prevent age-related diseases [28, 29]. These polyphenols exert their antioxidant activity through activation of various signaling pathways, including JAK-STAK, MAPK/ERK, NF-κB, and MAPK p38/JNK [30]. Several polyphenols (resveratrol, black chokeberry (*Aronia melanocarpa*) extract, and green tea polyphenols) have been shown to activate *SIRT3* [31–33]. In particular, decreased *SIRT3* expression and increased SOD2 acetylation resulting from a high-fat diet in Wister rats were attenuated by administration of green tea polyphenols, which likely explains the reduction in oxidative stress observed in the kidney tissues of the rats [32]. In the present study, we

FIGURE 2: Effects of SIRT3-augmenting polyphenols on SOD2 activity. Caco-2 cells were treated with SIRT3-augmenting polyphenols (10 μ M) for 48 h and SOD2 activity was measured using the SOD Assay Kit-WST (a). Deacetylation of SOD2 by pomegranin A; Caco-2 cells were transduced with Flag-SOD2 and treated with 10 μ M of eucalbanin B or pomegranin A for 48 h. Flag-SOD2 was immunoprecipitated with anti-Flag M2 antibodies immobilized to Protein G and analyzed by immunoblot using anti-acetylated-lysine antibodies (b and c). Band intensity corresponding to acetylated SOD2 was quantitatively determined using Image J software. The results are expressed as mean $\pm$ standard deviation. Control Caco-2 cells were treated with the same volume of DMSO used for treating with 10 μ M of polyphenols. Statistical significance between the respective polyphenol and control was determined by one-way ANOVA with Tukey's *post hoc* test. Statistical significance was defined as $P < 0.05$ (* $P < 0.05$; ** $P < 0.01$).

clarified the SIRT3-augmenting activity of the polyphenol pomegranin A, which activates SOD2 and reduces intracellular ROS in an *in vitro* system. These data provide direct evidence for the functionality of SIRT3-augmenting polyphenols [33]. Furthermore, these results confirm the utility of the model described here for identifying foods/food ingredients that increase SIRT3 expression and consequently reduce intracellular ROS. Although eucalbanin B was also identified as a SIRT3-augmenting polyphenol in this system, SOD2 activity increased in eucalbanin B-treated cells was not attenuated in the SIRT3-silenced Caco-2 cells. These results suggest that eucalbanin B activates SOD2 via a pathway that is distinct from SIRT3 activation, which will be pursued further in a future study.

In the present study, we identified 6 polyphenols among 22 polyphenols that augment SIRT3 expression and activate SOD2. Although we focused on eucalbanin B and pomegranin A as SIRT3-augmenting polyphenols, other polyphenols may elicit their effects by mechanisms similar to pomegranin A or by activating individual pathways. Until now, a limited number of polyphenols and food derived extracts were known to activate SIRT3 and showed distinct functions. Black chokeberry extract regulates lipid metabolism and lipoprotein assembly through the activation of SIRT3 [31]. NAD⁺ and grape wine polyphenols prevent axonal apoptosis through SIRT3-dependent mechanisms [34]. Resveratrol activates SIRT3, which then ameliorates cardiac fibrosis and improves cardiac function via

FIGURE 3: SIRT3-dependency of pomegranin A-induced SOD2 activation. Effects of eucalbanin B (a) and pomegranin A (b) on SOD2 activity in *SIRT1*- and *SIRT3*-silenced Caco-2 cells were evaluated using the SOD Assay Kit-WST. The results are expressed as mean $\pm$ standard deviation. Control Caco-2 cells were treated with the same volume of DMSO used for treating with $10 \mu\text{M}$ of polyphenols. Statistical significance between the respective polyphenol and control was determined by one-way ANOVA with Tukey's *post hoc* test. Statistical significance was defined as $P < 0.05$ (* $P < 0.05$; ** $P < 0.01$). SOD2, superoxide dismutase 2.

FIGURE 4: Pomegranin A reduces intracellular ROS levels via SIRT3. Effects of pomegranin A on intracellular ROS levels in Caco-2 cells; cells were treated with $10 \mu\text{M}$ pomegranin A for 48 h or 10 mM NAC for 2 h prior to detection, and intracellular ROS levels were analyzed with an IN Cell Analyzer 1000. The relative number of the cells with high ROS levels was shown in the scatter plot (a). Effects of pomegranin A on intracellular ROS levels in *SIRT3*-silenced Caco-2; cells were treated with $10 \mu\text{M}$ pomegranin A for 48 h, followed by detection of ROS levels with an IN Cell Analyzer 1000. The relative number of the cells with high ROS levels was shown in the scatter plot (b). Control Caco-2 cells were treated with the same volume of DMSO used for treating with $10 \mu\text{M}$ of polyphenols. Statistical significance between samples and control was determined by one-way ANOVA with Tukey's *post hoc* test. Statistical significance was defined as $P < 0.05$ (** $P < 0.01$). NAC, N-acetyl cysteine; ROS, reactive oxygen species; DMSO, dimethyl sulfoxide. Experiments were repeated three times, and representative was shown.

FIGURE 5: Effects of SIRT3-augmenting polyphenols on glutathione antioxidant system. Caco-2 cells were treated with SIRT3-augmenting polyphenols (10 μ M) for 48 h, and the amount of reduced glutathione (GSH) and oxidized glutathione (GSSG) was measured with the GSSG/GSH Quantification Kit, following the manufacturer's instructions. Control Caco-2 cells were treated with the same volume of DMSO used for treating with 10 μ M of polyphenols. Statistical significance between the respective polyphenol and control was determined by one-way ANOVA with Tukey's *post hoc* test. Statistical significance was defined as $P < 0.05$ (* $P < 0.05$; ** $P < 0.01$).

the transforming growth factor- β /Smad3 pathway [35]. Thus, other SIRT3-augmenting polyphenols have distinct functions that must be clarified in future studies.

Although pomegranin A activation of SOD2 occurs only via SIRT3, several polyphenols, such as kaempferol, have been reported to activate both SIRT1 and SIRT3 [36]. Furthermore, activation of SIRT1 is known to promote the nuclear factor-E2 related factor 2- (Nrf2-) antioxidant response element (ARE) pathway [37]. Here, we determined that the antioxidant activity of pomegranin A occurs through activation of the SIRT3-SOD2 pathway. Although pomegranin A and eucalbanin B are ellagitannin tetramer and dimer, respectively [9], with similar structural features, SOD2 activation by these ellagitannins showed distinct dependency on SIRT3. SOD2 activation by pomegranin A highly depended on SIRT3, but that by eucalbanin B did not, suggesting that eucalbanin B activates SOD2 by not only SIRT3, but also other molecules such as SIRT1 as described previously [38]. In future studies, we will clarify additional mechanisms associated with the antioxidant activity of pomegranin A.

Superoxide is one of the most abundant ROS produced by mitochondria. The extent of ROS-induced cellular damage accumulates with age. Thus, SOD-catalyzed reductions in ROS levels function to regulate intracellular ROS levels. In particular, overexpression of SOD in model organisms, including *Saccharomyces cerevisiae* and *Drosophila*, reduces oxidative damage and extends lifespan [39]. Currently, the potential for pomegranate-derived polyphenols to prevent and treat life-style related diseases has been extensively reported [10–12]. SOD2 activation by pomegranin A likely

provides a molecular basis for the broad spectrum of antioxidant activity observed for pomegranate juice. However, polyphenols having strong antioxidant activities do not always show similar biological activities *in vivo* due to their bioavailability, uptake, and metabolism. Thus, we must clarify biological activity of pomegranin A *in vivo* as well as its intestinal uptake, metabolism, and kinetics in blood by using animal models and reveal the molecular basis of *in vivo* function of pomegranin A.

SOD2 and isocitrate dehydrogenase 2, which regulates the ratio of GSH to GSSG, were reported to be targets of SIRT3 [40]; thus we investigated the effects of these polyphenols on SOD2 activity and GSH/GSSG ratio. Among SIRT3-augmenting polyphenols, 3 polyphenols including fisetin, eucalbanin B, and pomegranin A activated both SOD2 and glutathione antioxidant system, while other polyphenols including resveratrol, eugenin, and kaempferol only activated SOD2. These results suggest that there exist polyphenol-dependent activation mechanisms for cellular antioxidant system. Furthermore, we would investigate the effects of these polyphenols on other antioxidant enzymes including SOD1, glutathione peroxidase, glutathione reductase, and catalase in the future study.

In summary, we identified a pomegranate-derived polyphenol, pomegranin A, that eliminates ROS through SIRT3-dependent SOD2 activation in Caco-2 cells, which would ameliorate intestinal injury induced by hemorrhagic shock as reported previously [41]. Furthermore, this effect of pomegranin A on intestinal cells might lead to the activation of interorgan interaction and prevention of age-related disorders, the possibility of which must be evaluated in the future study [23]. However, the homogenate of fresh arils (2 kg) of pomegranate contains only 11 mg (3.5 μ mol) of pomegranin A. Thus, daily intake of pomegranin A as a supplement is thought to be more desirable.

Disclosure

The funder had no role in study design, data collection, and analysis.

Competing Interests

The authors declare that they have no competing interests.

Acknowledgments

The authors would like to thank N. Oshima (GE Healthcare) for her expert assistance with the IN Cell Analyzer 1000. This work was supported by MAFF Grant.

References

- [1] R. J. Colman, T. M. Beasley, J. W. Kemnitz, S. C. Johnson, R. Weindruch, and R. M. Anderson, "Caloric restriction reduces age-related and all-cause mortality in rhesus monkeys," *Nature Communications*, vol. 5, article 3557, 2014.
- [2] G. Donmez and L. Guarente, "Aging and disease: connections to sirtuins," *Aging Cell*, vol. 9, no. 2, pp. 285–290, 2010.

- [3] T. Finkel and N. J. Holbrook, "Oxidants, oxidative stress and the biology of ageing," *Nature*, vol. 408, no. 6809, pp. 239–247, 2000.
- [4] L. C. Heather, X. Wang, J. A. West, and J. L. Griffin, "A practical guide to metabolomic profiling as a discovery tool for human heart disease," *Journal of Molecular and Cellular Cardiology*, vol. 55, no. 1, pp. 2–11, 2013.
- [5] R. S. Balaban, S. Nemoto, and T. Finkel, "Mitochondria, oxidants, and aging," *Cell*, vol. 120, no. 4, pp. 483–495, 2005.
- [6] X. Qiu, K. Brown, M. D. Hirschey, E. Verdin, and D. Chen, "Calorie restriction reduces oxidative stress by SIRT3-mediated SOD2 activation," *Cell Metabolism*, vol. 12, no. 6, pp. 662–667, 2010.
- [7] S. Someya, W. Yu, W. C. Hallows et al., "Sirt3 mediates reduction of oxidative damage and prevention of age-related hearing loss under caloric restriction," *Cell*, vol. 143, no. 5, pp. 802–812, 2010.
- [8] J. Jurenka, "Therapeutic applications of pomegranate (*Punica granatum* L.): a review," *Alternative Medicine Review*, vol. 13, no. 2, pp. 128–144, 2008.
- [9] H. Ito, P. Li, M. Koreishi, A. Nagatomo, N. Nishida, and T. Yoshida, "Ellagitannin oligomers and a neolignan from pomegranate arils and their inhibitory effects on the formation of advanced glycation end products," *Food Chemistry*, vol. 152, pp. 323–330, 2014.
- [10] E. Turrini, L. Ferruzzi, and C. Fimognari, "Potential effects of pomegranate polyphenols in cancer prevention and therapy," *Oxidative Medicine and Cellular Longevity*, vol. 2015, Article ID 938475, 19 pages, 2015.
- [11] A. Zarfeshany, S. Asgary, and S. H. Javanmard, "Potent health effects of pomegranate," *Advanced Biomedical Research*, vol. 3, article 100, 2014.
- [12] S. Banihani, S. Swedan, and Z. Alguraan, "Pomegranate and type 2 diabetes," *Nutrition Research*, vol. 33, no. 5, pp. 341–348, 2013.
- [13] K. N. Chidambara Murthy, G. K. Jayaprakasha, and R. P. Singh, "Studies on antioxidant activity of pomegranate (*Punica granatum*) peel extract using in vivo models," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 17, pp. 4791–4795, 2002.
- [14] H. Z. Toklu, M. U. Dumlu, Ö. Sehirli et al., "Pomegranate peel extract prevents liver fibrosis in biliary-obstructed rats," *Journal of Pharmacy and Pharmacology*, vol. 59, no. 9, pp. 1287–1295, 2007.
- [15] R. Bachoual, W. Talmoudi, T. Boussetta, F. Braut, and J. El-Benna, "An aqueous pomegranate peel extract inhibits neutrophil myeloperoxidase in vitro and attenuates lung inflammation in mice," *Food and Chemical Toxicology*, vol. 49, no. 6, pp. 1224–1228, 2011.
- [16] M. I. Waly, A. Ali, N. Guizani, A. S. Al-Rawahi, S. A. Farooq, and M. S. Rahman, "Pomegranate (*Punica granatum*) peel extract efficacy as a dietary antioxidant against azoxymethane-induced colon cancer in rat," *Asian Pacific Journal of Cancer Prevention*, vol. 13, no. 8, pp. 4051–4055, 2012.
- [17] F. E. Agha, M. M. Hassannane, E. A. Omara, A. M. Hasan, and S. A. El-Toumy, "Protective effect of Punica granatum peel extract against pentachlorophenol-induced oxidative stress, cytogenetic toxicity and hepatic damage in rats," *Australian Journal of Basic and Applied Sciences*, vol. 7, pp. 853–864, 2013.
- [18] O. S. O. Amer, M. A. Dkhil, W. M. Hikil, and S. Al-Quraishy, "Antioxidant and anti-inflammatory activities of pomegranate (*Punica granatum*) on eimeria papillata-induced infection in mice," *BioMed Research International*, vol. 2015, Article ID 219670, 7 pages, 2015.
- [19] K. H. Al-Gubory, F. Blachier, P. Faure, and C. Garrel, "Pomegranate peel extract decreases small intestine lipid peroxidation by enhancing activities of major antioxidant enzymes," *Journal of the Science of Food and Agriculture*, vol. 96, pp. 3462–3468, 2016.
- [20] S. N. Al-Shaib, M. I. Waly, L. Al-Subhi, M. H. Tageldin, N. M. Al-Balushi, and M. S. Rahman, "Ameliorative effects of pomegranate peel extract against dietary-induced nonalcoholic fatty liver in rats," *Preventive Nutrition and Food Science*, vol. 21, no. 1, pp. 14–23, 2016.
- [21] R. E. Hartman, A. Shah, A. M. Fagan et al., "Pomegranate juice decreases amyloid load and improves behavior in a mouse model of Alzheimer's disease," *Neurobiology of Disease*, vol. 24, no. 3, pp. 506–515, 2006.
- [22] V. Tapias, J. R. Cannon, and J. T. Greenamyre, "Pomegranate juice exacerbates oxidative stress and nigrostriatal degeneration in Parkinson's disease," *Neurobiology of Aging*, vol. 35, no. 5, pp. 1162–1176, 2014.
- [23] K. Kadooka, K. Fujii, T. Matsumoto et al., "Mechanisms and consequences of carnosine-induced activation of intestinal epithelial cells," *Journal of Functional Foods*, vol. 13, pp. 32–37, 2015.
- [24] T. Fujiki, T. Miura, M. Maura et al., "TAK1 represses transcription of the human telomerase reverse transcriptase gene," *Oncogene*, vol. 26, no. 36, pp. 5258–5266, 2007.
- [25] S. Yamashita, K. Ogawa, T. Ikei, T. Fujiki, and Y. Katakura, "FOXO3a potentiates hTERT gene expression by activating c-MYC and extends the replicative life-span of human fibroblast," *PLoS ONE*, vol. 9, no. 7, Article ID e101864, 2014.
- [26] G. Harada, P. Pattarawat, K. Ito, T. Matsumoto, T. Hasegawa, and Y. Katakura, "Lactobacillus brevis T2102 suppresses the growth of colorectal cancer cells by activating SIRT1," *Journal of Functional Foods*, vol. 23, pp. 444–452, 2016.
- [27] Y. Noda, T. Kaneyuki, A. Mori, and L. Packer, "Antioxidant activities of pomegranate fruit extract and its anthocyanidins: delphinidin, cyanidin, and pelargonidin," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 1, pp. 166–171, 2002.
- [28] M. Reinisalo, A. Kärklund, A. Koskela, K. Kaarniranta, and R. O. Karjalainen, "Polyphenol stilbenes: molecular mechanisms of defence against oxidative stress and aging-related diseases," *Oxidative Medicine and Cellular Longevity*, vol. 2015, Article ID 340520, 24 pages, 2015.
- [29] J. M. Landete, "Ellagitannins, ellagic acid and their derived metabolites: a review about source, metabolism, functions and health," *Food Research International*, vol. 44, no. 5, pp. 1150–1160, 2011.
- [30] A. Varela-López, P. Bullón, F. Giampieri, and J. L. Quiles, "Non-nutrient, naturally occurring phenolic compounds with antioxidant activity for the prevention and treatment of periodontal diseases," *Antioxidants*, vol. 4, no. 3, pp. 447–481, 2015.
- [31] B. Kim, Y. Park, C. J. Wegner, B. W. Bolling, and J. Lee, "Polyphenol-rich black chokeberry (*Aronia melanocarpa*) extract regulates the expression of genes critical for intestinal cholesterol flux in Caco-2 cells," *Journal of Nutritional Biochemistry*, vol. 24, no. 9, pp. 1564–1570, 2013.
- [32] H. Yang, X. Z. Zuo, C. Tian et al., "Green tea polyphenols attenuate high-fat diet-induced renal oxidative stress through SIRT3-dependent deacetylation," *Biomedical and Environmental Sciences*, vol. 28, no. 6, pp. 455–459, 2015.
- [33] X. Zhou, M. Chen, X. Zeng et al., "Resveratrol regulates mitochondrial reactive oxygen species homeostasis through

- Sirt3 signaling pathway in human vascular endothelial cells,” *Cell Death and Disease*, vol. 5, no. 12, Article ID e1576, 2014.
- [34] S. Magnifico, L. Saias, B. Deleglise et al., “NAD⁺ acts on mitochondrial Sirt3 to prevent axonal caspase activation and axonal degeneration,” *The FASEB Journal*, vol. 27, no. 12, pp. 4712–4722, 2013.
- [35] T. Chen, J. Li, J. Liu et al., “Activation of SIRT3 by resveratrol ameliorates cardiac fibrosis and improves cardiac function via the TGF- β /Smad3 pathway,” *American Journal of Physiology. Heart and Circulatory Physiology*, vol. 308, no. 5, pp. H424–H434, 2015.
- [36] Z. Guo, Z. Liao, L. Huang, D. Liu, D. Yin, and M. He, “Kaempferol protects cardiomyocytes against anoxia/reoxygenation injury via mitochondrial pathway mediated by SIRT1,” *European Journal of Pharmacology*, vol. 761, pp. 245–253, 2015.
- [37] K. Huang, C. Chen, J. Hao et al., “Polydatin promotes Nrf2-ARE anti-oxidative pathway through activating Sirt1 to resist AGEs-induced upregulation of fibronectin and transforming growth factor- β 1 in rat glomerular mesangial cells,” *Molecular and Cellular Endocrinology*, vol. 399, pp. 178–189, 2015.
- [38] Z. Zeng, Z. Chen, S. Xu, R. Song, H. Yang, and K.-S. Zhao, “Polydatin alleviates small intestine injury during hemorrhagic shock as a SIRT1 activator,” *Oxidative Medicine and Cellular Longevity*, vol. 2015, Article ID 965961, 12 pages, 2015.
- [39] G. N. Landis and J. Tower, “Superoxide dismutase evolution and life span regulation,” *Mechanisms of Ageing and Development*, vol. 126, no. 3, pp. 365–379, 2005.
- [40] P. Parihar, I. Solanki, M. L. Mansuri, and M. S. Parihar, “Mitochondrial sirtuins: emerging roles in metabolic regulations, energy homeostasis and diseases,” *Experimental Gerontology*, vol. 61, pp. 130–141, 2015.
- [41] Z. Zeng, Y. Yang, X. Dai et al., “Polydatin ameliorates injury to the small intestine induced by hemorrhagic shock via SIRT3 activation-mediated mitochondrial protection,” *Expert Opinion on Therapeutic Targets*, vol. 20, no. 6, pp. 645–652, 2016.