

Research Article

Overexpression of Brg1 Alleviates Hepatic Ischemia/Reperfusion-Induced Acute Lung Injury through Antioxidative Stress Effects

Mian Ge, Chaojin Chen, Weifeng Yao, Shaoli Zhou, Fei Huang, Jun Cai, and Ziqing Hei

Department of Anesthesiology, The Third Affiliated Hospital of Sun Yat-sen University, Guangzhou, Guangdong 510630, China

Correspondence should be addressed to Jun Cai; gzcailun@hotmail.com and Ziqing Hei; heiziqing@sina.com

Received 1 March 2017; Revised 19 May 2017; Accepted 13 June 2017; Published 16 July 2017

Academic Editor: Serafina Perrone

Copyright © 2017 Mian Ge et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Aim. To investigate whether overexpression of Brahma-related gene-1 (Brg1) can alleviate lung injury induced by hepatic ischemia/reperfusion (HIR) and its precise mechanism. **Methods.** Cytomegalovirus-transgenic Brg1-overexpressing (CMV-Brg1) mice and wild-type (WT) C57BL/6 mice underwent HIR. Lung histology, oxidative injury markers, and antioxidant enzyme concentrations in the lung were assessed. The protein expression levels of Brg1, nuclear factor erythroid 2-related factor 2 (Nrf2), heme oxygenase-1 (HO-1), and NAD(P)H:quinone oxidoreductase 1 (NQO1) in the lung were analyzed by Western blotting. **Results.** In the WT group, histopathological analysis revealed that lung damage peaked at 6 h after HIR. Meanwhile, the lung reactive oxygen species (ROS) and 8-isoprostane levels were significantly increased. The protein expression of Brg1 in lung tissue decreased to a minimum at 6 h. Overexpression of Brg1 alleviated lung injury and decreased the amounts of oxidative products, including the levels of 8-isoprostane and ROS, as well as the percentage of positive cells for 4-hydroxynonenal (4-HNE) and 8-oxo-2'-deoxyguanosine (8-OHdG). Brg1 overexpression increased the expression and nuclear translocation of Nrf2 as well as activated the antioxidant enzymes. In addition, it decreased the expression of inflammatory factors. **Conclusion.** Overexpression of Brg1 alleviates oxidative lung injury induced by HIR, likely through the Nrf2 pathway.

1. Introduction

Hepatic ischemia/reperfusion injury (HIRI) occurs in many surgeries, including liver transplantation, liver resection, and cardiovascular surgery [1, 2]. HIRI not only leads to liver injury but also causes remote organ damage, especially lung injury [3, 4]. We have reported previously that up to 58.2% of patients developed pulmonary complications after liver transplantation; among them, 27.5% suffered from an acute lung injury (ALI) [5, 6]. Given that numerous risk factors are involved in ALI induced by hepatic ischemia/reperfusion (HIR), effective and preventive strategies are needed [7, 8].

Multiple factors are reported to take part in the pathogenesis of post-HIR ALI [9]. Excessive reactive oxygen species (ROS) produced in the liver after ischemia/reperfusion is one of the main mechanisms involved in HIR-induced ALI [10–12]. Excessive production of ROS causes ALI by

oxidative stress [13, 14], apoptosis [15–17], and inflammatory responses [18]. Thus, clearance of ROS appears to be a potential therapeutic strategy for post-HIR ALI.

Physiologically, every cell is able to defend itself against the oxidative damage caused by excessive ROS triggering the delivery of antioxidant enzymes [19]. Cellular antioxidant enzymes, including catalase, glutathione S-transferase (GST), thioredoxin 2, superoxide dismutase (SOD), glutathione peroxidase (GPx), heme oxygenase-1 (HO-1), and NAD(P)H:quinone oxidoreductase 1 (NQO1), constitute the first line of defense against the toxic effects of ROS [20, 21]. Nuclear factor erythroid 2-related factor 2 (Nrf2) is an important transcription factor that is known to act as a master regulator of the antioxidant response [21]. Following exposure of cells to oxidative stress, Nrf2 is induced and translocated into the nucleus to activate the expression of a series of antioxidative and cytoprotective proteins that

include GPx, glutamate-cysteine ligase (GCL), NQO1, and HO-1 [22, 23]. However, we have found previously that an increased expression of Nrf2 in liver tissue after liver transplantation did not effectively activate the expression and activity of antioxidant enzymes at an early reperfusion time. This finding indicated that the Nrf2-induced antioxidant response was restricted by post liver transplantation mechanisms. However, those mechanisms inhibiting Nrf2 induction after reperfusion of the liver and their importance in combating ALI remain unclear.

Being a catalytic subunit of SWI2/SNF2-like chromatin-remodeling complexes, Brahma-related gene-1 (Brg1) disrupts the chromatin architecture of target promoters and has been reported not only to play an important role in gene transcription and expression but also to regulate stem cell proliferation and differentiation in different tissues and cells in humans and animals, as well as the process of disease development [24]. Zhang et al. have reported that Brg1 can interact with Nrf2 and is specifically required for Nrf2-mediated activation of HO-1 gene transcription in multiple tumor cells [25]. However, whether Brg1 interacts with Nrf2 and mediates the activation of other antioxidant enzymes in the development of HIR-induced ALI is still unclear. Based on the above-mentioned findings, we hypothesized that upregulation of Brg1 expression could promote the activation of Nrf2-mediated antioxidant enzymes and alleviate lung injury induced by HIR.

2. Methods

2.1. Animals and Treatment. The experimental protocols and design were approved by the Institutional Animal Care and Use Committee at Sun Yat-sen University. Eight- to ten-week-old C57BL/6 male mice were used. Animal care was conducted in accordance with the Guidelines of Sun Yat-sen University for Animal Experimentation. Cytomegalovirus-transgenic Brg1-overexpressing (CMV-Brg1) mice (Cyagen Biosciences, ID: TGBS130618AG1, USA) with the C57BL/6 background were used to generate CMV-Brg1 and wild-type (WT) littermates. As described previously [26], genotyping was performed by polymerase chain reaction using genomic DNA extracted from a tail snip. The mice were randomly assigned into five groups ($n = 8$ per group) in the initial intervention model establishment study, which included sham-operated as well as 3, 6, 12, and 24 h reperfusion groups. Subsequent in vivo studies were performed using the 6 h reperfusion HIR model. Both the CMV-Brg1 and WT mice were then randomly assigned into two groups ($n = 5$ per group): sham-operated group (sham) and HIR group.

2.2. HIR Model. In order to minimize investigator bias, the operator was blinded to the treatment conditions. All the mice were anesthetized by an intraperitoneal injection of ketamine (60 mg/kg) and then received left and median liver lobe clamping for 60 min and reperfusion for the designated time. The sham group was treated in an identical manner to the surgery group except it was not subjected to renal pedicle occlusion. After the operation, the mice were housed in

microisolator cages and allowed free access to water and chow. After a designated reperfusion time, mouse blood and lungs were collected for further experiments.

2.3. Hematoxylin-Eosin Staining and Histopathological Analysis. At the end of the experiment, the mice were anesthetized by an intraperitoneal injection of 60 mg/kg ketamine. Following exposure, the left lung was removed and perfused with PBS and then fixed in 10% buffered formalin overnight at 4°C. The lung tissues were then embedded with paraffin, sectioned at 5 μ m thickness, and stained with hematoxylin and eosin. Histological severity of lung injury was graded using a 0 to 4 point scale according to combined assessments of edema, neutrophil infiltration, hemorrhage, bronchiole epithelial desquamation, and hyaline membrane formation as follows: 0, no injury; 1, modest injury (including limited interstitial edema and congestion, but no interstitial neutrophilic infiltration with inflammatory cells in the alveolar spaces); 2, intermediate injury (including mild interstitial edema, congestion, and interstitial neutrophilic infiltration with inflammatory cells in the alveolar spaces); 3, widespread injury (including more prominent congestion and interstitial edema with neutrophils partially filling the alveolar spaces); and 4, widespread injury (including most prominent, marked congestion and interstitial edema with neutrophilic infiltration nearly filling the alveolar spaces or with pulmonary consolidation). A total of six fields were randomly selected for each slide, and the average was used as the histopathological score [27].

2.4. ROS Detection. As reported previously [28, 29], the accumulation of ROS in lung tissue was estimated by using the in vitro reactive nitrogen species assay kit OxiSelect (Cell Biolabs, San Diego, CA, USA), according to the manufacturer's protocol.

2.5. Detection of 8-Isoprostane, Glutamate-Cysteine Ligase Catalytic Subunit (GCLC), Superoxide Dismutase (SOD), Glutathione S-Transferase Alpha 1 (GST α 1), Interleukin-6 (IL-6), and Tumor Necrosis Factor- α (TNF- α). Lung tissue was transformed into 10% homogenates with frozen normal saline and spun at 3000 rpm for 10 min. The pulmonary protein content was measured using a Bicinchoninic Acid Protein Assay Kit (KeyGen Biotech Company, Nanjing, China). The concentrations of 8-isoprostane (Cayman Chemical Company, USA), GCLC (USCN Business Co. Ltd., Wuhan, China), GST α 1 (USCN Business Co. Ltd., Wuhan, China.), SOD (KeyGen Biotech Company, Nanjing, China.), IL-6 (USCN Business Co. Ltd., Wuhan, China), and TNF- α (USCN Business Co. Ltd., Wuhan, China) were measured by enzyme-linked immunosorbent assay kits. All detections were performed according to the assay kit manufacturer's instructions.

2.6. Immunofluorescence and Immunohistochemistry. 5 μ m thick HIR-injured lung sections embedded in paraffin and cells grown on sterile coverslips and then subjected to the HIR protocol were prepared for immunofluorescence assays. The samples were stained with anti-8-OHdG antibody (1:200; LS-C415095, LSBio) at 4°C overnight, followed by

anti-rabbit or anti-mouse IgG (1:1000), according to the manufacturer's instructions. Cell nuclei were stained with DAPI. All images were captured using an EVOS FL fluorescence microscope (EVOS FL, Life Technology), and 10 randomly selected fields of each sample were semiquantified.

Immunohistochemical staining was performed to detect 4-hydroxynonenal (4-HNE) and Nrf2 expression in lung sections embedded in paraffin with anti-4-HNE antibody (1:100; ab46545, Abcam) and anti-Nrf2 antibody (1:100, AP52269, Abgent), respectively.

2.7. Western Blot Analysis. Western blotting was performed following standard procedures. Anti-Nrf2 antibody (1:100, AP52269, Abgent), anti-Brg1 antibody (1:100; ab110641, Abcam), anti-NQO1 antibody (1:100; ab28947, Abcam), anti-HO-1 antibody (1:100; sc-10,789, Santa Cruz), and secondary antibody (1:2000; Millipore) were used to detect protein expression. Anti- β -actin and anti-lamin B2 (Proteintech) were used at 1:5000. Images were acquired by a Tanon 5500 imaging system (Tanon, Shanghai). The images were scanned with the ImageJ scanning software, and the data were expressed as relative values to sham or control values.

2.8. Statistical Analysis. Statistical analysis was performed using SPSS 13.0 (SPSS Inc., Chicago, IL, USA) and SigmaPlot 10.0 (Systat Software Inc., Chicago, IL, USA). Data normality was tested using the Kolmogorov-Smirnov test. Multiple comparisons among different groups were analyzed using one-way analysis of variance, followed by Tukey's post hoc test. Quantitative data are presented as mean \pm standard error of the mean. *P* values less than 0.05 were considered statistically significant.

3. Results

3.1. HIR Induces Oxidative Stress and ALI in the WT Mice. After the process of HIR, pathological damage was detected in the lungs (Figure 1(a)), including increased alveolar damage, perivascular and peribronchial edema, alveolar wall capillary hyperemia, and pulmonary interstitial inflammatory cell infiltration. It reached a peak at 6 h after reperfusion (Figure 1(a)), as reflected by the significant increase in pathological scores (Figure 1(b)), and then was restored gradually to normal.

As shown in Figures 1(c) and 1(d), the changes of ROS and 8-isoprostane levels in lung tissues were consistent with lung pathological damage. Similarly, the Brg1 protein expression in lung tissue began to decline after reperfusion, significantly reduced at 3 h and 6 h ($P < 0.01$ versus the Sham group), reached a minimum at 6 h, and then gradually increased at 12 h and 24 h (Figure 1(e)). Nrf2 protein expression in the lung tissue did not significantly change within the first 6 h after reperfusion, but it increased significantly at 12 h and 24 h after reperfusion (Figure 1(f)).

3.2. Brg1 Overexpression Protects against ALI and Oxidative Stress Induced by HIR. HIR resulted in severe lung pathological injury that was associated with significant oxidative damage and a significant decrease in Brg1 expression, suggesting that there might be an important role for Brg1

in ALI induced by HIR. To further explore this role, the CMV-Brg1 and WT mice were used to develop an appropriate HIR model.

As shown in Figure 2(a), the Brg1 expression level of the CMV-Brg1 mice was increased by 2.2-fold, compared with the WT mice. The pathological results showed that increased alveolar damage, perivascular and peribronchial edema, alveolar wall capillary hyperemia, and pulmonary interstitial inflammatory cell infiltration were present after HIR in the WT mice (Figure 3(a)). In contrast, after HIR, the CMV-Brg1 mice suffered only mild alveolar damage, mild hyperemia in the alveolar wall capillaries, and less infiltration of inflammatory cells than the WT mice (Figure 3(a)). Compared with the WT mice, the pathological damage, as further assessed by the pathological scores, was therefore significantly reduced in the CMV-Brg1 mice (Figures 3(a) and 3(b)).

As shown in Figures 4(a) and 4(b), compared with the sham group, the levels of 8-isoprostane and ROS in the lung tissues of the WT mice were significantly increased after HIR; similarly, the relative fluorescence intensity of 8-OHdG (a marker of DNA oxidative damage) and the number of cells positive for 4-HNE (a marker of lipid peroxidation) were significantly increased in the WT HIR group (Figures 4(c), 4(d), 4(e), and 4(f)). However, compared with the WT mice undergoing HIR, the expression of 8-isoprostaglandin, ROS, 4-HNE, and 8-OHdG in the CMV-Brg1 mice was significantly decreased after HIR, showing that Brg1 overexpression can effectively suppress the oxidative damage induced by HIR.

3.3. Association of Brg1 Overexpression with Increased Nuclear Transfer of Nrf2. Next, we explored the role of Brg1 on Nrf2 in ALI induced by HIR. As shown in Figures 2(a) and 2(c), after the process of HIR, Nrf2 protein expression in the WT mice was not significantly increased, compared with the sham group. In contrast, Nrf2 protein expression in the CMV-Brg1 mice after the process of HIR was significantly increased.

The CMV-Brg1 mice expressed significantly more Nrf2 protein after HIR than the WT mice (Figures 2(a) and 2(c)). As shown by the immunohistochemistry results presented in Figures 2(d) and 2(e), HIR significantly increased the immunohistochemical staining intensity of Nrf2-positive cells in lung tissue of the CMV-Brg1 mice at 6 h after HIR.

3.4. Association of Brg1 Overexpression with Activated Antioxidant and Decreased Inflammatory Factors. As Nrf2 is known as a regulator of antioxidant enzymes, we compared the levels of enzymes between the WT and CMV-Brg1 mice. Compared with the sham group, the protein expressions of NQO1 and HO-1 in the WT mice after HIR were not significantly increased. In the CMV-Brg1 mice after HIR, both NQO1 and HO-1 protein expressions were significantly increased (Figures 5(a), 5(b), and 5(c)). Moreover, compared with the HIR WT mice, the HO-1 and NQO1 protein expressions in the CMV-Brg1 mice were significantly increased after HIR (Figures 5(a) and 5(b)). These data indicate that




FIGURE 1: Hepatic ischemia/reperfusion (HIR) induces oxidative stress and acute lung injury at different time points (3, 6, 12, and 24 h) after HIR in the wild-type mice. (a) Hematoxylin-eosin staining of lung sections ($\times 200$); (b) lung pathological scores; (c) and (d) levels of 8-isoprostane and reactive oxygen species (ROS); (e) Brg1 protein expression in lung tissue; and (f) Nrf2 protein expression in lung tissue; $*P < 0.05$; $**P < 0.01$.

the protein expressions of HO-1 and Nrf2 after HIR might be regulated by Brg1, but with little relation to NQO1.

At the same time, we tested the changes in antioxidant enzyme levels, including GST α 1, SOD, and GCLC. Compared with the sham groups, SOD, GCLC, and GST α 1 in the WT mice after HIR were reduced significantly. However, the decreased magnitude of SOD, GCLC, and GST α 1 in

the CMV-Brg1 mice was less than that in the WT mice after HIR, indicating that Brg1 overexpression might play a crucial role in prompting the activation and expression of antioxidant enzymes (Figures 5(d), 5(e), and 5(f)). Moreover, we observed an inflammatory response. As shown in Figures 5(g) and 5(h), compared with the sham groups, the levels of IL-6 and TNF- α were significantly increased after




FIGURE 2: Association of Brg1 overexpression with increased nuclear transfer of Nrf2. (a–c) The protein expression levels of Brg1 and Nrf2 were detected by Western blot in the CMV-Brg1 and WT mice after HIR. (d, e) Immunohistochemical staining ($\times 200$) of lung Nrf2 and densitometric analysis showed that the Nrf2-positive cells in lung tissue of the CMV-Brg1 mice were more pronounced. * $P < 0.05$; ** $P < 0.01$.




FIGURE 3: Overexpression of Brg1 protects against acute lung injury and oxidative stress induced by HIR. Cyomegalovirus-transgenic Brahma-related gene-1-overexpressing (CMV-Brg1) mice and wild-type (WT) mice were subjected to sham operation or HIR. (a) Hematoxylin-eosin staining of lung sections ($\times 200$); (b) pathological scores of lung tissue. * $P < 0.05$; ** $P < 0.01$.

HIR in the WT mice, but not in the CMV-Brg1 mice. After HIR, the TNF- α level was significantly decreased in the CMV-Brg1 mice ($P < 0.05$ versus the WT mice).

4. Discussion

In the current study, we demonstrated that HIR can induce remote ALI accompanied by a significant increase of oxidative damage and a significant decrease of Brg1 expression. Furthermore, we revealed that overexpression of Brg1 attenuated lung injury after HIR, promoted Nrf2 nuclear

translocation, and attenuated oxidative stress by activating the antioxidase system including HO-1, NQO1, SOD, GCLC, and GST α 1.

It is well known that excessive ROS production is one of the main mechanisms involved in ALI after HIR [12, 18] and that antioxidant enzyme activation alleviates ALI induced by HIR [10, 30]. Acting like a transcriptional factor and cell defense response regulator, Nrf2 translocates into the nucleus following exposure to oxidative stress and activates the gene expression of many antioxidase and cytoprotective proteins, including GCL, HO-1, GPx, and NQO1 [22]. We




FIGURE 4: Overexpression of Brg1 protects against oxidative stress induced by HIR. (a, b) Lung 8-isoprostane and ROS levels. (c, d) Immunohistochemical staining ($\times 200$) of lung 4-hydroxynonenal (4-HNE) (brown) and densitometric analysis showed that the 4-HNE-positive cells were significantly reduced in the CMV-Brg1 mice than in the WT mice at 6 h after HIR. (e, f) Immunofluorescence staining ($\times 200$) of lung 8-hydroxy-2-deoxyguanosine (8-OHdG) and densitometric analysis showed that the relative fluorescence intensity was significantly decreased in the CMV-Brg1 mice compared to the WT mice at 6 h after HIR. The 8-OHdG-positive cells were stained red, and the nuclei of tissue sections were counterstained blue by 4,6-diamidino-2-phenylindole. * $P < 0.05$; ** $P < 0.01$.




FIGURE 5: Association of Brg1 overexpression with activated antioxidant and decreased inflammatory factors. (a–c) the protein expression of HO-1 and NQO1 in the CMV-Brg1 and WT mice after HIR. (d–h) the concentrations of glutamate-cysteine ligase catalytic subunit (GCLC), glutathione S-transferase alpha 1 (GST α 1), superoxide dismutase (SOD), interleukin-6 (IL-6), and tumor necrosis factor- α (TNF- α) in the CMV-Brg1 and WT mice after HIR. * $P < 0.05$; ** $P < 0.01$.

have reported previously that HO-1 induction protects the lungs against ALI induced by liver transplantation in rats [31]. Recently, several studies have documented that upregulation of Nrf2/HO-1 plays a critical protective role in ALI induced by lipopolysaccharides or intestinal ischemia/reperfusion [23, 32, 33]. However, no studies have yet reported that Nrf2 and its downstream antioxidant enzymes can confer a protective effect against lung injury induced by HIR. Similar to earlier reports, we demonstrated that HIR induces significant oxidative stress by producing large amounts of ROS, 8-isoprostane, 4-HNE, and 8-OHdG. However, Nrf2 expression did not significantly

increase until the ROS levels and lung pathological damage reached a peak at 6 h after reperfusion. Interestingly, once significantly activated, Nrf2 quickly promoted the expression of antioxidant enzymes and protected the lung from oxidative injury. Considering both the immediate oxidative stress and the inhibited induction of Nrf2, we assumed that upregulation of Nrf2 expression could play a critical role in alleviating lung injury after HIR.

Brg1, a subunit of the chromatin remodeling complex, has been demonstrated not only to regulate gene transcription and expression but also to play an important role in stem cell proliferation and differentiation in different tissues and

cells in humans and animals [24, 34]. In addition, it has been shown to contribute to the process of disease development, such as lung tumorigenesis and cytokine response [24].

Reduction and loss of Brg1 expression have been related to dedifferentiation in lung cancers [35, 36]. Brg1 also has been found to act in concert with Fanconi anemia proteins to protect the promoters of antioxidant defense genes from oxidative damage [37]. In our experiments, we found that Brg1 was reduced after reperfusion, reaching a minimum at 6 h, while oxidative damage peaked. Interestingly, using transgenic CMV-Brg1 mice, we found that Brg1 overexpression promoted Nrf2 nuclear translocation, activated the downstream antioxidant system, and then alleviated post-HIR lung oxidative damage. Originally, Brg-1 was described as the factor responsible for selective HO-1 enzyme induction triggered by Nrf2 [25]. However, our findings indicated a more universal effect for Brg1: Brg1-promoted Nrf2 nuclear translocation mediates HO-1 induction; but the precise underlying mechanism remains to be determined. Therefore, the current study provides new potential therapeutic insights for scavenging ROS to combat HIR-mediated ALI.

It is worth noting that Brg1 interacted with Nrf2 to mediate HO-1 and NQO1 induction in order to counteract oxidative stress and activated antioxidant enzymes including GST, GCLC, and SOD. This finding was inconsistent with earlier reports showing that BRG1-mediated chromatin remodeling is essential for RNA polymerase II recruitment to the HO-1 promoter but not to the NQO1 promoter when subjected to oxidative stress in streptozotocin-induced diabetic rats [25, 38]. However, whether lung injury induced by HIR is aggravated in Brg1 gene deficiency still needs to be explored, especially by using Brg1 gene-deficient mice. These results will help to elucidate the role of Brg1 in lung protection. Moreover, whether other pathways can mediate Brg1 involvement in lung injury after HIR needs to be further investigated.

5. Conclusion

In conclusion, the present results demonstrated that HIR could induce remote lung injury accompanied by a significant increase of oxidative damage and a significant decrease of Brg1 expression at the early reperfusion time. Brg1 overexpression significantly protected the lung from HIR injury through Nrf2 activation and its downstream antioxidant activity, including HO-1, NQO1, SOD, GCLC, and GST α 1. Though the exact mechanisms of the interaction of Brg1 and Nrf2-induced downstream antioxidant enzymes under HIR settings remain to be studied, upregulation of Brg1 may be a potential therapy to alleviate remote lung injury after HIR.

Conflicts of Interest

There are no conflicts of interest for any of the authors.

Authors' Contributions

Mian Ge and Ziqing Hei conceived and designed the research; Chaojin Chen performed the experiments; Shaoli

Zhou analyzed the data; Weifeng Yao interpreted the experimental results and prepared the figures; Mian Ge and Chaojin Chen drafted the manuscript; Fei Huang and Jun Cai edited and revised the manuscript; Ziqing Hei approved the final version of the manuscript, and Mian Ge, Chaojin Chen, and Weifeng Yao contributed equally to this work.

Acknowledgments

The authors acknowledge Medjaden Bioscience Limited for the scientific editing and proofreading of this manuscript. This study was partly supported by grants from the Natural Science Foundation of China (no. 81372090, no. 81571926, no. 81501695, and no. 81601724) and key project of the Natural Science Foundation of Guangdong Province, China (Grant no. S2011020002780).

References

- [1] J. U. Bascom, P. Gosling, and B. A. Zikria, "Hepatic ischemia-reperfusion injury," *American Journal of Surgery*, vol. 184, no. 1, pp. 84-85, 2002.
- [2] M. Cannistra, M. Ruggiero, A. Zullo et al., "Hepatic ischemia reperfusion injury: a systematic review of literature and the role of current drugs and biomarkers," *International Journal of Surgery (London, England)*, vol. 33, Supplement 1, pp. S57-S70, 2016.
- [3] C. Nastos, K. Kalimeris, N. Papoutsidakis et al., "Global consequences of liver ischemia/reperfusion injury," *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 906965, 13 pages, 2014.
- [4] T. Brenner, J. Motsch, J. Werner, L. Grenacher, E. Martin, and S. Hofer, "Rapid-onset acute respiratory distress syndrome (ARDS) in a patient undergoing metastatic liver resection: a case report and review of the literature," *Anesthesiology Research and Practice*, vol. 2010, Article ID 586425, 9 pages, 2010.
- [5] A. Zhang, X. Chi, G. Luo et al., "Mast cell stabilization alleviates acute lung injury after orthotopic autologous liver transplantation in rats by downregulating inflammation," *PLoS One*, vol. 8, no. 10, article e75262, 2013.
- [6] S. R. Li, N. Shen, and Z. Q. Hei, "Early risk factors of acute lung injury following orthotopic liver transplantation," *Zhonghua Yi Xue Za Zhi*, vol. 88, no. 43, pp. 3049-3052, 2008.
- [7] J. F. Aduen, W. H. Stapelfeldt, M. M. Johnson et al., "Clinical relevance of time of onset, duration, and type of pulmonary edema after liver transplantation," *Liver Transplantation*, vol. 9, no. 7, pp. 764-771, 2003.
- [8] A. Jiang, C. Liu, F. Liu et al., "Liver cold preservation induce lung surfactant changes and acute lung injury in rat liver transplantation," *World Journal of Gastroenterology*, vol. 18, no. 4, pp. 323-330, 2012.
- [9] K. J. Kang, "Mechanism of hepatic ischemia/reperfusion injury and protection against reperfusion injury," *Transplantation Proceedings*, vol. 34, no. 7, pp. 2659-2661, 2002.
- [10] K. C. Chan, C. J. Lin, P. H. Lee et al., "Propofol attenuates the decrease of dynamic compliance and water content in the lung by decreasing oxidative radicals released from the reperfused liver," *Anesthesia and Analgesia*, vol. 107, no. 4, pp. 1284-1289, 2008.

- [11] A. P. Castro, M. A. Castro Junior, S. Lauz, E. Facin, J. Simoes Mde, and D. J. Fagundes, "The role of N-acetyl-cysteine in the lung remote injury after hepatic ischemia and reperfusion in rabbits," *Acta Cirurgica Brasileira*, vol. 27, no. 1, pp. 49–55, 2012.
- [12] H. Jaeschke and A. Ramachandran, "Reactive oxygen species in the normal and acutely injured liver," *Journal of Hepatology*, vol. 55, no. 1, pp. 227–228, 2011.
- [13] V. G. Nielsen, S. Tan, A. Weinbroum et al., "Lung injury after hepatoenteric ischemia-reperfusion: role of xanthine oxidase," *American Journal of Respiratory and Critical Care Medicine*, vol. 154, no. 5, pp. 1364–1369, 1996.
- [14] L. M. Colletti, S. L. Kunkel, A. Walz et al., "Chemokine expression during hepatic ischemia/reperfusion-induced lung injury in the rat. The role of epithelial neutrophil activating protein," *Journal of Clinical Investigation*, vol. 95, no. 1, pp. 134–141, 1995.
- [15] H. W. Chen, C. T. Chien, S. L. Yu, Y. T. Lee, and W. J. Chen, "Cyclosporine a regulate oxidative stress-induced apoptosis in cardiomyocytes: mechanisms via ROS generation, iNOS and Hsp70," *British Journal of Pharmacology*, vol. 137, no. 6, pp. 771–781, 2002.
- [16] X. Wang, N. Takahashi, H. Uramoto, and Y. Okada, "Chloride channel inhibition prevents ROS-dependent apoptosis induced by ischemia-reperfusion in mouse cardiomyocytes," *Cellular Physiology and Biochemistry*, vol. 16, no. 4–6, pp. 147–154, 2005.
- [17] C. Wang, K. Chen, Y. Xia et al., "N-acetylcysteine attenuates ischemia-reperfusion-induced apoptosis and autophagy in mouse liver via regulation of the ROS/JNK/Bcl-2 pathway," *PLoS One*, vol. 9, no. 9, article e108855, 2014.
- [18] L. Zhou, D. Zhao, H. An, H. Zhang, C. Jiang, and B. Yang, "Melatonin prevents lung injury induced by hepatic ischemia-reperfusion through anti-inflammatory and anti-apoptosis effects," *International Immunopharmacology*, vol. 29, no. 2, pp. 462–467, 2015.
- [19] I. Ben Mosbah, Y. Mouchel, J. Pajaud et al., "Pretreatment with mangafodipir improves liver graft tolerance to ischemia/reperfusion injury in rat," *PLoS One*, vol. 7, no. 11, article e50235, 2012.
- [20] K. H. Al-Gubory, C. Garrel, P. Faure, and N. Sugino, "Roles of antioxidant enzymes in corpus luteum rescue from reactive oxygen species-induced oxidative stress," *Reproductive Biomedicine Online*, vol. 25, no. 6, pp. 551–560, 2012.
- [21] M. Kobayashi and M. Yamamoto, "Molecular mechanisms activating the Nrf2-Keap1 pathway of antioxidant gene regulation," *Antioxidants and Redox Signaling*, vol. 7, no. 3–4, pp. 385–394, 2005.
- [22] S. Magesh, Y. Chen, and L. Hu, "Small molecule modulators of Keap1-Nrf2-ARE pathway as potential preventive and therapeutic agents," *Medicinal Research Reviews*, vol. 32, no. 4, pp. 687–726, 2012.
- [23] Q. T. Meng, C. Cao, Y. Wu et al., "Ischemic post-conditioning attenuates acute lung injury induced by intestinal ischemia-reperfusion in mice: role of Nrf2," *Laboratory Investigation*, vol. 96, no. 10, pp. 1087–1104, 2016.
- [24] W. Tian, H. Xu, F. Fang, Q. Chen, Y. Xu, and A. Shen, "Brahma-related gene 1 bridges epigenetic regulation of proinflammatory cytokine production to steatohepatitis in mice," *Hepatology*, vol. 58, no. 2, pp. 576–588, 2013.
- [25] J. Zhang, T. Ohta, A. Maruyama et al., "BRG1 interacts with Nrf2 to selectively mediate HO-1 induction in response to oxidative stress," *Molecular and Cellular Biology*, vol. 26, no. 21, pp. 7942–7952, 2006.
- [26] M. Malumbres, R. Mangués, N. Ferrer, S. Lu, and A. Pellicer, "Isolation of high molecular weight DNA for reliable genotyping of transgenic mice," *BioTechniques*, vol. 22, no. 6, pp. 1114–1119, 1997.
- [27] T. Zhu, D. X. Wang, W. Zhang et al., "Andrographolide protects against LPS-induced acute lung injury by inactivation of NF-kappaB," *PLoS One*, vol. 8, no. 2, article e56407, 2013.
- [28] S. Ravassa, J. Beaumont, A. Huerta et al., "Association of low GLP-1 with oxidative stress is related to cardiac disease and outcome in patients with type 2 diabetes mellitus: a pilot study," *Free Radical Biology and Medicine*, vol. 81, pp. 1–12, 2015.
- [29] A. Iglesias-Ara, O. Zenarruzabeitia, L. Buelta, J. Merino, and A. M. Zubiaga, "E2F1 and E2F2 prevent replicative stress and subsequent p53-dependent organ involution," *Cell Death and Differentiation*, vol. 22, no. 10, pp. 1577–1589, 2015.
- [30] V. Rocha-Santos, E. R. Figueira, J. A. Rocha-Filho et al., "Pentoxifylline enhances the protective effects of hypertonic saline solution on liver ischemia reperfusion injury through inhibition of oxidative stress," *Hepatobiliary and Pancreatic Diseases International*, vol. 14, no. 2, pp. 194–200, 2015.
- [31] X. Chi, N. Guo, W. Yao et al., "Induction of heme oxygenase-1 by hemin protects lung against orthotopic autologous liver transplantation-induced acute lung injury in rats," *Journal of Translational Medicine*, vol. 14, p. 35, 2016.
- [32] G. F. Zhu, H. J. Guo, Y. Huang, C. T. Wu, and X. F. Zhang, "Eriodictyol, a plant flavonoid, attenuates LPS-induced acute lung injury through its antioxidative and anti-inflammatory activity," *Experimental and Therapeutic Medicine*, vol. 10, no. 6, pp. 2259–2266, 2015.
- [33] Y. Jiang, Z. Zhou, Q. T. Meng et al., "Ginsenoside Rb1 treatment attenuates pulmonary inflammatory cytokine release and tissue injury following intestinal ischemia reperfusion injury in mice," *Oxidative Medicine and Cellular Longevity*, vol. 2015, Article ID 843721, 12 pages, 2015.
- [34] X. Zhang, B. Li, W. Li et al., "Transcriptional repression by the BRG1-SWI/SNF complex affects the pluripotency of human embryonic stem cells," *Stem Cell Reports*, vol. 3, no. 3, pp. 460–474, 2014.
- [35] T. Yoshimoto, D. Matsubara, T. Nakano et al., "Frequent loss of the expression of multiple subunits of the SWI/SNF complex in large cell carcinoma and pleomorphic carcinoma of the lung," *Pathology International*, vol. 65, no. 11, pp. 595–602, 2015.
- [36] D. Matsubara, Y. Kishaba, S. Ishikawa et al., "Lung cancer with loss of BRG1/BRM, shows epithelial mesenchymal transition phenotype and distinct histologic and genetic features," *Cancer Science*, vol. 104, no. 2, pp. 266–273, 2013.
- [37] W. Du, R. Rani, J. Sipple et al., "The FA pathway counteracts oxidative stress through selective protection of antioxidant defense gene promoters," *Blood*, vol. 119, no. 18, pp. 4142–4151, 2012.
- [38] H. Li, W. Yao, M. G. Irwin et al., "Adiponectin ameliorates hyperglycemia-induced cardiac hypertrophy and dysfunction by concomitantly activating Nrf2 and Brg1," *Free Radical Biology and Medicine*, vol. 84, pp. 311–321, 2015.



Hindawi
Submit your manuscripts at
<https://www.hindawi.com>

