Corrigendum

Corrigendum to “Perillaldehyde Inhibits AHR Signaling and Activates NRF2 Antioxidant Pathway in Human Keratinocytes”

Yoko Fuyuno ¹, Hiroshi Uchi,¹ Mao Yasumatsu,² Saori Morino-Koga,³ Yuka Tanaka ²,
Chikage Mitoma,^{1,2} and Masutaka Furue^{1,2}

¹Department of Dermatology, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashiku, Fukuoka 812-8582, Japan

²Research and Clinical Center for Yusho and Dioxin, Kyushu University Hospital, 3-1-1 Maidashi, Higashiku, Fukuoka 812-8582, Japan

³Division of Statistics, Center for Cohort Studies, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashiku, Fukuoka 812-8582, Japan

Correspondence should be addressed to Yoko Fuyuno; y-kuba@dermatol.med.kyushu-u.ac.jp

Received 25 March 2018; Accepted 8 April 2018; Published 22 May 2018

Copyright © 2018 Yoko Fuyuno et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Perillaldehyde Inhibits AHR Signaling and Activates NRF2 Antioxidant Pathway in Human Keratinocytes” [1], there was an error in Supplementary Figure S3. The figure incorrectly showed that PAH induced ROS expression and BaP suppressed ROS expression induced by PAH, whereas the corrected figure shows that BaP induced ROS expression and PAH suppressed ROS expression induced by BaP. The correct supplementary figure is available here.

Supplementary Materials

Supplementary 3. Figure S3: flow cytometric analysis of ROS confirmed the inhibitory action of PAH against BaP-induced ROS generation. $**p < 0.01$. (*Supplementary Materials*)

References

- [1] Y. Fuyuno, H. Uchi, M. Yasumatsu et al., “Perillaldehyde inhibits AHR signaling and activates NRF2 antioxidant pathway in human keratinocytes,” *Oxidative Medicine and Cellular Longevity*, vol. 2018, Article ID 9524657, 9 pages, 2018.