

Research Article

Flavonoids from *Apios americana* Medikus Leaves Protect RAW264.7 Cells against Inflammation via Inhibition of MAPKs, Akt-mTOR Pathways, and Nfr2 Activation

Qiang Chu,^{1,2} Xin Yu,¹ Ruoyi Jia,¹ Yaxuan Wang,¹ Yiru Zhang,¹ Shuang Zhang,³ Yangyang Liu,¹ Yonglu Li,¹ Wen Chen,¹ Xiang Ye,¹ and Xiaodong Zheng¹ 

¹Department of Food Science and Nutrition, National Engineering Laboratory of Intelligent Food Technology and Equipment, National-Local Joint Engineering Laboratory of Intelligent Food Technology and Equipment, Key Laboratory for Agro-Products Postharvest Handling of Ministry of Agriculture, Zhejiang Key Laboratory for Agro-Food Processing, Fuli Institute of Food Science, Zhejiang University, Hangzhou 310058, China

²State Key Laboratory of Silicon Materials, School of Materials Science and Engineering, Zhejiang University, Hangzhou 310027, China

³College of Environmental and Resource Sciences, Zhejiang University, Hangzhou 310058, China

Correspondence should be addressed to Xiaodong Zheng; xdzhengzju@163.com

Received 26 August 2019; Revised 20 October 2019; Accepted 15 November 2019; Published 13 December 2019

Guest Editor: Adam P. Lightfoot

Copyright © 2019 Qiang Chu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Apios americana Medikus was once widely accepted as staple food in India for a long time, and the tuber of which possesses high nutrients. During the past decades, most of the research has focused on the biological activity in the tubers of *Apios americana* Medikus whereas the leaves were ignored. In this study, the *Apios americana* Medikus leaf extract (ALE) was obtained and seven compounds were identified. LPS-induced RAW264.7 cells were used to study the anti-inflammation activity of ALE. As expected, ALE reduced the secretion of nitric oxide (NO) and inflammatory cytokines via inhibition of NF- κ B and MAPK signaling together with activation of Nrf2-Keap1 and FOXO pathways, as well as alleviating the oxidative stress and mitochondrial dysfunction. In addition, ALE could activate HMGB1-Beclin1 and Sirt1-FoxO1 pathways and inhibit the Akt-mTOR signaling pathway to activate autophagy, protecting RAW264.7 cells from inflammation. In summary, our results suggested that ALE might help activate the anti-inflammation system, resulting in the prevention of LPS-induced damage in RAW264.7 cells.

1. Introduction

Inflammation is a natural defensive response of the organisms to stimulation. Many factors could induce inflammation, which are involved in pathogens, chemical immune responses, damaged cells, irritants, etc. Excessive inflammation response would lead to disorder in organisms together with the overproduction of oxygen free radicals, eventually breaking the dynamic balance [1], which tends to cause diseases like fever, asthma, psoriasis, arthritis, and even cancer [2]. Meanwhile, macrophages play a vital part during the process of inflammation and many cytokines are generated, among which interleukin-1 (IL-1), interleukin-6 (IL-6), and tumor

necrosis factor α (TNF- α) could further promote inflammation. Therefore, the key to treat inflammation is to suppress the aberrant macrophage activation.

Multiple studies have demonstrated that lipopolysaccharide (LPS) induces the generation of inflammatory cytokines via NF- κ B and MAPK pathways [3]. RAW264.7 cells induced by LPS have widely been accepted as the classic inflammation model. Macrophages will produce a huge number of inflammatory cytokines like NO, TNF- α , IL- β , and COX under LPS conditions [4]. However, the incessant inflammation will cause the overproduction of reactive oxygen/nitrogen species including O₂⁻, H₂O₂, OH⁻, ONOO⁻, and HOONO, leading to DNA damage, mitochondrial dysfunction, and cell death.

The mitochondrion is the site where reactive oxygen/nitrogen species generate; meanwhile, mitochondrial dysfunction is closely connected with the overproduction reactive oxygen/nitrogen species [5]. In addition, as a self-protective mechanism of cells, autophagy could lead to mitochondrial transformation and the decomposition of impaired mitochondria. It has been proven that cell autophagy facilitates in scavenging reactive oxygen species, maintaining mitochondrial function, and attenuating inflammation [6, 7]. The balance between inflammation and autophagy is associated with many pathways such as the FoxO1 pathway, Akt-mTOR pathway, and HMGB1 pathway [8, 9].

Existing anti-inflammatory drugs always produce a side effect on the gastrointestinal tract and cardiovascular systems, suggesting that seeking other safe, efficient, and noncytotoxic anti-inflammatory drugs is necessary [10]. In the last few decades, more and more scientists have paid attention to the functional factors derived from plants for their favorable function of regulating the immune system and inflammation. *Apios americana* Medikus is a nitrogen-fixing legume native to North America, and the tubers of which are abundant in fatty acid, lipids, carbohydrate, and amino acids [10, 11]. However, most of studies on *Apios americana* Medikus focused on the biological activities of its tubers, and the leaves were ignored. In this study, flavonoids were extracted from dried *Apios americana* Medikus leaves and the main components were identified by liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis. Subsequently, we aimed to investigate the potential protective effect of ALE against LPS-induced inflammation in RAW264.7 cells and explored the underlying mechanism.

2. Materials and Methods

2.1. Materials and Reagents. RAW264.7 cells were purchased from the Wuhan Cell Bank of Chinese Academy of Sciences. Absolute ethyl alcohol, methanol, ethyl acetate, potassium bromide, formic acid, glucan, dimethyl sulfoxide, hydrochloric acid, and acetonitrile were obtained from Aladdin (Shanghai, China). Trypsin and MTT were purchased from Sigma-Aldrich (St. Louis, MO, USA). Fetal bovine serum and DMEM high glucose medium were obtained from Jinnuo Biotechnology (Hangzhou, China). 2',7'-Dichlorofluorescein diacetate (DCFH-DA), 3-amino,4-aminomethyl-2',7'-difluorescein diacetate (DAF-FMDA), dihydroethidium (DHE), naphthalene-2,3-dicarboxaldehyde (NDA), Nonyl Acridine Orange (NAO), Rhodamine 123, and Hoechst 33258 were purchased from Yusheng Biotechnology (Shanghai, China). ELISA kits were obtained from Jiyinmie Biotechnology (Wuhan, China). Primary antibodies were purchased from Abcam (Shanghai, China). NO assay kit, β -actin, the ECL Western blotting system, WB/IP lysis buffer, and BCA protein assay kit were purchased from Beyotime Biotechnology (Jiangsu, China). Other chemicals were of reagent grade from Aladdin (Shanghai, China).

2.2. Purification and Identification of ALE. Fresh *Apios* leaves were weighed, extracted, and mixed with water at a 1:10 ratio and treated by ultrasonic extraction at room tempera-

ture for 2 h. Then, the extract was centrifuged at 4000 r/min for 30 min, and centrifugation is repeated 3 times. The extract (1 g) was dissolved in 50 mL purified water, applied to a DM301 microporous resin column ($\varnothing 3.2 \times 60$ cm), washed with 1 L of water, and eluted at a flow rate of 1.0 mL/min with 20%, 40%, 60%, and 80% ethanol, respectively. After collection, the darker part of the eluent was merged, freeze-dried, and purified as ALE. ALE was stored at -80°C for subsequent experiments. The freeze-dried extract samples were dissolved in ultrapure water and prepared into a solution of 0.5 mg/mL. The main components were analyzed by LC-MS/MS after filtration with a 0.22-micron filter membrane. An Ultra Performance Liquid Chromatography (UPLC) system (Waters, Milford, MA, USA) was used for qualitative analysis. The flow rate was set at 0.8 mL/min and the temperature was maintained at 40°C . Solvent A was 15% (v/v) acetonitrile + 0.05 mol/L phosphate buffer (KH_2PO_4 -NaOH, pH 6.9), and solvent B was 40% (v/v) acetonitrile + 0.05 mol/L phosphate buffer. The solvent gradient of phase A was as follows: 0–15 min, 0–95% B; 15–21 min, 95–85% B; 21–22 min, 85–72% B; 22–24 min, 72–60% B; 24–27 min, 60–40% B; and 27–30 min, 40–95% B.

2.3. Cell Culture and Treatment. RAW264.7 cells were cultured in DMEM containing 10% fetal bovine serum, at 37°C , in a 5% CO_2 incubator. After 80% confluence, the cells were inoculated with ALE of different concentrations for 24 hours, and then, 250 ng/mL LPS was added for a total of 12 hours. Cells without LPS and ALE were used as a negative control group. Cells treated with LPS alone were used as a model group.

2.4. MTT Assays. RAW264.7 cells were seeded into a 96-well plate for 24 h. After treatments with ALE or LPS at different concentrations for 24 h, 100 μL (0.5 mg/mL) 3-(4,5-dimethylthazol-2-yl)-2,5-diphenyltetrazolium ammonium bromide (MTT) was added into each well, which was diluted with DMEM. After incubation at 37°C for 4 hours, formazan was precipitated and dissolved in 150 μL dimethyl sulfoxide (DMSO). The absorbance was measured at 570 nm with a spectrophotometer [12–14].

2.5. Nitrite Analysis. Nitric oxide levels were detected by the Griess reagent method. After treatment, RAW264.7 cell lysate and supernatant were harvested and the products from NO reactions were measured according to the instruction of the NO assay kit. The level of intracellular NO was quantified by total protein concentration. The protein concentration in the cell lysates was detected with the BCA protein concentration assay kit.

2.6. Detection of the Secretion of Cytokines. After treatment, the supernatant was collected, and the concentrations of TNF- α , IFN- γ , IL-1 β , IL-6, and IL-10 in the medium were determined according to ELISA kit instructions.

2.7. Fluorescent Staining. Fluorescent probes were used to detect intracellular ROS (DCFH, 10 mM), NO (DAF-FMDA, 5 μM), superoxide radical generation (DHE, 10 μM), GSH consumption (NDA, 50 μM), DNA damage

(Hoechst 33342, 10 μM), mitochondrial membrane potential alteration (RH123, 5 μM), mitochondrial membrane lipid peroxidation (NAO, 20 μM), and the number of mitochondria (MitoTracker Green, 20 μM). After treatment, the cells were incubated in phenol-free red medium containing different probes for 30 minutes. Then, the cells were washed with PBS 2 times and analyzed with a fluorescence microscope. The results were expressed as fluorescence intensity calculated by the image analysis software Image-Pro Plus 6.0.

2.8. Transmission Electron Microscopy (TEM). After grouping and treatment, cells in each group were collected and centrifuged at 1000 rpm for 5 min. Then, the ultrastructural changes were observed with a Hitachi H-7650 transmission electron microscope (Hitachi, Japan) operated at 15000 and 30000 magnifications after the process of fixation, rinsing, dehydration, embedding, ultrathin sectioning, and staining.

2.9. Western Blot. Cells were collected and lysed in a WB/IP lysis buffer. The lysate was centrifuged at 4°C and 12000 g for 10 min. The protein concentration was measured and leveled with the BCA protein assay kit. Equal amounts of protein (40 mg) are separated on SDS-PAGE and transferred to PVDF membranes. Then, the membrane was blocked with appropriate 5% nonfat milk and probed with 4 mL primary antibodies at 4°C overnight. The primary antibodies included iNOS, COX2, NF- κB , Keap1, Nrf2, p-JNK, JNK, p-ERK, ERK, p-p38, p38, Rab5, Rab7, Atg4, Atg5, Beclin1, LC3B, p62, HMGB1, p-FOXO1, FOXO1, p-Akt, Akt, p-mTOR, mTOR, and β -actin antibodies (Abcam). Horseradish peroxidase-(HRP-) conjugated secondary antibodies were probed to the membrane for 2 h at room temperature. The ECL luminescent liquid was added to the PVDF film. The protein bands were exposed to an Invitrogen iBright 1500 gel imaging system, and the densitometry of them was analyzed by the ImageJ software while β -actin was used as an internal reference [12].

2.10. Data Statistics and Analysis. In the study, every experiment contained at least 3 biological repetitions and every biological repetition was conducted with at least 3 technical repeats. Fluorescence images were analyzed by Image-Pro Plus for average optical density. Western blot results were quantified by ImageJ. Experimental data were expressed as the means and SD and analyzed by SPSS (Statistical Product and Service Solutions) 19.0 statistical software. Variance analysis was performed by one-way analysis of variance (ANOVA) followed by Duncan's multiple range test by Duncan's multiple comparisons. $p < 0.05$ was considered significant, and data were plotted with GraphPad Prism 5 [15, 16].

3. Results

3.1. Composition and Identification of ALE. According to the HPLC-MS/MS (Figure 1(a)) result, there are seven main components in ALE (Table 1): vicenin-2 (peak 1) [17], schaftoside (peak 2) [18], baimaside (peak 3), apigenin (6-C- α -L-arabinopyranosyl)-8-C- β -D-glucopyranoside (peak 4) [12], vitexin (peak 5), isquercetin (peak 6), and 6-C- β -D-glucopyranosyl-7-O-methyluteolin (peak 7). The highest

content of the component in ALE was vitexin, followed by schaftoside, and the proportion of other components was low. Besides, Table 1 showed the peak identification, retention times, MS/MS date, and compound identification of the components.

3.2. ALE Attenuates LPS-Induced Cytotoxicity and Inflammation in RAW264.7 Cells. LPS was applied to induce the inflammation of RAW264.7. The results in Figures 1(b)–1(d) showed that LPS stimulation induced the decline in cell proliferation activity and led to morphological changes of RAW264.7 cells. However, the pretreatment of ALE at different concentrations could reverse the situation induced by LPS (Figures 1(b) and 1(c)). It could be observed that ALE at concentrations below 100 $\mu\text{g}/\text{mL}$ had no toxicity on RAW264.7 cells and could alleviate the cytotoxicity induced by LPS. Then, we used the Hoechst 33258 fluorescence probe to measure the protective effect of ALE [19]. Compared with the LPS-induced group, the number of bright blue points in the ALE-treated group was significantly reduced, and the LPS-induced chromatin condensation was attenuated, suggesting that ALE had a positive effect on LPS-induced cell apoptosis (Figure 1(e)). Besides, we used the DAF-FMDA fluorescence probe to examine intracellular NO levels (Figure 1(f)). The ALE-treated group significantly reduced the content of intracellular NO (Figure 1(g)). In addition, Figures 1(h) and 1(i) show that the stimulation of LPS could significantly increase the level of NO in cells and in culture medium, while ALE inhibited such increases (Figures 1(h) and 1(i)). Herein, the above results showed that ALE helped relieve inflammation on LPS-induced RAW264.7 cells.

3.3. ALE Decreases LPS-Induced Overproduction of Inflammatory Cytokines in RAW264.7 Cells. Cytokines such as IL-1 β , IL-2, IL-6, TNF- α , and IFN- γ are involved in the inflammation process [20]. Therefore, we further detected the production of inflammatory cytokines in cell culture medium. The results showed that LPS stimulated the elevation of inflammatory cytokines. However, the treatment of ALE at different concentrations suppressed the secretion of inflammatory cytokines in RAW264.7 cells (Figure 2), suggesting the alleviation of inflammation induced by LPS.

3.4. ALE Ameliorates LPS-Induced Oxidative Damage in RAW264.7 Cells. LPS-induced inflammation is accompanied by oxidative stress [21]. Natural antioxidants, such as flavonoids, have potential ability to reduce oxidative stress. As shown in Figure 3, LPS significantly increased the levels of intracellular ROS (Figures 3(a) and 3(d)) and $\text{O}_2^{\cdot-}$ contents (Figures 3(a) and 3(e)) and decreased the level of glutathione (GSH) (Figures 3(c) and 3(f)). ALE treatment significantly inhibited the formation of ROS and $\text{O}_2^{\cdot-}$ and recover the level of GSH content, indicating the relief of oxidative stress induced by LPS. Studies have reported that natural products could inhibit the production of ROS and $\text{O}_2^{\cdot-}$ and increase the production of GSH [21]. Meanwhile, our previous studies have shown that extracts from the flowers and leaves of *Apios americana* Medikus




FIGURE 1: HPLC elution profiles of compounds in *Apios americana* Medik leaves and effects of ALE on LPS-induced cytotoxicity and inflammation in RAW264.7 cells. (a) The liquid chromatography profile of ALE in *Apios americana* Medik leaves. The peak numbers were labeled according to the retention times. (b) RAW264.7 cell viability was measured by the MTT method after treatment with ALE at different concentrations for 24h with/without LPS inducement ($n = 6$). (c) Cell proliferation assay via BeyoClick™ EdU Cell Proliferation Kit with Alexa Fluor 488. (d) Morphological changes of RAW264.7 cells after ALE treatment ($n = 3$). (e, f) Cell apoptosis, NO staining with Hoechst 33258, and DAF-FMDA in the presence of LPS and ALE ($n = 3$). (g) The quantitative data of (f). (h) NO production in cell culture medium according to the Griess reagent method ($n = 4$). (i) Intracellular concentration of NO according to the Griess reagent method quantified by total protein concentration ($n = 4$). ALE-treated cells were inoculated in different concentrations of ALE for 24 hours, and then, 250 ng/mL LPS was added for a total of 12 hours. Cells without LPS and ALE were used as a negative control group. Cells treated with LPS alone were used as a model group. Images were captured with a fluorescence microscope in the same settings. All the fluorescence images were quantified in the whole field with the background removed and represented by normalized fluorescence (y-axes) via Image-Pro Plus 6.0 ($n = 3$). Significance analysis was carried out according to the one-way ANOVA test, and different letters in figures mean statistically significant differences among the groups (A, B, C, and D were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).

TABLE 1: Identification of ALE.

No.	RT (min)	Molecular ion (M+)	MS/MS (M+)	Tentative identification
1	20.063	593.1555	297.0759, 325.0710, 353.0671, 383.0772, 473.1107, 503.1217	Vicenin-2
2	21.737	563.1427	297.0763, 325.0713, 353.0668, 383.0776, 413.0880, 443.0995, 473.1106	Schaftoside
3	22.033	625.1432	151.0029, 178.9978, 271.0238, 300.0272, 301.0343, 343.0454	Baimaside
4	22.217	563.1422	149.0240, 191.0342, 221.0454, 297.0770, 353.0668, 383.0776, 413.0878, 433.0986, 473.1100, 545.1307	Apigenin (6-C- α -L-arabinopyranosyl)-8-C- β -D-glucopyranoside
5	23.653	431.0994	161.0238, 268.0368, 281.0449, 283.0607, 311.0557, 341.0667	Vitexin
6	24.133	463.0883	151.0030, 243.0296, 255.0293, 271.0247, 300.0265, 301.0345	Isquercetin
7	24.397	461.1116	297.0397, 298.0478, 341.0674, 371.07788, 461.1116	6-C- β -D-Glucopyranosyl-7-O-methyluteolin




FIGURE 2: Effects of ALE on LPS-induced secretion of inflammatory factors in RAW264.7 cells. (a–e) IL-1 β , IL-2, IL-6, TNF- α , and IFN- γ production in cell culture medium according to the ELISA method ($n = 4$). ALE-treated cells were inoculated in different concentrations of ALE for 24 hours, and then, 250 ng/mL LPS was added for a total of 12 hours. Cells without LPS and ALE were used as a negative control group. Cells treated with LPS alone were used as a model group. Images were captured with a fluorescence microscope in the same settings. Significance analysis was carried out according to the one-way ANOVA test, and different letters in figures mean statistically significant differences among the groups (A, B, C, and D were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).

could also increase the levels of GSH in cells, which is beneficial to HepG2 cells' resistance to H₂O₂-induced oxidative stress [22].

3.5. ALE Attenuates LPS-Induced Mitochondrial Dysfunction in RAW264.7 Cells. Numerous studies have shown that LPS can induce inflammation, leading to oxidative stress and mitochondrial dysfunction, as well as the induction of autophagy [23, 24]. Mitochondria play a vital role in this process, which secrete mtDNA and ROS to activate the inflammasome and induce autophagy. Reports have shown that the overgeneration of ROS exerted a huge impact on

mitochondrial metabolism [25]. Additionally, mitochondrial dysfunction, characterized by the decrease of the mitochondria and the imbalance of the lipid peroxidation of the mitochondrial membrane (MMMP) and mitochondrial membrane potential (MMP), is correlative to oxidative stress caused by inflammation [26]. As shown in Figure 4, LPS treatment declined the mean fluorescence intensity of MitoTracker Green and NAO, while increasing the mean RH123 fluorescence intensity. However, these trends were suppressed by ALE, indicating that the reduction of mitochondria was inhibited after ALE pretreatment, while MMMP and MMP were significantly alleviated (Figures 4(b), 4(e),




FIGURE 3: Effects of ALE on LPS-induced oxidative damage in RAW264.7 cells ($n = 3$). (a) DCFH-DA staining for ROS. (b) DHE staining for $O_2^{\bullet-}$. (c) NDA staining for GSH. (d) The quantitative data of (a). (e) The quantitative data of (b). (f) The quantitative data of (c). ALE-treated cells were inoculated in different concentrations of ALE for 24 hours, and then, 250 ng/mL LPS was added for a total of 12 hours. Cells without LPS and ALE were used as a negative control group. Cells treated with LPS alone were used as a model group. Images were captured with a fluorescence microscope in the same settings. All the fluorescence images were quantified in the whole field with the background removed and represented by normalized fluorescence (y -axes) via Image-Pro Plus 6.0 ($n = 3$). Significance analysis was carried out according to the one-way ANOVA test, and different letters in figures mean statistically significant differences among the groups (A, B, C, and D were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).

5(c), and 5(f)). It has been reported that flavonoids can protect mitochondria from oxidative damage and remove excessive free radicals both *in vitro* and *in vivo*, which is consistent with our results [27, 28].

3.6. ALE Ameliorates Oxidative Stress and Inflammation via $NF-\kappa B$, $Nrf2$ -Keap1, and MAPKs in RAW264.7 Cells. $NF-\kappa B$ is an important factor related to the regulation of inflammatory mediators. The transport of $NF-\kappa B$ is involved in the expression of iNOS and COX2 [28, 29]. The above studies have confirmed that ALE can significantly inhibit the overproduction of NO; thus, we further detect the effects of ALE on expressions of iNOS, COX2, and $NF-\kappa B$. The results showed that LPS upregulated the expressions of iNOS, COX2, and $NF-\kappa B$, while the treatment of ALE could

suppress such upregulation triggered by LPS (Figures 5(a) and 5(c)–5(e)) thereby inhibiting the production of NO in cells. It has also been confirmed in above results that ALE could also attenuate the oxidative damage induced by LPS (Figure 3). As an essential protein in antioxidative stress, $Nrf2$ is able to help regulate the oxidative stress [30]. In order to investigate whether the anti-inflammatory effect of ALE is related to the $Nrf2$ pathway, we examined the expression of $Nrf2$ and Keap1 [30]. ALE significantly activated the expression of these two proteins in the presence of LPS, indicating that ALE promoted the dissociation of $Nrf2$ /Keap1 and activated the $Nrf2$ -mediated antioxidant cascade in RAW264.7 cells (Figures 5(a), 5(f), and 5(g)). MAPKs, including JNK1/2, ERK1/2, and p38 MAPK, are crucial in adjusting the inflammation induced by external factors [31]. ERK,




FIGURE 4: Effect of ALE on LPS-induced mitochondrial dysfunction in RAW264.7 cells ($n = 3$). (a–c) Mitochondria, mitochondrial membrane potential, and mitochondrial membrane lipid peroxidation alterations of RAW164.7 cells in the presence of LPS and ALE under different treatments which were incubated with MitoTracker, RH123, and NAO. (d–f) The quantitative data of (a), (b), and (c). Images were captured with a fluorescence microscope in the same settings. All the fluorescence images were quantified in the whole field with the background removed and represented by normalized fluorescence (y -axes) via Image-Pro Plus 6.0 ($n = 3$). Significance analysis was carried out according to the one-way ANOVA test, and different letters in figures mean statistically significant differences among the groups (A, B, C, and D were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).

which exists widely in various tissues, is involved in the regulation of cell proliferation and differentiation. JNK is a key in cell signal transduction induced by various factors and related to cell stress responses such as radiation, osmotic pressure, and temperature [32]. p38 is involved in inflammation and apoptosis [4]. The phosphorylation of MAPKs induced by LPS was involved in the activations of iNOS, COX2, and proinflammatory factors [32]. In order to confirm the effect of ALE on the expressions of MAPKs in RAW264.7 cells, we detected these proteins. Results showed that ALE inhibited the phosphorylation of MAPKs induced by LPS (Figures 5(b) and 5(h)–5(j)), indicating that ALE could relieve inflammation through MAPK pathways. The results above manifested that ALE ameliorated oxidative stress and inflammation via NF- κ B, Nrf2-Keap1, and MAPKs in RAW264.7 cells.

3.7. ALE Activates Autophagy to Resist Inflammation in RAW264.7 Cells. Redundant ROS/RNS will induce the mitochondrial dysfunction and the activation of mitochondrial autophagy against inflammation [31]. Autophagy is an intracellular self-protection mechanism that degrades misfolded proteins and dysfunctional organelles by lysosomes. Autophagy is involved in a variety of diseases including cancer, diabetes, aging, neurodegenerative diseases, inflammation, and liver diseases. In most cases, autophagy is both cytoprotective and destructive [32].

In our study, ALE could reduce oxidative stress, mitochondrial damage, and inflammation induced by LPS, which was consistent with previous studies [4]. So, we speculated that ALE could activate the autophagy in macrophages to resist inflammation. The result of TEM showed that the number of autophagosomes increased after ALE treatment,





FIGURE 5: ALE treatment altered expressions of proteins associated with inflammation and oxidative stress in RAW264.7 cells. (a) Western blot bands of iNOS, COX2, NF- κ B, and Nrf2-Keap1 ($n = 3$). (b) Western blot bands of MAPKs. (c–j) Relative expressions of iNOS, COX2, NF- κ B, Keap1, Nrf2, p-JNK/JNK, p-ERK/ERK, and p-p38/p38. The images were quantified by the ImageJ software, and the intensity of bands was corrected with β -actin ($n = 3$); vertical lines in the histogram represent SDs of three replicates. Significance analysis was carried out according to the one-way ANOVA test, and different letters in the histogram mean statistically significant differences among the groups (A, B, C, D, and E were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).


suggesting that ALE could enhance autophagy in RAW264.7 cells (Figure 6(a)). Western blot results showed that compared with the LPS-treated group, the expressions of Beclin1,

Atg4, and Atg5 were upregulated in the ALE-treated group, while the expression level of p62 was significantly inhibited in a dose-dependent manner (Figures 6(c)–6(i)). Meanwhile,




Control LPS ALE (40 µg/ml)


(a)




(b)




(c)



(d)



(e)



(f)

FIGURE 6: Continued.




FIGURE 6: ALE activated autophagy in LPS-induced RAW264.7 cells. (a) Transmission electron microscopy features of macrophages ($n = 3$). (b) Western blot bands of Rab5, Rab7, Atg4, Atg5, Beclin1, LC3B, p62, and β -actin ($n = 3$). (c–i) Relative expression levels of Atg4, Atg5, Beclin1, p62, LC3BII/LC3BI, Rab5, and Rab7. The images were quantified by the ImageJ software, and the intensity of bands was corrected with β -actin ($n = 3$); vertical lines in the histogram represent SDs of three replicates. Significance analysis was carried out according to the one-way ANOVA test, and different letters mean statistically significant differences among the groups (A, B, C, D, and E were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).

the expression of LC3II/LC3I was enhanced after the treatment of ALE. These results together confirmed that ALE could activate autophagy to resist inflammation in LPS-induced RAW264.7 cells [32].

3.8. ALE Induces Cell Autophagy via HMGB1, FoxO1, and Akt-mTOR Pathways in RAW264.7 Cells. ALE can induce autophagy, which facilitates in reducing inflammation. There are many signal pathways concerned with the regulation of autophagy as previously reported [33]. HMGB1, FoxO1, and Akt-mTOR are the key regulators to control autophagy [22, 34–36]. Researchers have proven that HMGB1 facilitates the phosphorylation of Bcl-2 to promote the dissociation of Beclin1 from the Beclin1-Bcl2 compound and the combination to the autophagy regulator Beclin1, thereby contributing to the assembly of the PI3KC3-Beclin1 compound and autophagy [37]. We found that ALE triggered the expression of HMGB1 (Figures 7(a) and 7(b)) and the delivery of Beclin1 (Figures 6(b)–6(i)). Under stress, FoxO1 could combine with the promoter of Atg12, which led to the activation of autophagy. Our research confirmed that ALE contributed to the phosphorylation of FoxO1 and promoted the expression of Rab5 and Rab7 (Figures 6(b)–6(i)), suggesting that ALE played an important role in the fusion of autophagosomes and lysosomes. In addition, the Akt-mTOR pathway is important in cellular inflammation and autophagy. The intervention of ALE caused the downregulation of relative expressions of p-Akt/Akt and p-mTOR/mTOR (Figure 7(c)), thereby contributing to autophagy, which was shown in the results of western blot. In conclusion, ALE induces cell autophagy via activation of HMGB1 and FoxO1 and inhibition of Akt-mTOR pathways in RAW264.7 cells.

4. Discussion

Flavonoids from natural plants have attracted extensive attention due to their excellent biological activities (such as antioxidant, anti-inflammatory, antitumor, and antibacterial). It is generally accepted that plant-derived functional

factors are nontoxic and have side effects [38]. Many studies have shown that the biological activities of flavonoids depend on their structural characteristics, and different flavonoids have different activities [39]. A large number of studies have shown that natural flavonoids have good anti-inflammatory activities. For example, apigenin can prevent colon inflammation and motor dysfunction in obese mice induced by a high-fat diet [40]. Naringin can reduce the cell viability, fat content, and inflammatory response of breast tumor cells in ovariectomized obese mice [41]. Liquiritigenin can reduce pulmonary inflammation and incidence of viral infection in mice [42]. The citrus flavonoid hesperidin can alleviate testicular injury in diabetic rats by inhibiting oxidative stress, inflammation, and apoptosis [43]. Baicalin can regulate the expression of inflammation-related genes in vivo to relieve skin inflammation [44].

In this study, a series of flavonoids were identified in ALE and the anti-inflammation effect of ALE was explored. Furthermore, ALE altered NF- κ B, MAPK, and Nrf2-Keap1 pathways to activate the anti-inflammatory system in RAW264.7 cells, at the same time via activating HMGB1-Beclin1 and FoxO1 pathways and inhibiting Akt-mTOR signaling pathways to induce cell autophagy, which could degrade damaged mitochondria and macromolecular substances, thereby lowering the secretion of NO and inflammatory cytokines. The anti-inflammatory processes of ALE involved in NF- κ B, MAPK, Nrf2-keap1, HMGB1-Beclin1, FoxO1, and Akt-mTOR signaling pathways and multiple pathways were linked to resist lipopolysaccharide-induced inflammatory response (Figure 7(e)).

During the anti-inflammation process, autophagy plays a vital role. Autophagy is an intracellular self-protective mechanism that degrades misfolded proteins and dysfunctional organelles through lysosomes [45]. Autophagy is also involved in a variety of diseases, including cancer, diabetes, aging, neurodegenerative diseases, inflammation, and liver diseases [46]. In most cases, autophagy is both cell protective and destructive. Numerous studies have shown that stimulation of LPS can induce inflammatory responses, leading to oxidative stress and mitochondrial dysfunction, as well as




FIGURE 7: Effect of ALE on expression levels of HMGB1, FoxO1, and Akt-mTOR pathways in RAW264.7 cells. (a) Western blot bands of HMGB1, p-FOXO1, FOXO1, p-Akt, Akt, p-mTOR, mTOR, and β -actin ($n = 3$). (b–e) Relative expression levels of HMGB1, p-FOXO1/FOXO1, p-Akt/Akt, and p-mTOR/mTOR. The images were quantified by the ImageJ software, and the intensity of bands was corrected with β -actin ($n = 3$); vertical lines in the histogram represent SDs of three replicates. (f) The schematic of the mechanism of impact of ALE on LPS-induced RAW264.7 cells. Significance analysis was carried out according to the one-way ANOVA test, and different letters mean statistically significant differences among the groups (A, B, C, D, and E were labeled from large to small and the same letters in columns mean statistical insignificance; otherwise, they mean statistical significance ($p < 0.05$)).

inducing autophagy. Mitochondria play a vital role in this process [47]. The secretion of mtDNA and ROS from mitochondria activates the inflammasome and induces impaired autophagy clearance, which is essential for activating the inflammatory response. In addition, oxidative stress can accelerate the formation of autophagy, and autophagy can reduce oxidative damage by degrading oxidative substances and damaged mitochondria [48, 49]. Our results confirm that ALE can induce autophagy of macrophages by activating the Hmgb1-Beclin1 and Sirt1-FOXO1 pathway and inhibiting the Akt-mTOR signaling pathway, thereby reducing LPS-induced oxidative stress, mitochondrial dysfunction, and inflammation in RAW264.7 cells. However, it is still unclear whether other pathways are also involved in regulating autophagy, and more studies are needed to demonstrate the interaction between ALE and autophagy in LPS-induced RAW264.7 cells.

5. Conclusion

Flavonoids derived from natural sources play significant roles in anti-inflammatory, antitumor, and antioxidant processes [50]. In this study, seven compounds were extracted from *Apios americana* Medik leaves and identified by LC-TOF-MS. Using *in vitro* studies, we first demonstrated that ALE could protect RAW264.7 cells from cytotoxicity, inflammation, oxidative damage, and mitochondrial dysfunction against LPS. Meanwhile, ALE reduced the overproduction of NO and inflammatory cytokines via inhibition of NF- κ B and MAPK signaling and activation of Nrf2-Keap1 and FOXO pathways [51]. In addition, ALE could activate HMGB1-Beclin1 and Sirt1-FoxO1 pathways and inhibit the Akt-mTOR signaling pathway to induce autophagy in RAW264.7 cells, thereby contributing to the decomposition of damaged mitochondria and maintaining ROS homeostasis. However, it is unclear whether other pathways are involved in the regulation of autophagy activation, and further research is needed to demonstrate the interaction between the protection effect of ALE and autophagy in RAW264.7 cells. Overall, this finding manifests that the leaf of *Apios americana* Medik has potential to be the source of an antioxidant and anti-inflammatory drug, which means *Apios americana* Medik leaves can be applied to promote human health in future.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare no competing interests.

Acknowledgments

This work was supported by the Science and Technology Department of Zhejiang Province (No. 2018C02045).

References

- [1] Q. Chu, R. Jia, M. Chen et al., "Tetrastigma hemsleyanum tubers polysaccharide ameliorates LPS-induced inflammation in macrophages and *Caenorhabditis elegans*," *International Journal of Biological Macromolecules*, vol. 141, pp. 611–621, 2019.
- [2] G. S. Hotamisligil, "Inflammation, metaflammation and immunometabolic disorders," *Nature*, vol. 542, no. 7640, pp. 177–185, 2017.
- [3] G. M. O. Irene, C. Marta, Á. C. Ana, Á. V. Luis, and S. Laura, "Lipopolysaccharide activates Toll-like receptor 4 (TLR4)-mediated NF- κ B signaling pathway and proinflammatory response in human reryocytes," *Journal of Biological Chemistry*, vol. 289, no. 4, pp. 2457–2468, 2014.
- [4] M. T. Islam, "Oxidative stress and mitochondrial dysfunction-linked neurodegenerative disorders," *Neurological Research*, vol. 39, no. 1, pp. 73–82, 2017.
- [5] A. Bhattacharyya, R. Chattopadhyay, S. Mitra, and S. E. Crowe, "Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases," *Physiological Reviews*, vol. 94, no. 2, pp. 329–354, 2014.
- [6] L. Li, J. Tan, Y. Miao, P. Lei, and Q. Zhang, "ROS and autophagy: interactions and molecular regulatory mechanisms," *Cellular and Molecular Neurobiology*, vol. 35, no. 5, pp. 615–621, 2015.
- [7] X. Zhu, J. S. Messer, Y. Wang et al., "Cytosolic HMGB1 controls the cellular autophagy/apoptosis checkpoint during inflammation," *Journal of Clinical Investigation*, vol. 125, no. 3, pp. 1098–1110, 2015.
- [8] S. Wang, P. Xia, G. Huang et al., "FoxO1-mediated autophagy is required for NK cell development and innate immunity," *Nature Communications*, vol. 7, article 11023, 2016.
- [9] M. W. James and C. J. Hawkey, "Assessment of non-steroidal anti-inflammatory drug (NSAID) damage in the human gastrointestinal tract," *British Journal of Clinical Pharmacology*, vol. 56, no. 2, pp. 146–155, 2015.
- [10] Q. Chu, S. Zhang, L. Yu et al., "*Apios americana* Medikus tuber polysaccharide exerts anti-inflammatory effects by activating autophagy," *International Journal of Biological Macromolecules*, vol. 130, pp. 892–902, 2019.
- [11] Q. Chu, S. Zhang, M. Chen et al., "Cherry anthocyanins regulate NAFLD by promoting autophagy pathway," *Oxidative Medicine and Cellular Longevity*, vol. 2019, Article ID 4825949, 16 pages, 2019.
- [12] C. Qiang, Y. Zhao, X. Shi et al., "In vivo-like 3-D model for sodium nitrite- and acrylamide-induced hepatotoxicity tests utilizing HepG2 cells entrapped in micro-hollow fibers," *Scientific Reports*, vol. 7, no. 1, article 14837, 2017.
- [13] F. Ferreres, B. M. Silva, P. B. Andrade, R. M. Seabra, and M. A. Ferreira, "Approach to the study of C-glycosyl flavones by ion trap HPLC-PAD-ESI/MS/MS: application to seeds of quince (*Cydonia oblonga*)," *Phytochemical Analysis*, vol. 14, no. 6, pp. 352–359, 2010.
- [14] Y. Li, Q. Chu, Y. Liu, X. Ye, Y. Jiang, and X. Zheng, "Radix Tetrastigma flavonoid ameliorates inflammation and prolongs the lifespan of *Caenorhabditis elegans* through JNK, P38 and Nrf2 pathways," *Free Radical Research*, vol. 53, no. 5, pp. 562–573, 2019.
- [15] Q. Chu, M. Chen, D. Song et al., "*Apios americana* Medik flowers polysaccharide (AFP-2) attenuates H₂O₂ induced neurotoxicity in PC12 cells," *International Journal of*

- Biological Macromolecules*, vol. 123, article S0141813018331532, pp. 1115–1124, 2019.
- [16] Y. Y. Zhao, Q. Chu, X. E. Shi, X. D. Zheng, X. T. Shen, and Y. Z. Zhang, “Toxicity testing of four silver nanoparticle-coated dental castings in 3-D LO2 cell cultures,” *Journal of Zhejiang University-SCIENCE B*, vol. 19, no. 2, pp. 159–167, 2018.
- [17] R. Colombo, J. H. Yariwake, E. F. Queiroz, K. Ndjoko, and K. Hostettmann, “On-line identification of further flavone C- and O-glycosides from sugarcane (*Saccharum officinarum* L., Gramineae) by HPLC-UV-MS,” *Phytochemical Analysis*, vol. 17, no. 5, pp. 337–343, 2010.
- [18] P. Anna Lisa, G. M. Milagros, A. Dulce María et al., “HPLC-PDA-MS and NMR characterization of C-glycosyl flavones in a hydroalcoholic extract of *Citrus aurantifolia* leaves with antiplatelet activity,” *Journal of Agricultural and Food Chemistry*, vol. 56, no. 5, pp. 1574–1581, 2008.
- [19] H. J. Young, P. Yoon Jeong, H. Mira, and K. Jae Bum, “Cross-talk between adipocytes and immune cells in adipose tissue inflammation and metabolic dysregulation in obesity,” *Molecules and Cells*, vol. 37, no. 5, pp. 365–371, 2014.
- [20] X. Xu, H. Li, X. Hou et al., “Punicalagin induces Nrf2/HO-1 expression via upregulation of PI3K/AKT pathway and inhibits LPS-induced oxidative stress in RAW264.7 macrophages,” *Mediators of Inflammation*, vol. 2015, Article ID 2015, 11 pages.
- [21] V. I. Lushchak, “Free radicals, reactive oxygen species, oxidative stress and its classification,” *Chemico-Biological Interactions*, vol. 224, pp. 164–175, 2014.
- [22] Q. Chu, L. Yu, Z. Zheng et al., “*Apios americana* Medik flowers extract protects PC12 cells against H₂O₂ induced neurotoxicity via regulating autophagy,” *Food and Chemical Toxicology*, vol. 124, pp. 231–238, 2019.
- [23] H. Blaser, C. Dostert, T. W. Mak, and D. Brenner, “TNF and ROS crosstalk in inflammation,” *Trends in Cell Biology*, vol. 26, no. 4, pp. 249–261, 2016.
- [24] N. A. Panat, D. K. Maurya, S. S. Ghaskadbi, and S. K. Sandur, “Trolox, a plant flavonoid, protects cells against oxidative stress-induced cell death through radical scavenging mechanism,” *Food Chemistry*, vol. 194, pp. 32–45, 2016.
- [25] T. Liu, L. Zhang, D. Joo, and S. C. Sun, “NF- κ B signaling in inflammation,” *Signal Transduction and Targeted Therapy*, vol. 2, no. 1, article 17023, 2017.
- [26] S. Lee, C. Kwak, S. Lee et al., “Anti-inflammatory effect of ascochlorin in LPS-stimulated RAW 264.7 macrophage cells is accompanied with the down-regulation of iNOS, COX-2 and proinflammatory cytokines through NF- κ B, ERK1/2 and P38 signaling pathway,” *Journal of Cellular Biochemistry*, vol. 117, no. 4, pp. 978–987, 2016.
- [27] S. Lee, C. Kwak, S. Lee et al., “Anti-inflammatory effect of ascochlorin in LPS-stimulated RAW 264.7 macrophage cells is accompanied with the down-regulation of iNOS, COX-2 and proinflammatory cytokines through NF- κ B, ERK1/2 and P38 signaling pathway,” *Journal of Cellular Biochemistry*, vol. 117, no. 4, pp. 978–987, 2016.
- [28] A. Loboda, M. Damulewicz, E. Pyza, A. Jozkowicz, and J. Dulak, “Role of Nrf2/HO-1 system in development, oxidative stress response and diseases: an evolutionarily conserved mechanism,” *Cellular and Molecular Life Sciences*, vol. 73, no. 17, pp. 3221–3247, 2016.
- [29] M. Chunfen, W. Ling, Z. Jie et al., “The crosstalk between Nrf2 and AMPK signal pathways is important for the anti-inflammatory effect of berberine in LPS-stimulated macrophages and endotoxin-shocked mice,” *Antioxidants & Redox Signaling*, vol. 20, no. 4, pp. 574–588, 2014.
- [30] Y. S. Kim, C. B. Ahn, and J. Y. Je, “Anti-inflammatory action of high molecular weight *Mytilus edulis* hydrolysates fraction in LPS-induced RAW264.7 macrophage via NF- κ B and MAPK pathways,” *Food Chemistry*, vol. 202, pp. 9–14, 2016.
- [31] B. Ahn, W. K. Jung, S. J. Park, Y. T. Kim, W. S. Kim, and J. Y. Je, “Gallic acid-g-chitosan modulates inflammatory responses in LPS-stimulated RAW264.7 cells via NF- κ B, AP-1, and MAPK pathways,” *Inflammation*, vol. 39, no. 1, pp. 366–374, 2016.
- [32] Y. Yu, M. Shen, Q. Song, and J. Xie, “Biological activities and pharmaceutical applications of polysaccharide from natural resources: a review,” *Carbohydrate Polymers*, vol. 183, pp. 91–101, 2018.
- [33] D. Wu, P. Zhong, J. Wang, and H. Wang, “Exogenous hydrogen sulfide mitigates LPS+ATP-induced inflammation by inhibiting NLRP3 inflammasome activation and promoting autophagy in L02 cells,” *Molecular and Cellular Biochemistry*, vol. 457, no. 1–2, pp. 145–156, 2019.
- [34] N. Wang, Q. Zhang, L. Luo, B. Ning, and Y. Fang, “ β -Asarone inhibited cell growth and promoted autophagy via P53/Bcl-2/Bcl-1 and P53/AMPK/mTOR pathways in human glioma U251 cells,” *Journal of Cellular Physiology*, vol. 233, no. 3, pp. 2434–2443, 2018.
- [35] N. K. Zenkov, A. V. Chechushkov, P. M. Kozhin, N. V. Kandalintseva, G. G. Martinovich, and E. B. Menshchikova, “Plant phenols and autophagy,” *Biochemistry*, vol. 81, no. 4, pp. 297–314, 2016.
- [36] L. Li, J. Tan, Y. Miao, P. Lei, and Q. Zhang, “ROS and autophagy: interactions and molecular regulatory mechanisms,” *Cellular and Molecular Neurobiology*, vol. 35, no. 5, pp. 615–621, 2015.
- [37] W. Martina, J. Justin, and S. A. Tooze, “Autophagosome formation—the role of ULK1 and Beclin1–PI3KC3 complexes in setting the stage,” *Seminars in Cancer Biology*, vol. 23, no. 5, pp. 301–309, 2013.
- [38] A. Zong, H. Cao, and F. Wang, “Anticancer polysaccharides from natural resources: a review of recent research,” *Carbohydrate Polymers*, vol. 90, no. 4, pp. 1395–1410, 2012.
- [39] T. Y. Wang, Q. Li, and K. S. Bi, “Bioactive flavonoids in medicinal plants: structure, activity and biological fate,” *Asian Journal of Pharmaceutical Sciences*, vol. 13, no. 1, pp. 12–23, 2018.
- [40] D. Gentile, M. Fornai, R. Colucci et al., “The flavonoid compound apigenin prevents colonic inflammation and motor dysfunctions associated with high fat diet-induced obesity,” *PLoS One*, vol. 13, no. 4, article e0195502, 2018.
- [41] J. Y. Ke, T. Banh, Y. H. Hsiao et al., “Citrus flavonoid naringenin reduces mammary tumor cell viability, adipose mass, and adipose inflammation in obese ovariectomized mice,” *Molecular Nutrition & Food Research*, vol. 61, no. 9, article 1600934, 2017.
- [42] H. Traboulsi, A. Cloutier, K. Boyapelly et al., “The flavonoid isoliquiritigenin reduces lung inflammation and mouse morbidity during influenza virus infection,” *Antimicrobial Agents and Chemotherapy*, vol. 59, no. 10, pp. 6317–6327, 2015.
- [43] A. Samie, R. Sedaghat, T. Baluchnejadmojarad, and M. Roghani, “Hesperetin, a citrus flavonoid, attenuates

- testicular damage in diabetic rats via inhibition of oxidative stress, inflammation, and apoptosis," *Life Sciences*, vol. 210, pp. 132–139, 2018.
- [44] S. J. van Dijk, P. L. Molloy, H. Varinli, J. L. Morrison, and B. S. Muhlhausler, "Epigenetics and human obesity," *International Journal of Obesity*, vol. 39, no. 1, pp. 85–97, 2015.
- [45] L. Galluzzi, E. H. Baehrecke, A. Ballabio et al., "Molecular definitions of autophagy and related processes," *The EMBO Journal*, vol. 36, no. 13, pp. 1811–1836, 2017.
- [46] J. A. Rodrigueznavarro and A. M. Cuervo, "Autophagy and lipids: tightening the knot," *Seminars in Immunopathology*, vol. 32, no. 4, pp. 343–353, 2010.
- [47] S. Shabalala, J. Louw, C. J. F. Muller, J. Louw, and R. Johnson, "Polyphenols, autophagy and doxorubicin-induced cardiotoxicity," *Life Sciences*, vol. 180, pp. 160–170, 2017.
- [48] K. Nakahira, J. A. Haspel, V. A. K. Rathinam et al., "Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome," *Nature Immunology*, vol. 12, no. 3, pp. 222–230, 2011.
- [49] G. Kroemer, "Autophagy and the integrated stress response," *Molecular Cell*, vol. 40, no. 2, pp. 280–293, 2010.
- [50] S. H. Nile, Y. S. Keum, A. S. Nile, S. S. Jalde, and R. V. Patel, "Antioxidant, anti-inflammatory, and enzyme inhibitory activity of natural plant flavonoids and their synthesized derivatives," *Journal of Biochemical and Molecular Toxicology*, vol. 32, no. 1, article e22002, 2018.
- [51] H. Yan, Y. Yuan, Z. Xi, Z. Kun, C. Shaohua, and D. Zhiyun, "Curcumin, inflammation, and chronic diseases: how are they linked?," *Molecules*, vol. 20, no. 5, pp. 9183–9213, 2015.



Hindawi

Submit your manuscripts at
www.hindawi.com

