Review Article

ROS-Mediated Cancer Cell Killing through Dietary Phytochemicals

Saranya NavaneethaKrishnan, Jesusa L. Rosales, and Ki-Young Lee

Department of Cell Biology and Anatomy, and Arnie Charbonneau Cancer and Alberta Children's Hospital Research Institutes, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada T2N 4N1

Correspondence should be addressed to Ki-Young Lee; kylee@ucalgary.ca

Received 27 February 2019; Accepted 18 April 2019; Published 14 May 2019

Guest Editor: Stefano Falone

Copyright © 2019 Saranya NavaneethaKrishnan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Reactive oxygen species (ROS) promote carcinogenesis by inducing genetic mutations, activating oncogenes, and raising oxidative stress, which all influence cell proliferation, survival, and apoptosis. Cancer cells display redox imbalance due to increased ROS level compared to normal cells. This unique feature in cancer cells may, therefore, be exploited for targeted therapy. Over the past few decades, natural compounds have attracted attention as potential cancer therapies because of their ability to maintain cellular redox homeostasis with minimal toxicity. Preclinical studies show that bioactive dietary polyphenols exert antitumor effects by inducing ROS-mediated cytotoxicity in cancer cells. These bioactive compounds also regulate cell proliferation, survival, and apoptotic and antiapoptotic signalling pathways. In this review, we discuss (i) how ROS is generated and (ii) regulated and (iii) the cell signalling pathways affected by ROS. We also discuss (iv) the various dietary phytochemicals that have been implicated to have cancer therapeutic effects through their ROS-related functions.

1. Introduction

Reactive oxygen species (ROS) are highly reactive metabolic by-products that cause both deleterious and beneficial effects. Cellular ROS act as secondary messengers in signalling cascades that are critical for normal physiological functions such as differentiation and development [1, 2]. However, overproduction of ROS can cause damage to biomolecules such as DNA, lipids, carbohydrates, and proteins [3, 4], leading to loss of cell integrity and subsequently cell pathology (Figure 1). For example, ROS is now recognized to promote tumorigenesis, metastasis, and angiogenesis [5]. But then again, in cancer, excessive accumulation of ROS induces cell death [6]. Studies have shown that cancer cells have increased ROS level compared to normal cells due to high metabolic rate and mitochondrial dysfunction, which render increased susceptibility to oxidative stress [7, 8]. Thus, additional surge in ROS level is likely to cause cancer cells to reach their oxidative stress threshold sooner than normal cells, resulting in oxidative stress-induced cancer cell death [7, 8]. Therefore, it

is not surprising that several natural dietary bioactive compounds that cause increased ROS levels have been shown to selectively target cancer cells [9]. For instance, dietary phytochemicals such as polyphenols, flavonoids, and stilbenes have the capacity to inhibit cancer cell proliferation and induce apoptosis and autophagy [10]. While most dietary bioactive compounds possess antioxidant capacity at low doses, high doses induce prooxidant activity that leads to cancer cell death. These compounds also influence mitochondrial functions by altering mitochondrial enzymes, oxidative phosphorylation, and mitochondrial pathways [11]. In this review, we focus on ROS regulation, ROS-mediated signalling pathways, and the contemporary use of dietary phytochemicals for cancer therapy.

2. ROS Regulation

ROS production is affected by both external factors such as tobacco smoke and ionizing radiation and intracellular factors such as the endoplasmic reticulum (ER), mitochondria,

FIGURE 1: Intracellular redox homeostasis and imbalance and their effects on cellular functions. SOD: superoxide dismutase; CAT: catalase; OH: hydroxyl radical; GPX: glutathione peroxidase; GSSG: glutathione disulfide; GR: GSSG reductase; GSH: glutathione.

and peroxisomes [12] (Figure 2). Endogenous ROS are mainly produced in mitochondria during oxidative phosphorylation. Superoxide anions are generated through the electron transport chain complexes I and III localized in the inner mitochondrial membrane, and superoxide dismutase (SOD) converts superoxide ions into hydrogen peroxide (H_2O_2), which is subsequently catalyzed by glutathione peroxidase (GPX) to generate H_2O . Catalase (CAT) also converts H_2O_2 to water (Figure 1) [13]. Other intracellular enzymes such as NADPH oxidase, lipoxygenases, and xanthine oxidase are also capable of ROS production [14]. Although intracellular redox homeostasis is well controlled by the enzymatic antioxidants, SOD, GPX, and CAT, it is also regulated by nonenzymatic antioxidants such as ascorbic acid (vitamin C) and glutathione (GSH) [15] (Figure 2).

Besides these antioxidants, the transcription factor, nuclear factor erythroid 2- (NFE2-) related factor 2 (Nrf2), also contributes in controlling oxidative stress. Activation of Nrf2 requires inhibition of its negative regulator Keap1, which results in Nrf2 nuclear translocation [16]. This leads to the expression and production of the antioxidant enzymes, CAT, GPX, heme oxygenase-1 (HO-1), and peroxiredoxin (PRX), and maintenance of redox balance [16]. We note, however, that intracellular oxidative stress induces activation of hypoxia-inducible factors (HIFs), resulting in the transcription of genes that promote survival and proliferation of cancer cells [17].

3. ROS in Cancer Signalling Pathways

ROS serve a crucial role in the regulation of a number of cellular processes such as cell proliferation and differentiation and cell death. Therefore, it is critical that a delicate balance in ROS level is maintained. ROS level is regulated by redox homeostasis via ROS elimination through antioxidants. Within the threshold limit of redox homeostasis, a regulated ROS increase could serve as a signal for H_2O_2 -

mediated oxidation of protein cysteine residues, triggering specific cellular events such as proliferation [18]. Conversely, disturbance of redox homeostasis in the direction of ROS overload leads to deleterious outcomes such as irreversible oxidative DNA damage that could trigger cell death. It is now known that metabolically transformed and fast-growing cancer cells have higher ROS levels than neighboring normal cells, placing cancer cells at a greater risk of reaching the ROS threshold to induce apoptosis. This infers that promoting further ROS production in cancer cells may be utilized as a strategy to induce cancer cell death.

ROS play an important role in tumor initiation, promotion, and progression [19]. At levels below the ROS threshold, ROS activate oncogenes such as Ras and c-Myc [20] and induce p53-mediated DNA repair and survival [21] in cancer cells. At levels above the ROS threshold, ROS trigger apoptotic signals [6]. These cellular processes are controlled by ROS through its regulation of various signalling pathways (Figure 3), including the mitogen-activated protein kinase (MAPK)/extracellular-signal-regulated kinase (ERK), the phosphoinositide-3-kinase (PI3K)/protein kinase B (AKT), the inhibitor of kappa B ($\text{I}\kappa\text{B}$) kinase (IKK)/nuclear factor κB ($\text{NF}\kappa\text{B}$), and the protein kinase D (PKD) signalling pathways [22, 23]. For example, ROS-dependent ERK activation controls the expression of proapoptotic genes by phosphorylation of transcription factors [23, 24]. Conversely, ROS-induced JNK activation results in phosphorylation and downregulation of antiapoptotic proteins such as BCL-2 and BCL-X_L [25]. In response to ROS, $\text{I}\kappa\text{B}$ phosphorylation by IKK and subsequently ubiquitination lead to activation and translocation of $\text{NF}\kappa\text{B}$ into the nucleus to stimulate the expression of antiapoptotic genes [26]. ROS directly activates PI3K subsequently converting phosphatidylinositol 4,5-bisphosphate (PIP_2) to phosphatidylinositol 3,4,5-triphosphate (PIP_3) and resulting in transcriptional inhibition of the AKT target genes, glycogen synthase kinase 3 (GSK3), forkhead box O (FOXO), and BCL-2-associated death

FIGURE 2: Exogenous and endogenous sources of ROS and enzymatic and nonenzymatic antioxidants.

FIGURE 3: ROS-mediated intracellular cell signalling pathways. The indicated signalling pathways regulate molecules associated with angiogenesis, survival, proliferation, apoptosis, and invasion and the expression of antioxidant enzymes. NRF2: nuclear factor erythroid 2-related factor 2; KEAP1: Kelch-like ECH-associated protein 1; HIF1 α/β : hypoxia inducing factor 1 α/β ; HRE: HIF-responsive elements; p38 MAPK: p38 mitogen-activated protein kinase; ERK: extracellular signal-related kinases; MEK: MAPK kinase; JNK: c-Jun N-terminal kinase; PTEN: phosphatase and tensin homolog; PI3K: phosphoinositide-3-kinase; AKT: protein kinase B; IKK: I κ B kinase; NF κ B: nuclear factor kappa-light-chain-enhancer of activated B cells; FOXO: forkhead box protein O; mTOR1: mechanistic target of rapamycin 1; ATF2: activating transcription factor 2; CAT: catalase; HO-1: heme oxygenase-1; GPX: glutathione peroxidase; VEGF: vascular endothelial growth factor; TGF β ₃: transforming growth factor beta 3; NOS₂: nitric oxide synthase 2; BCL-2: B-cell lymphoma 2; BCL-X_L: B-cell lymphoma-extra large; BAD: BCL2-associated agonist of cell death; TRAIL: TNF-related apoptosis-inducing ligand; MMP: matrix metalloproteinase; ICAM: intercellular adhesion molecule-1.

promoter (BAD) and activation of mammalian target of rapamycin (mTOR1) [27].

ROS-mediated apoptosis can be initiated by mitochondrial intrinsic apoptotic signalling or by extrinsic apoptotic signalling through death receptor pathways (Figure 4). Increased production of ROS depolarizes the mitochondrial membrane, releasing cytochrome C from the mitochondria. Cytochrome C induces activation of caspase-9 by promoting nucleotide binding to apoptotic protein-activating factor 1 (APAF-1), which leads to activation of caspase-3 [28].

Antiapoptotic (BCL-2 and BCL-X_L) and proapoptotic (BAD, BAK, BAX, BID, and BIM) proteins also contribute to the formation of distinct channels for mitochondrial membrane permeabilization [29]. Elevated ROS levels have also been implicated in the activation of death receptors and in triggering caspase 8-mediated cleavage of caspase 3 [6]. In addition, ROS modulates the TRAIL- and Fas-mediated apoptosis through p53-mediated upregulation of death receptors. p53 regulates such apoptosis by controlling the expression of anti- and proapoptotic (e.g., PUMA and

FIGURE 4: ROS-mediated extrinsic and intrinsic apoptotic pathways. TRAIL: TNF-related apoptosis-inducing ligand; FADD: Fas-associated death domain; $[Ca^{2+}]_i$: intracellular calcium concentration; mPTP: mitochondrial permeability transition pore; $\Delta\Psi_m$: mitochondrial membrane potential.

NOXA) proteins [30, 31]. ROS further promotes apoptosis by inducing increased Ca^{2+} -mediated mitochondrial permeability transition pore opening [32].

4. Dietary Polyphenols

There is increasing claim that certain natural bioactive compounds can maintain redox homeostasis and hold promise as anticancer therapeutics due to their biocompatibility, biodegradability, comparatively less toxicity, and reduced side effects. The polyphenol bioactive compounds are secondary metabolites found in plants [33]. The most abundantly occurring plant polyphenols are phenolic acids and flavonoids which account for 30% and 60%, respectively, of dietary polyphenols [33]. Interestingly, they have both antioxidant and prooxidant properties that modulate cell proliferation and apoptotic pathways [34]. Some of the most common bioactive compounds that were suggested to have cancer therapeutic effects through their ROS-related activities are discussed below.

4.1. Quercetin. Quercetin (3,5,7,3',4'-pentahydroxyflavone) is a flavonoid, present in numerous vegetables and fruits [34, 35]. Quercetin (Qu) displays neuroprotective, chemopreventive, and anticancer activities [36, 37], and these have been attributed to their anti- and prooxidative capacities. Qu efficiently scavenges mitochondrial superoxide anions (O_2^-) and subsequently generates semiquinone, Qu radicals, and H_2O_2 [11, 34, 38]. While, H_2O_2 is eliminated by peroxidase, semiquinone and Qu radicals alter intracellular ROS metabolism by depleting the intracellular GSH pool in a concentration-dependent manner [39–41] and inhibiting thioredoxin reductase activity [42]. *In vitro* and *in vivo* studies (Table 1) show that Qu promotes ROS-induced apoptosis, necrosis, and autophagy [43] at a range of 10–100 μM in a

variety of cancers, including glioma [43], osteosarcoma [44], and cervical [45] and breast cancer [46]. Qu induces apoptosis through distinct mechanisms: (i) via the mitochondrial pathway through activation of caspase-3. Qu reduces the mitochondrial membrane potential (MMP), inducing cytochrome C release and subsequent activation of caspase-3. This mechanism was observed in MDA MB-231 breast cancer cells [47], U937 promonocytic leukemia cells [48], HL-60 promyelocytic leukemia cells [49], HepG2 hepatocellular carcinoma cells [50], and oral cancer cells [51]. (ii) Qu alters the expression of the antiapoptotic BCL-2 and BCL-X_L and proapoptotic BAX and BAD proteins [47, 48]. Leukemic cells treated with Qu showed upregulation of BAX and increased phosphorylation of BCL-2 [52]. Similar results were observed in osteosarcoma [44] and breast cancer cells [46]. (iii) Qu induces the expression of death receptor-(DR-) 5, enhancing TNF-related apoptosis-inducing ligand-(TRAIL-) induced apoptosis [53–55] either by accumulating death receptors in lipid rafts [56] or inhibiting survivin in the ERK signalling pathway [57]. In addition to its proapoptotic capacity, Qu also promotes cell cycle arrest [58] by modulating p21^{WAF1}, cyclin B, and p27^{KIP1} in squamous cell carcinoma [59] and breast [60], lung [61], and hepatoma cancer cells [62].

4.2. Curcumin. Curcumin (1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione) is the principal polyphenol derived from turmeric (*Curcuma longa*). Various pharmacological activities have been attributed to curcumin, including its anti-inflammatory and anticarcinogenic properties which are triggered at 25 μM [63]. Its anticancer effect is currently being evaluated in clinical trials for a variety of cancers [64–66] (Table 2). In normal cells, curcumin acts as a potent antioxidant. It scavenges hydroxyl radicals, superoxide, nitric oxide, H_2O_2 , and peroxynitrite [11, 67–69] and modulates the expression of SOD, HO-1, and GPX through an indirect mechanism [11, 70–72]. In contrast, curcumin's anticancer properties rely on its prooxidative capacity to induce apoptosis, likely via the mitochondria-mediated pathway [73–75]. Curcumin oxidizes thiols in the mitochondrial membrane, leading to mitochondrial permeability transition pore (mPTP) opening, mitochondrial swelling, mitochondrial depolarization, and inhibition of ATP synthesis, resulting in apoptosis [76]. Evidence shows that curcumin increases ROS levels, including superoxides, hydroxyl radicals, and H_2O_2 [77–79]. Indeed, in human hepatoma cells, curcumin causes cell death by ROS-induced mitochondrial DNA damage and impairment of OXPHOS [80, 81]. Curcumin also activates TRAIL-induced apoptosis by ROS-mediated upregulation of DR5 in renal cancer cells and colon cancer cells [82, 83]. Curcumin further induces autophagy in colon cancer cells through ROS-dependent activation of the ERK1/2 and the p38 MAPK pathway [84]. In glioblastoma [85] and liver cancer [86], curcumin decreases cancer stem cell viability and proliferation by ROS-mediated inhibition of NF κ B and signal transducer and activator of transcription 3 (STAT3). As with Qu, curcumin promotes cancer cell apoptosis by upregulating proapoptotic proteins (BAX, BIM, BAK, and NOXA) [87, 88] and downregulating antiapoptotic

TABLE 1: *In vivo* dosages and mechanistic effects of known natural bioactive compounds.

Compound	Animals	Cancer model	Dose	Mechanism
Quercetin	Male Sprague-Dawley rats	Glioma	100 mg/kg, every other day for 15 days, i.v	Autophagy and apoptosis [43]
	Female BALB/c mice	Colon & breast cancer	100 and 200 mg/kg for 36 days, i.p	Apoptosis [178]
	Male BALB/cA nude mice	Prostate cancer	20 mg/kg for 16 days, i.p	Antiangiogenesis [179]
	Female BALB/c/nude mice	Hepatic cancer	10 mg/kg for 7 days, i.p	Necrosis and antiproliferation [180]
Curcumin	Female NOD.CB17-Prkdcscid/J lineage	Acute myelogenous leukemia	120 mg/kg, once every 4 days for 21 days, i.p	Apoptosis, autophagy, and cell cycle arrest [181]
	Female BALB/c/nude mice	Colon cancer (multidrug resistance)	50 mg/kg, 2x/day for 14 days, peritumoral	Reduced expression of MDR1 and survivin [182]
	Male BALB/c/nude mice	Prostate cancer	25, 50, and 100 mg/kg, every 2 days for 30 days, abdominal cavity injection	Apoptosis [89]
	Female athymic nude mice	Breast cancer	45 mg/kg, 2x/week for 4 consecutive weeks, i.p	Antiproliferation [183]
Capsaicin	Female athymic nude mice	Pancreatic cancer	2.5 & 5 mg/kg, 5x/week, gavage	Activation of JNK and apoptosis [99]
	Female BALB/c nude mice	Colon cancer	1 & 3 mg/kg, 3 days once for 40 days, i.p	Apoptosis [95]
	Male BNX nu/nu mice	Prostate cancer	5 mg/kg, 3x/week for 4 weeks, gavage	Antiproliferation and apoptosis [184]
	Female BNX nu/nu	Breast cancer	5 mg/kg, 3x/week for 4 weeks, gavage	Reduced EGFR/HER2 activation and apoptosis [185]
ECGC	Female C3H/HeJ syngeneic mice	Squamous cell carcinoma	50 mg/kg, 5 days/week, i.p	Apoptosis [186]
	NOD/SCID mice	Myeloid leukemia	10 mM, oral drinking fluid	Antiproliferation [110]
	Female BALB/c mice	Bladder cancer	100 mg/kg for 4 weeks, i.p	Antiproliferation and migration [187]
	Male BALB/c/nude mice	Lung cancer	0.05% in drinking water for 21 days	Angiogenesis [188]
PEITC	Male BALB/c/nude mice	Adrenal pheochromocytoma	15 mg/kg, every other day for 15 days, i.p	Apoptosis [189]
	Male athymic nude mice	Glioblastoma	20 μ mol/100 μ l PBS for 21 days, gavage	Apoptosis [190]
	Male athymic mice	Prostate cancer	12 μ mol/100 μ l PBS for 5 days, oral	Apoptosis [191]
	Female BALB/c/nude mice	Lung cancer	25 mg/kg, 3x/week, i.p	Antiproliferation, reduced cancer stem cells [128]
Piperine	Female SCID/NOD mice	Breast cancer	81 mg/kg for 35 days, oral gavage	Apoptosis [192]
	Female athymic nude mice	Ovarian cancer	12 μ mol for 42 days, oral gavage	EGFR-AKT pathway inhibition, antiproliferation, and apoptosis [193]
	Female BALB/c mice	Mouse 4T1 mammary carcinoma	2.5 and 5 mg/kg, every 3 days for 3 times, intratumoral	Cell cycle arrest and apoptosis [194]
	Male nude mice	Prostate cancer	100 mg/kg/day for 1 month, i.p 10 mg/kg for 1 month, gavage	Antiproliferation and apoptosis [195]
Piperine	Male albino Wistar rats	Hepatocellular carcinoma (diethylnitrosamine-induced)	5 mg/kg, 3x/week for 6 weeks, oral	Apoptosis [138]

TABLE 1: Continued.

Compound	Animals	Cancer model	Dose	Mechanism
Resveratrol	Male nude mice	Lung cancer	20 mg/kg, every other day for 25 days, i.p	Reduce metastasis [196]
	Male BALB/c/nude mice	Bladder cancer	20 mg/kg/day for 4 weeks, i.p	Decreased VEGF and FGF-2 level, cell cycle arrest, and apoptosis [197]
	Female athymic mice	Breast cancer	25 mg/kg/day for 3 weeks, i.p	Apoptosis [198]
	BALB/c/nude mice	Pancreatic cancer	20, 40, and 60 mg/kg, 5 days/week for 6 weeks, gavage	Inhibition of FOXO transcription factors and apoptosis [199]
	Male athymic nude mice	Prostate cancer	50 mg/kg, every other day for 2 weeks, gavage	Antiproliferation [200]

i.p: intraperitoneal; i.v: intravenous.

TABLE 2: Clinical trials of natural phytochemicals.

Bioactive compounds (Clinicaltrials.gov identifier)	Disease condition	Phase	Dosage	Study goal
<i>Quercetin</i> (NCT03476330)	Squamous cell carcinoma	II	4 g/day	Efficacy in reducing buccal micronuclei in patients with Fanconi anemia
<i>Curcumin</i> (NCT03769766)	Prostate cancer	III	500 mg, 2x/day	Effect on prostate cancer progression
(NCT00094445)	Pancreatic cancer	II	8 g/day	Effect in pancreatic cancer growth and the safety of treatment
(NCT01246973)	Radiation dermatitis	III	500 mg, 3x/day	Effect on dermatitis caused by radiation therapy in breast cancer patients
With piperine (NCT02598726)	Neoplasms	I	A dose escalation study	Optimal biological dose in cancer patients
<i>Capsaicin</i> (NCT02037464)	Prostate cancer	II	2 capsules/day for 6 months	Expression of Ki67 and p27 in a posttreatment biopsy
(NCT00003610)	Head & neck cancer, mucositis	III	4 lozenges/day up to 2 weeks after radiation therapy	Efficacy of lozenges in patients with mucositis caused by radiation therapy
Patch (Qutenza) (NCT03317613)	Cancer	II	Qutenza (8% capsaicin patch) for every 3 months	Efficacy in peripheral neuropathic pain in cancer patients
<i>EGCG</i> (NCT02891538)	Colon cancer	Early I	450 mg, 2x/day	Chemopreventive effects
(NCT01317953)	Small cell lung carcinoma	I	2 × 450 mg/day to 5 × 450 mg/day	Side effects and best dose
<i>PEITC</i> (NCT00691132)	Lung cancer	II	4x/day for 5 days in week 4	Effect in preventing lung cancer in smokers
(NCT01790204)	Oral cancer	I & II		Effect on oral cells with mutant p53
Nutri-PEITC jelly (NCT03034603)	Head & neck neoplasms		200 mg/day, 5 days/week for 3 months	Safety and efficacy
<i>Resveratrol</i> (NCT00256334)	Colon cancer	I	20 mg/day	Modulation of Wnt signalling <i>in vivo</i>
(NCT01476592)	Neuroendocrine tumor		5 g/day	Effect on Notch-1 signalling
<i>SRT501</i> (NCT00920803)	Colorectal cancer	I	5 g/day	Safety and tolerability

proteins (BCL-2 and BCL-X_L) [89, 90]. In addition, curcumin can impede tumor angiogenesis by downregulating the expression of the vascular endothelial growth factor (VEGF) and matrix metalloproteinases (MMPs) [91, 92].

4.3. Capsaicin. Capsaicin (trans-8-methyl-N-vanillyl-6-nonenamide), the major component of *Capsicum* [93], has been implicated to have anticarcinogenic properties [94–96]. However, the mechanisms by which capsaicin induces cancer cell death are still unclear. The proposed anticancer mechanisms of capsaicin include promotion of ROS accumulation, mitochondria-mediated apoptosis, cell cycle arrest, and impairment of endoplasmic reticulum (ER) calcium homeostasis [97]. Capsaicin induces a rapid rise of ROS level followed by a disruption of mitochondrial membrane potential and subsequent activation of downstream caspase-3 in human colon cancer [98], pancreatic cancer [99], glioma [100], and prostate cancer [101]. In transformed T-cells, capsaicin inhibits the plasma membrane NADH-oxidoreductase

(PMOR) electron transport chain, causing an increase in ROS level and subsequent disruption of the mitochondrial membrane potential [102]. Capsaicin at 150 μ M also blocks complexes I and III of the respiratory chain and decreases SOD activity in pancreatic cancer [103]. Interestingly, binding of capsaicin to the transient receptor potential vanilloid type 1 (TRPV1) results in an increase in intracellular calcium level and activation of the apoptotic pathway [104–106]. Besides its proapoptotic effects, capsaicin can also induce cell cycle arrest through inhibition of the cyclin-dependent kinases, Cdk2, Cdk4, and Cdk6 [107, 108].

4.4. Epigallocatechin-3-Gallate (EGCG). Epigallocatechin-3-gallate ((2R,3R)-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)-3,4-dihydro-2H-1-benzopyran-3-yl 3,4,5-trihydroxybenzoate) is a prominent catechin polyphenol in green tea. EGCG has dual antioxidant and prooxidant roles. It produces ROS by autooxidation [109] and its ability to modulate ROS level accounts for its chemopreventive property. EGCG induces

apoptosis in various cancer cell types, including myeloid leukemia cells [110], human lymphoblastoid B cells [111], and hepatocarcinoma cells [112]. In pancreatic carcinoma [113] and lung cancer cells [114], EGCG-induced apoptosis occurs through inhibition of the PI3K/AKT signalling pathway. EGCG also decreases the mitochondrial membrane potential, increasing the intracellular free Ca^{2+} level and causing activation of the intrinsic apoptotic pathway. EGCG further decreases the expression of the antiapoptotic BCL-2, BCL-X_L, xIAP, and cIAP and increases the expression of the proapoptotic BAD, BAX, and FAS/CD95 [115]. In pancreatic [116] and bladder cancer cells [117], EGCG also induces G₀/G₁ cell cycle arrest through regulation of cyclin D1, Cdk4, Cdk6, p21^{WAF1}, and p27^{KIP1} via the ERK, IKK, and PI3K signalling pathways. A combination of EGCG (10 μM) and curcumin (10 μM) inhibits breast cancer stem cell growth by inactivating the NF κ B-STAT3 pathway [118].

4.5. PEITC and BITC. Phenethyl isothiocyanate (PEITC) and benzyl isothiocyanate (BITC) are abundant in cruciferous vegetables that have been implicated to have anticancer properties [119–122]. Epidemiological studies show that increased intake of dietary isothiocyanates (ITC) reduces cancer risk [123] and increases cancer patient survival [124]. Both PEITC and BITC induce ROS production in many cancer cells [125–127]. IC₅₀ value of PEITC is at the range of 3–14 μM in various human cancer cells [128]. PEITC increases ROS level by decreasing intracellular GSH level, leading to mitochondrial dysfunction as observed in ovarian [126, 129] and non-small-cell lung cancer [128] cells but not in normal cells. PEITC-induced ROS production correlates with inhibition of complex III activity, inhibition of OXPHOS, and ATP depletion in prostate cancer [125]. PEITC also inhibits HO-1 and subsequently induces the ROS-mediated mitochondrial apoptotic pathway, which was noted in human chronic myeloid leukemia [130]. Conversely, BITC causes oxidative stress in pancreatic [131], glioma [122], and prostate cancer [132] cells by depleting SOD and GSH, which is accompanied by the induction of caspase-mediated apoptosis [121, 133]. BITC also activates the ERK/JNK/p38MAPK pathway in pancreatic cancer [134]. Both PEITC and BITC induce G₂/M cell cycle arrest by downregulating cyclin B1, Cdc2, and Cdc25C [135, 136].

4.6. Piperine. Piperine ([5-(1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]piperidine) is the most abundant natural alkaloid found in long pepper (*Piper longum* L.). Recently, it was determined to be a promising anticancer compound [137]. Piperine suppresses tumor growth in vitro and in vivo by modulating the ROS-induced oxidative stress response pathway, cell cycle arrest, and ER stress. In hepatocellular carcinoma, piperine treatment initiates ROS-induced mitochondria-mediated apoptosis by inhibiting catalase activity [138]. In human oral squamous cells exposed to high concentrations of piperine, ROS elevation is associated with mitochondrial depolarization and activation of caspase-mediated apoptosis. Piperine also induces nuclear condensation and cell cycle arrest in these cells [139].

4.7. Resveratrol. Resveratrol (3,4',5-trihydroxystilbene), a polyphenol that is found in grapes and berries, effectively prevents tumor initiation and progression by stimulating apoptosis at 10 to 100 μM [140] in prostate [141] and neuroblastoma cells [142]. Resveratrol has been shown to promote apoptosis by activating p53, ROS-dependent caspases, and death receptors for TRAIL and FasL [143]. Resveratrol-mediated apoptosis is mainly associated with the inhibition of the PI3K/AKT, MAPK, and NF κ B pathways [144] and STAT3 [145]. Moreover, resveratrol suppresses the expression of antiapoptotic proteins such as survivin, xIAP, and BCL-X_L and increases BAX/caspase-3-associated apoptosis [146]. Resveratrol further binds to F₁-ATPase, inhibiting mitochondrial ATP synthesis [147, 148]. It triggers cell cycle arrest by upregulating p21^{WAF1} and p27^{KIP1} and downregulating cyclins D1, D2, and E and Cdks 2, 4, and 6 [149, 150].

4.8. Others. Peanuts, tomatoes, and carrots are rich in *p-Coumaric acid* (*p-CoA*), an isomer of cinnamic acid [151]. In colon cancer cells, *p-CoA* triggers apoptosis by increasing ROS generation and mitochondrial depolarization, resulting in p53-mediated upregulation of BAX and downregulation of BCL-2 [151, 152]. In addition, *p-CoA* treatment of these cells in vitro and in vivo induces apoptosis mediated by the unfolded protein response [153].

The naturally occurring quinone compounds have potent cytotoxicity against cancer cells. In lung adenocarcinoma cells, *2-methoxy-1,4-naphthoquinone* (MNQ) and *8-hydroxy-2-methoxy-1,4-naphthoquinone* (HMNQ) elicit ROS production and induce apoptosis via the JNK/p38 MAPK pathway [154–156].

Naringenin, a citrus flavonoid, triggers ROS-induced apoptosis and stimulates p38MAPK-mediated caspase activation [157, 158].

Gallic acid (3,4,5-trihydroxy-benzoic acid; GA), which is widely present in grapes and red wine, inhibits lung cancer cell growth by increasing ROS level and depleting GSH [159]. In prostate cancer cells, autooxidation of GA produces H₂O₂ and O₂^{•−}, leading to mitochondria-dependent apoptosis [160]. GA also induces apoptosis via ROS-dependent activation of the ATM/p53 [161] and JNK pathways [162].

5. Limitations

Poor bioavailability is a major obstacle for natural bioactive compounds, especially for Qu, curcumin, and resveratrol, which are associated with poor absorption and fast metabolism in the liver and intestine. Pharmacokinetic profile analysis of Qu revealed that about 93% of the compound is metabolised after oral administration (10 mg/kg) in male Sprague-Dawley rats [163]. On the other hand, people taking high oral doses (10 or 12 g) of curcumin attained limited availability of this compound in the plasma and other tissues [164]. Similarly, oral bioavailability of resveratrol is low at less than 1% [165]. Thus, the cytotoxic concentration of these compounds appears to be difficult to achieve by oral administration in cancer patients [166]. Several strategies have been proposed to overcome the problem of low oral bioavailability. One approach is to use a combination of phytochemicals.

For example, a combination of piperine and curcumin [167] (in rats: 20 mg/kg piperine + 2 g/kg curcumin; in humans: 20 mg piperine + 2 g curcumin) or piperine and resveratrol [168] (in mice: 10 mg/kg piperine + 100 mg/kg resveratrol) showed increased bioavailability of curcumin and resveratrol, respectively. Other promising approaches include the use of novel formulations, synthetic analogues, prodrugs, and different drug delivery systems (e.g., via liposomes, phospholipid complexes, micelles, and nanoparticles). These methods could increase bioavailability as well as solubility and/or metabolic stability [169, 170]. Some studies have also shown that natural bioactive compounds may promote carcinogenesis by inducing ROS-mediated chromosome aberrations and DNA damage [80, 171, 172]. For example, an *in vivo* study showed that curcumin promotes lung cancer [173] and topical application of capsaicin causes skin cancer in mice [174], suggesting that these natural compounds must be carefully assessed for safety prior to clinical application.

As dietary phytochemicals lack mechanistic selectivity, these natural compounds display a variety of effects in different cancer cell types and thus the discrepancies in results among separate studies. Other possible reasons for divergent findings in different studies include changes or differences in (i) stability of the bioactive compounds in cell culture medium, for example, stability of Qu decreases at pH 7 or 8 [175]; (ii) release of bioactive compounds under different conditions, for example, the maximum release of curcumin occurs in phosphate buffered saline at pH 6.4 [176]; (iii) sensitivity of different cell types to bioactive compounds; (iv) cellular permeability of bioactive compounds; (v) presence or contamination by metal ions [177]; (vi) number of hydroxyl groups present in a molecule [177]; and (vii) *in vivo* biodistribution.

6. Conclusion

Natural phytochemicals have been associated with anticancer properties through their ability to modulate oxidative stress, cell cycle regulators, and proapoptotic, antiapoptotic, and survival signalling pathways. In preclinical and clinical trials, bioactive compounds show a promising and wide therapeutic window against various malignancies, including glioblastoma and breast, colon, and prostate cancers where phytochemical-induced cancer cell death was observed. However, certain attributes such as poor solubility and bioavailability of these bioactive compounds limit their clinical application. Thus, further studies are required to identify ways for effective biological delivery of these compounds in different cancer cell types. It is also critical that detailed studies are conducted in large cohorts to establish the pharmacokinetic profile of these compounds alone and in combination with other chemotherapeutic agents to determine dosage, tissue targets, and toxicity. Indeed, natural phytochemicals may serve as future therapy for specific types of cancer.

Conflicts of Interest

The authors declare no conflict of interest.

Authors' Contributions

SN wrote the draft. JR and KYL revised the manuscript.

Acknowledgments

This work was supported in part by grants from the CIHR (MOP-123400) and NSERC (RGPIN/312985-2011) to KYL and an Alberta Cancer Foundation graduate studentship to SN.

References

- [1] L. A. Sena and N. S. Chandel, "Physiological roles of mitochondrial reactive oxygen species," *Molecular Cell*, vol. 48, no. 2, pp. 158–167, 2012.
- [2] L. Covarrubias, D. Hernandez-Garcia, D. Schnabel, E. Salas-Vidal, and S. Castro-Obregon, "Function of reactive oxygen species during animal development: passive or active?," *Developmental Biology*, vol. 320, no. 1, pp. 1–11, 2008.
- [3] K. R. Martin and J. C. Barrett, "Reactive oxygen species as double-edged swords in cellular processes: low-dose cell signaling versus high-dose toxicity," *Human & Experimental Toxicology*, vol. 21, no. 2, pp. 71–75, 2002.
- [4] N. Bellance, P. Lestienne, and R. Rossignol, "Mitochondria: from bioenergetics to the metabolic regulation of carcinogenesis," *Frontiers in Bioscience*, vol. 14, pp. 4015–4034, 2009.
- [5] L. B. Sullivan and N. S. Chandel, "Mitochondrial reactive oxygen species and cancer," *Cancer & Metabolism*, vol. 2, no. 1, 2014.
- [6] M. Redza-Dutordoir and D. A. Averill-Bates, "Activation of apoptosis signalling pathways by reactive oxygen species," *Biochimica et Biophysica Acta*, vol. 1863, no. 12, pp. 2977–2992, 2016.
- [7] D. Trachootham, J. Alexandre, and P. Huang, "Targeting cancer cells by ROS-mediated mechanisms: a radical therapeutic approach?," *Nature Reviews Drug Discovery*, vol. 8, no. 7, pp. 579–591, 2009.
- [8] S. NavaneethaKrishnan, J. L. Rosales, and K. Y. Lee, "Targeting Cdk5 for killing of breast cancer cells via perturbation of redox homeostasis," *Oncoscience*, vol. 5, no. 5-6, pp. 152–154, 2018.
- [9] A. M. Mileo and S. Miccadei, "Polyphenols as modulator of oxidative stress in cancer disease: new therapeutic strategies," *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 6475624, 17 pages, 2016.
- [10] M. Gonzalez-Vallinas, M. Gonzalez-Castejon, A. Rodriguez-Casado, and A. Ramirez de Molina, "Dietary phytochemicals in cancer prevention and therapy: a complementary approach with promising perspectives," *Nutrition Reviews*, vol. 71, no. 9, pp. 585–599, 2013.
- [11] L. Gibellini, E. Bianchini, S. De Biasi, M. Nasi, A. Cossarizza, and M. Pinti, "Natural compounds modulating mitochondrial functions," *Evidence-Based Complementary and Alternative Medicine*, vol. 2015, Article ID 527209, 13 pages, 2015.
- [12] A. Phaniendra, D. B. Jestadi, and L. Periyasamy, "Free radicals: properties, sources, targets, and their implication in various diseases," *Indian Journal of Clinical Biochemistry*, vol. 30, no. 1, pp. 11–26, 2015.

- [13] S. S. Sabharwal and P. T. Schumacker, "Mitochondrial ROS in cancer: initiators, amplifiers or an Achilles' heel?," *Nature Reviews Cancer*, vol. 14, no. 11, pp. 709–721, 2014.
- [14] D. I. Brown and K. K. Griendling, "Regulation of signal transduction by reactive oxygen species in the cardiovascular system," *Circulation Research*, vol. 116, no. 3, pp. 531–549, 2015.
- [15] I. Mironczuk-Chodakowska, A. M. Witkowska, and M. E. Zujko, "Endogenous non-enzymatic antioxidants in the human body," *Advances in Medical Sciences*, vol. 63, no. 1, pp. 68–78, 2018.
- [16] B. D'Autreaux and M. B. Toledano, "ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis," *Nature Reviews Molecular Cell Biology*, vol. 8, no. 10, pp. 813–824, 2007.
- [17] A. Galanis, A. Pappa, A. Giannakakis, E. Lanitis, D. Dangaj, and R. Sandaltzopoulos, "Reactive oxygen species and HIF-1 signalling in cancer," *Cancer Letters*, vol. 266, no. 1, pp. 12–20, 2008.
- [18] M. Schieber and N. S. Chandel, "ROS function in redox signaling and oxidative stress," *Current Biology*, vol. 24, no. 10, pp. R453–R462, 2014.
- [19] G. Y. Liou and P. Storz, "Reactive oxygen species in cancer," *Free Radical Research*, vol. 44, no. 5, pp. 479–496, 2010.
- [20] S. Kumari, A. K. Badana, M. M. G. S. G., and R. R. Malla, "Reactive oxygen species: a key constituent in cancer survival," *Biomarker Insights*, vol. 13, p. 117727191875539, 2018.
- [21] B. Liu, Y. Chen, and D. K. St. Clair, "ROS and p53: a versatile partnership," *Free Radical Biology & Medicine*, vol. 44, no. 8, pp. 1529–1535, 2008.
- [22] J. N. Moloney and T. G. Cotter, "ROS signalling in the biology of cancer," *Seminars in Cell & Developmental Biology*, vol. 80, pp. 50–64, 2018.
- [23] J. Zhang, X. Wang, V. Vikash et al., "ROS and ROS-mediated cellular signaling," *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 4350965, 18 pages, 2016.
- [24] Z. Lu and S. Xu, "ERK1/2 MAP kinases in cell survival and apoptosis," *IUBMB Life*, vol. 58, no. 11, pp. 621–631, 2006.
- [25] D. N. Dhanasekaran and E. P. Reddy, "JNK signaling in apoptosis," *Oncogene*, vol. 27, no. 48, pp. 6245–6251, 2008.
- [26] S. Mincheva-Tasheva and R. M. Soler, "NF- κ B Signaling Pathways," *The Neuroscientist*, vol. 19, pp. 175–194, 2013.
- [27] B. D. Manning and L. C. Cantley, "AKT/PKB signaling: navigating downstream," *Cell*, vol. 129, no. 7, pp. 1261–1274, 2007.
- [28] S. NavaneethaKrishnan, J. L. Rosales, and K. Y. Lee, "Loss of Cdk5 in breast cancer cells promotes ROS-mediated cell death through dysregulation of the mitochondrial permeability transition pore," *Oncogene*, vol. 37, no. 13, pp. 1788–1804, 2018.
- [29] R. J. Youle and A. Strasser, "The BCL-2 protein family: opposing activities that mediate cell death," *Nature reviews Molecular Cell Biology*, vol. 9, no. 1, pp. 47–59, 2008.
- [30] D. H. Lee, J. G. Rhee, and Y. J. Lee, "Reactive oxygen species up-regulate p53 and Puma; a possible mechanism for apoptosis during combined treatment with TRAIL and wogonin," *British Journal of Pharmacology*, vol. 157, no. 7, pp. 1189–1202, 2009.
- [31] K. Yoshida and Y. Miki, "The cell death machinery governed by the p53 tumor suppressor in response to DNA damage," *Cancer Science*, vol. 101, no. 4, pp. 831–835, 2010.
- [32] N. Hempel and M. Trebak, "Crosstalk between calcium and reactive oxygen species signaling in cancer," *Cell Calcium*, vol. 63, pp. 70–96, 2017.
- [33] S. Ramos, "Effects of dietary flavonoids on apoptotic pathways related to cancer chemoprevention," *The Journal of Nutritional Biochemistry*, vol. 18, no. 7, pp. 427–442, 2007.
- [34] L. Gibellini, M. Pinti, M. Nasi et al., "Interfering with ROS metabolism in cancer cells: the potential role of quercetin," *Cancers*, vol. 2, no. 2, pp. 1288–1311, 2010.
- [35] C. H. Jeong and S. H. Joo, "Downregulation of reactive oxygen species in apoptosis," *Journal of Cancer Prevention*, vol. 21, no. 1, pp. 13–20, 2016.
- [36] F. Dajas, "Life or death: neuroprotective and anticancer effects of quercetin," *Journal of Ethnopharmacology*, vol. 143, no. 2, pp. 383–396, 2012.
- [37] C. Chen, J. Zhou, and C. Ji, "Quercetin: a potential drug to reverse multidrug resistance," *Life Sciences*, vol. 87, no. 11–12, pp. 333–338, 2010.
- [38] D. Metodiewa, A. K. Jaiswal, N. Cenas, E. Dickancaite, and J. Segura-Aguilar, "Quercetin may act as a cytotoxic prooxidant after its metabolic activation to semiquinone and quinoidal product," *Free Radical Biology & Medicine*, vol. 26, no. 1–2, pp. 107–116, 1999.
- [39] R. Kachadourian and B. J. Day, "Flavonoid-induced glutathione depletion: potential implications for cancer treatment," *Free radical Biology & Medicine*, vol. 41, no. 1, pp. 65–76, 2006.
- [40] R. Ferraresi, L. Troiano, E. Roat et al., "Essential requirement of reduced glutathione (GSH) for the anti-oxidant effect of the flavonoid quercetin," *Free Radical Research*, vol. 39, no. 11, pp. 1249–1258, 2005.
- [41] A. M. Ramos and P. Aller, "Quercetin decreases intracellular GSH content and potentiates the apoptotic action of the anti-leukemic drug arsenic trioxide in human leukemia cell lines," *Biochemical Pharmacology*, vol. 75, no. 10, pp. 1912–1923, 2008.
- [42] J. Lu, L. V. Papp, J. Fang, S. Rodriguez-Nieto, B. Zhivotovsky, and A. Holmgren, "Inhibition of mammalian thioredoxin reductase by some flavonoids: implications for myricetin and quercetin anticancer activity," *Cancer Research*, vol. 66, no. 8, pp. 4410–4418, 2006.
- [43] Y. Bi, C. Shen, C. Li et al., "Inhibition of autophagy induced by quercetin at a late stage enhances cytotoxic effects on glioma cells," *Tumor Biology*, vol. 37, no. 3, pp. 3549–3560, 2016.
- [44] X. Xie, J. Yin, Q. Jia et al., "Quercetin induces apoptosis in the methotrexate-resistant osteosarcoma cell line U2-OS/MTX300 via mitochondrial dysfunction and dephosphorylation of Akt," *Oncology Reports*, vol. 26, no. 3, pp. 687–693, 2011.
- [45] R. Vidya Priyadarsini, R. Senthil Murugan, S. Maitreyi, K. Ramalingam, D. Karunakaran, and S. Nagini, "The flavonoid quercetin induces cell cycle arrest and mitochondria-mediated apoptosis in human cervical cancer (HeLa) cells through p53 induction and NF- κ B inhibition," *European Journal of Pharmacology*, vol. 649, no. 1–3, pp. 84–91, 2010.
- [46] E. J. Choi, S. M. Bae, and W. S. Ahn, "Antiproliferative effects of quercetin through cell cycle arrest and apoptosis in human breast cancer MDA-MB-453 cells," *Archives of Pharmacological Research*, vol. 31, no. 10, pp. 1281–1285, 2008.

- [47] S. Y. Chien, Y. C. Wu, J. G. Chung et al., "Quercetin-induced apoptosis acts through mitochondrial- and caspase-3-dependent pathways in human breast cancer MDA-MB-231 cells," *Human & Experimental Toxicology*, vol. 28, no. 8, pp. 493–503, 2009.
- [48] E. Lugli, L. Troiano, R. Ferraresi et al., "Characterization of cells with different mitochondrial membrane potential during apoptosis," *Cytometry Part A*, vol. 68, no. 1, pp. 28–35, 2005.
- [49] I. K. Wang, S. Y. Lin-Shiau, and J. K. Lin, "Induction of apoptosis by apigenin and related flavonoids through cytochrome c release and activation of caspase-9 and caspase-3 in leukaemia HL-60 cells," *European journal of cancer*, vol. 35, no. 10, pp. 1517–1525, 1999.
- [50] A. B. Granado-Serrano, M. A. Martin, L. Bravo, L. Goya, and S. Ramos, "Quercetin induces apoptosis via caspase activation, regulation of Bcl-2, and inhibition of PI-3-kinase/Akt and ERK pathways in a human hepatoma cell line (HepG2)," *The Journal of Nutrition*, vol. 136, no. 11, pp. 2715–2721, 2006.
- [51] J. W. Kang, J. H. Kim, K. Song, S. H. Kim, J. H. Yoon, and K. S. Kim, "Kaempferol and quercetin, components of Ginkgo biloba extract (EGb 761), induce caspase-3-dependent apoptosis in oral cavity cancer cells," *Phytotherapy Research*, vol. 24, Suppl 1, pp. S77–S82, 2010.
- [52] J. Duraj, K. Zazrivcova, J. Bodo, M. Sulikova, and J. Sedlak, "Flavonoid quercetin, but not apigenin or luteolin, induced apoptosis in human myeloid leukemia cells and their resistant variants," *Neoplasma*, vol. 52, no. 4, pp. 273–279, 2005.
- [53] Y. H. Jung, J. Heo, Y. J. Lee, T. K. Kwon, and Y. H. Kim, "Quercetin enhances TRAIL-induced apoptosis in prostate cancer cells via increased protein stability of death receptor 5," *Life Sciences*, vol. 86, no. 9–10, pp. 351–357, 2010.
- [54] J. Y. Kim, E. H. Kim, S. S. Park, J. H. Lim, T. K. Kwon, and K. S. Choi, "Quercetin sensitizes human hepatoma cells to TRAIL-induced apoptosis via Sp1-mediated DR5 up-regulation and proteasome-mediated c-FLIPS down-regulation," *Journal of Cellular Biochemistry*, vol. 105, no. 6, pp. 1386–1398, 2008.
- [55] W. Chen, X. Wang, J. Zhuang, L. Zhang, and Y. Lin, "Induction of death receptor 5 and suppression of survivin contribute to sensitization of TRAIL-induced cytotoxicity by quercetin in non-small cell lung cancer cells," *Carcinogenesis*, vol. 28, no. 10, pp. 2114–2121, 2007.
- [56] F. H. Psahoulia, K. G. Drosopoulos, L. Doubavaska, L. Andera, and A. Pintzas, "Quercetin enhances TRAIL-mediated apoptosis in colon cancer cells by inducing the accumulation of death receptors in lipid rafts," *Molecular Cancer Therapeutics*, vol. 6, no. 9, pp. 2591–2599, 2007.
- [57] Y. H. Kim, D. H. Lee, J. H. Jeong, Z. S. Guo, and Y. J. Lee, "Quercetin augments TRAIL-induced apoptotic death: involvement of the ERK signal transduction pathway," *Biochemical Pharmacology*, vol. 75, no. 10, pp. 1946–1958, 2008.
- [58] L. Gibellini, M. Pinti, M. Nasi et al., "Quercetin and cancer chemoprevention," *Evidence-Based Complementary and Alternative Medicine*, vol. 2011, Article ID 591356, 15 pages, 2011.
- [59] Q. Zhang, X. H. Zhao, and Z. J. Wang, "Cytotoxicity of flavones and flavonols to a human esophageal squamous cell carcinoma cell line (KYSE-510) by induction of G2/M arrest and apoptosis," *Toxicology in Vitro*, vol. 23, no. 5, pp. 797–807, 2009.
- [60] J. H. Jeong, J. Y. An, Y. T. Kwon, J. G. Rhee, and Y. J. Lee, "Effects of low dose quercetin: cancer cell-specific inhibition of cell cycle progression," *Journal of Cellular Biochemistry*, vol. 106, no. 1, pp. 73–82, 2009.
- [61] J. H. Yang, T. C. Hsia, H. M. Kuo et al., "Inhibition of lung cancer cell growth by quercetin glucuronides via G2/M arrest and induction of apoptosis," *Drug Metabolism and Disposition*, vol. 34, no. 2, pp. 296–304, 2006.
- [62] C. Mu, P. Jia, Z. Yan, X. Liu, X. Li, and H. Liu, "Quercetin induces cell cycle G1 arrest through elevating Cdk inhibitors p21 and p27 in human hepatoma cell line (HepG2)," *Methods and Findings in Experimental and Clinical Pharmacology*, vol. 29, no. 3, pp. 179–183, 2007.
- [63] B. B. Aggarwal and B. Sung, "Pharmacological basis for the role of curcumin in chronic diseases: an age-old spice with modern targets," *Trends in Pharmacological Sciences*, vol. 30, no. 2, pp. 85–94, 2009.
- [64] A. S. Strimpakos and R. A. Sharma, "Curcumin: preventive and therapeutic properties in laboratory studies and clinical trials," *Antioxidants & Redox Signaling*, vol. 10, no. 3, pp. 511–546, 2008.
- [65] B. B. Aggarwal, A. Kumar, and A. C. Bharti, "Anticancer potential of curcumin: preclinical and clinical studies," *Anti-cancer Research*, vol. 23, no. 1A, pp. 363–398, 2003.
- [66] K. M. Terlikowska, A. M. Witkowska, M. E. Zujko, B. Dobrzycka, and S. J. Terlikowski, "Potential application of curcumin and its analogues in the treatment strategy of patients with primary epithelial ovarian cancer," *International Journal of Molecular Sciences*, vol. 15, no. 12, pp. 21703–21722, 2014.
- [67] T. Ak and I. Gulcin, "Antioxidant and radical scavenging properties of curcumin," *Chemico-Biological Interactions*, vol. 174, no. 1, pp. 27–37, 2008.
- [68] A. Barzegar and A. A. Moosavi-Movahedi, "Intracellular ROS protection efficiency and free radical-scavenging activity of curcumin," *PloS ONE*, vol. 6, no. 10, article e26012, 2011.
- [69] J. E. Kim, A. R. Kim, H. Y. Chung, S. Y. Han, B. S. Kim, and J. S. Choi, "In vitro peroxynitrite scavenging activity of diarylheptanoids from *Curcuma longa*," *Phytotherapy Research*, vol. 17, no. 5, pp. 481–484, 2003.
- [70] S. J. McNally, E. M. Harrison, J. A. Ross, O. J. Garden, and S. J. Wigmore, "Curcumin induces heme oxygenase 1 through generation of reactive oxygen species, p38 activation and phosphatase inhibition," *International Journal of Molecular Medicine*, vol. 19, no. 1, pp. 165–172, 2007.
- [71] H. D. Panchal, K. Vranizan, C. Y. Lee, J. Ho, J. Ngai, and P. S. Timiras, "Early anti-oxidative and anti-proliferative curcumin effects on neuroglioma cells suggest therapeutic targets," *Neurochemical Research*, vol. 33, no. 9, pp. 1701–1710, 2008.
- [72] E. Balogun, M. Hoque, P. Gong et al., "Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive element," *The Biochemical Journal*, vol. 371, no. 3, pp. 887–895, 2003.
- [73] S. Karmakar, N. L. Banik, S. J. Patel, and S. K. Ray, "Curcumin activated both receptor-mediated and mitochondria-mediated proteolytic pathways for apoptosis in human glioblastoma T98G cells," *Neuroscience Letters*, vol. 407, no. 1, pp. 53–58, 2006.
- [74] R. Gogada, M. Amadori, H. Zhang et al., "Curcumin induces Apaf-1-dependent, p21-mediated caspase activation and apoptosis," *Cell Cycle*, vol. 10, no. 23, pp. 4128–4137, 2011.

- [75] N. R. Jana, P. Dikshit, A. Goswami, and N. Nukina, "Inhibition of proteasomal function by curcumin induces apoptosis through mitochondrial pathway," *The Journal of Biological Chemistry*, vol. 279, no. 12, pp. 11680–11685, 2004.
- [76] D. Morin, S. Barthelemy, R. Zini, S. Labidalle, and J. P. Tillement, "Curcumin induces the mitochondrial permeability transition pore mediated by membrane protein thiol oxidation," *FEBS Letters*, vol. 495, no. 1-2, pp. 131–136, 2001.
- [77] F. Thayyullathil, S. Chathoth, A. Hago, M. Patel, and S. Galadari, "Rapid reactive oxygen species (ROS) generation induced by curcumin leads to caspase-dependent and -independent apoptosis in L929 cells," *Free Radical Biology & Medicine*, vol. 45, no. 10, pp. 1403–1412, 2008.
- [78] S. Bhaumik, R. Anjum, N. Rangaraj, B. V. V. Pardhasaradhi, and A. Khar, "Curcumin mediated apoptosis in AK-5 tumor cells involves the production of reactive oxygen intermediates," *FEBS Letters*, vol. 456, no. 2, pp. 311–314, 1999.
- [79] J. L. Watson, R. Hill, P. B. Yaffe et al., "Curcumin causes superoxide anion production and p53-independent apoptosis in human colon cancer cells," *Cancer Letters*, vol. 297, no. 1, pp. 1–8, 2010.
- [80] J. Cao, L. Jia, H. M. Zhou, Y. Liu, and L. F. Zhong, "Mitochondrial and nuclear DNA damage induced by curcumin in human hepatoma G2 cells," *Toxicological Sciences*, vol. 91, no. 2, pp. 476–483, 2006.
- [81] J. Cao, Y. Liu, L. Jia et al., "Curcumin induces apoptosis through mitochondrial hyperpolarization and mtDNA damage in human hepatoma G2 cells," *Free Radical Biology & Medicine*, vol. 43, no. 6, pp. 968–975, 2007.
- [82] E. M. Jung, J. H. Lim, T. J. Lee, J. W. Park, K. S. Choi, and T. K. Kwon, "Curcumin sensitizes tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-induced apoptosis through reactive oxygen species-mediated upregulation of death receptor 5 (DR5)," *Carcinogenesis*, vol. 26, no. 11, pp. 1905–1913, 2005.
- [83] X. Yang, Z. Li, Q. Wu, S. Chen, C. Yi, and C. Gong, "TRAIL and curcumin codelivery nanoparticles enhance TRAIL-induced apoptosis through upregulation of death receptors," *Drug Delivery*, vol. 24, no. 1, pp. 1526–1536, 2017.
- [84] Y. J. Lee, N. Y. Kim, Y. A. Suh, and C. Lee, "Involvement of ROS in curcumin-induced autophagic cell death," *The Korean Journal of Physiology and Pharmacology*, vol. 15, no. 1, pp. 1–7, 2011.
- [85] Z. C. Gersey, G. A. Rodriguez, E. Barbarite et al., "Curcumin decreases malignant characteristics of glioblastoma stem cells via induction of reactive oxygen species," *BMC Cancer*, vol. 17, no. 1, p. 99, 2017.
- [86] J. U. Marquardt, L. Gomez-Quiroz, L. O. Arreguin Camacho et al., "Curcumin effectively inhibits oncogenic NF- κ B signaling and restrains stemness features in liver cancer," *Journal of Hepatology*, vol. 63, no. 3, pp. 661–669, 2015.
- [87] T. Choudhuri, S. Pal, M. L. Agwarwal, T. Das, and G. Sa, "Curcumin induces apoptosis in human breast cancer cells through p53-dependent Bax induction," *FEBS Letters*, vol. 512, no. 1-3, pp. 334–340, 2002.
- [88] S. Shankar and R. K. Srivastava, "Bax and Bak genes are essential for maximum apoptotic response by curcumin, a polyphenolic compound and cancer chemopreventive agent derived from turmeric, *Curcuma longa*," *Carcinogenesis*, vol. 28, no. 6, pp. 1277–1286, 2007.
- [89] J. Yang, J. Ning, L. Peng, and D. He, "Effect of curcumin on Bcl-2 and Bax expression in nude mice prostate cancer," *International Journal of Clinical and Experimental Pathology*, vol. 8, no. 8, pp. 9272–9278, 2015.
- [90] J. Yu, X. Zhou, X. He, M. Dai, and Q. Zhang, "Curcumin induces apoptosis involving bax/bcl-2 in human hepatoma SMMC-7721 cells," *Asian Pacific Journal of Cancer Prevention*, vol. 12, no. 8, pp. 1925–1929, 2011.
- [91] S. Y. Kim, S. H. Jung, and H. S. Kim, "Curcumin is a potent broad spectrum inhibitor of matrix metalloproteinase gene expression in human astrogloma cells," *Biochemical and Biophysical Research Communications*, vol. 337, no. 2, pp. 510–516, 2005.
- [92] D. Kumar, M. Kumar, C. Saravanan, and S. K. Singh, "Curcumin: a potential candidate for matrix metalloproteinase inhibitors," *Expert Opinion on Therapeutic Targets*, vol. 16, no. 10, pp. 959–972, 2012.
- [93] C. Aza-Gonzalez, H. G. Nunez-Palenius, and N. Ochoa-Alejo, "Molecular biology of capsaicinoid biosynthesis in chili pepper (*Capsicum* spp.)," *Plant Cell Reports*, vol. 30, no. 5, pp. 695–706, 2011.
- [94] K. Bley, G. Boorman, B. Mohammad, D. McKenzie, and S. Babbar, "A comprehensive review of the carcinogenic and anticarcinogenic potential of capsaicin," *Toxicologic Pathology*, vol. 40, no. 6, pp. 847–873, 2012.
- [95] H. F. Lu, Y. L. Chen, J. S. Yang et al., "Antitumor activity of capsaicin on human colon cancer cells in vitro and colo 205 tumor xenografts in vivo," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 24, pp. 12999–13005, 2010.
- [96] R. Clark and S. H. Lee, "Anticancer properties of capsaicin against human cancer," *Anticancer Research*, vol. 36, no. 3, pp. 837–843, 2016.
- [97] H. Impheng, S. Pongcharoen, L. Richert, D. Pekthong, and P. Srisawang, "The selective target of capsaicin on FASN expression and de novo fatty acid synthesis mediated through ROS generation triggers apoptosis in HepG2 cells," *PLoS ONE*, vol. 9, no. 9, article e107842, 2014.
- [98] K. M. Yang, J. O. Pyo, G. Y. Kim et al., "Capsaicin induces apoptosis by generating reactive oxygen species and disrupting mitochondrial transmembrane potential in human colon cancer cell lines," *Cellular & Molecular Biology Letters*, vol. 14, no. 3, pp. 497–510, 2009.
- [99] R. Zhang, I. Humphreys, R. P. Sahu, Y. Shi, and S. K. Srivastava, "In vitro and in vivo induction of apoptosis by capsaicin in pancreatic cancer cells is mediated through ROS generation and mitochondrial death pathway," *Apoptosis*, vol. 13, no. 12, pp. 1465–1478, 2008.
- [100] L. Xie, G. H. Xiang, T. Tang et al., "Capsaicin and dihydrocapsaicin induce apoptosis in human glioma cells via ROS and Ca²⁺-mediated mitochondrial pathway," *Molecular Medicine Reports*, vol. 14, no. 5, pp. 4198–4208, 2016.
- [101] A. M. Sanchez, M. G. Sanchez, S. Malagarie-Cazenave, N. Olea, and I. Diaz-Laviada, "Induction of apoptosis in prostate tumor PC-3 cells and inhibition of xenograft prostate tumor growth by the vanilloid capsaicin," *Apoptosis*, vol. 11, no. 1, pp. 89–99, 2006.
- [102] A. Macho, M. A. Calzado, J. Muñoz-Blanco et al., "Selective induction of apoptosis by capsaicin in transformed cells: the role of reactive oxygen species and calcium," *Cell Death and Differentiation*, vol. 6, no. 2, pp. 155–165, 1999.

- [103] K. C. Pramanik, S. R. Boreddy, and S. K. Srivastava, "Role of mitochondrial electron transport chain complexes in capsaicin mediated oxidative stress leading to apoptosis in pancreatic cancer cells," *PLoS ONE*, vol. 6, no. 5, article e20151, 2011.
- [104] C. Amantini, M. Mosca, M. Nabissi et al., "Capsaicin-induced apoptosis of glioma cells is mediated by TRPV1 vanilloid receptor and requires p38 MAPK activation," *Journal of Neurochemistry*, vol. 102, no. 3, pp. 977–990, 2007.
- [105] C. Amantini, P. Ballarini, S. Caprodossi et al., "Triggering of transient receptor potential vanilloid type 1 (TRPV1) by capsaicin induces Fas/CD95-mediated apoptosis of urothelial cancer cells in an ATM-dependent manner," *Carcinogenesis*, vol. 30, no. 8, pp. 1320–1329, 2009.
- [106] S. R. Kim, S. U. Kim, U. Oh, and B. K. Jin, "Transient receptor potential vanilloid subtype 1 mediates microglial cell death in vivo and in vitro via Ca²⁺-mediated mitochondrial damage and cytochrome c release," *Journal of Immunology*, vol. 177, no. 7, pp. 4322–4329, 2006.
- [107] D. Chen, Z. Yang, Y. Wang, G. Zhu, and X. Wang, "Capsaicin induces cycle arrest by inhibiting cyclin-dependent-kinase in bladder carcinoma cells," *International Journal of Urology*, vol. 19, no. 7, pp. 662–668, 2012.
- [108] C. H. Lin, W. C. Lu, C. W. Wang, Y. C. Chan, and M. K. Chen, "Capsaicin induces cell cycle arrest and apoptosis in human KB cancer cells," *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, 2013.
- [109] K. J. Min and T. K. Kwon, "Anticancer effects and molecular mechanisms of epigallocatechin-3-gallate," *Integrative Medicine Research*, vol. 3, no. 1, pp. 16–24, 2014.
- [110] T. Nakazato, K. Ito, Y. Miyakawa et al., "Catechin, a green tea component, rapidly induces apoptosis of myeloid leukemic cells via modulation of reactive oxygen species production in vitro and inhibits tumor growth in vivo," *Haematologica*, vol. 90, no. 3, pp. 317–325, 2005.
- [111] C. Noda, J. He, T. Takano et al., "Induction of apoptosis by epigallocatechin-3-gallate in human lymphoblastoid B cells," *Biochemical and biophysical research communications*, vol. 362, no. 4, pp. 951–957, 2007.
- [112] W. Li, S. Nie, Q. Yu, and M. Xie, "(-)-Epigallocatechin-3-gallate induces apoptosis of human hepatoma cells by mitochondrial pathways related to reactive oxygen species," *Journal of agricultural and food chemistry*, vol. 57, no. 15, pp. 6685–6691, 2009.
- [113] S. Liu, X. J. Wang, Y. Liu, and Y. F. Cui, "PI3K/AKT/mTOR signaling is involved in (-)-epigallocatechin-3-gallate-induced apoptosis of human pancreatic carcinoma cells," *The American journal of Chinese medicine*, vol. 41, no. 03, pp. 629–642, 2013.
- [114] J. J. Gu, K. S. Qiao, P. Sun, P. Chen, and Q. Li, "Study of EGCG induced apoptosis in lung cancer cells by inhibiting PI3K/Akt signaling pathway," *European review for medical and pharmacological sciences*, vol. 22, no. 14, pp. 4557–4563, 2018.
- [115] P. P. Wu, S. C. Kuo, W. W. Huang et al., "Epigallocatechin gallate induced apoptosis in human adrenal cancer NCI-H295 cells through caspase-dependent and caspase-independent pathway," *Anticancer research*, vol. 29, no. 4, pp. 1435–1442, 2009.
- [116] S. Shankar, G. Suthakar, and R. K. Srivastava, "Epigallocatechin-3-gallate inhibits cell cycle and induces apoptosis in pancreatic cancer," *Frontiers in Bioscience*, vol. 12, no. 12, pp. 5039–5051, 2007.
- [117] J. Qin, L. P. Xie, X. Y. Zheng et al., "A component of green tea, (-)-epigallocatechin-3-gallate, promotes apoptosis in T24 human bladder cancer cells via modulation of the PI3K/Akt pathway and Bcl-2 family proteins," *Biochemical and biophysical research communications*, vol. 354, no. 4, pp. 852–857, 2007.
- [118] S. S. Chung and J. V. Vadgama, "Curcumin and Epigallocatechin Gallate Inhibit the Cancer Stem Cell Phenotype via Down-regulation of STAT3–NF κ B Signaling," *Anticancer Research*, vol. 35, no. 1, pp. 39–46, 2015.
- [119] Y. C. Chou, M. Y. Chang, M. J. Wang et al., "PEITC induces apoptosis of human brain glioblastoma GBM8401 cells through the extrinsic- and intrinsic -signaling pathways," *Neurochemistry international*, vol. 81, pp. 32–40, 2015.
- [120] P. Gupta, C. Adkins, P. Lockman, and S. K. Srivastava, "Metastasis of breast tumor cells to brain is suppressed by phenethyl isothiocyanate in a novel in vivo metastasis model," *PLoS ONE*, vol. 8, no. 6, article e67278, 2013.
- [121] S. Kalkunte, N. Swamy, D. S. Dizon, and L. Brard, "Benzyl isothiocyanate (BITC) induces apoptosis in ovarian cancer cells in vitro," *Journal of experimental therapeutics & oncology*, vol. 5, no. 4, pp. 287–300, 2006.
- [122] Y. Zhu, J. X. Zhuang, Q. Wang, H. Y. Zhang, and P. Yang, "Inhibitory effect of benzyl isothiocyanate on proliferation in vitro of human glioma cells," *Asian Pacific journal of cancer prevention*, vol. 14, no. 4, pp. 2607–2610, 2013.
- [123] J. V. Higdon, B. Delage, D. E. Williams, and R. H. Dashwood, "Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis," *Pharmacological Research*, vol. 55, no. 3, pp. 224–236, 2007.
- [124] L. Tang, G. R. Zirpoli, K. Guru et al., "Intake of cruciferous vegetables modifies bladder cancer survival," *Cancer Epidemiology Biomarkers & Prevention*, vol. 19, no. 7, pp. 1806–1811, 2010.
- [125] D. Xiao, A. A. Powolny, M. B. Moura et al., "Phenethyl isothiocyanate inhibits oxidative phosphorylation to trigger reactive oxygen species-mediated death of human prostate cancer cells," *The Journal of biological chemistry*, vol. 285, no. 34, pp. 26558–26569, 2010.
- [126] D. Trachootham, Y. Zhou, H. Zhang et al., "Selective killing of oncogenically transformed cells through a ROS-mediated mechanism by beta-phenylethyl isothiocyanate," *Cancer Cell*, vol. 10, no. 3, pp. 241–252, 2006.
- [127] D. Xiao, V. Vogel, and S. V. Singh, "Benzyl isothiocyanate-induced apoptosis in human breast cancer cells is initiated by reactive oxygen species and regulated by Bax and Bak," *Molecular Cancer Therapeutics*, vol. 5, no. 11, pp. 2931–2945, 2006.
- [128] J. Wang, B. Luo, X. Li et al., "Inhibition of cancer growth in vitro and in vivo by a novel ROS-modulating agent with ability to eliminate stem-like cancer cells," *Cell death & disease*, vol. 8, no. 6, article e2887, 2017.
- [129] Y.-H. Hong, H. Uddin, U. Jo et al., "ROS accumulation by PEITC selectively kills ovarian cancer cells via UPR-mediated apoptosis," *Frontiers in oncology*, vol. 5, p. 167, 2015.
- [130] Y. Wang, S. Wei, J. Wang, Q. Fang, and Q. Chai, "Phenethyl isothiocyanate inhibits growth of human chronic myeloid leukemia K562 cells via reactive oxygen species generation

- and caspases," *Molecular medicine reports*, vol. 10, no. 1, pp. 543–549, 2014.
- [131] R. Kasiappan, I. Jutooru, K. Karki, E. Hedrick, and S. Safe, "Benzyl isothiocyanate (BITC) induces reactive oxygen species-dependent repression of STAT3 protein by down-regulation of specificity proteins in pancreatic cancer," *The Journal of biological chemistry*, vol. 291, no. 53, pp. 27122–27133, 2016.
 - [132] J. F. Lin, T. F. Tsai, S. C. Yang et al., "Benzyl isothiocyanate induces reactive oxygen species-initiated autophagy and apoptosis in human prostate cancer cells," *Oncotarget*, vol. 8, no. 12, pp. 20220–20234, 2017.
 - [133] A. Ranjan, N. M. Fofaria, S. H. Kim, and S. K. Srivastava, "Modulation of signal transduction pathways by natural compounds in cancer," *Chinese journal of natural medicines*, vol. 13, no. 10, pp. 730–742, 2015.
 - [134] R. P. Sahu, R. Zhang, S. Batra, Y. Shi, and S. K. Srivastava, "Benzyl isothiocyanate-mediated generation of reactive oxygen species causes cell cycle arrest and induces apoptosis via activation of MAPK in human pancreatic cancer cells," *Carcinogenesis*, vol. 30, no. 10, pp. 1744–1753, 2009.
 - [135] Y. T. Yeh, H. Yeh, S. H. Su et al., "Phenethyl isothiocyanate induces DNA damage-associated G2/M arrest and subsequent apoptosis in oral cancer cells with varying p53 mutations," *Free radical biology & medicine*, vol. 74, pp. 1–13, 2014.
 - [136] R. Zhang, S. Loganathan, I. Humphreys, and S. K. Srivastava, "Benzyl isothiocyanate-induced DNA damage causes G2/M cell cycle arrest and apoptosis in human pancreatic cancer cells," *The Journal of Nutrition*, vol. 136, no. 11, pp. 2728–2734, 2006.
 - [137] R. A. Rather and M. Bhagat, "Cancer chemoprevention and piperine: molecular mechanisms and therapeutic opportunities," *Frontiers in cell and developmental biology*, vol. 6, p. 10, 2018.
 - [138] V. Gunasekaran, K. Elangovan, and S. Niranjali Devaraj, "Targeting hepatocellular carcinoma with piperine by radical-mediated mitochondrial pathway of apoptosis: an in vitro and in vivo study," *Food and Chemical Toxicology*, vol. 105, pp. 106–118, 2017.
 - [139] S. Siddiqui, M. S. Ahamad, A. Jafri, M. Afzal, and M. Arshad, "Piperine triggers apoptosis of human oral squamous carcinoma through cell cycle arrest and mitochondrial oxidative stress," *Nutrition and Cancer*, vol. 69, no. 5, pp. 791–799, 2017.
 - [140] M. Takashina, S. Inoue, K. Tomihara et al., "Different effect of resveratrol to induction of apoptosis depending on the type of human cancer cells," *International Journal of Oncology*, vol. 50, no. 3, pp. 787–797, 2017.
 - [141] T. C. Hsieh and J. M. Wu, "Differential effects on growth, cell cycle arrest, and induction of apoptosis by resveratrol in human prostate cancer cell lines," *Experimental Cell Research*, vol. 249, no. 1, pp. 109–115, 1999.
 - [142] Y. Chen, S. H. Tseng, H. S. Lai, and W. J. Chen, "Resveratrol-induced cellular apoptosis and cell cycle arrest in neuroblastoma cells and antitumor effects on neuroblastoma in mice," *Surgery*, vol. 136, no. 1, pp. 57–66, 2004.
 - [143] S. Shankar, Q. Chen, I. Siddiqui, K. Sarva, and R. K. Srivastava, "Sensitization of TRAIL-resistant LNCaP cells by resveratrol (3, 4', 5 tri-hydroxystilbene): molecular mechanisms and therapeutic potential," *Journal of Molecular Signaling*, vol. 2, p. 7, 2007.
 - [144] E. M. Varoni, A. F. Lo Faro, J. Sharifi-Rad, and M. Iriti, "Anti-cancer molecular mechanisms of resveratrol," *Frontiers in Nutrition*, vol. 3, 2016.
 - [145] K. B. Harikumar and B. B. Aggarwal, "Resveratrol: A multi-targeted agent for age-associated chronic diseases," *Cell Cycle*, vol. 7, no. 8, pp. 1020–1035, 2008.
 - [146] N. C. Whitlock and S. J. Baek, "The anticancer effects of resveratrol: modulation of transcription factors," *Nutrition and Cancer*, vol. 64, no. 4, pp. 493–502, 2012.
 - [147] J. R. Gledhill, M. G. Montgomery, A. G. W. Leslie, and J. E. Walker, "Mechanism of inhibition of bovine F1-ATPase by resveratrol and related polyphenols," *Proceedings of the National Academy of Sciences*, vol. 104, no. 34, pp. 13632–13637, 2007.
 - [148] J. Zheng and V. D. Ramirez, "Inhibition of mitochondrial proton F0F1-ATPase/ATP synthase by polyphenolic phytochemicals," *British Journal of Pharmacology*, vol. 130, no. 5, pp. 1115–1123, 2000.
 - [149] S. K. Singh, S. Banerjee, E. P. Acosta, J. W. Lillard, and R. Singh, "Resveratrol induces cell cycle arrest and apoptosis with docetaxel in prostate cancer cells via a p53/ p21WAF1/ CIP1 and p27KIP1 pathway," *Oncotarget*, vol. 8, no. 10, pp. 17216–17228, 2017.
 - [150] M. Athar, J. H. Back, L. Kopelovich, D. R. Bickers, and A. L. Kim, "Multiple molecular targets of resveratrol: anti-carcinogenic mechanisms," *Archives of Biochemistry and Biophysics*, vol. 486, no. 2, pp. 95–102, 2009.
 - [151] L. D. S. Rosa, N. A. Jordão, N. D. C. P. Soares, J. F. DeMesquita, M. Monteiro, and A. J. Teodoro, "Pharmacokinetic, Antiproliferative and Apoptotic Effects of Phenolic Acids in Human Colon Adenocarcinoma Cells Using In Vitro and In Silico Approaches," *Molecules*, vol. 23, no. 10, p. 2569, 2018.
 - [152] S. K. Jaganathan, E. Supriyanto, and M. Mandal, "Events associated with apoptotic effect of p-Coumaric acid in HCT-15 colon cancer cells," *World Journal of Gastroenterology*, vol. 19, no. 43, pp. 7726–7734, 2013.
 - [153] S. H. Sharma, V. Rajamanickam, and S. Nagarajan, "Antiproliferative effect of p-Coumaric acid targets UPR activation by downregulating Grp78 in colon cancer," *Chemico-Biological Interactions*, vol. 291, pp. 16–28, 2018.
 - [154] J. Y. H. Ong, P. V. C. Yong, Y. M. Lim, and A. S. H. Ho, "2-Methoxy-1,4-naphthoquinone (MNQ) induces apoptosis of A549 lung adenocarcinoma cells via oxidation-triggered JNK and p38 MAPK signaling pathways," *Life Sciences*, vol. 135, pp. 158–164, 2015.
 - [155] J. Ignacio Aguiló, M. Iturralde, I. Monleón et al., "Cytotoxicity of quinone drugs on highly proliferative human leukemia T cells: reactive oxygen species generation and inactive shortened SOD1 isoform implications," *Chemico-Biological Interactions*, vol. 198, no. 1–3, pp. 18–28, 2012.
 - [156] E. B. Lee, M. G. Cheon, J. Cui, Y. J. Lee, E. K. Seo, and H. H. Jang, "The quinone-based derivative, HMNQ induces apoptotic and autophagic cell death by modulating reactive oxygen species in cancer cells," *Oncotarget*, vol. 8, no. 59, pp. 99637–99648, 2017.
 - [157] E. R. Lee, Y. J. Kang, H. J. Kim et al., "Regulation of apoptosis by modified naringenin derivatives in human colorectal carcinoma RKO cells," *Journal of Cellular Biochemistry*, vol. 104, no. 1, pp. 259–273, 2008.
 - [158] P. Totta, F. Acconcia, S. Leone, I. Cardillo, and M. Marino, "Mechanisms of Naringenin-induced Apoptotic Cascade in

- Cancer Cells: Involvement of Estrogen Receptor α and β Signaling," *IUBMB Life*, vol. 56, no. 8, pp. 491–499, 2004.
- [159] B. R. You, S. Z. Kim, S. H. Kim, and W. H. Park, "Gallic acid-induced lung cancer cell death is accompanied by ROS increase and glutathione depletion," *Molecular and Cellular Biochemistry*, vol. 357, no. 1-2, pp. 295–303, 2011.
- [160] L. H. Russell JR, E. Mazzio, R. B. Badisa et al., "Autooxidation of gallic acid induces ROS-dependent death in human prostate cancer LNCaP cells," *Anticancer Research*, vol. 32, no. 5, pp. 1595–1602, 2012.
- [161] C. Y. Chuang, H. C. Liu, L. C. Wu, C. Y. Chen, J. T. Chang, and S. L. Hsu, "Gallic acid induces apoptosis of lung fibroblasts via a reactive oxygen species-dependent ataxia telangiectasia mutated-p53 activation pathway," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 5, pp. 2943–2951, 2010.
- [162] C. Y. Chen, K. C. Chen, T. Y. Yang, H. C. Liu, and S. L. Hsu, "Gallic acid induces a reactive oxygen species-provoked c-Jun NH2-terminal kinase-dependent apoptosis in lung fibroblasts," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 613950, 12 pages, 2013.
- [163] X. Chen, O. Q. P. Yin, Z. Zuo, and M. S. S. Chow, "Pharmacokinetics and modeling of quercetin and metabolites," *Pharmaceutical Research*, vol. 22, no. 6, pp. 892–901, 2005.
- [164] S. K. Vareed, M. Kakarala, M. T. Ruffin et al., "Pharmacokinetics of curcumin conjugate metabolites in healthy human subjects," *Cancer Epidemiology Biomarkers & Prevention*, vol. 17, no. 6, pp. 1411–1417, 2008.
- [165] T. Walle, "Bioavailability of resveratrol," *Annals of the New York Academy of Sciences*, vol. 1215, no. 1, pp. 9–15, 2011.
- [166] E. Burgos-Moron, J. M. Calderon-Montano, J. Salvador, A. Robles, and M. Lopez-Lazaro, "The dark side of curcumin," *International Journal of Cancer*, vol. 126, no. 7, pp. 1771–1775, 2010.
- [167] G. Shoba, D. Joy, T. Joseph, M. Majeed, R. Rajendran, and P. S. Srinivas, "Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers," *Planta Medica*, vol. 64, no. 04, pp. 353–356, 1998.
- [168] J. J. Johnson, M. Nihal, I. A. Siddiqui et al., "Enhancing the bioavailability of resveratrol by combining it with piperine," *Molecular Nutrition & Food Research*, vol. 55, no. 8, pp. 1169–1176, 2011.
- [169] A. Vyas, P. Dandawate, S. Padhye, A. Ahmad, and F. Sarkar, "Perspectives on new synthetic curcumin analogs and their potential anticancer properties," *Current Pharmaceutical Design*, vol. 19, no. 11, pp. 2047–2069, 2013.
- [170] X. Cai, Z. Fang, J. Dou, A. Yu, and G. Zhai, "Bioavailability of quercetin: problems and promises," *Current Medicinal Chemistry*, vol. 20, no. 20, pp. 2572–2582, 2013.
- [171] P. Urbina-Cano, L. Bobadilla-Morales, M. A. Ramírez-Herrera et al., "DNA damage in mouse lymphocytes exposed to curcumin and copper," *Journal of Applied Genetics*, vol. 47, no. 4, pp. 377–382, 2006.
- [172] J. Blasiak, A. Trzeciak, and J. Kowalik, "Curcumin damages DNA in human gastric mucosa cells and lymphocytes," *Journal of Environmental Pathology, Toxicology and Oncology*, vol. 18, pp. 271–276, 1999.
- [173] S. T. Dance-Barnes, N. D. Kock, J. E. Moore et al., "Lung tumor promotion by curcumin," *Carcinogenesis*, vol. 30, no. 6, pp. 1016–1023, 2009.
- [174] A. M. Bode and Z. Dong, "The two faces of capsaicin," *Cancer Research*, vol. 71, no. 8, pp. 2809–2814, 2011.
- [175] J. Hu, L. Chen, F. Lei et al., "Investigation of quercetin stability in cell culture medium: role in in vitro experiment," *African Journal of Pharmacy and Pharmacology*, vol. 6, pp. 1069–1076, 2012.
- [176] S. K. Savale, "Curcumin as a model drug: conformation, solubility estimation, morphological, in vitro and in vivo bio-distribution study," *Journal of PharmaSciTech*, vol. 7, pp. 31–35, 2017.
- [177] A. Kocyigit, E. M. Guler, and M. Dikilitas, "Role of antioxidant phytochemicals in prevention, formation and treatment of cancer," in *Reactive Oxygen Species (ROS) in Living Cells*, IntechOpen, 2017.
- [178] M. Hashemzaei, A. D. Far, A. Yari et al., "Anticancer and apoptosis-inducing effects of quercetin in vitro and in vivo," *Oncology Reports*, vol. 38, no. 2, pp. 819–828, 2017.
- [179] P. Pratheeshkumar, A. Budhraj, Y. O. Son et al., "Quercetin inhibits angiogenesis mediated human prostate tumor growth by targeting VEGFR-2 regulated AKT/mTOR/P70S6K signaling pathways," *PLoS ONE*, vol. 7, no. 10, article e47516, 2012.
- [180] J. Zhou, L. Fang, J. Liao et al., "Investigation of the anti-cancer effect of quercetin on HepG2 cells in vivo," *PLoS ONE*, vol. 12, no. 3, article e0172838, 2017.
- [181] V. Maso, A. K. Calgarotto, G. C. Franchi et al., "Multitarget effects of quercetin in leukemia," *Cancer Prevention Research*, vol. 7, no. 12, pp. 1240–1250, 2014.
- [182] W. D. Lu, Y. Qin, C. Yang, L. Li, and Z. X. Fu, "Effect of curcumin on human colon cancer multidrug resistance in vitro and in vivo," *Clinics*, vol. 68, no. 5, pp. 694–701, 2013.
- [183] H. W. Lai, S. Y. Chien, S. J. Kuo et al., "The Potential Utility of Curcumin in the Treatment of HER-2-Overexpressed Breast Cancer: An In Vitro and In Vivo Comparison Study with Herceptin," *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 486568, 12 pages, 2012.
- [184] A. Mori, S. Lehmann, J. O'Kelly et al., "Capsaicin, a component of red peppers, inhibits the growth of androgen-independent, p53 mutant prostate cancer cells," *Cancer Research*, vol. 66, no. 6, pp. 3222–3229, 2006.
- [185] N. H. Thoenissen, J. O'Kelly, D. Lu et al., "Capsaicin causes cell-cycle arrest and apoptosis in ER-positive and -negative breast cancer cells by modulating the EGFR/HER-2 pathway," *Oncogene*, vol. 29, no. 2, pp. 285–296, 2010.
- [186] Y. S. Shin, S. U. Kang, J. K. Park et al., "Anti-cancer effect of (-)-epigallocatechin-3-gallate (EGCG) in head and neck cancer through repression of transactivation and enhanced degradation of β -catenin," *Phytomedicine*, vol. 23, no. 12, pp. 1344–1355, 2016.
- [187] K. W. Luo, W. Y. Lung, C. X., X. L. Luo, and W. R. Huang, "EGCG inhibited bladder cancer T24 and 5637 cell proliferation and migration via PI3K/AKT pathway," *Oncotarget*, vol. 9, no. 15, pp. 12261–12272, 2018.
- [188] Y. Sakamoto, N. Terashita, T. Muraguchi, T. Fukusato, and S. Kubota, "Effects of epigallocatechin-3-gallate (EGCG) on A549 lung cancer tumor growth and angiogenesis," *Bioscience, Biotechnology, and Biochemistry*, vol. 77, no. 9, pp. 1799–1803, 2013.
- [189] Q. Hu, X. Chang, R. Yan et al., "(-)-Epigallocatechin-3-gallate induces cancer cell apoptosis via acetylation of amyloid precursor protein," *Medical Oncology*, vol. 32, no. 1, p. 390, 2015.

- [190] Y. C. Chou, M. Y. Chang, H. T. Lee et al., "Phenethyl isothiocyanate inhibits in vivo growth of xenograft tumors of human glioblastoma cells," *Molecules*, vol. 23, no. 9, p. 2305, 2018.
- [191] D. Xiao, K. L. Lew, Y. Zeng et al., "Phenethyl isothiocyanate-induced apoptosis in PC-3 human prostate cancer cells is mediated by reactive oxygen species-dependent disruption of the mitochondrial membrane potential," *Carcinogenesis*, vol. 27, no. 11, pp. 2223–2234, 2006.
- [192] P. Gupta and S. K. Srivastava, "Antitumor activity of phenethyl isothiocyanate in HER2-positive breast cancer models," *BMC Medicine*, vol. 10, no. 1, 2012.
- [193] S. Loganathan, P. K. Kandala, P. Gupta, and S. K. Srivastava, "Inhibition of EGFR-AKT axis results in the suppression of ovarian tumors in vitro and in preclinical mouse model," *PloS ONE*, vol. 7, no. 8, article e43577, 2012.
- [194] L. H. Lai, Q. H. Fu, Y. Liu et al., "Piperine suppresses tumor growth and metastasis in vitro and in vivo in a 4T1 murine breast cancer model," *Acta Pharmacologica Sinica*, vol. 33, no. 4, pp. 523–530, 2012.
- [195] A. Samykutty, A. V. Shetty, G. Dakshinamoorthy et al., "Piperine, a bioactive component of pepper spice exerts therapeutic effects on androgen dependent and androgen independent prostate cancer cells," *PloS ONE*, vol. 8, no. 6, article e65889, 2013.
- [196] L. Sun, H. Li, J. Chen et al., "A SUMOylation-dependent pathway regulates SIRT1 transcription and lung cancer metastasis," *Journal of the National Cancer Institute*, vol. 105, no. 12, pp. 887–898, 2013.
- [197] Y. Bai, Q. Q. Mao, J. Qin et al., "Resveratrol induces apoptosis and cell cycle arrest of human T24 bladder cancer cells in vitro and inhibits tumor growth in vivo," *Cancer Science*, vol. 101, no. 2, pp. 488–493, 2010.
- [198] S. Garvin, K. Ollinger, and C. Dabrosin, "Resveratrol induces apoptosis and inhibits angiogenesis in human breast cancer xenografts in vivo," *Cancer Letters*, vol. 231, no. 1, pp. 113–122, 2006.
- [199] S. K. Roy, Q. Chen, J. Fu, S. Shankar, and R. K. Srivastava, "Resveratrol inhibits growth of orthotopic pancreatic tumors through activation of FOXO transcription factors," *PloS ONE*, vol. 6, no. 9, article e25166, 2011.
- [200] S. J. Dias, K. Li, A. M. Rimando et al., "Trimethoxy-resveratrol and piceatannol administered orally suppress and inhibit tumor formation and growth in prostate cancer xenografts," *The Prostate*, vol. 73, no. 11, pp. 1135–1146, 2013.