

Research Article

Activation of the NF- κ B and MAPK Signaling Pathways Contributes to the Inflammatory Responses, but Not Cell Injury, in IPEC-1 Cells Challenged with Hydrogen Peroxide

Kan Xiao,¹ Congcong Liu,¹ Zhixiao Tu,¹ Qiao Xu,¹ Shaokui Chen,¹ Yang Zhang,¹ Xiuying Wang,¹ Jing Zhang,¹ Chien-An Andy Hu,^{1,2} and Yulan Liu¹ 

¹Hubei Key Laboratory of Animal Nutrition and Feed Science, Hubei Collaborative Innovation Center for Animal Nutrition and Feed Safety, Wuhan Polytechnic University, Wuhan, China

²Department of Biochemistry and Molecular Biology, University of New Mexico School of Medicine, Albuquerque, NM 87131, USA

Correspondence should be addressed to Yulan Liu; yulanflower@126.com

Received 28 May 2019; Revised 20 November 2019; Accepted 10 December 2019; Published 21 January 2020

Academic Editor: Nicoletta Guaragnella

Copyright © 2020 Kan Xiao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oxidative stress can lead to intestinal cell injury as well as the induction of inflammation. It is not clear whether inflammation is an important factor leading to cell injury caused by oxidative stress. The purpose of this study was to investigate the role of inflammation in intestinal injury caused by hydrogen peroxide (H₂O₂). Our results revealed that H₂O₂ stimulation significantly decreased the viability of intestinal porcine epithelial cells (IPEC-1), increased lactate dehydrogenase (LDH) activity, and disrupted the distribution of the tight junction protein claudin-1. H₂O₂ significantly increased the mRNA expression of interleukin-6 (IL-6), IL-8, and tumor necrosis factor- α (TNF- α). H₂O₂ stimulation also led to increased phosphorylation of p38 and jun N-terminal kinase (JNK), and p65 NF- κ B protein translocation into the nucleus of IPEC-1 cells. Cells treated with the NF- κ B inhibitor (BAY11-7082), the p38 inhibitor (SB202190), or the JNK inhibitor (PD98059) significantly decreased mRNA and protein expression of IL-6, IL-8, and TNF- α . However, treatment with mitogen-activated protein kinase (MAPK) or NF- κ B inhibitors did not prevent the damage effect on cell viability, LDH activity, or the distribution of claudin-1 in cells challenged with H₂O₂. In summary, our data demonstrate that activation of the NF- κ B and MAPK signaling pathways can contribute to the inflammatory response, but not cell injury, in IPEC-1 cells challenged with H₂O₂.

1. Introduction

The intestinal barrier is the first line of defense inside the body against harmful antigens and pathogens, within the intestinal lumen [1, 2]. It is a single layer of cells lining the gut, consisting of enterocyte membranes and tight junctions between enterocytes. The integrity of the intestinal barrier is essential for the digestion and absorption of nutrients, both in humans and animals [3–5]. Intestinal epithelial cells establish a barrier between the hostile external environment and the internal milieu. However, numerous factors such as infection, inflammation, and oxidative stress can result in intestinal epithelia injury, and dysfunction [1].

Hydrogen peroxide (H₂O₂) is a highly reactive oxygen specie and serves as a signaling molecule in the regulation

of a wide variety of biological processes [6–8]. Excessive H₂O₂ is capable of diffusing throughout mitochondria and across cell membranes, where it can cause multiple forms of cellular oxidative injury [9, 10]. Oxidative stress can lead to intestinal cell injury by various means [11, 12]. It can cause cell damage by free radicals or by induction of inflammation [13]. Oxidative stress appears to be an important driving force for an enhanced cytokine production in intestinal epithelial cells, resulting in gut mucosal inflammation [14]. However, it is not clear whether inflammation is an important factor leading to cell injury caused by oxidative stress. Mitogen-activated protein kinase (MAPK) and NF- κ B pathways are common inflammatory signaling pathways. Activation of MAPKs and NF- κ B pathways, caused proinflammatory cytokines such as interleukin-6

(IL-6), IL-8, and tumor necrosis factor- α (TNF- α) to be released, which could lead to the inflammatory response [15]. Traditionally, it is thought that cell death is the most extreme form of injury produced by inflammation [16]. Inflammation and cell death are distinct but highly interconnected and mutually regulated host-mediated mechanisms, which are important for maintaining homeostasis in organisms [17].

Therefore, the aim of the present study was to investigate the role of inflammation in intestinal injury caused by H₂O₂. It was found that H₂O₂ induced cell injury, upregulated proinflammatory cytokine-related gene expression, and activated NF- κ B, and MAPK signaling pathways in intestinal porcine epithelial cells (IPEC-1). In addition, the inhibition of p38, JNK, and NF- κ B with selective inhibitors downregulated the inflammatory response but did not prevent cell injury in IPEC-1 cells. Our data demonstrated that activation of the NF- κ B and MAPK signaling pathways could contribute to the inflammatory responses but not cell injury in IPEC-1 cells challenged with H₂O₂.

2. Materials and Methods

2.1. Cell Reagents and Antibodies. IPEC-1 cells were derived from the jejunum/ileum of a neonatal unsuckled piglet and were kindly provided by Dr. Guoyao Wu's laboratory at Texas A&M University. These cells were used for all experiments and were cultured in Dulbecco's Modified Eagle's Medium-F12 (HyClone, St. Louis, MO, USA), supplemented with 5% fetal bovine serum (Sigma-Aldrich, St. Louis, MO, USA), 1% insulin-transferrin-selenium, 1% penicillin/streptomycin (Gibco, NY, USA), and epidermal growth factor (5 ng/mL) (Gibco, NY, USA). Cells were split into 75 cm² culture flasks when cell monolayers reached 70%-80% confluency using trypsin/EDTA (0.25%, 0.9 mM EDTA) (Gibco, NY, USA). The cells were incubated at 37°C in a humidified atmosphere of 5% CO₂, and culture medium was changed every second day, according to standard culture protocols. The MAPK inhibitors (p38 inhibitor SB202190 and JNK inhibitor PD98059) and NF- κ B inhibitor (Bay11-7082) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The MAPK antibodies and anti-phospho-MAPK antibodies were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA). The secondary antibody was horseradish peroxidase-conjugated anti-rabbit IgG (Santa Cruz Biotechnology, Santa Cruz, CA, USA).

2.2. Cell Viability Assays. Cell viability was detected using the Cell Counting Kit-8 detection kit (CCK-8, Beyotime Institute of Biotechnology, Wuhan, CH). Cells were cultured in 96-well plates. When they were cultured to 70%-80% confluency, medium was removed and after washing two times with phosphate-buffered saline (PBS), cells were treated with different concentrations of H₂O₂ (0, 0.2, and 0.5 mM) for 3 h, or pretreated with different inhibitors (Bay11-7082, SB202190, and PD98059) for 1 h, in the presence or absence of 0.5 mM H₂O₂ for 3 h. Dulbecco's Modified Eagle's Medium-F12 medium was added to the control group. After a 3 h incubation, media was removed and 10% CCK-8

reagent (100 μ L per well) was added and incubated at 37°C for 1 h according to the manufacturer's instructions. The absorbance was quickly measured at a wavelength of 450 nm using a microplate reader (Bio-Rad, CA, USA). The viability of the cells was recorded as the percentage relative to untreated controls. All assays were performed in triplicate and in at least three independent experiments.

2.3. Lactate Dehydrogenase (LDH) Activity Measurement. The concentration of LDH released into the culture medium through damaged cell membranes was measured spectrophotometrically using an LDH assay kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, China). IPEC-1 cells were cultured in 12-well plates until 80% confluent. Cells were then exposed to H₂O₂ (0, 0.2, and 0.5 mM) or pretreated with different inhibitors (Bay11-7082, SB202190, or PD98059) for 1 h, in the presence or absence of 0.5 mM H₂O₂ for 3 h. Subsequently, cell supernatants were collected for LDH measurement according to the manufacturer's protocol. The absorbance was read at a wavelength of 450 nm using an automated microplate reader (Bio-Rad, CA, USA).

2.4. Quantitative Real-Time PCR. To investigate the effects of H₂O₂ or the inhibitors on proinflammatory cytokine expression, we used 0.5 mM H₂O₂ to treat the cells or cells were pretreated with different inhibitors (Bay11-7082, SB202190, or PD98059) for 1 h, in the presence or absence of 0.5 mM H₂O₂ for 3 h; cells were then collected for mRNA expression analysis. Messenger RNA levels were determined by real-time PCR as described by [16]. After washing IPEC-1 cells two times with ice-cold PBS, total RNA was extracted using the RNAiso Plus Kit (TaKaRa Biotechnology (Dalian) Co., Ltd., Dalian, China) following the manufacturer's instructions. After purification and quantitation, reverse transcription was performed using the PrimeScript[®] RT Reagent Kit (TaKaRa Biotechnology (Dalian) Co., Ltd., Dalian, China) following the manufacturer's instructions. Quantitative analysis of PCR was carried out on a ABI 7500 Real-Time PCR System (Applied Biosystems, Life Technologies) using a SYBR[®] Premix Ex Taq[™] qPCR Kit (TaKaRa Biotechnology (Dalian) Co., Ltd., Dalian, China). The primer pairs used are listed in Table 1. Results were analyzed by the 2^{- $\Delta\Delta$ CT} method with GAPDH as the housekeeping gene as GAPDH displayed no variation among all groups [16]. All samples were run in triplicate. Relative mRNA abundance of each target gene was normalized to the control group.

2.5. Cytokine Production Measurement. IPEC-1 cells were seeded in 12-well plates and treated with different concentrations of H₂O₂ (0 and 0.5 mM) for 3 h or pretreated with different inhibitors (Bay11-7082, SB202190, and PD98059) for 1 h, in the presence or absence of 0.5 mM H₂O₂ for 3 h. Subsequently, cell supernatants were collected for cytokine IL-6, IL-8, and TNF- α measurement by ELISA using commercial kits (4A Biotech, Beijing, China) according to the manufacturer's protocol.

2.6. Western Blot. To evaluate the effects of H₂O₂ on MAPK signaling activation, we used the 0.5 mM concentrations of

TABLE 1: Primer sequences used for real-time PCR.

Primer name	Primer sequence (5'-3')	Product size (bp)	Accession numbers
<i>IL-6</i>	F: AAGGTGATGCCACCTCAGAC R: TCTGCCAGTACCTCCTTGCT	151	JQ839263.1
<i>IL-8</i>	F: ACAGCAGTAACAACAACAAG R: GACCAGCACAGGAATGAG	117	NM_213867.1
<i>TNF-α</i>	F: TCCAATGGCAGAGTGGGTATG R: AGCTGGTTGTCTTTCAGCTTCAC	67	NM_214022.1
<i>β-Actin</i>	F: TCGCGGACATCAAGGAGAAG R: AGTTGAAGGTGGTCTCGTGG	194	AF017079.1




FIGURE 1: Effects of H₂O₂ on cell viability and lactate dehydrogenase activity in IPEC-1 cells. (a) Cell viability. (b) LDH activity. ^{AB}Means within a figure with different letters indicate H₂O₂ treatment and the control group differs significantly ($P \leq 0.05$).

H₂O₂ to treat IPEC-1 cells. After incubation for 3 h, we collected the cells for Western blot analysis. Cells were then lysed and centrifuged, and the supernatants were collected for western blot and protein assay according to our previous study [16]. After transmembrane electrophoresis, blots were incubated with primary antibodies against ERK, p38, JNK, phospho-ERK (p-ERK), phospho-p38 (p-p38), and phospho-JNK1/2 (p-JNK1/2) (Cell Signaling Technology, Inc., Danvers, MA, USA) overnight at 4°C and then incubated with a secondary antibody (Santa Cruz Biotechnology, Santa Cruz, CA, USA) at 25°C for 2 h. The blots were visualized with an enhanced Chemiluminescence Western Blot Kit (Amersham Biosciences) and processed with the Quantity One® software (Bio-Rad, CA, USA). The results were expressed as the abundance of p-ERK, p-p38, and p-JNK1/2 relative to the total protein content of ERK, p38, and JNK, respectively.

2.7. Immunofluorescence. IPEC-1 cells were seeded onto 12-well glass coverslips (Corning, MA, USA). Cells were then exposed to H₂O₂ (0 or 0.5 mM) for 3 h when the cells reached 70%-80% confluency. After fixing and blocking, the cells were then incubated with primary rabbit anti-NF- κ B p65 antibody overnight at 4°C, followed by incubation with the secondary Cy3-labelled antibody in a humidified chamber for 1 h at room temperature. Subsequently, the cells were

stained with 2-(4-amidinophenyl)-6-indolecarbamide dihydrochloride solution. Each of the steps above was followed by washing three times for 5 mins each in cold PBS. Finally, the IPEC-1 cells on coverslips were preserved in antifade mounting medium. The activation and nuclear translocation of NF- κ B were observed using a fluorescent microscope (Olympus, Tokyo, Japan).

2.8. Confocal Laser Scanning Microscope Analysis. IPEC-1 cells were seeded onto glass microscope coverslips (Corning, MA, USA). To assess the effect of inhibitors on H₂O₂-induced barrier function, cells were pretreated with SB202190, PD98059, or Bay11-7082 for 1 h and then challenged with H₂O₂ when cells reached 70%-80% confluency. All cells were fixed with 4% paraformaldehyde for 10 min at room temperature (RT) and permeabilized with 0.2% Triton X-100 in PBS at RT for 10 min. The cells were blocked with 1% bovine serum albumin in PBS for 30 min at RT and then incubated with anti-claudin-1 antibody overnight. After being washed five times in PBS, the cells were incubated for 2 h at RT with a goat anti-rabbit secondary antibody conjugated to Alexa Fluor 488 (Invitrogen, CA, USA), followed by counterstaining with 4,6-diamidino-2-phenylindole (Sigma-Aldrich, St. Louis, MO, USA). The coverslips were mounted onto glass microscope slides using mounting buffer, and the stained cells were visualized using a fluorescent




FIGURE 2: Effect of H₂O₂ on tight junction protein claudin-1 distribution in IPEC-1 cells. (a) Representative picture of claudin-1 protein distribution. (b) Quantification of claudin-1. DAPI: 4',6-diamidino-2-phenylindole (blue); FITC: fluorescein isothiocyanate (green). The cells were examined using a confocal laser microscope at 60x magnification. ^{AB}Means within a figure with different letters indicate H₂O₂ treatment and the control group differs significantly ($P \leq 0.05$).

TABLE 2: Effect of H₂O₂ on the mRNA expression of proinflammatory cytokines in IPEC-1 cells.

Items ¹	Control	0.5 mM H ₂ O ₂	<i>P</i> value
IL-6	1.00 ± 0.05 ^a	4.89 ± 0.51 ^b	<0.001
IL-8	1.00 ± 0.05 ^a	4.19 ± 0.14 ^b	<0.001
TNF- α	1.00 ± 0.03 ^a	6.51 ± 0.10 ^b	<0.001

^{ab}Means within a row differ significantly ($P \leq 0.05$). ¹The $2^{-\Delta\Delta Ct}$ method was used to analyze the relative gene expression (fold changes), which was calculated relative to the values in samples from the control group.

confocal laser scanning microscope (Carl Zeiss, Gottingen, Germany).

2.9. Statistical Analysis. Data were analyzed by Student's *t*-test or ANOVA (IBM Institute, Chicago, IL, USA). For the inhibitor trial, the effects of inhibitors and H₂O₂ were analyzed as a 2 × 2 factorial arrangement

TABLE 3: Effect of H₂O₂ on the protein expression of proinflammatory cytokines in IPEC-1 cells.

Items	Control	0.5 mM H ₂ O ₂	<i>P</i> value
IL-6 (pg/mL)	10.04 ± 1.14 ^a	13.40 ± 1.49 ^b	0.017
IL-8 (pg/mL)	15.85 ± 0.67 ^a	219.31 ± 2.14 ^b	<0.001
TNF- α (pg/mL)	7.88 ± 0.38 ^a	10.99 ± 0.49 ^b	<0.001

^{ab}Means within a row differ significantly ($P \leq 0.05$).

using the general linear model (GLM) procedures. All data were presented as means with standard errors of means. The statistical model included the effects of inhibitors, H₂O₂, and their interactions. When significant inhibitor × H₂O₂ interaction or a trend for inhibitor × H₂O₂ interaction occurred, multiple comparison tests were performed using ANOVA. All data were represented by at least three independent experiments. Differences were considered significant for values of $P \leq 0.05$.




FIGURE 3: Effect of H₂O₂ on p38 (a), JNK (b), and ERK (c) signaling pathway activation (phosphorylation) in IPEC-1 cells. P-MAPK/total MAPK was the ratio reflected by the bars. Values are means (six replicates) with standard errors of means represented by vertical bars. ^{AB}Means within a figure with different letters indicate H₂O₂ treatment and the control group differs significantly ($P \leq 0.05$).

3. Results

3.1. H₂O₂ Induces Cell Injury in IPEC-1 Cells

3.1.1. Cell Viability. To explore the effect of H₂O₂ on cell injury, we used the CCK-8 kit to determine cell viability after treatment with 0, 0.2 mM, and 0.5 mM H₂O₂ for 3 h. The dose and time point used were determined based on our preliminary experiments (data not shown). Our results demonstrated that 0.2 mM and 0.5 mM H₂O₂ stimulation significantly decreased cell viability in IPEC-1 cells (Figure 1(a)).

3.1.2. LDH. This enzyme is found in virtually all living cells, is released extracellularly when cells are damaged, and is commonly used as a marker of cell injury. In our study, compared with control cells, 0.5 mM H₂O₂ stimulation significantly increased the LDH activity in the cell supernatant (Figure 1(b)). This data, in combination with a cell viability indicator, led us to use 0.5 mM of H₂O₂ in all subsequent experiments.

3.1.3. Tight Junction Protein. To explore the effect of H₂O₂ on cell barrier function, we also assessed the distribution of the tight junction protein claudin-1 in IPEC-1 cells, using confocal microscopy. Claudin-1 protein is one of the most important proteins associated with tight junctions. In the control cells, it is normally distributed uniformly across the cell membrane. Under external stimulation, however, it becomes nonuniformly distributed, both inter and extracellularly. Our results demonstrated that 0.5 mM H₂O₂ stimulation for 3 h significantly altered the distribution of claudin-1 in IPEC-1 cells (Figures 2(a) and 2(b)), suggesting a role for H₂O₂ in the destruction of the cell barrier.

3.2. H₂O₂ Upregulates mRNA and Protein Expression of Inflammatory-Related Genes in IPEC-1 Cells. To verify the

effect of H₂O₂ on the inflammatory response, we measured the gene and protein expression levels of the proinflammatory cytokines IL-6, IL-8, and TNF- α using RT-PCR. Our results demonstrated that compared with the control cells, 0.5 mM H₂O₂ significantly increased the mRNA and protein expression of IL-6, IL-8, and TNF- α when compared to control (Tables 2 and 3).

3.3. H₂O₂ Activates MAPK (p38 and JNK Phosphorylation) and NF- κ B p65 Signaling in IPEC-1 Cells. To evaluate the activation of MAPK and NF- κ B signaling pathways after H₂O₂ stimulation, we measured p38, ERK, and JNK phosphorylation and NF- κ B p65 protein translocation by western blot analysis and immunofluorescence after incubation with 0.5 mM H₂O₂ for 3 h. As demonstrated in Figures 3(a) and 3(b), H₂O₂ stimulation significantly increased the phosphorylation of p38 and JNK in IPEC-1 cells compared with control cells. However, H₂O₂ challenge had no significant effect on the phosphorylation of ERK and total protein levels for the MAPKs (p38, ERK, and JNK) (Figure 3(c)). Our data also demonstrated that H₂O₂ treatment increased NF- κ B p65 protein entering the nucleus of IPEC-1 cells (Figures 4(a) and 4(b)). These data demonstrated that H₂O₂ activated MAPK (p38 and JNK phosphorylation) and NF- κ B p65 signaling pathways in IPEC-1 cells after 3 h incubation.

3.4. Inhibition of p38, JNK, and NF- κ B with Bay11-7082, SB202190, and PD98059 Downregulates mRNA and Protein Expression of Proinflammatory Cytokine-Related Genes in IPEC-1 Cells. To determine the role of p38, JNK, and NF- κ B signaling in inducing the inflammatory response after H₂O₂ stimulation, cells were pretreated with specific inhibitors (SB202190, PD98059, or Bay11-7082) or DMSO as control. The time points and concentrations of inhibitors were selected after preliminary time and dose experiments (data




FIGURE 4: Effect of H₂O₂ on NF-κB p65 protein translocation in IPEC-1 cells. NF-κB p65 protein translocation in the IPEC-1 cells challenged with H₂O₂. (a) Representative picture of p65 protein translocation (immunofluorescence; ×100). (b) Quantification of p65 translocation. DAPI: 4',6-diamidino-2-phenylindole; cy3: cyanines. ^{AB}Means within a figure with different letters indicate H₂O₂ treatment and the control group differs significantly ($P \leq 0.05$).

TABLE 4: Effect of Bay11-7082 on mRNA expression of proinflammatory cytokines in IPEC-1 cells stimulated with H₂O₂.

Items ¹	-H ₂ O ₂		+H ₂ O ₂			P value H ₂ O ₂	Interactions
	Control	Bay11-7082	Control	Bay11-7082	Bay11-7082		
IL-6	1.00 ± 0.10	0.50 ± 0.07	1.27 ± 0.02	0.78 ± 0.03	<0.001	0.002	0.898
IL-8	1.00 ± 0.03 ^b	0.63 ± 0.02 ^a	2.48 ± 0.15 ^c	0.52 ± 0.06 ^a	<0.001	<0.001	<0.001
TNF-α	1.00 ± 0.05 ^a	0.80 ± 0.08 ^a	15.75 ± 0.7 ^c	3.05 ± 0.18 ^b	<0.001	<0.001	<0.001

^{a-c}Means within a row with different letters differ significantly ($P \leq 0.05$). ¹The $2^{-\Delta\Delta Ct}$ method was used to analyze the relative mRNA expression (fold changes), calculated relative to the values in the control group.

not shown). Cells were pretreated with the p38 inhibitor SB202190 (20 μM), JNK inhibitor PD98059 (50 μM), or the NF-κB inhibitor Bay11-7082 (10 μM) for 1 h prior to H₂O₂ stimulation. As depicted in Tables 4, 5, and 6, after 3 h incubation, H₂O₂ challenge significantly increased the mRNA expression of IL-6, IL-8, and TNF-α. However, inhibition of p38, JNK, and NF-κB signaling pathways

with Bay11-7082 (Tables 4 and 7), SB202190 (Tables 5 and 8), or PD98059 (Tables 6 and 9) downregulated the mRNA and protein expression of the inflammatory-related genes IL-6, IL-8, and TNF-α in IPEC-1 cells. Collectively, the data suggested that the p38, JNK, and NF-κB signaling pathways played an important role in the induction of cell inflammation.

TABLE 5: Effect of SB202190 on the mRNA expression of proinflammatory cytokines in IPEC-1 cells stimulated with H₂O₂.

Items ¹	-H ₂ O ₂		+H ₂ O ₂			P value H ₂ O ₂	Interactions
	Control	SB202190	Control	SB202190	SB202190		
IL-6	1.00 ± 0.05 ^b	0.29 ± 0.01 ^a	5.33 ± 0.28 ^d	2.31 ± 0.14 ^c	<0.001	<0.001	<0.001
IL-8	1.00 ± 0.09	0.54 ± 0.01	4.61 ± 0.58	3.04 ± 0.22	0.012	<0.001	0.112
TNF- α	1.00 ± 0.10 ^a	0.86 ± 0.14 ^a	6.09 ± 0.35 ^c	2.84 ± 0.21 ^b	<0.001	<0.001	<0.001

^{a-d}Means within a row with different letters differ significantly ($P \leq 0.05$). ¹The 2^{- $\Delta\Delta C_t$} method was used to analyze the relative mRNA expression (fold changes), calculated relative to the values in the control group.

TABLE 6: Effect of PD98059 on the mRNA expression of proinflammatory cytokines in IPEC-1 cells stimulated with H₂O₂.

Items ¹	-H ₂ O ₂		+H ₂ O ₂			P value H ₂ O ₂	Interactions
	Control	PD98059	Control	PD98059	SB202190		
IL-6	1.00 ± 0.11 ^a	0.82 ± 0.1 ^a	1.69 ± 0.11 ^b	1.24 ± 0.14 ^a	0.001	<0.001	<0.042
IL-8	1.00 ± 0.24 ^a	0.88 ± 0.03 ^a	27.90 ± 1.72 ^c	22.64 ± 2.85 ^b	0.024	<0.001	0.029
TNF- α	1.00 ± 0.10 ^a	0.85 ± 0.23 ^a	41.63 ± 5.69 ^c	10.86 ± 1.74 ^b	<0.001	<0.001	<0.001

^{a-c}Means within a row with different letters differ significantly ($P \leq 0.05$). ¹The 2^{- $\Delta\Delta C_t$} method was used to analyze the relative mRNA expression (fold changes), calculated relative to the values in the control group.

TABLE 7: Effect of Bay11-7082 on protein expression of proinflammatory cytokines in IPEC-1 cells stimulated with H₂O₂.

Items	-H ₂ O ₂		+H ₂ O ₂			P value H ₂ O ₂	Interactions
	Control	Bay11-7082	Control	Bay11-7082	Bay11-7082		
IL-6 (pg/mL)	11.54 ± 0.82 ^b	12.03 ± 1.27 ^b	225.26 ± 2.02 ^a	14.91 ± 2.08 ^b	0.008	<0.001	0.004
IL-8 (pg/mL)	14.90 ± 0.97 ^d	61.05 ± 5.89 ^c	2222.20 ± 7.75 ^a	119.30 ± 4.58 ^b	<0.001	<0.001	<0.001
TNF- α (pg/mL)	5.55 ± 0.35 ^{ab}	3.13 ± 0.32 ^b	8.65 ± 1.95 ^a	5.93 ± 1.26 ^{ab}	0.047	0.025	0.917

^{abcd}Means within a row differ significantly ($P \leq 0.05$).

TABLE 8: Effect of SB202190 on the protein expression of proinflammatory cytokines in IPEC-1 cells stimulated with H₂O₂.

Items	-H ₂ O ₂		+H ₂ O ₂			P value H ₂ O ₂	Interactions
	Control	SB202190	Control	SB202190	SB202190		
IL-6 (pg/mL)	12.29 ± 1.55 ^b	14.34 ± 1.28 ^b	26.11 ± 2.27 ^a	14.91 ± 1.08 ^b	0.012	<0.001	0.001
IL-8 (pg/mL)	17.80 ± 3.16 ^c	25.53 ± 2.39 ^c	228.08 ± 7.45 ^a	71.83 ± 2.72 ^b	<0.001	<0.001	<0.001
TNF- α (pg/mL)	4.93 ± 0.58 ^b	2.60 ± 0.23 ^b	8.40 ± 1.62 ^a	2.88 ± 0.54 ^b	0.001	0.026	0.049

^{ab}Means within a row differ significantly ($P \leq 0.05$).

TABLE 9: Effect of PD98059 on the protein expression of proinflammatory cytokines in IPEC-1 cells stimulated with H₂O₂.

Items	-H ₂ O ₂		+H ₂ O ₂			P value H ₂ O ₂	Interactions
	Control	PD98059	Control	PD98059	SB202190		
IL-6 (pg/mL)	11.91 ± 1.04 ^b	15.34 ± 1.42 ^b	25.69 ± 1.60 ^a	16.96 ± 0.82 ^b	0.047	<0.001	<0.001
IL-8 (pg/mL)	16.35 ± 1.78 ^c	18.02 ± 0.78 ^c	224.54 ± 8.28 ^a	76.97 ± 3.64 ^b	<0.001	<0.001	<0.001
TNF- α (pg/mL)	5.24 ± 0.36 ^b	3.40 ± 0.38 ^b	8.50 ± 1.33 ^a	4.51 ± 0.42 ^b	<0.001	<0.001	<0.001

^{abc}Means within a row differ significantly ($P \leq 0.05$).

3.5. Inhibition of p38, JNK, and NF- κ B with Bay11-7082, SB202190, and PD98059 Does Not Prevent Cell Injury in IPEC-1 Cells

3.5.1. Cell Viability. To further determine a role for p38, JNK, and NF- κ B signaling in the induction of the cell injury after H₂O₂ stimulation, cells were pretreated with inhibitors (SB202190, PD98059, or Bay11-7082) 1 h prior to a 3 h incubation with H₂O₂. Our data demonstrated that H₂O₂ stimulation significantly decreased the viability of IPEC-1 cells

when compared to control. The inhibitors SB202190, PD98059, and Bay11-7082 did not have a protective effect on cell viability, suggesting that the inhibition of MAPK (p38 and JNK) and NF- κ B pathways did not prevent cell injury induced by H₂O₂ challenge in IPEC-1 cells (Figures 5(a), 5(b), and 5(c)). Bay11-7082 further decreased the cell viability after H₂O₂ stimulation.

3.5.2. LDH. To investigate whether p38, JNK, and NF- κ B signaling pathways influenced H₂O₂-induced LDH activity in




FIGURE 5: Effects of Bay11-7082, SB202190, and PD98059 on cell viability and LDH activity in the supernatant of IPEC-1 cells stimulated with H₂O₂. (a and d) Effects of Bay11-7082 on cell viability and LDH activity in the supernatant of IPEC-1 cells stimulated with H₂O₂. (b and e) Effects of SB202190 on cell viability and LDH activity in the supernatant of IPEC-1 cells stimulated with H₂O₂. (c and f) Effects of PD98059 on cell viability and LDH activity in the supernatant of IPEC-1 cells stimulated with H₂O₂. Values are means with standard errors of means represented by vertical bars. ^{ABC}Means within a figure with different letters indicate H₂O₂ treatment and the inhibitor group differs significantly ($P \leq 0.05$).

IPEC-1 cells, we pretreated cells with inhibitors (SB202190, PD98059, or Bay11-7082) for 1 h prior to the H₂O₂ incubation for 3 h. The results demonstrated that H₂O₂ stimulation significantly increased the activity of LDH in the IPEC-1 cell supernatant, and pretreatment with inhibitors (SB202190, PD98059, or Bay11-7082) did not alleviate the increase of LDH activity, indicating that inhibition of the MAPKs (p38 and JNK) and NF- κ B signaling pathways did not prevent the H₂O₂-induced LDH activity in the IPEC-1 cell supernatant (Figures 5(d), 5(e), and 5(f)). Bay11-7082 further increased LDH activity after H₂O₂ stimulation.

3.5.3. Tight Junction Protein. Next, we sought to identify the role of the p38, JNK, and NF- κ B signaling pathways in cell barrier function by using confocal microscopic analysis. As presented in Figures 6, 7, 8, and 9, H₂O₂ treatment significantly destroyed the normal distribution of claudin-1 protein in the cells. Pretreating cells with inhibitors (SB202190, PD98059, or Bay11-7082) 1 h prior to the H₂O₂ incubation had no effect on the distribution of claudin-1 after H₂O₂ challenge. This suggested that inhibition of the MAPKs (p38 and JNK) and NF- κ B signaling pathways did not prevent barrier function injury.

4. Discussion

The intestinal epithelium performs a dual function as it can absorb dietary nutrients and it can form a physical barrier

against noxious stimuli. As the first line of defense, intestinal epithelial cells also participate in multiple physiological activities including immune response and tissue renewal. Oxidative stress has been identified as a key factor in the breakdown of intestinal function in pigs [18]. H₂O₂ is a highly reactive oxygen species and serves as a signaling molecule in the regulation of a wide variety of biological processes [6–8]. H₂O₂ is often used in *in vitro* studies of redox-regulated processes because it is relatively stable *in vivo* compared to other reactive oxygen species.

Our results indicated that H₂O₂ stimulation significantly decreased cell viability and increased LDH activity in the IPEC-1 cell supernatant, demonstrating a role for H₂O₂ in cell injury. In agreement with our study, Zheng et al. found that incubation with H₂O₂ caused significant damage to nucleus pulposus cells, as indicated by an increase in apoptotic cells [19]. Similarly, H₂O₂ has been reported to decrease the cell viability and increase cell apoptosis in human umbilical vein endothelial cells [20]. H₂O₂ can also cause high death rates and induce chronic intestinal injury in mice, as demonstrated by increased apoptotic intestinal epithelial cells and damaged intestinal morphology [21]. The intestinal barrier is composed of a layer of columnar epithelium and intraepithelial tight junctions. Tight junction proteins, such as claudins, occludins, and zonula occludins, form the tight junction, work as a rate-limiting step in the paracellular pathway, and form a selectively permeable barrier [22]. We determined the distribution of tight junction claudin-1 and found




FIGURE 6: Effect of Bay11-7082 on the protein distribution of claudin-1 in IPEC-1 cells stimulated with H_2O_2 . DAPI: 4',6-diamidino-2-phenylindole (blue); FITC: fluorescein isothiocyanate (green). The cells were examined using a confocal laser microscope at 60x magnification.

that H_2O_2 challenge disrupted the tight junction distribution when compared to control, demonstrating a role for H_2O_2 in severe intestinal barrier dysfunction. Our study was in agreement with Ma et al. [23] who found H_2O_2 -induced barrier disruption in a monolayer of Caco-2 cells. An H_2O_2 -induced increase in renal epithelial cell paracellular permeability was also demonstrated, mediated by occludin protein, possibly by a reduction in the rate of occludin movement into the tight junction region [24]. It is known that H_2O_2 could cause the production of reactive oxygen species, resulting in oxidative damage of the intestinal barrier and cell apoptosis and necrosis [25, 26]. Therefore, it is possible that stimulation destroyed the IPEC-1 cell structure and then disrupted the distribution of the tight junction protein in the cell membrane.

Oxidative stress is an important driving force for an enhanced proinflammatory cytokine production in intestinal epithelial cells, causing gut mucosal inflammation. Several

cytokines are now known to contribute to tissue injury and cell death, including $\text{TNF-}\alpha$, interferon- γ , $\text{IL-1}\beta$, and IL-18 [27]. In the present study, we found that 0.5 mM H_2O_2 significantly increased the mRNA and protein expression of IL-6 , IL-8 , and $\text{TNF-}\alpha$. Previous studies have reported that acute oxidative stress affected IL-8 and $\text{TNF-}\alpha$ expression in IPEC-J2 porcine epithelial cells [28]. It has been found that short-term H_2O_2 and $\text{TNF-}\alpha$ incubation induced IL-8 secretion in Caco-2 and HT-29 cells [29].

We further explored the signaling pathways involved in proinflammatory cytokine regulation. It is well known that inflammation can result in tissue injury through multiple signaling pathways. $\text{NF-}\kappa\text{B}$ signaling is a key pathway whose activation can lead to proinflammatory cytokine production. MAPKs are a family of serine threonine kinases and are able to transduce signals from a diverse array of extracellular stimuli including oxidative stress and cytotoxic factors [30]. The extracellular regulated kinases (ERK1/2), p38 MAPK,




FIGURE 7: Effect of SB202190 on the protein distribution of claudin-1 in IPEC-1 cells stimulated with H_2O_2 . DAPI: 4',6-diamidino-2-phenylindole (blue); FITC: fluorescein isothiocyanate (green). The cells were examined using a confocal laser microscope at 60x magnification.

and JNK represent the three primary MAPK signaling pathways. We found that H_2O_2 stimulation significantly enhanced NF- κB pathway activation and increased its translocation to the nucleus. As well as canonical NF- κB pathways, MAPK pathways have been reported to be involved in proinflammatory cytokine regulation [31]. We also detected MAPK activation in this study. Our results demonstrated that H_2O_2 stimulation activated the phosphorylation of p38 and JNK in IPEC-1 cells but had no significant effect on phosphorylation of ERK. In agreement with our results, Zhou et al. [32] reported that H_2O_2 stimulation activated the p38 signaling pathway in mouse intestinal epithelial cells, potentially causing apoptosis. H_2O_2 stimulation has been reported to activate JNK and nuclear NF- κB phosphorylation in human HepG2 cells [33]. The majority of studies have demonstrated that JNK activation ultimately leads to cellular damage. Therefore, it may be possible that activation of

MAPKs and NF- κB signaling pathways contribute to the cell damage associated with H_2O_2 .

To further explore the underlying mechanism of cell injury induced by H_2O_2 , we used MAPK and NF- κB inhibitors to block their signaling pathways in order to evaluate their roles in cell injury. As expected, pretreatment with the p38 inhibitor SB202190, JNK inhibitor PD98059, and NF- κB inhibitor (Bay11-7082) significantly decreased IL-6, IL-8, and TNF- α gene expression. Similarly, other studies have demonstrated that p38 and JNK signaling pathways could activate proinflammatory cytokine release in numerous cell types [34–37]. It was found by Kim and Lee that inhibition of p38 and JNK signaling reduced the lipopolysaccharide-induced production of inflammatory mediators in keratinocytes [37]. It was also reported that inhibition of the p38 signaling pathway could suppress apoptosis and expression of proinflammatory cytokines in human chondrocytes [38].




FIGURE 8: Effect of PD98059 on the protein distribution of claudin-1 in IPEC-1 cells stimulated with H₂O₂. DAPI: 4',6-diamidino-2-phenylindole (blue); FITC: fluorescein isothiocyanate (green). The cells were examined using a confocal laser microscope at 60x magnification.




FIGURE 9: Quantitation of fluorescence intensity of claudin-1 in IPEC-1 cells after NF- κ B and MAPK inhibitor treatment. (a) Effect of Bay11-7082 on the protein quantitation of claudin-1 in IPEC-1 cells stimulated with H₂O₂. (b) Effect of SB202190 on the protein quantitation of claudin-1 in IPEC-1 cells stimulated with H₂O₂. (c) Effect of PD98059 on the protein quantitation of claudin-1 in IPEC-1 cells stimulated with H₂O₂. ^{ABC}Means within a figure with different letters indicate H₂O₂ treatment and the inhibitor group differs significantly ($P \leq 0.05$).

Our results demonstrated that inhibition of MAPKs and NF- κ B signaling pathways could prevent the cell inflammatory response induced by H₂O₂ stimulation.

To further determine whether p38, JNK, and NF- κ B signaling pathways are important for the induction of the cell injury after H₂O₂ stimulation, we measured cell injury after treatment with MAPK and NF- κ B inhibitors. Unexpectedly, we found that inhibiting MAPK and NF- κ B signaling pathways did not attenuate the cell injury as indicated by decreased cell viability and increased LDH in the cell supernatant. Bay11-7082 further decreased the cell viability after H₂O₂ stimulation. That is because Bay11-7082 can inhibit the cell viability and promote apoptosis. These data suggested that p38, JNK, and NF- κ B signaling pathways were not the principle pathways mediating cell injury in IPEC-1 cells. In fact, cell death *in vivo* can trigger an inflammatory response. Damaged cells release “danger signals” that alert the host to cell death. Some of these molecules are recognized by cellular receptors that can stimulate the generation of proinflammatory mediators [39]. Dead cells can also release danger signals activating dendritic cells and promoting the generation of an immune response to antigens in and around the dying cells. For example, factor-related apoptosis suicide (FAS) is a prototypical death receptor which stimulates cells to undergo apoptosis when activated [40]. Injection of an agonistic form of an anti-FAS antibody into mice causes hepatocytes (which express Fas) to undergo apoptosis and generate a very strong inflammatory response [37]. If apoptosis is blocked, however, this inflammatory response is inhibited [41]. It is possible that the stressful stimuli (oxidative stress) firstly lead to cell damage and then induce an inflammatory response. Our results led us to speculate that proinflammatory cytokines were not the factors leading to cell injury in IPEC-1 cells after H₂O₂ stimulation. It is also known that dead cells can induce inflammation by releasing proinflammatory signal(s). Taking the inflammation index into consideration, it is possible that, in our study, H₂O₂ first induced cell death and then induced a secondary release of proinflammatory cytokines.

From our results, H₂O₂ caused a cell inflammatory response through the activation of MAPK and NF- κ B pathways in IPEC-1 cells. However, MAPK and NF- κ B pathways were not the specific pathways involved in the mediation of cell injury. Many other pathways such as those involved in apoptosis, necrosis, and autophagy may be involved in the cell damage induced by H₂O₂ [42]. There is a need for further research to determine the contribution of cell death in the regulation of inflammation and to improve our ability to distinguish these cytotoxic mechanisms from other pathways by which death-related molecules regulate inflammation. Such progress will provide us with better tools with which to clarify the role of cell death in inflammation. It will also pave the way for the development of new therapies whereby cell death can be targeted and manipulated, ultimately modulating the course of inflammation.

5. Conclusion

In conclusion, the present study demonstrates that H₂O₂ stimulation can cause IPEC-1 cell injury and alter the distri-

bution of tight junction proteins, leading to the inflammatory response and activation of p38, JNK, and NF- κ B signaling pathways. Inhibiting p38, JNK, and NF- κ B signaling pathways prevents the cell inflammatory response but does not alleviate the cell injury. Therefore, it is suggested that H₂O₂ stimulation firstly leads to cell injury, followed by an inflammatory response in IPEC-1 cells.

Data Availability

The data used to support the findings of this study are included within the article.

Disclosure

Kan Xiao and Congcong Liu are co-first authors.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Authors' Contributions

The authors' contributions are as follows: Yulan Liu designed the research; Yulan Liu, Kan Xiao, Zhixiao Tu, Congcong Liu, Qiao Xu, Yang Zhang, and Xiuying Wang conducted the research; Yulan Liu, Kan Xiao, Zhixiao Tu, and Jing Zhang analyzed the data; Yulan Liu and Kan Xiao wrote the article; Yulan Liu, Kan Xiao, and Chien-An Andy Hu edited and revised the manuscript; and Yulan Liu had the primary responsibility for the final content. All authors read and approved the final manuscript.

Acknowledgments




The present study was supported by the Project of Innovative Research Groups of the Natural Science Foundation of Hubei Province (2019CFA015) and the Project of the Hubei Provincial Department of Education (T201508).

References

- [1] A. T. Blikslager, A. J. Moeser, J. L. Gookin, S. L. Jones, and J. Odle, “Restoration of barrier function in injured intestinal mucosa,” *Physiological Reviews*, vol. 87, no. 2, pp. 545–564, 2007.
- [2] P. J. A. Wijtten, J. . . Meulen, and M. W. A. Verstegen, “Intestinal barrier function and absorption in pigs after weaning: a review,” *British Journal of Nutrition*, vol. 105, no. 7, pp. 967–981, 2011.
- [3] B. Wang, G. Wu, Z. Zhou et al., “Glutamine and intestinal barrier function,” *Amino Acids*, vol. 47, no. 10, pp. 2143–2154, 2015.
- [4] S. T. Ballard, J. H. Hunter, and A. E. Taylor, “Regulation of tight-junction permeability during nutrient absorption across the intestinal epithelium,” *Annual Review of Nutrition*, vol. 15, no. 1, pp. 35–55, 1995.
- [5] J. R. Turner, “Intestinal mucosal barrier function in health and disease,” *Nature Reviews Immunology*, vol. 9, no. 11, pp. 799–809, 2009.

- [6] A. D. Bokare and W. Choi, "Review of iron-free Fenton-like systems for activating H_2O_2 in advanced oxidation processes," *Journal of Hazardous Materials*, vol. 275, pp. 121–135, 2014.
- [7] I. A. Pomytkin, " H_2O_2 signalling pathway: a possible bridge between insulin receptor and mitochondria," *Current Neuropharmacology*, vol. 10, no. 4, pp. 311–320, 2012.
- [8] H. Sies, "Role of metabolic H_2O_2 generation: redox signaling and oxidative stress," *Journal of Biological Chemistry*, vol. 289, no. 13, pp. 8735–8741, 2014.
- [9] J. Zhou, J. Wang, X. Li et al., " H_2O_2 mediates the crosstalk of brassinosteroid and abscisic acid in tomato responses to heat and oxidative stresses," *Journal of Experimental Botany*, vol. 65, no. 15, pp. 4371–4383, 2014.
- [10] J. R. Burgoyne, S. Oka, N. Ale-Agha, and P. Eaton, "Hydrogen peroxide sensing and signaling by protein kinases in the cardiovascular system," *Antioxidants & Redox Signaling*, vol. 18, no. 9, pp. 1042–1052, 2013.
- [11] Y. Saito, K. Nishio, Y. Ogawa et al., "Turning point in apoptosis/necrosis induced by hydrogen peroxide," *Free Radical Research*, vol. 40, no. 6, pp. 619–630, 2006.
- [12] M. Strus, A. Janczyk, A. Gonet-Surówka et al., "Effect of hydrogen peroxide of bacterial origin on apoptosis and necrosis of gut mucosa epithelial cells as a possible pathomechanism of inflammatory bowel disease and cancer," *Journal of Physiology and Pharmacology*, vol. 60, no. 6, pp. 55–60, 2009.
- [13] X. Cai, X. Chen, X. Wang et al., "Pre-protective effect of lipoic acid on injury induced by H_2O_2 in IPEC-J2 cells," *Molecular and Cellular Biochemistry*, vol. 378, no. 1–2, pp. 73–81, 2013.
- [14] A. Bhattacharyya, R. Chattopadhyay, S. Mitra, and S. E. Crowe, "Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases," *Physiological Reviews*, vol. 94, no. 2, pp. 329–354, 2014.
- [15] J. Park, J. S. Min, B. Kim et al., "Mitochondrial ROS govern the LPS-induced pro-inflammatory response in microglia cells by regulating MAPK and NF- κ B pathways," *Neuroscience Letters*, vol. 584, pp. 191–196, 2015.
- [16] Y. Liu, F. Chen, J. Odle et al., "Fish oil enhances intestinal integrity and inhibits TLR4 and NOD2 signaling pathways in weaned pigs after LPS challenge," *The Journal of Nutrition*, vol. 142, no. 11, pp. 2017–2024, 2012.
- [17] I. Jorgensen, M. Rayamajhi, and E. A. Miao, "Programmed cell death as a defence against infection," *Nature Reviews Immunology*, vol. 17, no. 3, pp. 151–164, 2017.
- [18] L. H. Zhu, K. L. Zhao, X. L. Chen, and J. X. Xu, "Impact of weaning and an antioxidant blend on intestinal barrier function and antioxidant status in pigs," *Journal of Animal Science*, vol. 90, no. 8, pp. 2581–2589, 2012.
- [19] W. W. Zheng, M. Yang, C. Wu, and X. L. Su, "Protective effects of puerarin on injury of nucleus pulposus cells induced by hydrogen peroxide through inhibiting inflammatory response, oxidative stress and anti-apoptotic pathway," *International Journal of Clinical and Experimental Medicine*, vol. 9, no. 9, pp. 17258–17266, 2016.
- [20] R. Han, F. Tang, M. Lu et al., "Astragalus polysaccharide ameliorates H_2O_2 -induced human umbilical vein endothelial cell injury," *Molecular Medicine Reports*, vol. 15, no. 6, pp. 4027–4034, 2017.
- [21] Z. J. Cui, J. Yin, L. Wang, L. He, Y. Yin, and T. Li, "Effects of pro-inflammatory cytokines and antioxidants expression in the jejunum of mice induced by hydrogen peroxide," *International Immunopharmacology*, vol. 31, pp. 9–14, 2016.
- [22] X. Li, S. Akhtar, and M. A. Choudhry, "Alteration in intestine tight junction protein phosphorylation and apoptosis is associated with increase in IL-18 levels following alcohol intoxication and burn injury," *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, vol. 1822, no. 2, pp. 196–203, 2012.
- [23] C. Ma, L. G. Shi, H. S. Zhang et al., "Salvianolic acid B ameliorates hydrogen peroxide-induced barrier disruption in monolayer Caco-2 cells," *International Journal of Clinical and Experimental Pathology*, vol. 9, no. 9, pp. 9018–9027, 2016.
- [24] D. Janosevic, J. Axis, R. L. Bacallao, and K. Amsler, "Occludin content modulates hydrogen peroxide-induced increase in renal epithelial paracellular permeability," *Journal of Cellular Biochemistry*, vol. 117, no. 3, pp. 769–779, 2016.
- [25] R. Yang, Q. Hui, Q. Jiang et al., "Effect of manitoba-grown red-osier dogwood extracts on recovering Caco-2 cells from H_2O_2 -induced oxidative damage," *Antioxidants*, vol. 8, no. 8, p. 250, 2019.
- [26] A. Iraha, H. Chinen, A. Hokama et al., "Fucoidan enhances intestinal barrier function by upregulating the expression of claudin-1," *World Journal of Gastroenterology*, vol. 19, no. 33, pp. 5500–5507, 2013.
- [27] D. Wallach, A. Kovalenko, and M. Feldmann, "Advances in TNF family research," in *Proceedings of the 12th International TNF Conference*, New York, NY, USA, 2011.
- [28] E. Paszti-Gere, E. Csibrik-Nemeth, K. Szeker, R. Csizinszky, C. Jakab, and P. Galfi, "Acute oxidative stress affects IL-8 and TNF- α expression in IPEC-J2 porcine epithelial cells," *Inflammation*, vol. 35, no. 3, pp. 994–1004, 2012.
- [29] D. O. Son, H. Satsu, and M. Shimizu, "Histidine inhibits oxidative stress- and TNF- α -induced interleukin-8 secretion in intestinal epithelial cells," *FEBS Letters*, vol. 579, no. 21, pp. 4671–4677, 2005.
- [30] D. E. Shifflett, S. L. Jones, A. J. Moeser, and A. T. Blikslager, "Mitogen-activated protein kinases regulate COX-2 and mucosal recovery in ischemic-injured porcine ileum," *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 286, no. 6, pp. G906–G913, 2004.
- [31] H. K. Koul, M. Pal, and S. Koul, "Role of p38 MAP kinase signal transduction in solid tumors," *Genes & Cancer*, vol. 4, no. 9–10, pp. 342–359, 2013.
- [32] Y. Zhou, Q. Wang, B. Mark Evers, and D. H. Chung, "Oxidative stress-induced intestinal epithelial cell apoptosis is mediated by p38 MAPK," *Biochemical and Biophysical Research Communications*, vol. 350, no. 4, pp. 860–865, 2006.
- [33] X. He, C. Y. Liu, J. F. Yin, A. H. Jin, X. Z. Yin, and J. S. Quan, "Rhaponticum uniflorum inhibits H_2O_2 -induced apoptosis of liver cells via JNK and NF- κ B pathways," *China Journal of Chinese Materia Medica*, vol. 42, no. 6, pp. 1189–1193, 2017.
- [34] D. Couto, M. Freitas, G. Porto et al., "Polyacrylic acid-coated and non-coated iron oxide nanoparticles induce cytokine activation in human blood cells through TAK1, p38 MAPK and JNK pro-inflammatory pathways," *Archives of Toxicology*, vol. 89, no. 10, pp. 1759–1769, 2015.
- [35] X. T. Wu, Z. Yang, A. R. Ansari et al., "Visfatin regulates the production of lipopolysaccharide-induced inflammatory cytokines through p38 signaling in murine macrophages," *Microbial Pathogenesis*, vol. 117, pp. 55–59, 2018.
- [36] Y. T. Yeung, N. S. Bryce, S. Adams et al., "p38 MAPK inhibitors attenuate pro-inflammatory cytokine production and the invasiveness of human U251 glioblastoma cells," *Journal of Neuro-Oncology*, vol. 109, no. 1, pp. 35–44, 2012.

- [37] A. Kim and C. S. Lee, "Apigenin reduces the Toll-like receptor-4-dependent activation of NF- κ B by suppressing the Akt, mTOR, JNK, and p38-MAPK," *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 391, no. 3, pp. 271–283, 2018.
- [38] H. Y. Sun, K. Z. Hu, and Z. S. Yin, "Inhibition of the p38-MAPK signaling pathway suppresses the apoptosis and expression of proinflammatory cytokines in human osteoarthritis chondrocytes," *Cytokine*, vol. 90, pp. 135–143, 2017.
- [39] D. Wallach, T. B. Kang, and A. Kovalenko, "Concepts of tissue injury and cell death in inflammation: a historical perspective," *Nature Reviews Immunology*, vol. 14, no. 1, pp. 51–59, 2014.
- [40] T. Nunes, C. Bernardazzi, and H. S. de Souza, "Cell death and inflammatory bowel diseases: apoptosis, necrosis, and autophagy in the intestinal epithelium," *BioMed Research International*, vol. 2014, 12 pages, 2014.
- [41] S. Faouzi, B. E. Burckhardt, J. C. Hanson et al., "Anti-Fas induces hepatic chemokines and promotes inflammation by an NF- κ B-independent, caspase-3-dependent pathway," *Journal of Biological Chemistry*, vol. 276, no. 52, pp. 49077–49082, 2001.
- [42] M. Sendler, J. Mayerle, and M. M. Lerch, "Necrosis, Apoptosis, Necroptosis, Pyroptosis: It Matters How Acinar Cells Die During Pancreatitis," *Cellular and Molecular Gastroenterology and Hepatology*, vol. 2, no. 4, pp. 407–408, 2016.



Hindawi

Submit your manuscripts at
www.hindawi.com

