

## *Retraction*

# **Retracted: MKP-1 Overexpression Reduces Postischemic Myocardial Damage through Attenuation of ER Stress and Mitochondrial Damage**

### **Oxidative Medicine and Cellular Longevity**

Received 10 October 2023; Accepted 10 October 2023; Published 11 October 2023

Copyright © 2023 Oxidative Medicine and Cellular Longevity. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been retracted by Hindawi following an investigation undertaken by the publisher [1]. This investigation has uncovered evidence of one or more of the following indicators of systematic manipulation of the publication process:

- (1) Discrepancies in scope
- (2) Discrepancies in the description of the research reported
- (3) Discrepancies between the availability of data and the research described
- (4) Inappropriate citations
- (5) Incoherent, meaningless and/or irrelevant content included in the article
- (6) Peer-review manipulation

The presence of these indicators undermines our confidence in the integrity of the article's content and we cannot, therefore, vouch for its reliability. Please note that this notice is intended solely to alert readers that the content of this article is unreliable. We have not investigated whether authors were aware of or involved in the systematic manipulation of the publication process.

Wiley and Hindawi regrets that the usual quality checks did not identify these issues before publication and have since put additional measures in place to safeguard research integrity.

We wish to credit our own Research Integrity and Research Publishing teams and anonymous and named external researchers and research integrity experts for contributing to this investigation.

The corresponding author, as the representative of all authors, has been given the opportunity to register their agreement or disagreement to this retraction. We have kept a record of any response received.

### **References**

- [1] X. Hou, L. Li, S. Chen, C. Ge, M. Shen, and Z. Fu, "MKP-1 Overexpression Reduces Postischemic Myocardial Damage through Attenuation of ER Stress and Mitochondrial Damage," *Oxidative Medicine and Cellular Longevity*, vol. 2021, Article ID 8905578, 12 pages, 2021.

## Research Article

# MKP-1 Overexpression Reduces Postischemic Myocardial Damage through Attenuation of ER Stress and Mitochondrial Damage

Xiaoling Hou <sup>1</sup>, Lijun Li,<sup>1</sup> Si Chen,<sup>2</sup> Cheng Ge,<sup>2</sup> Mingzhi Shen <sup>3</sup>, and Zhenhong Fu <sup>1</sup>

<sup>1</sup>Senior Department of Cardiology, The Sixth Medical Center of People's Liberation Army General Hospital, Beijing, China

<sup>2</sup>Department of Cardiology, The First Medical Center of People's Liberation Army General Hospital, China

<sup>3</sup>Department of Cardiology, Hainan Hospital of Chinese People's Liberation Army General Hospital, Sanya, Hainan 572013, China

Correspondence should be addressed to Mingzhi Shen; [shenmz301@163.com](mailto:shenmz301@163.com) and Zhenhong Fu; [fuzhenh@126.com](mailto:fuzhenh@126.com)

Received 19 July 2021; Revised 4 August 2021; Accepted 16 August 2021; Published 3 September 2021

Academic Editor: Jin Wang

Copyright © 2021 Xiaoling Hou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mitochondrial dysfunction and endoplasmic reticulum (ER) stress contribute to postischemic myocardial damage, but the upstream regulatory mechanisms have not been identified. In this study, we analyzed the role of mitogen-activated protein kinase (MAPK) phosphatase 1 (MKP-1) in the regulation of mitochondrial function and ER stress in hypoxic cardiomyocytes. Our results show that MKP-1 overexpression sustains viability and reduces hypoxia-induced apoptosis among H9C2 cardiomyocytes. MKP-1 overexpression attenuates ER stress and expression of ER stress genes and improves mitochondrial function in hypoxia-treated H9C2 cells. MKP-1 overexpression also increases ATP production and mitochondrial respiration and attenuates mitochondrial oxidative damage in hypoxic cardiomyocytes. Moreover, our results demonstrate that ERK and JNK are the downstream signaling targets of MKP-1 and that MKP-1 overexpression activates ERK, while it inhibits JNK. Inhibition of ERK reduces the ability of MKP-1 to preserve mitochondrial function and ER homeostasis in hypoxic cardiomyocytes. These results show that MKP-1 plays an essential role in the regulation of mitochondrial function and ER stress in hypoxic H9C2 cardiomyocytes through normalization of the ERK pathway and suggest that MKP-1 may serve as a novel target for the treatment of postischemic myocardial injury.

## 1. Introduction

Myocardial ischemia and/or hypoxia are common clinical diseases, which cause serious harm to the body, tissues, and organs [1, 2]. Myocardial damage, necrosis, and arrhythmia caused by myocardial ischemia and hypoxia rank first in the incidence of heart disease, which is also one of the main problems facing the world today [3, 4]. Hypoxia is the main factor of myocardial ischemia and plays a vital role in causing ischemic heart disease damage [5, 6].

Myocardial postischemia injury refers to the interruption of myocardial blood supply after partial or complete coronary artery obstruction [7, 8]. After the blood supply is restored, the ischemic myocardium regains normal perfusion, but additional damage occurs to the postischemic myocardium tissues [9, 10]. A series of pathological changes

occur during the ischemic period including changes in myocardial ultrastructure, energy metabolism, malignant arrhythmia, mitochondria damage, and endoplasmic reticulum (ER) stress [11–15]. However, the mechanisms underlying the pathogenesis of myocardial postischemia damage are not fully understood.

Myocardial postischemia damage has been associated with reactive oxygen species (ROS) overload, excessive production of calcium, granulocyte activation, and energy metabolism disorders in cardiomyocytes [16–19]. Interestingly, these molecular events are closely associated with mitochondrial dysfunction and ER stress [20]. ER is an important organelle that regulates calcium homeostasis, protein synthesis and processing, and protein transport in eukaryotic cells [21]. An excessive accumulation of unfolded or misfolded proteins or calcium ion disbalance in ER leads

to ER dysfunction, resulting in cardiomyocyte death [22]. Abnormal or excessive accumulation of unfolded proteins in the ER activates three main signaling pathways of ER stress: IRE1-XBP1, PERK-eIF2 $\alpha$ , and ATF6 [23]. Activation of IRE1 activates the transcription factor XBP1, which regulates the expression of downstream ER stress proteins [22]. Activation of PERK causes the phosphorylation of eIF2 $\alpha$ , leading to caspase-12 and caspase-3 activation, and irreversible cardiomyocyte damage [24].

Mitochondria are organelles with a three-dimensional network structure [20, 23]. Their structure is not fixed but is always in a dynamic balance of fusion and division [24, 25]. The continuous movement and structural changes of mitochondria are regulated by the GTPase protein family [26–28]. The main proteins regulating mitochondrial fission are Drp1 and Fis1 [29, 30]. The main proteins regulating mitochondrial fusion are Mfn1/2 and Opa1 [31, 32]. An increase in ROS caused by mitochondria dysfunction can downregulate the expression of Mfn1/2 [33, 34], thus shifting the balance in favor of mitochondrial division. The increased mitochondrial division then leads to the production of fragmented mitochondria [35]. Abnormal mitochondrial fission has been linked to cardiomyocyte apoptosis through a mechanism involving mitochondrial outer membrane hyperpermeability [36, 37].

Earlier studies suggested that mitochondria are independent organelles performing specific cellular and metabolic functions [38]. However, recent studies indicate that mitochondria and ER are closely related under physiological conditions [39]. Calcium signal transduction, lipid transport, and energy metabolism are regulated by mitochondria-ER interactions, although detailed molecular patterns have not been identified [40]. Recent studies have identified mitogen-activated protein kinase (MAPK) phosphatase 1 (MKP-1) as a novel regulator of ER function and mitochondrial homeostasis in neuroinflammation [41] and diabetic nephropathy [42]. In this study, we asked whether MKP-1 affects myocardial postischemia damage through regulating mitochondrial homeostasis and ER function.

## 2. Materials and Methods

**2.1. Cell Culture and H/R Model Establishment.** H9C2 myocardial cells were derived from rat embryo fetal heart tissues and cultured in a high-sugar DMEM medium containing 10% fetal calf serum (FCS) at 37°C and 5% CO<sub>2</sub>. The control group was cultured 12 h in a complete DMEM, while the hypoxia group was cultured in DMEM containing 600  $\mu$ mol/L CoCl<sub>2</sub> to induce hypoxia [43].

**2.2. Determination of Intracellular ROS Levels.** Lysine-coated coverslips were placed in a 6-well culture plate, and H9C2 cardiomyocytes were overlaid on the coverslips [44]. When cells grew to about 80% confluence, they were treated 12 h with the following: (1) control group, high-sugar DMEM containing 10% FCS at 37°C and 5% CO<sub>2</sub>, and (2) hypoxia group, DMEM containing 600  $\mu$ mol/L CoCl<sub>2</sub>. Each group consisted of 3 replicate wells. After treatment, cells were rinsed with PBS and washed with 10  $\mu$ mol/L green fluores-

cent probe DCFH-DA. Cells were observed under a fluorescent microscope, and 5 randomly selected nonduplicated areas were analyzed by ImageJ software to determine the mean fluorescence intensity (MFI) [45].

**2.3. Western Blotting.** Cells were lysed and centrifuged, and the protein lysates were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred to a PVDF membrane. After blocking with 2% serum albumin for 2 hours [46], the membranes were incubated with primary antibodies overnight at 4°C. After washing with TBST, the membranes were incubated with HRP-labeled goat anti-rabbit antibody (1:5000) at 37°C for 1 h. The membranes were developed with luminescent reagent ECL and exposed to X-ray film [47].

**2.4. Cell Viability MTT Assay.** After treatment, 200  $\mu$ L of 10% MTT was added to each well, and cells were incubated in a culture box for 4 hours. After adding 150  $\mu$ L of DMSO and shaking for 10 minutes, absorbance (OD value) was measured at 490 nm [48]. The cell survival rate was calculated according to the formula: cell survival rate (%) = OD treatment group/OD control group  $\times$  100% [49].

**2.5. Calcium Fluo-3 AM Assay.** The Fluo-3 AM Calcium Ion Indicator Kit was used according to the manufacturer's instructions. In brief, Fluo-3 AM was incubated with cells at 37°C for 40 min in the dark [50]. After removal of the dye solution, cells were rinsed with PBS, and fluorescence was observed using a fluorescence microscope [51].

**2.6. Mitochondria Fluorescence Staining.** Mitochondria were stained with MitoTracker solution according to the manufacturer's instructions [52]. Briefly, cells were seeded on cell slides in 24-well plates and after treatment incubated with the MitoTracker solution for 30 minutes at 37°C in the dark. After washing, cells were fixed with 4% neutral paraformaldehyde at room temperature for 30 minutes. After rinsing, cells were stained with DAPI and observed with a laser confocal scanning microscope [53].

**2.7. MKP-1 Adenovirus Transfection.** H9C2 cells were transfected with MKP-1 overexpression adenovirus or empty vector control virus for 24 hours, and transfection efficiency was analyzed under a fluorescence microscope [54]. The transfection efficiency exceeding 70% was considered successful, and the MKP-1 overexpression was confirmed by western blotting [55].

**2.8. Statistics.** One-way analysis of variance (ANOVA) was performed to analyze statistically significant changes between two groups; *p* values of *p* < 0.05 were considered statistically significant.

## 3. Results

**3.1. MKP-1 Overexpression Reduces Hypoxia-Induced H9C2 Cardiomyocyte Apoptosis.** To investigate the possible cardioprotective function of MKP-1 in postischemic myocardial damage, we first analyzed cell viability in hypoxia-treated cardiomyocytes transfected with MKP-1 adenovirus. As

shown in Figure 1(a), hypoxia significantly reduced cardiomyocyte viability compared to the control group, but overexpression of MKP-1 preserved the H9C2 cardiomyocyte viability under hypoxia stress. In addition, hypoxia induced the release of lactate dehydrogenase (LDH) into the medium, suggesting damage to cell membrane integrity in the presence of hypoxia stress (Figure 1(b)). However, transfection of MKP-1 adenovirus prevented the LDH leakage (Figure 1(b)), indicating a cardioprotective action of MKP-1 in hypoxic cardiomyocytes. A drop in cell viability is followed by cell apoptosis activation. Indeed, the number of apoptotic cardiomyocytes rapidly increased after exposure to hypoxia (Figures 1(c) and 1(d)), but MKP-1 overexpression decreased the ratio of apoptotic cardiomyocytes (Figures 1(c) and 1(d)). Furthermore, hypoxia increased the activities of caspase-3 and caspase-9, but this increase was inhibited by MKP-1 overexpression (Figures 1(e) and 1(f)). Together, these results show that hypoxia-induced cardiomyocyte death can be inhibited by MKP-1 overexpression.

**3.2. MKP-1 Overexpression Attenuates Cardiomyocyte ER Stress.** Since hypoxia induces cardiomyocyte ER stress and mitochondrial damage, we asked whether the hypoxia-induced ER stress could be attenuated by MKP-1 overexpression. As shown in Figures 2(a)–2(c), the mRNA levels of PERK, CHOP, and GRP78, the markers of ER stress, were significantly increased in hypoxic cardiomyocytes. However, MKP-1 overexpression prevented the hypoxia-induced PERK, CHOP, and GRP78 upregulation (Figures 2(a)–2(c)), indicating that MKP-1 overexpression might suppress ER stress in hypoxic cardiomyocytes. ER stress induces ER damage that is associated with increased intracellular calcium and ER-mediated cell death. The Fluo-3 AM staining assay showed a significant increase in the levels of intracellular calcium in hypoxia-treated cardiomyocytes, but this increase was partly repressed in H9C2 cardiomyocytes overexpressing MKP-1 (Figures 2(d) and 2(e)). In addition, the marker of ER-related cell death, caspase-12 activity (Figure 2(f)), was rapidly upregulated by hypoxia but downregulated to near-normal levels after MKP-1 transfection, demonstrating that MKP-1 overexpression suppresses the ER stress in hypoxic cardiomyocytes.

**3.3. MKP-1 Overexpression Sustains Cardiomyocyte Mitochondrial Function.** Mitochondrial damage, characteristic of postischemic myocardial damage, functions upstream of ER stress. Thus, we asked whether MKP-1 could affect mitochondrial function in hypoxia-treated cardiomyocytes. As shown in Figure 3(a), compared to the control, hypoxia impaired the mitochondria function, as measured by ATP production. However, the ATP production was reversed to normal levels by MKP-1 overexpression. Considering the essential role of mitochondrial respiration in controlling ATP production, we measured baseline and maximal mitochondrial respiration. As shown in Figures 3(b) and 3(c), hypoxia markedly repressed the baseline and maximal mitochondrial respiration, whereas these parameters were normalized in H9C2 cardiomyocytes overexpressing MKP-1.

In addition, hypoxia increased mitochondrial ROS production, but this increase could be normalized by MKP-1 overexpression (Figures 3(d) and 3(e)). These results indicate that MKP-1 overexpression sustains the mitochondrial function in hypoxic cardiomyocytes.

**3.4. MKP-1 Overexpression Inhibits JNK Activity and Activates ERK in Cardiomyocytes.** Previous studies have identified three downstream signaling pathways of MKP-1: JNK, ERK, and p38. Interestingly, ample data have reported the regulatory role played by JNK and ERK in regulating postischemic myocardial injury. Therefore, we analyzed which pathway is controlled by MKP-1 in postischemic myocardial damage. The ELISA assay demonstrated that the activity of ERK was markedly downregulated in hypoxia-treated cardiomyocytes (Figures 4(a) and 4(b)). In contrast, JNK activity was rapidly increased in response to hypoxia treatment (Figures 4(a) and 4(b)). After transfection of MKP-1 adenovirus, the activity of ERK was increased, whereas the activity of JNK was decreased (Figures 4(a) and 4(b)), suggesting that MKP-1 overexpression increases the ERK activity but inhibits the JNK pathway. Similar changes were observed on protein levels; as shown in Figures 4(c)–4(e), the JNK protein levels rapidly increased, whereas the ERK expression decreased in hypoxia-treated cardiomyocytes. However, MKP-1 overexpression suppressed the JNK activity and increased the ERK expression (Figures 4(c)–4(e)). Together, these results indicate that the downstream pathways of MKP-1 include JNK and ERK.

**3.5. ERK Inhibition Decreases MKP-1-Mediated Protection of Mitochondria and ER.** To determine whether MKP-1 controls mitochondrial function and ER homeostasis through ERK, we used the ERK inhibitor SCH772984. MKP-1 transfection improved mitochondrial ATP production (Figure 5(a)) and baseline and maximal mitochondrial respiration (Figures 5(b) and 5(c)) in hypoxic cardiomyocytes. However, inhibition of ERK by SCH772984 significantly reduced the mitochondrial ATP production and ROS generation in MKP-1-overexpressing H9C2 cardiomyocytes (Figure 5(d)), indicating that MKP-1 modulates the mitochondrial function through ERK. In addition, MKP-1 transfection suppressed the expression of ER stress genes in hypoxic cardiomyocytes (Figures 5(e) and 5(f)), but ERK inhibition by SCH772984 increased the expression of ER stress genes, indicating the ERK involvement in MKP-1-mediated ER homeostasis.

## 4. Discussion

H9C2 cells are derived from rat embryonic heart tissue, have characteristics of mature myocardial cells, and are widely used in cell morphology, electrophysiology, and toxicology studies. *In vitro*, hypoxia can be induced physically and chemically. Physical hypoxia is often induced in nitrogen-filled hypoxic devices and normoxia devices. However, due to the need for an exact oxygen concentration detection, its application is limited. Chemical hypoxia is easy to induce, and  $\text{CoCl}_2$  is a commonly used chemical hypoxia stimulant.




FIGURE 1: MKP-1 overexpression reduces hypoxia-induced cardiomyocyte apoptosis. (a) Cell viability analyzed by MTT assay in hypoxia-treated cardiomyocytes transfected with MKP-1 adenovirus or a control vector. (b) LDH release assay was used to analyze cell death. (c, d) TUNEL apoptosis staining. (e, f) ELISA assays of caspase-3 and caspase-9. \* $p < 0.05$ . Bar:  $120 \mu\text{m}$ .




FIGURE 2: MKP-1 overexpression attenuates ER stress. (a–c) qRT-PCR of PERK, CHOP, and GRP78 mRNA levels in hypoxia-treated cardiomyocytes transfected with MKP-1 adenovirus or control vector. (d, e) Immunofluorescence assay of intracellular calcium using Fluo-3 AM probe. (f) ELISA was used to analyze the activity of caspase-12. \* $p < 0.05$ .

Cobalt can replace ferrous iron chelated in hemoglobin, preventing cell oxygen uptake and inducing oxidative stress damage. Removal of CoCl<sub>2</sub> can restore hemoglobin oxygen uptake and hypoxia-reoxygenation damage. Our results indeed show that CoCl<sub>2</sub> inhibits the survival of H9C2 cardiomyocytes in a dose- and time-dependent manner. The induced cardiomyocyte hypoxia was associated with increased levels of ROS, highlighting the important role of oxidative stress in aggravating myocardial postischemic damage. ROS include O<sup>2-</sup>, OH, ONOO<sup>-</sup>, and H<sub>2</sub>O<sub>2</sub> [56, 57]. It is believed that the hypoxia-induced ROS can be transported from the cell membrane to the cytoplasm where they act as a second messenger to modulate gene expression and DNA oxidative modification. Therefore, antioxidative drugs or compounds that enhance the activity of antioxidative enzymes (such as glutathione, superoxide dismutase,

coenzyme II, and heme oxygenase-1) may have additional benefits for a postischemic heart.

Apoptosis is an important mechanism that causes postischemic myocardial injury and heart failure. Recent studies have reported that ROS-related oxidative response is capable of interrupting the MAPK pathway, resulting in increased apoptosis. MAPKs are serine-threonine protein kinases that regulate cell growth, differentiation, and responses to environmental stress; they include extracellular signal-regulated protein kinase (ERK), N-terminal kinase (JNK), and p38. Our study shows that hypoxia activates JNK and inhibits ERK in H9C2 cardiomyocytes, and this is associated with mitochondrial damage and ER stress.

Previous studies have shown that myocardial ischemic damage is closely associated with cardiomyocyte mitochondrial dysfunction [15, 58]. A reduction of mitochondrial




FIGURE 3: MKP-1 overexpression sustains mitochondrial function. (a) ATP production analyzed by ELISA in hypoxia-treated cardiomyocytes transfected with MKP-1 adenovirus or control vector. (b, c) The baseline and maximal mitochondrial respiration was measured through ELISA. (d, e) Immunofluorescence assay of intracellular calcium using ROS probe. \* $p < 0.05$ .




FIGURE 4: MKP-1 overexpression inhibits JNK activity and activates the ERK pathway. (a, b) ELISA assay of activity of JNK and ERK in hypoxia-treated cardiomyocytes transfected with MKP-1 adenovirus or control vector. (c-e) Western analysis of JNK and ERK levels. \**p* < 0.05. Bar: 75 μm.






FIGURE 5: Continued.




FIGURE 5: ERK inhibition reduces MKP-1-mediated protection of mitochondria and ER. (a) ATP production was analyzed by ELISA in hypoxia-treated cardiomyocytes transfected with MKP-1 adenovirus or control vector and treated with the ERK inhibitor SCH772984. (b, c) The baseline and maximal mitochondrial respiration was measured through ELISA. (d) Immunofluorescence assay was used to measure intracellular ROS. (e, f) qRT-PCR analysis of PERK and CHOP mRNA levels. \* $p < 0.05$ .

membrane potential has been considered an important step inducing cell damage by increasing apoptosis [14, 59]. In our study, mitochondrial damage was characterized by decreased ATP production, impaired mitochondrial respiration, and reduced mitochondrial membrane potential. Interestingly, these alterations could be reversed by MKP-1 overexpression through normalization of the ERK pathway. During apoptosis, mitochondria are known to undergo dramatic structural changes. Indeed, our results showed that mitochondrial function [60], ER function, and ER structure were improved by MKP-1 overexpression in a manner dependent on ERK. These results are consistent with previous studies. For example, MKP-1 deletion caused mitochondrial damage, resulting in decreased insulin release in obesity [41, 61]. In diabetic rats, MKP-1 overexpression improved cardiac performance by promoting mitochondrial metabolism [62]. LPS-associated myocardial damage could be attenuated by MKP-1 through reducing inflammatory response [63]. Oxidative stress-mediated endothelial cell injury was associated with ER stress due to a loss of MKP-1 activity [64, 65]. Overexpression of MKP-1 reduced ER stress, resulting in increased neural cell survival [66]. In hypoxia-treated mesenchymal stem cells, overexpression of MKP-1 prevented activation of ER stress and promoted differentiation and proliferation of stem cells [67, 68]. Based on these data, MKP-1 could be considered a critical regulator of mitochondrial function and ER stress in the postischemic myocardium.

There are several issues that should be addressed in future studies. First, our MKP-1 overexpression results should be confirmed by loss-of-function experiments. *In*

*vivo* data will be needed to rule out a possible negative effect of MKP-1 on other tissues, such as the liver and lung. In addition, although adenovirus transfection is an effective way to enhance MKP-1 expression *in vitro*, novel chemotherapeutic drugs and gene therapeutic approaches targeting the MKP-1 expression will facilitate the development of better therapies for the treatment of postischemic myocardial injury.

## Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## Conflicts of Interest

The authors declare no conflict of interest, financial or otherwise.

## Authors' Contributions

Experiments were performed by XLH and LJL. The manuscript was written by SC and CG. The study was supervised by ZHF and MZS. The manuscript was corrected by ZHF. Xiaoling Hou and Lijun Li contributed equally to this work.

## Acknowledgments

This research was supported by the Hainan science and technology project (ZDYF2020123 and ZDYF2019188).

## References

- [1] B. R. Pagliaro, F. Cannata, G. G. Stefanini, and L. Bolognese, "Myocardial ischemia and coronary disease in heart failure," *Heart Failure Reviews*, vol. 25, no. 1, pp. 53–65, 2020.
- [2] T. Carbonell and A. V. Gomes, "MicroRNAs in the regulation of cellular redox status and its implications in myocardial ischemia-reperfusion injury," *Redox Biology*, vol. 36, p. 101607, 2020.
- [3] X. Wang, X. Li, S. Liu et al., "PCSK9 regulates pyroptosis via mtDNA damage in chronic myocardial ischemia," *Basic Research in Cardiology*, vol. 115, no. 6, p. 66, 2020.
- [4] J. Van De Wouw, O. Sorop, R. W. A. Van Drie et al., "Perturbations in myocardial perfusion and oxygen balance in swine with multiple risk factors: a novel model of ischemia and no obstructive coronary artery disease," *Basic Research in Cardiology*, vol. 115, no. 2, p. 21, 2020.
- [5] H. Feng, S. Q. Mou, W. J. Li et al., "Resveratrol inhibits ischemia-induced myocardial senescence signals and NLRP3 inflammasome activation," *Oxidative Medicine and Cellular Longevity*, vol. 2020, Article ID 2647807, 20 pages, 2020.
- [6] Y. Sawashita, N. Hirata, Y. Yoshikawa, H. Terada, Y. Tokinaga, and M. Yamakage, "Remote ischemic preconditioning reduces myocardial ischemia-reperfusion injury through unacylated ghrelin-induced activation of the JAK/STAT pathway," *Basic Research in Cardiology*, vol. 115, no. 4, p. 50, 2020.
- [7] J. Wang, S. Toan, and H. Zhou, "New insights into the role of mitochondria in cardiac microvascular ischemia/reperfusion injury," *Angiogenesis*, vol. 23, no. 3, pp. 299–314, 2020.
- [8] Y. Tan, D. Mui, S. Toan, P. Zhu, R. Li, and H. Zhou, "SERCA overexpression improves mitochondrial quality control and attenuates cardiac microvascular ischemia-reperfusion injury," *Mol Ther Nucleic Acids*, vol. 22, pp. 696–707, 2020.
- [9] J. Wang, S. Toan, and H. Zhou, "Mitochondrial quality control in cardiac microvascular ischemia-reperfusion injury: new insights into the mechanisms and therapeutic potentials," *Pharmacological Research*, vol. 156, p. 104771, 2020.
- [10] H. Zhou and S. Toan, "Pathological roles of mitochondrial oxidative stress and mitochondrial dynamics in cardiac microvascular ischemia/reperfusion injury," *Biomolecules*, vol. 10, no. 1, p. 85, 2020.
- [11] H. Zhou, S. Toan, P. Zhu, J. Wang, J. Ren, and Y. Zhang, "DNA-PKcs promotes cardiac ischemia reperfusion injury through mitigating BI-1-governed mitochondrial homeostasis," *Basic Research in Cardiology*, vol. 115, no. 2, p. 11, 2020.
- [12] H. Zhou, P. Zhu, J. Guo et al., "Ripk3 induces mitochondrial apoptosis via inhibition of FUNDC1 mitophagy in cardiac IR injury," *Redox Biology*, vol. 13, pp. 498–507, 2017.
- [13] X. Sun, Z. Wei, Y. Li et al., "Renal denervation restrains the inflammatory response in myocardial ischemia-reperfusion injury," *Basic Research in Cardiology*, vol. 115, no. 2, p. 15, 2020.
- [14] J. Martínez-Milla, C. Galán-Arriola, M. Carnero et al., "Translational large animal model of hibernating myocardium: characterization by serial multimodal imaging," *Basic Research in Cardiology*, vol. 115, no. 3, p. 33, 2020.
- [15] the EU-CardioRNA COST Action (CA17129), A. Jusic, and Y. Devaux, "Mitochondrial noncoding RNA-regulatory network in cardiovascular disease," *Basic Research in Cardiology*, vol. 115, no. 3, p. 23, 2020.
- [16] S. Hamilton, R. Terentyeva, B. Martin et al., "Increased RyR2 activity is exacerbated by calcium leak-induced mitochondrial ROS," *Basic Research in Cardiology*, vol. 115, no. 4, p. 38, 2020.
- [17] H. Zhou, P. Zhu, J. Wang, H. Zhu, J. Ren, and Y. Chen, "Pathogenesis of cardiac ischemia reperfusion injury is associated with CK2 $\alpha$ -disturbed mitochondrial homeostasis via suppression of FUNDC1-related mitophagy," *Cell Death and Differentiation*, vol. 25, no. 6, pp. 1080–1093, 2018.
- [18] H. Zhou, S. Wang, S. Hu, Y. Chen, and J. Ren, "ER-mitochondria microdomains in cardiac ischemia-reperfusion injury: a fresh perspective," *Frontiers in Physiology*, vol. 9, p. 755, 2018.
- [19] C. W. Vogel, "The role of complement in myocardial infarction reperfusion injury: an underappreciated therapeutic target," *Frontiers in Cell and Development Biology*, vol. 8, p. 606407, 2020.
- [20] S. Cadenas, "ROS and redox signaling in myocardial ischemia-reperfusion injury and cardioprotection," *Free Radical Biology & Medicine*, vol. 117, pp. 76–89, 2018.
- [21] H. Zhu and H. Zhou, "Novel insight into the role of endoplasmic reticulum stress in the pathogenesis of myocardial ischemia-reperfusion injury," *Oxidative Medicine and Cellular Longevity*, vol. 2021, Article ID 5529810, 2021.
- [22] B. Delgado-Valero, L. De La Fuente-Chávez, A. Romero-Miranda et al., "Role of endoplasmic reticulum stress in renal damage after myocardial infarction," *Clinical Science (London, England)*, vol. 135, no. 1, pp. 143–159, 2021.
- [23] J. L. Chiang, P. Shukla, K. Pagidas et al., "Mitochondria in ovarian aging and reproductive longevity," *Ageing Research Reviews*, vol. 63, p. 101168, 2020.
- [24] F. Di Lisa, M. Canton, R. Menabò, N. Kaludercic, and P. Bernardi, "Mitochondria and cardioprotection," *Heart Failure Reviews*, vol. 12, no. 3-4, pp. 249–260, 2007.
- [25] S. M. Davidson, "Endothelial mitochondria and heart disease," *Cardiovascular Research*, vol. 88, no. 1, pp. 58–66, 2010.
- [26] H. Zhou, C. Shi, S. Hu, H. Zhu, J. Ren, and Y. Chen, "BI1 is associated with microvascular protection in cardiac ischemia reperfusion injury via repressing Syk-Nox2-Drp1-mitochondrial fission pathways," *Angiogenesis*, vol. 21, no. 3, pp. 599–615, 2018.
- [27] J. Wang, P. Zhu, S. Toan, R. Li, J. Ren, and H. Zhou, "Pum2-Mff axis fine-tunes mitochondrial quality control in acute ischemic kidney injury," *Cell Biology and Toxicology*, vol. 36, no. 4, pp. 365–378, 2020.
- [28] H. Zhou, S. Wang, P. Zhu, S. Hu, Y. Chen, and J. Ren, "Empagliflozin rescues diabetic myocardial microvascular injury via AMPK-mediated inhibition of mitochondrial fission," *Redox Biology*, vol. 15, pp. 335–346, 2018.
- [29] H. Zhou, P. Zhu, J. Wang, S. Toan, and J. Ren, "DNA-PKcs promotes alcohol-related liver disease by activating Drp1-related mitochondrial fission and repressing FUNDC1-required mitophagy," *Signal Transduction and Targeted Therapy*, vol. 4, no. 1, p. 56, 2019.
- [30] H. Zhou, W. Du, Y. Li et al., "Effects of melatonin on fatty liver disease: the role of NR4A1/DNA-PKcs/p53 pathway, mitochondrial fission, and mitophagy," *Journal of Pineal Research*, vol. 64, no. 1, 2018.
- [31] J. Wang and H. Zhou, "Mitochondrial quality control mechanisms as molecular targets in cardiac ischemia \*\*\* reperfusion injury," *Acta Pharmaceutica Sinica B*, vol. 10, no. 10, pp. 1866–1879, 2020.

- [32] A. Del Campo, "Mitophagy as a new therapeutic target for sarcopenia," *Acta Physiologica*, vol. 225, no. 2, article e13219, 2019.
- [33] Y. Z. Li, X. D. Wu, X. H. Liu, and P. F. Li, "Mitophagy imbalance in cardiomyocyte ischaemia/reperfusion injury," *Acta Physiologica*, vol. 225, no. 4, article e13228, 2019.
- [34] M. Akbari, T. B. L. Kirkwood, and V. A. Bohr, "Mitochondria in the signaling pathways that control longevity and health span," *Ageing Research Reviews*, vol. 54, p. 100940, 2019.
- [35] H. Zhou, S. Hu, Q. Jin et al., "Mff-dependent mitochondrial fission contributes to the pathogenesis of cardiac microvasculature ischemia/reperfusion injury via induction of mROS-mediated cardiolipin oxidation and HK2/VDAC1 disassociation-involved mPTP opening," *Journal of the American Heart Association*, vol. 6, no. 3, 2017.
- [36] J. Wang, S. Toan, R. Li, and H. Zhou, "Melatonin fine-tunes intracellular calcium signals and eliminates myocardial damage through the IP3R/MCU pathways in cardiorenal syndrome type 3," *Biochemical Pharmacology*, vol. 174, p. 113832, 2020.
- [37] H. Zhou, J. Wang, S. Hu, H. Zhu, S. Toan, and J. Ren, "BI1 alleviates cardiac microvascular ischemia-reperfusion injury via modifying mitochondrial fission and inhibiting XO/ROS/F-actin pathways," *Journal of Cellular Physiology*, vol. 234, no. 4, pp. 5056–5069, 2019.
- [38] Z. Fu, Y. Jiao, J. Wang et al., "Cardioprotective role of melatonin in acute myocardial infarction," *Frontiers in Physiology*, vol. 11, p. 366, 2020.
- [39] S. Yang, R. Zhou, C. Zhang, S. He, and Z. Su, "Mitochondria-associated endoplasmic reticulum membranes in the pathogenesis of type 2 diabetes mellitus," *Frontiers in Cell and Development Biology*, vol. 8, p. 571554, 2020.
- [40] R. Khatoon, M. Pahuja, and S. Parvez, "Cross talk between mitochondria and other targets in Alzheimer's disease," *Journal of Environmental Pathology, Toxicology and Oncology*, vol. 39, no. 3, pp. 261–279, 2020.
- [41] D. Huang and Y. Jiang, "MKP1 reduces neuroinflammation via inhibiting endoplasmic reticulum stress and mitochondrial dysfunction," *Journal of Cellular Physiology*, vol. 235, no. 5, pp. 4316–4325, 2020.
- [42] Y. Zhang, J. Feng, Q. Wang et al., "Hyperglycaemia stress-induced renal injury is caused by extensive mitochondrial fragmentation, attenuated MKP1 signalling, and activated JNK-CaMKII-Fis1 biological axis," *Cellular Physiology and Biochemistry*, vol. 51, no. 4, pp. 1778–1798, 2018.
- [43] J. Bai, M. Khajavi, L. Sui et al., "Angiogenic responses in a 3D micro-engineered environment of primary endothelial cells and pericytes," *Angiogenesis*, vol. 24, no. 1, pp. 111–127, 2021.
- [44] S. Khadjeh, V. Hindmarsh, F. Weber et al., "CRISPLD1: a novel conserved target in the transition to human heart failure," *Basic Research in Cardiology*, vol. 115, no. 3, p. 27, 2020.
- [45] D. Bausch, S. Fritz, L. Bolm et al., "Hedgehog signaling promotes angiogenesis directly and indirectly in pancreatic cancer," *Angiogenesis*, vol. 23, no. 3, pp. 479–492, 2020.
- [46] P. Kleinbongard, "Cardioprotection by early metoprolol-attenuation of ischemic vs. reperfusion injury?," *Basic Research in Cardiology*, vol. 115, no. 5, p. 54, 2020.
- [47] A. L. Bayliss, A. Sundararaman, C. Granet, and H. Mellor, "Raflin is recruited by neuropilin-1 to the activated VEGFR2 complex to control proangiogenic signaling," *Angiogenesis*, vol. 23, no. 3, pp. 371–383, 2020.
- [48] S. Liu, J. Chen, J. Shi et al., "M1-like macrophage-derived exosomes suppress angiogenesis and exacerbate cardiac dysfunction in a myocardial infarction microenvironment," *Basic Research in Cardiology*, vol. 115, no. 2, p. 22, 2020.
- [49] E. Bridges, H. Sheldon, E. Kleibeuker et al., "RHOQ is induced by DLL4 and regulates angiogenesis by determining the intracellular route of the Notch intracellular domain," *Angiogenesis*, vol. 23, no. 3, pp. 493–513, 2020.
- [50] S. K. Lahiri, A. P. Quick, B. Samson-Couterie et al., "Nuclear localization of a novel calpain-2 mediated junctophilin-2 C-terminal cleavage peptide promotes cardiomyocyte remodeling," *Basic Research in Cardiology*, vol. 115, no. 4, p. 49, 2020.
- [51] D. B. Buglak, E. J. Kushner, A. P. Marvin, K. L. Davis, and V. L. Bautch, "Excess centrosomes disrupt vascular lumenization and endothelial cell adherens junctions," *Angiogenesis*, vol. 23, no. 4, pp. 567–575, 2020.
- [52] Y. Li, P. Liang, B. Jiang et al., "CARD9 promotes autophagy in cardiomyocytes in myocardial ischemia/reperfusion injury via interacting with Rubicon directly," *Basic Research in Cardiology*, vol. 115, no. 3, p. 29, 2020.
- [53] J. Cao, X. Liu, Y. Yang et al., "Decylubiquinone suppresses breast cancer growth and metastasis by inhibiting angiogenesis via the ROS/p53/ BAI1 signaling pathway," *Angiogenesis*, vol. 23, no. 3, pp. 325–338, 2020.
- [54] M. Lindner, H. Mehel, A. David et al., "Fibroblast growth factor 23 decreases PDE4 expression in heart increasing the risk of cardiac arrhythmia; Klotho opposes these effects," *Basic Research in Cardiology*, vol. 115, no. 5, p. 51, 2020.
- [55] T. L. Capasso, B. Li, H. J. Volek et al., "BMP10-mediated ALK1 signaling is continuously required for vascular development and maintenance," *Angiogenesis*, vol. 23, no. 2, pp. 203–220, 2020.
- [56] M. Neri, V. Fineschi, M. Di Paolo et al., "Cardiac oxidative stress and inflammatory cytokines response after myocardial infarction," *Current Vascular Pharmacology*, vol. 13, no. 1, pp. 26–36, 2015.
- [57] R. S. Gangwar, G. H. Bevan, R. Palanivel, L. Das, and S. Rajagopalan, "Oxidative stress pathways of air pollution mediated toxicity: recent insights," *Redox Biology*, vol. 34, p. 101545, 2020.
- [58] C. L. Depoix, A. Colson, C. Hubinont, and F. Debieve, "Impaired vascular endothelial growth factor expression and secretion during in vitro differentiation of human primary term cytotrophoblasts," *Angiogenesis*, vol. 23, no. 2, pp. 221–230, 2020.
- [59] X. Lu, Y. He, C. Tang et al., "Triad3A attenuates pathological cardiac hypertrophy involving the augmentation of ubiquitination-mediated degradation of TLR4 and TLR9," *Basic Research in Cardiology*, vol. 115, no. 2, p. 19, 2020.
- [60] S. Pabel, S. Ahmad, P. Tirilomis et al., "Inhibition of NaV1.8 prevents atrial arrhythmogenesis in human and mice," *Basic Research in Cardiology*, vol. 115, no. 2, p. 20, 2020.
- [61] M. Lobo-Gonzalez, C. Galán-Arriola, X. Rossello et al., "Metoprolol blunts the time-dependent progression of infarct size," *Basic Research in Cardiology*, vol. 115, no. 5, p. 55, 2020.
- [62] Y. Weng, F. Shen, J. Li, Y. Shen, and X. Zhang, "Expression changes of mitogen-activated protein kinase phosphatase-1 (MKP-1) in myocardium of streptozotocin-induced diabetic rats," *Experimental and Clinical Endocrinology & Diabetes*, vol. 115, no. 7, pp. 455–460, 2007.

- [63] Z. Tucsek, B. Radnai, B. Racz et al., "Suppressing LPS-induced early signal transduction in macrophages by a polyphenol degradation product: a critical role of MKP-1," *Journal of Leukocyte Biology*, vol. 89, no. 1, pp. 105–111, 2011.
- [64] W. J. Lee, W. Huey-Herng Sheu, S. H. Liu et al., "Nε-carboxymethyllysine-mediated endoplasmic reticulum stress promotes endothelial cell injury through Nox4/MKP-3 interaction," *Free Radical Biology & Medicine*, vol. 74, pp. 294–306, 2014.
- [65] J. Chen, F. L. Lin, J. Y. K. Leung et al., "A drug-tunable Flt23k gene therapy for controlled intervention in retinal neovascularization," *Angiogenesis*, vol. 24, no. 1, pp. 97–110, 2021.
- [66] B. Li, P. Yi, B. Zhang et al., "Differences in endoplasmic reticulum stress signalling kinetics determine cell survival outcome through activation of MKP-1," *Cellular Signalling*, vol. 23, no. 1, pp. 35–45, 2011.
- [67] Z. Li, H. Wei, X. Liu, S. Hu, X. Cong, and X. Chen, "LPA rescues ER stress-associated apoptosis in hypoxia and serum deprivation-stimulated mesenchymal stem cells," *Journal of Cellular Biochemistry*, vol. 111, no. 4, pp. 811–820, 2010.
- [68] D. V. C. De Jel, F. J. M. Disch, S. Kroon, J. J. Mager, and F. J. Verdam, "Intranasal Efudix reduces epistaxis in hereditary hemorrhagic telangiectasia," *Angiogenesis*, vol. 23, no. 3, pp. 271–274, 2020.