

Research Article

Sublethal Oxidative Stress Induces the Premature Senescence of Human Mesenchymal Stem Cells Derived from Endometrium

Elena Burova, Aleksandra Borodkina, Alla Shatrova, and Nikolay Nikolsky

Department of Intracellular Signaling and Transport, Institute of Cytology of Russian Academy of Sciences, St. Petersburg 194064, Russia

Correspondence should be addressed to Elena Burova; burova_elena@yahoo.com

Received 10 June 2013; Revised 19 July 2013; Accepted 19 July 2013

Academic Editor: Mohammad Abdollahi

Copyright © 2013 Elena Burova et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The specific responses of mesenchymal stem cells to oxidative stress may play a crucial role in regulation of tissue homeostasis as well as regeneration of organs after oxidative injury. The responses of human endometrium-derived mesenchymal stem cells (hMESC) to oxidative stress remain still unknown. Herein, we examined the impact of H_2O_2 on cell viability, induction of premature senescence, and apoptosis. hMESC were highly resistant to H_2O_2 compared with human diploid fibroblasts. To test a hypothesis whether hMESC may undergo oxidative stress-induced premature senescence, cells were briefly exposed to the sublethal H_2O_2 doses. H_2O_2 -treated cells were permanently arrested, lost Ki67 proliferation marker, and exhibited a senescent phenotype including cell hypertrophy and increased SA- β -Gal activity. Additionally, in stressed cells the expression levels of p21Cip1, SOD1, SOD2, and GPX1 were elevated. hMESC survived under stress were not able to resume proliferation, indicating the irreversible loss of proliferative potential. While the low H_2O_2 doses promoted senescence in hMESC, the higher H_2O_2 doses induced also apoptosis in a part of the cell population. Of note, senescent hMESC exhibited high resistance to apoptosis. Thus, we have demonstrated for the first time that hMESC may enter a state of premature senescence in response to sublethal oxidative stress.

1. Introduction

Stress responses of human embryonic and adult stem cells to γ -radiation, oxidative stress, heat shock, and so forth are widely researched to establish cell-based strategies of tissue repair, tissue engineering, and transplantation [1]. Human mesenchymal stem cells are adult multipotent stem cells with the capacity of self-renewal and undergoing adipogenic, osteogenic, chondrogenic, myogenic differentiation [2, 3]. They contribute to the homeostatic maintenance of many organs and tissues [4, 5]. Unlike some other adult stem cells (e.g., hematopoietic stem cells) human mesenchymal stem cells are not immortal. These cells exhibit *ex vivo* growth characteristics typical of the Hayflick model of cellular senescence with a limited life span [6]. Recently, it has been reported that the mesenchymal stem cells subjected to oxidative stress [7–9] or ionizing radiation [10–12] may undergo stress-induced premature senescence *in vitro*. Many types of normal and tumor cells also enter a state of premature senescence after exposure to radiation [13–15], H_2O_2 [16–19], or treatment

with histone deacetylase inhibitors [20, 21]. Prematurely senescent cells exhibit some of the characteristics inherent in replicatively senescent cells, including a large flat morphology, increased senescence-associated β -galactosidase (SA- β -Gal) activity, and permanent cell cycle arrest [14, 17]. Besides, cellular overactivation and hyperfunction, feedback signal resistance, and loss of regenerative potential are considered hallmarks of senescence [22]. Progress in understanding the causes and mechanisms of cellular senescence and significance of senescence for ageing and suppressing cancer has been reviewed [23–25].

In the current study, oxidative stress responses of human mesenchymal stem cells derived from endometrium (hMESC) were investigated. Our knowledge of specific responses of these cells to stress is very limited, though they prove to be useful in the treatment of pathologies in which tissue damage is linked to oxidative stress. Unlike most of the human mesenchymal stem cells, the isolation of which as a rule is complicated by invasive procedures, the mesenchymal stem cells produced from desquamated endometrium in

menstrual blood by a simple noninvasive way provide a good opportunity to explore the stress responses of hMESCs. Regarding hMESCs, phenomenon of premature senescence induced by oxidative stress remains still unknown. This study aimed to test a hypothesis whether hMESCs after exposure with sublethal doses of H_2O_2 may undergo the stress-induced premature senescence. In parallel, the impact of H_2O_2 on cell viability and development of apoptosis has been evaluated.

2. Materials and Methods

2.1. Cell Culture and Cell Treatment. Human mesenchymal stem cells isolated from desquamated endometrium in menstrual blood (hMESCs, line 2304), as described previously [26], as well as human embryonic lung-derived diploid fibroblasts (HDF, line FRL-9505) were cultured in complete medium (DMEM/F12 (Gibco BRL, Gaithersburg, MD, USA) supplemented with 10% FBS (HyClone, Waltham, MA, USA), 1% gentamycin, and 1% glutamax (Gibco BRL, Gaithersburg, MD, USA)) at 37°C in humidified incubator, containing 5% CO_2 . hMESCs have a positive expression of CD73, CD90, CD105, CD13, CD29, and CD44 markers and absence of expression of the hematopoietic cell surface antigens CD19, CD34, CD45, CD117, CD130, and HLA-DR (class II). Multipotency of isolated hMESCs is confirmed by their ability to differentiate into other mesodermal cell types, such as osteocytes and adipocytes. Besides, the isolated hMESCs partially (over 50%) express the pluripotency marker SSEA-4 but do not express Oct-4. Immunofluorescent analysis of the derived cells revealed the expression of the neural precursor markers nestin and beta-III-tubulin. This suggests a neural predisposition of the established hMESCs. These cells are characterized by high rate of cell proliferation (doubling time 22-23 h) and high cloning efficiency (about 60%). Cells at early passages (between 6 and 9 passages for hMESCs and between 16 and 21 passages for HDF) were used in all experiments to avoid complications of replicative senescence. Cells were harvested by trypsinization and plated at a density of 15×10^3 cells per cm^2 . For microscopy experiments, cells were grown on glass coverslips. H_2O_2 treatments were performed on subconfluent cells to avoid variability of H_2O_2 toxicity. H_2O_2 stock solution in serum-free medium was prepared from 30% H_2O_2 (Sigma, St. Louis, MO, USA) just before adding. Cells were treated with H_2O_2 in the range of concentrations from 200 μM to 2,000 μM for 1 h for MTT assay. Based on LD_{50} values 200 μM H_2O_2 was chosen as a sublethal concentration for the induction of premature senescence of hMESCs, whereas apoptosis was tested under higher concentrations of H_2O_2 (900 and 3,000 μM). The cells were washed twice with serum-free medium to remove H_2O_2 and then recultured in fresh complete medium for various durations as specified in individual experiments.

2.2. Assessment of Cell Viability. The cell viability after exposure to H_2O_2 for 1 h was evaluated by the enzymatic conversion of MTT (AppliChem, Darmstadt, Germany, number A2231) to formazan in live cells. The culture medium from the cells grown in plates was removed, and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide

(MTT; 0,715 mg/mL) in serum-containing growth medium was added to each well. In 2 h the solution was changed to DMSO to solve formazan produced. The plates were shaken for 15 min at room temperature; thereafter the absorbance was measured at 570 nm using microplate reader (Fluorofot "Charity," Russia). All points were read as parallels of 8 similar samples. The average absorbance at a given time point was normalized to the start time point.

2.3. Flow Cytometry. Adherent cells were rinsed twice with PBS and harvested by trypsinization. Detached cells were collected with supernatants, pelleted by centrifugation. Detached and adherent cells were finally pooled and resuspended in PBS. One part of each sample was used for propidium iodide (PI) staining to evaluate cell viability and another part for cell cycle phase distribution analysis that was performed as described previously [27]. 50 $\mu g/mL$ PI was added to each sample just before analysis and mixed gently. Samples were analyzed on a Coulter EPICS XL Flow Cytometer (Beckman Coulter, Brea, CA, USA). For cell cycle analysis, each cell sample was suspended in 300 μL PBS containing 200 $\mu g/mL$ of saponin (Fluka, NY, USA), used for cell permeabilization, 250 $\mu g/mL$ RNase A (Sigma, St. Louis, MO, USA, number R4642), and 50 $\mu g/mL$ PI, incubated for 30 min at room temperature and subjected to FACS analysis. At least 10,000 cells were measured per sample. Cell cycle analysis was performed using Win MDI program version 2.8 and ModFit LT software (Verity Software House, Topsham, ME, USA).

2.4. FACS Analysis of Cell Enlargement. The same procedure of sample preparation as described previously was done for light-scattering cytometry. As after H_2O_2 treatment live and dead cells were very close to each other on one-parameter histogram, two-parameter histogram was used (FL4LOG versus FSLOG) to discriminate live and dead cells. Analysis of each sample was performed for 100 sec with high sample delivery. Cell size of control and H_2O_2 -treated cells was measured by means of cytometric light scattering of PI-stained cells by using Win MDI program version 2.8.

2.5. SA- β -Gal Activity. Cells expressing senescent-associated β -galactosidase were detected with senescence β -galactosidase staining kit (Cell Signaling Technology, Beverly, MA, USA, number 9860) according to manufacturer's instructions. The kit detects β -galactosidase activity at pH 6 in cultured cells which is present only in senescent cells and is not found in presenescent, quiescent, or immortal cells. The percent of SA- β -Gal-positive cells was calculated by counting not less than 500 cells.

2.6. Immunofluorescence Staining. Cells cultured on coverslips were fixed with PBS/4% formalin for 15 min and then permeabilized with 0.1% Triton X-100. After blocking with 1% bovine serum albumin, they were incubated with a rabbit polyclonal antibody against Ki67 (Abcam, Cambridge, UK, number 15580) (1 : 1000) overnight at 4°C and then with Alexa Fluor 568 donkey anti-rabbit antibody (Invitrogen, Carlsbad,

USA, number A10042) (1:500) at room temperature for 1 h after extensive washing with PBS/0.1% Tween 20 between each step. The slides were counterstained with 1 $\mu\text{g}/\text{mL}$ DAPI (Sigma, St. Louis, MO, USA, number D9564) and mounted using 2% propyl gallate. A Zeiss Axiovert 200M fluorescence microscope (Carl Zeiss, Germany) equipped with a digital camera DFC 420C (Leica, Germany) utilizing Adobe Photoshop software was used to view and acquire images.

2.7. Apoptosis Detection. Apoptosis detection was performed by Annexin V/PI staining according to standard manufacture protocols (BD Pharmingen). H_2O_2 -treated and control cells were harvested as described previously. Then cells were washed twice with cold PBS, resuspended in 1X binding buffer at concentration 10^6 cells/mL. 100 μL of this suspension was transferred to 5 mL tube, and 5 μL Annexin V/FITC (Annexin V/FITC Apoptosis Detection Kit II, BD Biosciences, San Diego, CA, USA) and 10 μL PI were added; suspension was gently mixed and incubated for 15 min at room temperature in the dark. Just before flow cytometric analysis 400 μL of 1X binding buffer was added to each sample.

2.8. Western Blotting. Total cell lysates were prepared as described previously [27]. Protein content was determined by the method of Bradford. The cell lysates were dissolved in SDS sample buffer and separated on 8% or 12% SDS gel. SDS-PAGE electrophoresis, transfer to nitrocellulose membrane, and immunoblotting with ECL (Thermo Scientific, USA) detection were performed according to standard manufacturer's protocols (Bio-Rad Laboratories, USA). The following antibodies were used: rabbit monoclonal antibodies against $p21^{\text{Waf1/Cip1}}$ (12D1, number 2947S) (1:1000) and against glyceraldehyde-3-phosphate dehydrogenase (GAPDH, clone 14C10, number 2118S) (1:1000), as well as horseradish peroxidase-conjugated goat anti-rabbit IG (number 7074S) (1:10000). All antibodies were purchased from Cell Signaling, USA. Hyperfilm (CEA) was from Amersham (Sweden). Equal protein loading was confirmed by Ponceau S (Sigma, St. Louis, MO, USA, number P7170) staining.

2.9. RT-PCR Assay. To analyze gene expression, total RNA from cells was isolated with RNesy Micro Kit (Qiagen, USA) according to manufacturer's instructions. cDNA synthesis was performed with 1 μg of total RNA using RevertAid H Minus First Strand cDNA Synthesis Kit (Fermentas, Lithuania) according to manufacturer's instructions. Specific genes were amplified by Taq DNA polymerase (Fermentas, Lithuania) with C1000 Touch Thermal Cycler amplifier (Bio-Rad Laboratories, USA). The program was as follows: hot start-denaturation at 93°C for 3 min, primer annealing at $51\text{--}70^\circ\text{C}$ for 2 min, and then elongation at 72°C for 1 min 30 sec; thereafter, 25–27 cycles of denaturation at 93°C for 45 sec, primer annealing at $51\text{--}70^\circ\text{C}$ for 1 min, elongation at 72°C for 1 min 30 sec, and then final elongation at 72°C for 10 min; primers $p21^{\text{Waf1/Cip1}}$, sense 5' CCA CAT GGT CTT CCT CTG CTG 3', antisense 5' GAT GTC CGT CAG AAC CCA

TG 3', annealing temperature 55°C (316 bp); *SOD1*, sense 5' GGT CCT CAC TTT AAT CCT CTA T 3', antisense 5' CAT CTT TGT CAG CAG TCA CAT T 3', annealing temperature 62°C (96 bp); *SOD2*, sense 5' TGA CAA GTT TAA GGA GAA GC 3', antisense 5' GAA TAA GGC CTG TTG TTC C 3', annealing temperature 56°C (148 bp); *GPXI*, sense 5' CGC CAC CGC GCT TAT GAC CG 3', antisense 5' GCA GCA CTG CAA CTG CCA AGC AG 3', annealing temperature 68°C (238 bp); β -actin gene was used for RNA quantitative control and DNA contamination monitoring: sense primer 5'-GCC GAG CGG GAA ATC GTG CGT-3', antisense 5'-CGG TGG ACG ATG GAG GGG CCG-3', annealing temperature 70°C (507 bp). All primers were obtained from SYNTOL (Russia). The electrophoresis of amplified products was performed in 2% agarose gel with TAE buffer and ethidium bromide. 100 kb DNA ladder (Fermentas, Lithuania) was used as molecular weight markers. Amplified products were visualized in UV light (302 nm) with transilluminator and registered with a digital Canon camera.

2.10. Statistics. All data are presented as the mean and standard error of the mean from at least three separate experiments performed. Statistical differences were calculated using the Student's *t*-test and considered significant at $*P < 0.05$.

3. Results and Discussion

3.1. Cell Viability under Oxidative Stress. H_2O_2 treatment of cultured cells is a commonly used model to test oxidative stress susceptibility in different cell types. Accumulating evidence pointed to a high resistance of mesenchymal stem cells to oxidative stress caused by H_2O_2 [28, 29]. On the other hand, it has been recently reported that some types of human mesenchymal stem cells are very sensitive to H_2O_2 exposure [9]. Susceptibility of hMESC to oxidative stress remains unexplored up to date. We earlier reported that hMESC subjected to prolonged treatment (for 24 h) with H_2O_2 demonstrated a higher resistance compared with human diploid fibroblasts [30]. In this study, a pulse cell treatment with H_2O_2 in varying concentrations from 200 μM to 2 mM for 1 h was applied. Human diploid fibroblasts (HDF) were used as a H_2O_2 sensitive cell model to compare effect of H_2O_2 cytotoxicity on hMESC. Firstly, it was necessary to assess hMESC viability under oxidative stress to examine a sublethal H_2O_2 concentration required for further experiments. Previously we have found out that, at a fixed H_2O_2 concentration, both cytotoxicity and rate of degradation were dependent on the volume of H_2O_2 solution added to the culture medium; therefore the volume used was adjusted proportionally according to the surface area in order to obtain a consistent H_2O_2 cytotoxicity. In addition, it was very important to control plating cell density because the H_2O_2 effect (at equal volume and concentration) on cells was inversely related to cell density; that is, confluent cell cultures were more resistant to H_2O_2 than subconfluent ones. Cell viability was evaluated by MTT assay as a broad indicator of cellular activity that allows estimating a number of viable cells via monitoring mitochondrial dehydrogenase activity. As shown in Figure 1, H_2O_2 affected the cell viability of both cell




FIGURE 1: The viability of hMESC and HDF under oxidative stress. Cells were either treated or not with H₂O₂ at indicated concentrations for 1 h. The percentage of viable cells was evaluated in 24 h after treatment using MTT assay as described in Section 2. Results are shown as a percent of control. Data represent mean ± SEM of at least three independent experiments. ****P* < 0.001 significantly different from the untreated control cells. LD values were 600–700 and 370–400 μM for hMESC and HDF, respectively. Control (Ctr): untreated cells.




FIGURE 2: Gene expression of *SOD1*, *SOD2*, and *GPX1* is enhanced in hMESC in response to H₂O₂ treatment. Exponentially growing cells were treated with the sublethal dose (200 μM) of H₂O₂ for 1 h with following H₂O₂ replacement and then cultured under normal conditions for the indicated time. Total RNA isolated from hMESC was amplified with specific primers for referred genes. β-Actin was used as a loading control.

lines in a dose-dependent manner. LD₅₀ values characterizing cell viability corresponded to 600–700 μM (15–17.5 pmol/cell) for hMESC and 370–400 μM (3.5–4.0 pmol/cell) for HDF. Thus, hMESC were found to be very resistant to H₂O₂ compared to HDF.

3.2. Expression of Antioxidant Enzymes. To examine whether this high resistance to effect of H₂O₂ correlates with the ability of hMESC to effectively scavenge reactive oxygen species (ROS), expression of the genes coding for enzymes involved in the elimination of ROS such as *SOD1*, *SOD2*, and *GPX1* was tested. Previously, we have observed a rapid increase of the intracellular ROS levels in hMESC after exposure to H₂O₂ (data not presented). Here, it was shown that, in untreated cells, the basal levels of mRNA expression of all of the antioxidant enzymes were elevated, while after treatment with 200 μM H₂O₂ an expression level of each enzyme

was upregulated to different extent (Figure 2). Consequently, the significant insensitivity to H₂O₂ was consistent with the enhanced expression levels of the antioxidant enzymes. These findings are in agreement with previous report demonstrating a high resistance of human bone marrow-derived mesenchymal stem cells to oxidative stress [29]. The conflicting findings demonstrating a particular sensitivity of human umbilical cord blood-derived mesenchymal stem cells to H₂O₂ have correlated with the low levels of antioxidant enzyme activity [9].

3.3. A Sublethal Oxidative Stress Induces a Premature Senescence Phenotype in hMESC. In hMESC, phenomenon of H₂O₂-induced premature senescence was not so far described. In all experiments, early-passage cells were used to avoid undesirable replicative senescence of cells, because the major features of both replicative and stress-induced senescence are known to be alike [14, 17, 18]. To test whether hMESC after treatment with a sublethal H₂O₂ concentration (200 μM) could undergo SIPS, we assessed a variety of senescent-associated biomarkers: change of cell morphology, SA-β-Gal staining, increasing of cell size, loss of proliferative potential, cell cycle arrest, and p21^{Waf1/Cip1} (hereafter p21) status. H₂O₂ treatment was found to lead to development of senescent-like morphology: cells become enlarged, flattened, and heterogeneous. It should be noted that, in a part of cell population, we could see such morphological changes within 24 h after H₂O₂ treatment. In addition, senescent cells demonstrated SA-β-Gal staining which increased gradually, and the most remarkable effect was reached at 7 days after treatment: more 95% H₂O₂-treated cells were SA-β-Gal positive (Figures 3(a) and 3(b)). Exponentially growing control cells displayed very weak, if any, increase of SA-β-Gal staining. Importantly, a premature senescence phenotype was maintained during the follow-up period of 21 days (data not shown).

The increased heterogeneity of the cellular size of H₂O₂-treated hMESC was further confirmed and quantified by




FIGURE 3: The sublethal dose of H₂O₂ induces senescent phenotype in hMESC. Cells were treated as indicated in the legend of Figure 2. (a) SA-β-Gal staining. (b) Quantitative assay of SA-β-Gal-positive cells. (c) H₂O₂-induced cell size increase. Typical presentation of forward scatter (FS), reflecting the average cell size (left). Cell size was determined daily: H₂O₂-treated cells were either cultured for 5 days under standard conditions (middle) or were reseeded in 2 days and additionally cultivated for 3 days (right). Data were obtained by light-scattering cytometry with using Win MDI program version 2.8. Data represent mean ± SEM of at least three independent experiments. Significant difference was based on the Student's *t*-test (**P* < 0.05, ****P* < 0.001). Control (Ctr): untreated cells.

light-scattering cytometry of PI-stained cells. As indicated in Figure 3(c), H₂O₂ induced a 2-fold increase of cell size after 5 days compared with control, as measured by the shift in the mean value of the forward scatter. The elevated size of treated cells was sustained constant, at least, for 5 days, whereas the size of control cells was almost not changed. To

test whether treated hMESC retain their increased size being reseeded, after H₂O₂ treatment cells were cultured for 2 days and then reseeded and additionally cultured under normal cell culture conditions for 3 days. As a result, we observed the similar increase of cell size (1.8-fold) in both reseeded and cultured cells for 5 days without reseeding cells (Figure 3(c)).




FIGURE 4: Continued.




FIGURE 4: Induction of the premature senescence in hMESC cells under oxidative stress leads to the permanent arrest of cell cycle and irreversible loss of proliferative potential. Cell treatment was done as described in Figure 2. (a) Growth curve of both H_2O_2 -treated and untreated cells. Cell number was determined daily after cell exposure to H_2O_2 by FACS analysis ($M \pm SEM$, $n = 3$, $*P < 0.05$, $**P < 0.01$). (b) H_2O_2 -treated cells were either cultured for 5 days or were reseeded in 2 days and additionally cultured for 3 days. Flow cytometry analysis of cell cycle phase distribution: the percentage of cells in the G0/G1, S, and G2/M phases (upper panel) ($*P < 0.05$); visualization of phase distribution based on light-scattering analysis (lower panel); SS: side scattering, FL3: PI fluorescence. (c) The expression levels of p21 protein. Representative results of the three experiments are shown in the figure. (d) The levels of p21 mRNA expression. GAPDH and β -actin were used as loading controls. (e) The nuclear localization of Ki67 was tested in control or H_2O_2 -treated cells by immunofluorescence and DAPI staining. Representative photomicrographs of the staining are shown. Images were taken at magnification 100x. Control (Ctr): untreated cells.

Importantly, cell size increase was accompanied with protein content elevation, suggesting protein synthesis in H_2O_2 -treated cells. Together, the results obtained demonstrate cellular hypertrophy within hMESC population in response to H_2O_2 .

3.4. The Permanent Cell Cycle Arrest and Loss of the Proliferative Potential in hMESC Cells Subjected to Sublethal Oxidative Stress. In order to further characterize H_2O_2 -induced senescent-like state of hMESC cells, we analyzed their proliferative potential. Cell number in both untreated and H_2O_2 -treated cell cultures was counted during 5 days. As seen in Figure 4(a), the pattern of growth curves indicates a significant increase (more than two times in 5 days) in the number of proliferating control cells compared with H_2O_2 -treated cells. Consequently, 200 μM H_2O_2 caused a permanent growth arrest, that is, a permanent loss of the proliferative potential. Additionally, a proliferative status of cells was examined by staining with antibodies against proliferation marker Ki67. As seen in Figure 4(e), in 5 days

after H_2O_2 treatment, there were no Ki67-positive cells in the cell culture, while the proliferating control cells had a pronounced staining. As viability of hMESC cells in response to 200 μM H_2O_2 did not decrease appreciably (Figure 1), we suggested that H_2O_2 -induced growth inhibition of hMESC cells could be associated with rather the cell cycle arrest than promotion of cell death.

The analysis of the cell cycle phase distribution in hMESC cells showed that a pulse H_2O_2 treatment led to the arrest in all of the cycle phases (Figure 4(b), upper panel). Treated cells demonstrated the prolonged arrest, at least, for 5 days. The phase distribution of treated cells in each time point tested was characterized with a minor accumulation of cells in G0/G1 phase compared with control cells. The distribution analysis with using light scattering confirmed these findings (Figure 4(b), lower panel). To test whether arrested cells could recover their proliferative potentials, in 2 days after treatment cells were reseeded and cultivated for 3 more days. As expected, reseeded cells also displayed the cell cycle arrest. Moreover, cell cycle phase distributions of both reseeded




FIGURE 5: Dot plots of FITC-annexin V/PI flow cytometry. hMESC were subjected to 900 μM or 3,000 μM H_2O_2 for 1 h with following H_2O_2 replacement and cell cultivation under normal conditions. Apoptosis was detected at indicated time points. Control (Ctr): untreated cells.

and taken-before-reseeding cells were identical (Figure 4(b)). Consequently, the senescent cells were not able to resume proliferation even after being reseeded, indicating the irreversible growth arrest. These observations were confirmed by proliferation assay (Figure 4(a)), which demonstrated that H_2O_2 -treated cells were not able to proliferate normally for 5 days. Overall, these results suggest that cellular senescence in hMESC was induced through growth arrest by H_2O_2 .

3.5. A Permanent Loss of the Proliferative Potential in hMESC Is Accompanied with Elevated Levels of p21. In human mesenchymal stem cells, cyclin-dependent kinase inhibitor p21 was recently shown to be upregulated during H_2O_2 -induced premature senescence [7, 8]. Increased levels of p21 may mediate the initiation of H_2O_2 -induced cell cycle arrest by inhibiting various cyclin-dependent kinases that contribute cell cycle phase progression [14, 16]. To find out whether p21 could be involved in the regulation of H_2O_2 -induced senescence of hMESC, protein and mRNA expression levels of p21 were determined. 200 μM H_2O_2 promoted a significant elevation in protein (Figure 4(c)) and mRNA (Figure 4(d)) expression of p21 in 7 h after treatment. An inducible expression of p21 was upregulated during 1-2 days with a following decline to insignificant, but not control, levels and was accompanied with the cell cycle arrest at the same time (Figure 4(b)). Importantly, the arrested cells thereafter could acquire a senescent morphology (Figure 3(a)) but could not resume proliferation (Figure 4(a)). We assume

that the elevated p21 expression is essential to drive H_2O_2 -induced premature senescence in hMESC. In support of our findings, it has been reported that, in bone marrow-derived mesenchymal stem cells exposed to sublethal doses of H_2O_2 , a rapid decrease of proliferation rate was detected within 3 days and correlated with G1 phase arrest of the cell cycle when p21 was accumulated at the same time [8].

In summary, our findings strongly indicate that hMESC under a sublethal oxidative stress are able to undergo premature senescence.

3.6. Effect of High H_2O_2 Doses on hMESC. H_2O_2 is well documented to cause apoptosis by a dose-dependent manner in various cell types. According to our data, at 200 μM H_2O_2 , no apoptosis was detected in the cell population up to day 21; therefore we tested here whether H_2O_2 at a high concentration causes apoptosis in hMESC. In order to detect apoptosis, annexin V/PI staining was performed. As presented in Figure 5, upper panel, 900 μM H_2O_2 reduced the number of viable cells from 90.3% to 69.7% in 48 h after treatment by increasing the number of AnV/PI+ cells. Remarkably, no significant changes in the number of early apoptotic cells (AnV+) compared with control were observed for 48 h. By contrast, H_2O_2 at higher concentration (3,000 μM) caused apoptosis with no evidence of necrosis in a similar pattern but with much stronger effect (Figure 5, lower panel). Thus, the apoptotic levels in H_2O_2 -treated hMESC were regulated by a dose-dependent manner. According to our




FIGURE 6: The senescent phenotype in hMESC cells treated with 900 μM H₂O₂. (a) H₂O₂-induced inhibition of cell proliferation (M ± SEM, n = 3). (b) SA-β-Gal staining. (c) Quantitative assay of SA-β-Gal-positive cells. (d) Cell hypertrophy detected by light-scattering cytometry. Significant difference was based on the Student's *t*-test (**P* < 0.05, ***P* < 0.01, and ****P* < 0.001). Control (Ctr): untreated cells.

preliminary results, apoptosis was mediated by both caspase-8 and activated caspase-3; however, the exact mechanism of H₂O₂-induced apoptosis in hMESC cells remains to be further elucidated.

Interestingly, 900 μM H₂O₂ triggered not only delayed apoptosis but also led to the emergence of enlarged and flattened cells in the same cell cultures. Cell changes in the presence of both 200 μM and 900 μM H₂O₂ were similar in appearance. These observations prompted us to test if cell hypertrophy could be connected with premature senescence. As shown in Figure 6, 900 μM H₂O₂

actually promoted the senescent morphology and SA-β-Gal staining, permanent growth arrest, and approximately 2-fold increase of cell size, pointing to premature senescence of the main part of cell population. Notably, the major senescence features induced by both 200 μM and 900 μM H₂O₂ were found to be alike. Together, these findings demonstrate that hMESC cells exhibit high apoptosis resistance compared with human mesenchymal stem cells derived from both umbilical cord blood [9] and bone marrow [7], in which H₂O₂ above 200 μM triggered apoptosis, whereas 100–150 μM H₂O₂ induced senescence. Moreover, after exposure

to sublethal doses of H_2O_2 , senescent hMESC acquired the increased stability in culture and displayed enhanced resistance to H_2O_2 -induced apoptosis (data not shown). Many cell types acquire resistance to some apoptotic signals when they become senescent. So, senescent human fibroblasts resist apoptosis induced by oxidative stress or growth factor deprivation but do not resist Fas-mediated apoptosis [31, 32]. Resistance to apoptosis might in part explain the enhanced stability of senescent cells in culture. The mechanisms by which senescent cells resist apoptosis are poorly investigated. The senescence and apoptosis regulatory systems are supposed to communicate probably through their common regulator, p53 tumor suppressor protein [33].

In this study, we have provided the reliable evidence for our hypothesis that hMESC are able to undergo the premature senescence in response to oxidative stress induced by H_2O_2 in a wide range of concentrations from 200 to 900 μM . According to data obtained, entering senescence was accompanied with a rapid initiation of the cellular events, such as the changes of cell phenotype, the increase of SOD1, SOD2, and GPX1 expression, and upregulation of p21 without increase over time, leading to the irreversible cell cycle arrest and loss of proliferative potential. Since 2009, when phenomenon of stress-induced premature senescence in human mesenchymal stem cells was described for the first time, there were only a few publications concerning the oxidative stress-induced premature senescence of human mesenchymal cells derived from bone marrow [7, 8] and umbilical cord blood [9]. Even though both stem cell lines under sublethal stress respond with senescence, the major features of this process, in particular, dynamics of p21 accumulation and decline of cell proliferation rate, were extremely different, depending on the cell context. The precise molecular mechanism required to regulate the oxidative stress-induced premature senescence of human mesenchymal stem cells is far from understanding. Senescence program seems to develop in mesenchymal stem cells as a result of DNA damage response, leading to functional activation of either the p53/p21 or the p16INK4a (p16)/retinoblastoma protein (pRb) pathway, both of which can establish and maintain the growth arrest that is typical of senescence [23, 24]. The cyclin-dependent kinase inhibitors p16 and p21 may maintain pRb in active hypophosphorylated state [34, 35]. In turn, pRb halts cell proliferation by suppressing the activity of transcription factor E2F that regulates cell cycle progression. Our preliminary data, indicating a time- and dose-dependent formation of foci that contain phosphorylated histone H2AX (γ H2AX), activation of both ATM and p53, and upregulation of p21 expression, suggest that in hMESC subjected to sublethal doses of H_2O_2 the senescence process may be controlled by the p53 pathway. In parallel, by monitoring the kinetics of p38 mitogen-activated protein kinase (MAPK) activation in H_2O_2 -induced senescence of hMESC, we have revealed a rapid and continued phosphorylation of p38 MAPK, indicating its possible role in the regulation of the premature senescence [36]. On the other hand, pRb was reported to induce growth arrest as a downstream molecule of p38 MAPK [37]. Taking into consideration these results, we

cannot exclude the possibility that p16/pRb signaling cascade is also implicated in hMESC senescence promotion.

Understanding the mechanisms of senescence process will be of great importance in developing applications of hMESC in regenerative medicine to provide new strategies in autologous transplant and bioengineering. Primarily, hMESC may be applied for cell therapy of infertility associated with decidualization insufficiency. Decidualization of endometrium is known to be an essential process for embryo implantation, placenta forming, and maintenance of pregnancy [38]. A noninvasive and easily available source for isolation of hMESC, high proliferation activity during long-term cultivation, genetic stability, lack of tumorigenicity [39], and low immunogenicity make hMESC a promising source of stem cells for clinical applications, including reproduction technology.

In summary, we have displayed for the first time that hMESC in oxidative stress conditions undergo a premature senescence. Data obtained broaden a conception of mesenchymal stem cell senescence under oxidative stress. Taken together, the findings presented here and the data published allow us to assume that induction of premature senescence might be a common physiological response to sublethal oxidative stress in human mesenchymal stem cells of any origin.

Abbreviations

hMESC:	Human endometrium-derived mesenchymal stem cells
HDF:	Human diploid fibroblasts
SIPS:	Stress-induced premature senescence
H_2O_2 :	Hydrogen peroxide
FACS:	Flow-activated cell sorting
SA- β -Gal:	Senescence-associated β -galactosidase
SOD1:	Cytosolic superoxide dismutase
SOD2:	Mitochondrial superoxide dismutase
GPX1:	Glutathione peroxidase.

Acknowledgments





The authors thank Alekseenko L. L. (Institute of Cytology) for help with RT-PCR. This research was supported by Grants of MCB RAS and from the Russian Foundation for Basic Research (RFBR) (11-04-01581-a).

References

- [1] W. R. Otto and N. A. Wright, "Mesenchymal stem cells: from experiment to clinic," *Fibrogenesis and Tissue Repair*, vol. 4, no. 1, article 20, 2011.
- [2] M. F. Pittenger, A. M. Mackay, S. C. Beck et al., "Multilineage potential of adult human mesenchymal stem cells," *Science*, vol. 284, pp. 143–144, 1999.
- [3] C. Toma, M. F. Pittenger, K. S. Cahill, B. J. Byrne, and P. D. Kessler, "Human mesenchymal stem cells differentiate to a cardiomyocyte phenotype in the adult murine heart," *Circulation*, vol. 105, no. 1, pp. 93–98, 2002.
- [4] M. K. Majumdar, M. A. Thiede, J. D. Mosca, M. Moorman, and S. L. Gerson, "Phenotypic and functional comparison of

- cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells," *Journal of Cellular Physiology*, vol. 176, pp. 57–66, 1998.
- [5] N. E. Sharpless and R. A. DePinho, "How stem cells age and why this makes us grow old," *Nature Reviews Molecular Cell Biology*, vol. 8, no. 9, pp. 703–713, 2007.
 - [6] J. Campisi, S.-H. Kim, C.-S. Lim, and M. Rubio, "Cellular senescence, cancer and aging: the telomere connection," *Experimental Gerontology*, vol. 36, no. 10, pp. 1619–1637, 2001.
 - [7] A. Brandl, M. Meyer, V. Bechmann, M. Nerlich, and P. Angele, "Oxidative stress induces senescence in human mesenchymal stem cells," *Experimental Cell Research*, vol. 317, no. 11, pp. 1541–1547, 2011.
 - [8] J.-S. Kim, E.-J. Kim, H.-J. Kim, J.-Y. Yang, G.-S. Hwang, and C.-W. Kim, "Proteomic and metabolomic analysis of H₂O₂-induced premature senescent human mesenchymal stem cells," *Experimental Gerontology*, vol. 46, no. 6, pp. 500–510, 2011.
 - [9] E. Ko, K. Y. Lee, and D. S. Hwang, "Human umbilical cord blood-derived mesenchymal stem cells undergo cellular senescence in response to oxidative stress," *Stem Cells and Development*, vol. 21, pp. 1877–1886, 2012.
 - [10] D. Wang and D.-J. Jang, "Protein kinase CK2 regulates cytoskeletal reorganization during ionizing radiation-induced senescence of human mesenchymal stem cells," *Cancer Research*, vol. 69, no. 20, pp. 8200–8207, 2009.
 - [11] Á. M. Prendergast, S. Cruet-Hennequart, G. Shaw, F. P. Barry, and M. P. Carty, "Activation of DNA damage response pathways in human mesenchymal stem cells exposed to cisplatin or γ -irradiation," *Cell Cycle*, vol. 10, no. 21, pp. 3768–3777, 2011.
 - [12] J. Cmielova, R. Havelek, T. Soukup et al., "Gamma radiation induces senescence in human adult mesenchymal stem cells from bone marrow and periodontal ligaments," *International Journal of Radiation Biology*, vol. 88, no. 5, pp. 393–404, 2012.
 - [13] E. E. Medrano, S. Im, F. Yang, and Z. A. Abdel-Malek, "Ultraviolet B light induces G₁ arrest in human melanocytes by prolonged inhibition of retinoblastoma protein phosphorylation associated with long-term expression of the p21^(Waf-1/Sdi-1/Cip-1) protein," *Cancer Research*, vol. 55, no. 18, pp. 4047–4052, 1995.
 - [14] O. Toussaint, E. E. Medrano, and T. Von Zglinicki, "Cellular and molecular mechanisms of stress-induced premature senescence (SIPS) of human diploid fibroblasts and melanocytes," *Experimental Gerontology*, vol. 35, no. 8, pp. 927–945, 2000.
 - [15] C.-W. Oh, E. A. Bump, J.-S. Kim, D. Janigro, and M. R. Mayberg, "Induction of a senescence-like phenotype in bovine aortic endothelial cells by ionizing radiation," *Radiation Research*, vol. 156, no. 3, pp. 232–240, 2001.
 - [16] Q. M. Chen, J. C. Bartholomew, J. Campisi, M. Acosta, J. D. Reagan, and B. N. Ames, "Molecular analysis of H₂O₂-induced senescent-like growth arrest in normal human fibroblasts: p53 and Rb control G₁ arrest but not cell replication," *The Biochemical Journal*, vol. 332, pp. 43–50, 1998.
 - [17] P. Dumont, M. Burton, Q. M. Chen et al., "Induction of replicative senescence biomarkers by sublethal oxidative stresses in normal human fibroblast," *Free Radical Biology and Medicine*, vol. 28, no. 3, pp. 361–373, 2000.
 - [18] C. Fripiat, Q. M. Chen, S. Zdanov, J.-P. Magalhaes, J. Remacle, and O. Toussaint, "Subcytotoxic H₂O₂ stress triggers a release of transforming growth factor- β 1, which induces biomarkers of cellular senescence of human diploid fibroblasts," *Journal of Biological Chemistry*, vol. 276, no. 4, pp. 2531–2537, 2001.
 - [19] J. Duan, J. Duan, Z. Zhang, and T. Tong, "Irreversible cellular senescence induced by prolonged exposure to H₂O₂ involves DNA-damage-and-repair genes and telomere shortening," *International Journal of Biochemistry and Cell Biology*, vol. 37, no. 7, pp. 1407–1420, 2005.
 - [20] V. V. Ogryzko, T. H. Hirai, V. R. Russanova, D. A. Barbie, and B. H. Howard, "Human fibroblast commitment to a senescence-like state in response to histone deacetylase inhibitors is cell cycle dependent," *Molecular and Cellular Biology*, vol. 16, no. 9, pp. 5210–5218, 1996.
 - [21] V. S. Romanov, M. V. Abramova, S. B. Svetlikova et al., "p21Waf1 is required for cellular senescence but not for cell cycle arrest induced by the HDAC inhibitor sodium butyrate," *Cell Cycle*, vol. 9, no. 19, pp. 3945–3955, 2010.
 - [22] M. V. Blagosklonny, "Cell cycle arrest is not yet senescence, which is not just cell cycle arrest: terminology for TOR-driven aging," *Aging*, vol. 4, pp. 159–165, 2012.
 - [23] J. Campisi and F. D'Adda Di Fagagna, "Cellular senescence: when bad things happen to good cells," *Nature Reviews Molecular Cell Biology*, vol. 8, no. 9, pp. 729–740, 2007.
 - [24] F. Rodier and J. Campisi, "Four faces of cellular senescence," *Journal of Cell Biology*, vol. 192, no. 4, pp. 547–556, 2011.
 - [25] S. Momtaz and M. Abdollahi, "A comprehensive review of biochemical and molecular evidences from animal and human studies on the role of oxidative stress in aging: an epiphenomenon or the cause," *Asian Journal of Animal and Veterinary Advances*, vol. 7, no. 1, pp. 1–19, 2012.
 - [26] V. I. Zemel'ko, T. M. Grinchuk, A. P. Domnina et al., "Multipotent mesenchymal stem cells of desquamated endometrium: isolation, characterization and use as feeder layer for maintenance of human embryonic stem cell lines," *Tsitologiya*, vol. 53, no. 12, pp. 919–929, 2011.
 - [27] E. B. Burova, I. S. Smirnova, I. V. Gonchar, A. N. Shatrova, and N. N. Nikolsky, "Inhibition of the EGF receptor and ERK1/2 signaling pathways rescues the human epidermoid carcinoma A431 cells from IFN γ -induced apoptosis," *Cell Cycle*, vol. 10, no. 13, pp. 2197–2205, 2011.
 - [28] M.-F. Chen, C.-T. Lin, W.-C. Chen et al., "The sensitivity of human mesenchymal stem cells to ionizing radiation," *International Journal of Radiation Oncology Biology Physics*, vol. 66, no. 1, pp. 244–253, 2006.
 - [29] A. Valle-Prieto and P. A. Conget, "Human mesenchymal stem cells efficiently manage oxidative stress," *Stem Cells and Development*, vol. 19, no. 12, pp. 1885–1893, 2010.
 - [30] E. B. Burova, O. G. Lublinskaya, A. N. Shatrova, A. V. Borodkina, and N. N. Nikolsky, "Comparison of human endometrial stem cells and fibroblasts resistance to oxidative stress," *Tsitologiya*, vol. 54, pp. 478–483, 2012.
 - [31] Q. M. Chen, J. Liu, and J. B. Merrett, "Apoptosis or senescence-like growth arrest: influence of cell-cycle position, p53, p21 and bax in H₂O₂ response of normal human fibroblasts," *The Biochemical Journal*, vol. 347, no. 2, pp. 543–551, 2000.
 - [32] C. G. Tepper, M. F. Seldin, and M. Mudryj, "Fas-mediated apoptosis of proliferating, transiently growth-arrested, and senescent normal human fibroblasts," *Experimental Cell Research*, vol. 260, no. 1, pp. 9–19, 2000.
 - [33] A. Seluanov, V. Gorbunova, A. Falcovitz et al., "Change of the death pathway in senescent human fibroblasts in response to DNA damage is caused by an inability to stabilize p53," *Molecular and Cellular Biology*, vol. 21, no. 5, pp. 1552–1564, 2001.
 - [34] G. H. Stein, L. F. Drullinger, A. Soulard, and V. Dulić, "Differential roles for cyclin-dependent kinase inhibitors p21 and p16

- in the mechanisms of senescence and differentiation in human fibroblasts," *Molecular and Cellular Biology*, vol. 19, no. 3, pp. 2109–2117, 1999.
- [35] J. J. L. Jacobs and T. De Lange, "Significant role for p16INK4a in p53-independent telomere-directed senescence," *Current Biology*, vol. 14, no. 24, pp. 2302–2308, 2004.
- [36] J. Maruyama, I. Naguro, K. Takeda, and H. Ichijo, "Stress-activated MAP kinase cascades in cellular senescence," *Current Medicinal Chemistry*, vol. 16, no. 10, pp. 1229–1235, 2009.
- [37] H. Iwasa, J. Han, and F. Ishikawa, "Mitogen-activated protein kinase p38 defines the common senescence-signalling pathway," *Genes to Cells*, vol. 8, no. 2, pp. 131–144, 2003.
- [38] J. J. Kim, H. S. Taylor, Z. Lu et al., "Altered expression of HOXA10 in endometriosis: potential role in decidualization," *Molecular Human Reproduction*, vol. 13, no. 5, pp. 323–332, 2007.
- [39] A. P. Domnina, I. I. Fridlyanskaya, V. I. Zemelko et al., "Mesenchymal stem cells from human endometrium do not undergo spontaneous transformation during long-term cultivation," *Cell and Tissue Biology*, vol. 7, pp. 221–226, 2013.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

