Review Article

Clinical Aspects and Management of Levodopa-Induced Dyskinesia

Nicola Tambasco,¹ Simone Simoni,¹ Erica Marsili,¹ Elisa Sacchini,¹ Donatella Murasecco,¹ Gabriela Cardaioli,¹ Aroldo Rossi,¹ and Paolo Calabresi^{1,2}

¹*Clinica Neurologica, Azienda Ospedaliera—Università di Perugia, S. Andrea delle Fratte, 06156 Perugia, Italy*

²*IRCCS Fondazione S. Lucia, Via Ardeatina 306, 00179 Roma, Italy*

Correspondence should be addressed to Nicola Tambasco, n.tambasco@libero.it

Received 5 February 2012; Accepted 2 April 2012

Academic Editor: Gilberto Fisone

Copyright © 2012 Nicola Tambasco et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In Parkinson's disease, one of the most troublesome dilemmas is the treatment of levodopa-induced dyskinesia. After a few years, chronic treatment with levodopa is associated with the development of dyskinesias. Strategies to delay or to reduce dyskinesias are based on the change of levodopa dosing or the early use of dopamine agonists. Dopamine agonists with different pharmacological profile are available. Our paper was aimed to analyse the clinical impact and the management of dyskinesias with dopamine agonists.

1. Introduction

Four decades after its introduction, levodopa remains the most effective agent to improve motor symptoms in PD, but chronic use is associated with the emergence of motor fluctuations, defined as a loss of clinical benefit before the next levodopa dose (wearing off), abnormal involuntary movements (dystonia, chorea, and athetosis—collectively referred to as dyskinesia) [1, 2], and nonmotor complications, as behavioural and cognitive changes [3]. Levodopa is initially well tolerated in most of the cases and allows a substantial improvement of motor performances despite its erratic pharmacokinetics [1, 4]. With the disease progression, therapeutic window of levodopa narrows, and the duration of each dose shortens. Motor fluctuations usually precede dyskinesias [5], and it has been observed that the development of one is a risk factor for the development of the other [5].

Although more commonly associated with levodopa, dyskinesias can also occur with dopamine agonist monotherapy [6–8]. The development of dyskinesia in some patients treated with dopamine agonists that have relatively long half-lives (ropinirole, 6 h; pramipexole, 8 h) or very long half-lives (cabergoline, 68 h) suggests that, to some extent, even

dopamine stimulation provided in a continuous fashion can cause dyskinesias.

2. Epidemiology and Clinical Aspects of Motor Complications

The three most important risk factors positively associated with increased occurrence of dyskinesias are younger age at disease onset [9, 10], longer disease duration [11, 12], and longer duration of pulsatile dopaminergic treatment (typically, levodopa) [13, 14]. The first two factors are interrelated and almost all patients with early-onset PD [15] develop dyskinesias, whereas they are less frequent in patients with late-onset PD [16]. PD patients with early disease onset have a high probability to carry mutations for monogenic PD forms, and therefore, early onset and genetic predisposition are two overlapping and possibly interrelated risk factors. Other risk factors associated with increased risk of dyskinesias are female gender [17, 18] and the occurrence of specific polymorphisms for dopamine receptors or dopamine transporters [19–21].

Dyskinesias more commonly appear as choreiform, but in some cases, they may resemble dystonia, myoclonus, or

other movement disorders. Peak-dose dyskinesias are the most common type of dyskinesia. They occur during peaks of levodopa-derived dopamine in the brain, when the patient is otherwise experiencing a beneficial response (the “on” state). Peak-dose dyskinesias worsen with increases in dopaminergic dose and lessen with its reductions. In certain cases, dyskinesias seem to appear with a more particular pattern, as dyskinesia-improvement-dyskinesia. This is termed diphasic dyskinesia, and it tends to occur when levodopa-derived dopamine concentrations are increasing or decreasing, whereas the clinical condition of the patient turns “on” and “off” [22]. Diphasic dyskinesias are typically displayed with large-amplitude stereotypic, rhythmic, and repetitive movements, more often of the legs, that may be associated with Parkinsonian features in other body regions. In extreme cases, patients treated with levodopa can cycle between “on” periods, which are complicated by disabling dyskinesias, and “off” periods in which Parkinsonism is uncontrolled and the patient is akinetic and frozen.

Motor complications occur in about 50% of patients with PD who have been in therapy with levodopa for more than 5 years, and in almost 100% of patients with young-onset disease [23, 24]. Achieving an acceptable clinical control once these motor fluctuations have appeared is usually a relatively simple matter, nearing together the levodopa doses or adding medications that reduce “off” time. However, when a patient develops peak-dose dyskinesias too, it becomes difficult to smooth the clinical response. Although for many patients, dyskinesias are not disabling, they create a barrier to adequate treatment of fluctuations and Parkinsonian symptoms.

3. Pathophysiology of Dyskinesias

A primary condition in LID pathophysiology is the presence of dopaminergic cell loss in substantia nigra [25–27]. The nonappearance of dyskinesia in normal humans chronically treated with levodopa (i.e., mistaken diagnosis [28]) and its rapid emergence in PD patients either with late diagnosis or a young onset, where denervation is high at diagnosis [15, 29, 30], heavily support this theory. Moreover, the progression of nigral denervation seems to be closely related with the lowering of the dyskinesia onset threshold in MPTP-exposed primates [31]. Nonetheless, denervation cannot be the unique factor responsible for dyskinesia, whereas not all patients with advanced illness and extensive nigral denervation develop dyskinesia when treated with levodopa [32, 33]. Thus, a chronic dopaminergic stimulation on a denervated substantia nigra induces a process of sensitisation such that each following administration modifies the response to subsequent dopaminergic treatments. This process, called priming, increases over time of treatment the chance of eliciting dyskinesias and, once dyskinesias have been established, their severity. The priming process, which is responsible for the insidious evolution of dyskinesias over time of treatment, is associated with changes in receptors for dopamine or other neurotransmitters [34, 35]. A crucial role has been postulated for both dopamine receptors and NMDA glutamate receptors in the induction of priming; this mechanism could

be regarded as an increased responsiveness of postsynaptic striatal dopamine receptors (mainly D₁-like), which are activated in conjunction with glutamatergic inputs [1]. Dyskinesias are probably generated by a persistent enhancement of the responsiveness of striatal medium-sized spiny neurones to dopaminergic treatment. This is an aftermath of dopamine depletion and is associated with overexpression of specific components of the signal transduction machinery. If protracted, this condition may ultimately lead to long-term changes in gene expression, which will permanently affect the function of striatal medium spiny neurones [36]. Following priming, the development of dyskinesias largely depends on two additional factors, the pulsatile administration of levodopa (or another short-acting dopaminergic agent) and the severity of dopaminergic denervation in the striatum. The latter plays an important role in setting the threshold required in developing dyskinesias [37]. A direct relationship between the severity of striatal denervation and the time required to develop dyskinesias has been demonstrated in PD patients [38] and has been indirectly confirmed by the finding that patients with dopa-responsive dystonia, who have Parkinsonism without nigrostriatal denervation, uncommonly develop dyskinesias [39].

In early PD patients, levodopa-derived dopamine is packaged into synaptic vesicles by vesicular monoamine transporter 2 (VMAT-2), stored, and released in both tonic and phasic bursts in response to impulse flow [40, 41], in order to preserve dopamine receptors from levodopa plasma concentration fluctuations and, therefore, to maintain physiological dopaminergic transmission [42, 43]. With the progression of the disease, and the striatal dopaminergic cell loss, the formation of dopamine from levodopa and its storage capacity are increasingly compromised, and the response to levodopa becomes dominated by its pharmacokinetic characteristics and general bioavailability [4]. Thus, in advanced PD, peak concentrations of drug in plasma become coincidental with the expression of dyskinesia. As observed in animal models, the continuous release of dopamine leads to improvements in motor function and, together, to a marked reduction in the expression of involuntary movements [44]. These studies support the clinical findings that the continuous intravenous or intraduodenal administration of levodopa or the continuous subcutaneous or intravenous infusion of apomorphine results in improved motor response but also with a marked reduction of dyskinesia [45, 46].

Other mechanisms are involved to explain the underlying cause and expression of dyskinesia. Although dopamine agonists when used as monotherapy in early PD are associated to a lower incidence of dyskinesia, involuntary movements are still observed, reflecting some kind of activity at the postsynaptic dopamine receptor level, as dopamine agonists are not dependent on the presence of presynaptic terminals.

Subtle changes in D₁ and D₂ receptor density as well as the complex interaction between receptor activation and synaptic plasticity [1] have been proposed as playing significant roles in dyskinesia induction and expression. Although the exact molecular mechanisms of LID still remain to be fully elucidated, exaggerated signalling of the striatal D₁ [47–49], the reduction of the modulating function of D₂/D₃

TABLE 1: Pharmacological characteristics of dopamine agonists.

	Pramipexole	Ropinirole	Rotigotine	Pergolide	Bromocriptine	Cabergoline	Apomorphine	Lisuride
D ₁	0	0	+	+	–	0/+	+++	–
D ₂	+++	+++	+++	++++	++	+++	++	++++
D ₃	++++	++++	++++	+++	++	++	++++	+++
Type	Nonergot	Nonergot	Nonergot	Ergoline	Ergoline	Ergoline	Morphine deriv.	Ergoline
Routes	os	os	td	os	os	os	sc	sc
Metabolism	—	Hepatic	Hepatic	Hepatic	?	Hepatic	Hepatic	Hepatic
Elimin.	Urine	Urine	Urine/fecal	Urine/fecal	Fecal	Fecal/urine	Urine/fecal	Urine/fecal
Half-life (h)	8–12	5–6	5–7	27	12–14	63–69	40 min.	2

td: transdermal; sc: subcutaneous.

receptors [42, 43, 50–52], and the interaction between D₂ and A2A adenosine receptors [53] have been implicated in both rodents and primates, suggesting that a normalisation of signalling may be beneficial in the treatment of dyskinesia.

In clinical practice, postsynaptic mechanisms can be partially explained by the dopamine agonists capability to prime for involuntary movements. Switching from a chronic dopamine agonist administration that usually results in a low expression and intensity of dyskinesia to an equivalent dose of levodopa in fact immediately results in the appearance of dramatic involuntary movements [54, 55]. These findings suggest that dopamine agonists principally prime for, but less commonly express dyskinesia. Vice versa, when considering the expression of dyskinesia in patients with PD with a history of levodopa exposure, switching to a dopamine agonist after the introduction of levodopa, established dyskinesia still occurred [56]. Moreover, patients receiving a combination of levodopa and the dopamine D₂/D₃ agonist pramipexole showed a level of dyskinesia that was greater than the additive effect of the individual drug [57]. Once established, dopamine agonists produce the same pattern of dyskinesia although its intensity is reduced, suggesting that agonists do not express dyskinesia to the same extent as levodopa [54, 55, 58]. Both the lower priming for dyskinesia and the lower expression of involuntary movements by dopamine agonists may be a reflection of their more specific pharmacology compared to levodopa.

4. Reducing or Delaying Parkinsonian Dyskinesias

Any type of exogenous dopaminergic stimulation in a denervated striatum can cause dyskinesias [59], but pulsatile stimulation produced by short-acting drugs (as typically occurs with levodopa) particularly favours their occurrence [60]. The expression LID is still currently used, although levodopa is not the only drug causing dyskinesias in PD patients [61]. Based on published series, it has been estimated that PD patients treated for less than 5 years have a 11% risk of developing dyskinesias, those treated for 6–9 years have a risk of 32%, whereas patients treated for more than 10 years have a risk of 89% [13].

Levodopa, however, seems to be the most important factor in inducing dyskinesia expression in chronically treated

PD patients; therefore, it appears that the benefit of initial treatment with a dopamine agonist in lowering the incidence of dyskinesias is related to the ability of the agonist to delay the need for levodopa [12, 62]. Moreover, experimental data suggest that the administration of long-acting dopamine agonists results in significantly less dyskinesia than does levodopa [63, 64] and other short-acting agents administered in a pulsatile fashion [65]. However, once a long-acting agonist is administered to animals already primed to exhibit dyskinesias with levodopa, the resultant dyskinesias are comparable to those seen in the levodopa group [63]. Clinical studies randomly assigning patients to initial treatment with a dopamine agonist or levodopa have shown a lower risk for dyskinesias in the groups treated with pramipexole [7], ropinirole [8, 12], bromocriptine [66, 67], pergolide [68], and cabergoline [6]; nevertheless, once levodopa was added, the rate of development of dyskinesias was similar in both groups.

One therapeutic strategy that has been tried in this sense is to use higher doses of a dopamine agonist to reduce both the total daily levodopa dose and its frequency [69] or to gradually substitute a dopamine agonist for levodopa [70]. Unfortunately, these strategies are unsatisfactory and typically reduce dyskinesias at the expense of less control of Parkinsonian symptoms. Indeed, the evidence that early levodopa exposure adversely affects the course of disease and leads to disabling dyskinesias and motor fluctuations constituted the rationale for the initial treatment with dopamine agonist.

5. Different Profile and Efficacy of Dopamine Agonists in Reducing Dyskinesia

In order to create a valid alternative to levodopa, and with the aim of eliminating its related complications, many different drugs acting on dopaminergic receptors have been developed and studied during the last years. They have different metabolism, plasma half-life, affinity to receptors subtypes, excretion, and routes of administration (Table 1). Moreover, these drugs have different efficacies on reducing the incidence of dyskinesia, improving motor symptoms, and reducing the daily levodopa dose (Table 2, Figure 1).

Initially dopamine agonists have been used as adjuvant therapy to improve levodopa-induced complications, but

TABLE 2: Adjuvant therapy versus placebo.

	Pramipexole	Ropinirole	Pergolide*	Bromocriptine	Cabergoline
Off-time reduction (h/day)	-1.81	-0.93	-1.60	-1.78	-1.29
LEDD red (mg/day)	-114.82	-119.81	-183.90	-52.17	-149.60
UPDRS ADL reduction (pts)					-1.78
UPDRS III reduction (pts)		-4.80			-1.74
Incidence of dyskinesia (OR)	2.63	3.21	4.64	2.52	1.44

* Based on data from just one trial [71].

FIGURE 1: Effects of dopamine agonists on (a) reducing off time, (b) reducing levodopa daily dose, and (c) inducing dyskinesia.

once their effects on delaying the need for levodopa have been demonstrated, they have often been prescribed before the introduction of levodopa. Patients receiving dopamine agonists rather than levodopa as initial monotherapy showed a reduced risk for developing dyskinesias [7, 8, 12, 62, 72–76] (Table 3).

5.1. Dopamine Agonists Monotherapy and the Risk of Dyskinesia. The CALM-PD trial (Comparison of the Agonist Pramipexole versus Levodopa on Motor Complications of Parkinson's Disease) was a randomised controlled trial

evaluating the risk of developing dyskinesias in patients with early PD initially treated with either pramipexole or levodopa. After 24 months, pramipexole-treated patients were receiving pramipexole plus levodopa, compared with levodopa alone. A minority of pramipexole-treated patients reached the endpoint of time to first occurrence of wearing off, dyskinesias, or on-off motor fluctuations (27.8% versus 50.7%, $P < 0.001$). Moreover, a significantly lower incidence of dyskinesias (9.9% versus 30.7%, $P < 0.001$) also has been demonstrated in patients in the pramipexole group. However, after a mean 6-year followup, >90% of patients were receiving levodopa therapy regardless of their initial treatment assignment. Compared to those taking pramipexole, patients initially treated with levodopa had significantly more dyskinesias (20.4% versus 36.8%), but there was no difference between groups in the incidence of disabling or painful dyskinesias [62, 74]. Interestingly, 5 subjects taking pramipexole developed dyskinesias before the supplemental levodopa, and 4 of them had no prior levodopa exposure [73]. No significant difference in the Lang-Fahn activities of daily living dyskinesia score was observed (1.3 versus 1.1 with pramipexole, $P < 0.06$) [7, 62, 72–74].

In a randomised, double-blind 5-year study of patients with early PD, the risk of developing dyskinesias after initial monotherapy with ropinirole was less than with levodopabenserazide (hazard ratio (HR), 2.82 (1.78, 4.44); $P < 0.001$) [8]. However, many of these patients eventually required supplemental levodopa to control the symptoms of the disease [8, 12]. When patients receiving ropinirole monotherapy required the addition of levodopa, the risk for developing dyskinesias increased and then did not differ significantly from that associated with levodopa alone [12]. The use of ropinirole as monotherapy, with only later addition of levodopa, delayed the onset of dyskinesias by up to 3 years, although it was associated with a higher incidence of neuropsychiatric complications than levodopa monotherapy.

Apomorphine, a subcutaneous nonergolinic dopaminergic agent, has been studied in 2 retrospective chronic monotherapy trials in which no oral anti-parkinsonian therapies were permitted from the time the pump was turned on in the morning until it was turned off in the evening [77, 78]. The mean maximum reduction of dyskinesia per patient was 64% ($P < 0.005$), and the mean time to achieve maximal dyskinesia improvement was 12.1 months.

Lisuride, another subcutaneously administered dopamine agonist, given as a continuous daytime infusion via pump, has been utilised as a strategy for minimising dyskinesias in 40 patients with advanced, levodopa-responsive PD

TABLE 3: Series on adjuvant therapy with dopamine agonists*. In italic, dyskinesia evaluation.

Author	Duration	Characteristics of participants	Interventions	Primary outcomes	Secondary outcomes
Pewee <i>et al.</i> [79]	(6 months)	N: 302; MFs. Mean duration of PD: 8.5 y	<i>Pramipexole (n = 201) versus placebo (n = 101)</i>	<i>Disability; motor complications; on/off time</i>	SE
Pahwa <i>et al.</i> [80, 81]; Sethi <i>et al.</i> [82, 83]; Stacy <i>et al.</i> [84]; Stocchi <i>et al.</i> [85, 86]	(24 weeks)	N: 393; MFs. Mean duration of PD: 8.6 y	<i>Ropinirole (24-h) (n = 202) versus placebo (n = 191)</i>	<i>Disability; patient-rated QoL; on/off time; levodopa dose</i>	SE <i>Depression sleep scales</i>
Oertel <i>et al.</i> [87]; Pogarell <i>et al.</i> [88]	(32 weeks)	N: 363 (354 analysed); MFs. Mean duration of PD: 7.8 y	<i>Pramipexole (n = 180) versus placebo (n = 183)</i>	<i>Disability; off time; levodopa dose</i>	SE
Wong <i>et al.</i> [89]	(15 weeks)	N: 150; Mean duration of PD: 4.4 y	<i>Pramipexole (n = 73) versus placebo (n = 77)</i>	<i>Disability; off time</i>	SE
Musch and Bonura [90]	(24 weeks)	N: 218; on levodopa. Mean duration of PD: NA	<i>Cabergoline (n = 145) versus placebo (n = 73)</i>	<i>Disability; off time; levodopa dose</i>	SE
Pinter <i>et al.</i> [91]	(11 weeks)	N: 78; MFs. Mean duration of PD: 8.2 y	<i>Pramipexole (n = 34) versus placebo (n = 44)</i>	<i>Disability; off time; levodopa dose</i>	SE
Wermuth [92]	(12 weeks)	N: 69; MFs. Mean duration of PD: 10 y (range: 3–27 y)	<i>Pramipexole (n = 36) versus placebo (n = 33)</i>	<i>Disability; motor complications; off time; levodopa dose</i>	SE
Lieberman <i>et al.</i> [93]; Weiner <i>et al.</i> [94]	(32 weeks)	N: 360; MFs. Mean duration of PD: 9.2 y	<i>Pramipexole (n = 181) versus placebo (n = 179)</i>	<i>Disability; on/off time; levodopa dose</i>	SE
Guttman [95]	(9 months)	N: 247; MFs. Mean duration of PD: 7 y (range: 0.67–36 y)	<i>Pramipexole (n = 79) versus bromocriptine (n = 84) versus placebo (n = 83)</i>	<i>Disability; off time</i>	SE
Kreider <i>et al.</i> [96]; Lieberman <i>et al.</i> [97]	(6 months)	N: 149; predictable MFs. Mean duration of PD: 9 y	<i>Ropinirole (n = 95) versus placebo (n = 54)</i>	<i>Disability; motor complications; off time; levodopa dose</i>	SE
Rascol <i>et al.</i> [98, 99]	(12 weeks)	N: 46; not optimally controlled with levodopa. Mean duration of PD: 8 y	<i>Ropinirole (n = 23) versus placebo (n = 23)</i>	<i>Disability; motor complications; off time</i>	SE

TABLE 3: Continued.

Author	Duration	Characteristics of participants	Interventions	Primary outcomes	Secondary outcomes
Steiger et al. [100]	(6 months)	N: 37; MFs. Mean duration of PD: 12.8 y (range: 3–33 y)	Cabergoline (n = 19) versus placebo (n = 18)	Disability; motor complications; off time; levodopa dose	SE
Hutton et al. [101]; Lieberman and Hutton [102]; Schoenfelder et al. [103]	(24 weeks)	N: 188; MFs. Mean duration of PD: 10.6 y (range: 2–30 y)	Cabergoline (n = 123) versus placebo (n = 65)	Disability; on/off time; levodopa dose	SE
Olanow et al. [104]	(6 months)	N: 376; MFs. Mean duration of PD: 10.9 y	Pergolide (n = 189) versus placebo (n = 187)	Disability; motor complications; off time; levodopa dose	SE
Temlett et al. [105]	(5 weeks)	N: 44 (40 analysed); Mean duration of PD: 13.4 y	Bromocriptine (n = 22) versus placebo (n = 18)	Levodopa dose	SE
Toyokura et al. [106]	(8 weeks)	N: 222; not optimally controlled with levodopa. Mean duration of PD: 6.6 y	Bromocriptine (n = 114) versus placebo (n = 108)	Motor complications; on/off time	SE
Schneider and Fischer [107]	(4 weeks)	N: 40; not optimally controlled with levodopa. Mean duration of PD: 9.1 y	Bromocriptine (n = 20) versus placebo (n = 20)	On/off time; levodopa dose	
Jansen [108]	(5 months)	N: 23; not optimally controlled with levodopa. Mean duration of PD: 8.7 y	Bromocriptine (n = 12) versus placebo (n = 11)	Disability	

* Performed on PD patients, parallel groups, double blind.

MFs: motor fluctuations; SE: side effects.

characterised by motor fluctuations and dyskinesias [109]. After 4 years, the lisuride-treated patients had improved their baseline dyskinesia scores (measured by AIMS) by 49% ($P < 0.0001$), whereas the levodopa-treated patients had worsened their scores by 59% ($P < 0.0001$).

5.2. Long-Acting Dopamine Agonists and the Risk of Dyskinesia. In animal-model studies, the long-acting dopamine agonists have been demonstrated to prevent or reduce the onset time for LIDs. In a study of monkeys with MPTP-induced parkinsonism, small doses of subcutaneously administered cabergoline, a D_2 -selective dopamine agonist with a relatively long half-life, were added as adjuvant therapy to orally administered levodopa/benserazide (100/25 mg) for 1 month, showing significantly lower dyskinesia scores (sum for all body segments) than when levodopa/benserazide was given alone for 1 month ($P < 0.01$).

A report on the effect of cabergoline compared to levodopa showed a reduced incidence of dyskinesias [110]. Nevertheless, more recently, an increased incidence of dyskinesia and confusion in patients treated with bromocriptine was reported [111].

5.3. Differences among Drugs in Adjuvant Therapy. A recent systematic meta-analysis, which performs indirect comparisons among three classes of drugs, including nondopaminergic agents as catechol-O-methyl transferase inhibitors (COMTIs) or monoamine oxidase type B inhibitors (MAO-BIs), used as add-on (adjuvant) treatment to levodopa therapy in PD patients with motor complications, suggests that dopamine agonists may provide more effective symptomatic control [112].

5.3.1. Off-Time Reduction. There is no (or little) evidence of a difference across the different dopamine agonists for the overall reduction in off-time [pramipexole (-1.81 hours/day, CI -2.19 to -1.43 ; $P < 0.00001$); bromocriptine (-1.78 hours/day, CI -2.91 to -0.65 ; $P = 0.002$); pergolide (-1.60 hours/day, CI -2.57 to -0.63 ; $P = 0.001$); cabergoline (-1.29 hours/day, CI -1.89 to -0.69 ; $P < 0.0001$); ropinirole (-0.93 hours/day, CI -1.53 to -0.33 ; $P = 0.002$)] [112].

5.3.2. Levodopa Daily Dose Reduction. The largest reduction was with pergolide (-183.90 mg/day, CI -259.09 to -72.71 ; $P = 0.001$), though this was based on data from just one trial [71]. Cabergoline reduced the required levodopa dose by 149.60 mg/day (CI -208.79 to -90.41 ; $P < 0.00001$), ropinirole by 119.81 mg/day (CI -150.63 to -89.00 ; $P < 0.00001$), pramipexole by 114.82 mg/day (CI -143.01 to -86.64 ; $P < 0.00001$), and bromocriptine by 52.17 mg/day (CI -95.16 to -9.18 ; $P = 0.02$) [112].

5.3.3. UPDRS Scores Improvement. The agonist pramipexole appeared to produce larger improvements for UPDRS motor score (-6.31 points, CI -7.69 to -4.93 ; $P < 0.00001$) compared to ropinirole (UPDRS motor: -4.80 points, CI -7.32 to -2.28 ; $P = 0.0002$) and cabergoline (UPDRS motor: -1.74 points, CI -3.78 to 0.30 ; $P = 0.09$) [112].

5.3.4. Incidence of Dyskinesia. The analysis included 6476 participants, which represented 85% of the 7590 randomised participants included in the meta-analysis. Compared to placebo, the incidence of dyskinesia was increased with adjuvant therapy. The incidence of dyskinesia was greatest with pergolide (OR 4.64, CI 3.09 to 6.97; $P < 0.00001$), although the data were obtained from just one trial [71], followed by ropinirole (OR 3.21, CI 1.98 to 5.21; $P < 0.00001$), pramipexole (OR 2.63, CI 2.01 to 3.42; $P < 0.00001$), bromocriptine (OR 2.52, CI 1.42 to 4.48; $P = 0.002$), and cabergoline (OR 1.44, CI 0.96 to 2.16; $P = 0.08$) [112].

Though this meta-analysis indirectly compares several series on dopaminergic agents as adjuvant treatment, the need of large randomised studies that directly compare different agents administered as monotherapy with patient-rated overall quality of life and health economic measures as primary outcomes is recommended.

6. Alternative Treatments to Reduce Dyskinesia

As seen earlier, the primary therapeutic strategy for managing LIDs in PD patients is to delay their occurrence through delaying the introduction of levodopa therapy administering dopaminergic agents.

Once dyskinesias have occurred, other strategies should be attempted: (1) substitution of immediate release for controlled-release levodopa. The immediate-release preparation is easier to adjust, as onset of its effects is sooner, and duration of action (and dyskinesias) is shorter than with controlled-release preparations. For the same reason, agents that prolong the half-life of levodopa, such as entacapone, should be stopped; (2) discontinuation of other therapy that may embitter dyskinesias, as dopamine agonists or other factors delaying dopamine degradation as selegiline and rasagiline; (3) incrementation of the number of administrations of levodopa, in lower doses; (4) addition of an antidyskinetic agent as amantadine, an NMDA receptor antagonist. Diphasic dyskinesias that may manifest at the beginning and the end of a dosing cycle should be managed by utilising more frequent doses of levodopa, and the therapy should be sewed on the patient [113].

6.1. Amantadine. The NMDA receptor-binding and neurotoxic effects of excessive glutamate have led to the hypothesis that an NMDA antagonist may have antidyskinetic effects and reduce the severity of LIDs. Amantadine has been studied as adjuvant treatment in levodopa-treated patients experiencing motor complications, including dyskinesias, with the aim of reducing these effects without worsening Parkinsonian symptoms [114–117]. Three randomized placebo-controlled crossover clinical studies in a group of 53 PD patients showed a reduction (up to 60%) in the severity of LIDs after challenge with acute levodopa administration, without impacting the beneficial effects of levodopa on motor function.

6.2. Clozapine. It is an atypical antipsychotic that has been assessed for the treatment of drug-induced psychosis in PD.

It may also be effective in decreasing dyskinesias [70], and a few studies have focused on its antidyskinetic effect [118, 119].

6.3. Intraduodenal Levodopa. It provides direct delivery of levodopa to the duodenum and jejunum. The method involves insertion of a permanent access tube in the abdominal wall by percutaneous endoscopic gastrostomy. Several clinical studies have been conducted using this approach, demonstrating significant reductions in "off" time and dyskinesia after 6 months. It may be an option for patients with marked fluctuations and dyskinesia in whom deep-brain stimulation (DBS) is contraindicated or not possible due to advanced age, or it may provide an alternative to DBS.

6.4. Surgical Treatment. Patients with PD who may benefit from surgery include those who have substantial dyskinesias unresponsive to medication adjustments, are levodopa responsive, do not have dementia, and do not have neuropsychiatric impairment [80]. DBS is the most frequently performed surgery for PD in North America [80]. In patients with advanced PD, DBS of the globus pallidus interna (GPi) or the subthalamic nucleus (STN) has been shown to reduce dyskinesia severity by up to 89% [120, 121] and to reduce the duration of dyskinesias by 86% [122]. It provides significant improvement in Parkinsonian motor features and allows a reduction of dyskinesias, in part through the subsequent reduction of levodopa [123, 124].

References

- [1] P. Calabresi, M. D. Filippo, V. Ghiglieri, N. Tambasco, and B. Picconi, "Levodopa-induced dyskinesias in patients with Parkinson's disease: filling the bench-to-bedside gap," *The Lancet Neurology*, vol. 9, no. 11, pp. 1106–1117, 2010.
- [2] F. Stocchi, M. Tagliati, and C. W. Olanow, "Treatment of levodopa-induced motor complications," *Movement Disorders*, vol. 23, supplement 3, pp. S599–S612, 2008.
- [3] N. Tambasco, V. Belcastro, A. Gallina, A. Castrìto, P. Calabresi, and A. Rossi, "Levodopa-induced breathing, cognitive and behavioral changes in Parkinson's disease," *Journal of Neurology*, pp. 2296–2299, 2011.
- [4] J. G. Nutt, "Pharmacokinetics and pharmacodynamics of levodopa," *Movement Disorders*, vol. 23, supplement 3, pp. S580–S584, 2008.
- [5] R. A. Hauser, M. P. McDermott, and S. Messing, "Factors associated with the development of motor fluctuations and dyskinesias in Parkinson disease," *Archives of Neurology*, vol. 63, no. 12, pp. 1756–1760, 2006.
- [6] F. Bracco, A. Battaglia, C. Chouza et al., "The long-acting dopamine receptor agonist cabergoline in early Parkinson's disease: final results of a 5-year, double-blind, levodopa-controlled study," *CNS Drugs*, vol. 18, no. 11, pp. 733–746, 2004.
- [7] The Parkinson Study Group, "Pramipexole vs levodopa as initial treatment for Parkinson Disease: a 4-year randomized controlled trial," *Archives of Neurology*, vol. 61, no. 7, pp. 1044–1053, 2004.
- [8] O. Rascol, D. J. Brooks, A. D. Korczyn, P. P. De Deyn, C. E. Clarke, and A. E. Lang, "A five-year study of the incidence of dyskinesia in patients with early Parkinson's disease who were treated with ropinirole or levodopa," *The New England Journal of Medicine*, vol. 342, no. 20, pp. 1484–1491, 2000.
- [9] V. Kostic, S. Przedborski, E. Flaster, and N. Sternic, "Early development of levodopa-induced dyskinesias and response fluctuations in young-onset Parkinson's disease," *Neurology*, vol. 41, no. 2 I, pp. 202–205, 1991.
- [10] A. Schrag, A. Hovris, D. Morley, N. Quinn, and M. Jahan-shahi, "Young- versus older-onset Parkinson's disease: impact of disease and psychosocial consequences," *Movement Disorders*, vol. 18, no. 11, pp. 1250–1256, 2003.
- [11] The Parkinson Study Group, "Impact of deprenyl and tocopherol treatment on Parkinson's disease in DATATOP subjects not requiring levodopa," *Annals of Neurology*, vol. 39, no. 1, pp. 29–36, 1996.
- [12] O. Rascol, D. J. Brooks, A. D. Korczyn et al., "Development of dyskinesias in a 5-year trial and ropinirole and L-dopa," *Movement Disorders*, vol. 21, no. 11, pp. 1844–1850, 2006.
- [13] G. Fabbrini, J. M. Brotchie, F. Grandas, M. Nomoto, and C. G. Goetz, "Levodopa-induced dyskinesias," *Movement Disorders*, vol. 22, no. 10, pp. 1379–1389, 2007.
- [14] A. Schrag and N. Quinn, "Dyskinesias and motor fluctuations in Parkinson's disease: a community-based study," *Brain*, vol. 123, no. 11, pp. 2297–2305, 2000.
- [15] A. Schrag and J. M. Schott, "Epidemiological, clinical, and genetic characteristics of early-onset parkinsonism," *The Lancet Neurology*, vol. 5, no. 4, pp. 355–363, 2006.
- [16] N. Kumar, J. A. Van Gerpen, J. H. Bower, and J. E. Ahlskog, "Levodopa-dyskinesia incidence by age of Parkinson's disease onset," *Movement Disorders*, vol. 20, no. 3, pp. 342–344, 2005.
- [17] K. E. Lyons, J. P. Hubble, A. I. Tröster, R. Pahwa, and W. C. Koller, "Gender differences in Parkinson's disease," *Clinical Neuropharmacology*, vol. 21, no. 2, pp. 118–121, 1998.
- [18] M. Zappia, G. Annesi, G. Nicoletti et al., "Sex differences in clinical and genetic determinants of levodopa peak-dose dyskinesias in Parkinson disease: an exploratory study," *Archives of Neurology*, vol. 62, no. 4, pp. 601–605, 2005.
- [19] Y. Gilgun-Sherki, R. Djaldetti, E. Melamed, and D. Offen, "Polymorphism in candidate genes: implications for the risk and treatment of idiopathic Parkinson's disease," *Pharmacogenomics Journal*, vol. 4, no. 5, pp. 291–306, 2004.
- [20] R. Kaiser, A. Hofer, A. Grapengiesser et al., "L-Dopa-induced adverse effects in PD and dopamine transporter gene polymorphism," *Neurology*, vol. 60, no. 11, pp. 1750–1755, 2003.
- [21] R. L. Oliveri, G. Annesi, M. Zappia et al., "Dopamine D₂ receptor gene polymorphism and the risk of levodopa-induced dyskinesias in PD," *Neurology*, vol. 53, no. 7, pp. 1425–1430, 1999.
- [22] W. C. Koller, W. Tse, Olanow et al., "Unmet medical needs in Parkinson's disease," *Neurology*, vol. 62, supplement 1, pp. S1–S8, 2004.
- [23] S. Fahn, "The spectrum of levodopa-induced dyskinesias," *Annals of Neurology*, vol. 47, supplement 1, pp. S2–S11, 2000.
- [24] L. I. Globe, "Young-onset Parkinson's disease: a clinical review," *Neurology*, vol. 41, pp. 168–173, 1991.
- [25] S. Boyce, C. E. Clarke, R. Luquin et al., "Induction of chorea and dystonia in parkinsonian primates," *Movement Disorders*, vol. 5, no. 1, pp. 3–7, 1990.
- [26] F. Durif, "Treating and preventing levodopa-induced dyskinesias: current and future strategies," *Drugs and Aging*, vol. 14, no. 5, pp. 337–345, 1999.
- [27] T. Borau, E. Bezard, B. Bioulac, and C. E. Gross, "Dopamine agonist-induced dyskinesias are correlated to both firing pattern and frequency alterations of pallidal neurones in the MPTP-treated monkey," *Brain*, vol. 124, pp. 546–557, 2001.

- [28] J. Jankovic, A. H. Rajput, M. P. McDermott, and D. P. Perl, "The evolution of diagnosis in early Parkinson disease," *Archives of Neurology*, vol. 57, no. 3, pp. 369–372, 2000.
- [29] A. Schrag, Y. Ben-Shlomo, R. Brown, C. D. Marsden, and N. Quinn, "Young-onset Parkinson's disease revisited—clinical features, natural history, and mortality," *Movement Disorders*, vol. 13, no. 6, pp. 885–894, 1998.
- [30] V. Sossi, R. De La Fuente-Fernández, M. Schulzer, J. Adams, and J. Stoessl, "Age-related differences in levodopa dynamics in Parkinson's: implications for motor complications," *Brain*, vol. 129, pp. 1050–1058, 2006.
- [31] D. A. Di Monte, A. McCormack, G. Petzinger, A. M. Janson, M. Quik, and W. J. Langston, "Relationship among nigrostriatal denervation, parkinsonism, and dyskinesias in the MPTP primate model," *Movement Disorders*, vol. 15, no. 3, pp. 459–466, 2000.
- [32] G. Linazasoro, "New ideas on the origin of L-dopa-induced dyskinesias: age, genes and neural plasticity," *Trends in Pharmacological Sciences*, vol. 26, no. 8, pp. 391–397, 2005.
- [33] G. Linazasoro, N. Van Blercom, A. Bergaretxe, I. Fernández Manchola, E. Laborda, and J. A. Ruiz Ortega, "Levodopa-induced dyskinesias in parkinson disease are independent of the extent of striatal dopaminergic denervation: a pharmacological and SPECT study," *Clinical Neuropharmacology*, vol. 32, no. 6, pp. 326–329, 2009.
- [34] C. R. Gerfen, T. M. Engber, L. C. Mahan et al., "D1 and D2 dopamine receptor-regulated gene expression of striatonigral and striatopallidal neurons," *Science*, vol. 250, no. 4986, pp. 1429–1432, 1990.
- [35] J. E. Nash and J. M. Brotchie, "A common signaling pathway for striatal NMDA and adenosine A(2a) receptors: implications for the treatment of Parkinson's disease," *Journal of Neuroscience*, vol. 20, no. 20, pp. 7782–7789, 2000.
- [36] E. Santini, E. Valjent, and G. Fisone, "Parkinson's disease: levodopa-induced dyskinesia and signal transduction," *FEBS Journal*, vol. 275, no. 7, pp. 1392–1399, 2008.
- [37] S. Boyce, N. M. J. Rupniak, M. J. Steventon, and S. D. Iversen, "Nigrostriatal damage is required for induction of dyskinesias by L-DOPA in squirrel monkeys," *Clinical Neuropharmacology*, vol. 13, no. 5, pp. 448–458, 1990.
- [38] A. Kumar, S. Mann, V. Sossi et al., "[11C]DTBZ-PET correlates of levodopa responses in asymmetric Parkinson's disease," *Brain*, vol. 126, no. 12, pp. 2648–2655, 2003.
- [39] R. de la Fuente-Fernández, "Drug-induced motor complications in dopa-responsive dystonia: implications for the pathogenesis of dyskinesias and motor fluctuations," *Clinical Neuropharmacology*, vol. 22, no. 4, pp. 216–219, 1999.
- [40] A. E. Fleckenstein, T. J. Volz, and G. R. Hanson, "Psychostimulant-induced alterations in vesicular monoamine transporter-2 function: neurotoxic and therapeutic implications," *Neuropharmacology*, vol. 56, supplement 1, pp. 133–138, 2009.
- [41] J. K. Dreyer, K. F. Herrik, R. W. Berg, and J. D. Hounsgaard, "Influence of phasic and tonic dopamine release on receptor activation," *Journal of Neuroscience*, vol. 30, no. 42, pp. 14273–14283, 2010.
- [42] J. A. Obeso, C. W. Olanow, and J. G. Nutt, "Levodopa motor complications in Parkinson's disease," *Trends in Neurosciences*, vol. 23, no. 10, pp. S2–S7, 2000.
- [43] J. A. Obeso, M. C. Rodríguez-Oroz, M. Rodríguez, M. R. DeLong, and C. W. Olanow, "Pathophysiology of levodopa-induced dyskinesias in Parkinson's disease: problems with the current model," *Annals of Neurology*, vol. 47, supplement 1, pp. S22–S34, 2000.
- [44] T. Björklund, T. Carlsson, E. A. Cederfjäll, M. Carta, and D. Kirik, "Optimized adeno-associated viral vector-mediated striatal DOPA delivery restores sensorimotor function and prevents dyskinesias in a model of advanced Parkinson's disease," *Brain*, vol. 133, pp. 496–511, 2010.
- [45] M. M. Mouradian, J. L. Juncos, G. Fabbrini, and T. N. Chase, "Motor fluctuations in Parkinson's disease: pathogenetic and therapeutic studies," *Annals of Neurology*, vol. 22, no. 4, pp. 475–479, 1987.
- [46] F. Stocchi, "The therapeutic concept of continuous dopaminergic stimulation (CDS) in the treatment of Parkinson's disease," *Parkinsonism and Related Disorders*, vol. 15, supplement 3, pp. S68–S71, 2009.
- [47] E. Bezard, J. M. Brotchie, and C. E. Gross, "Pathophysiology of levodopa-induced dyskinesia: potential for new therapies," *Nature Reviews Neuroscience*, vol. 2, no. 8, pp. 577–588, 2001.
- [48] M. A. Cenci, C. S. Lee, and A. Björklund, "L-DOPA-induced dyskinesia in the rat is associated with striatal overexpression of prodynorphin- and glutamic acid decarboxylase mRNA," *European Journal of Neuroscience*, vol. 10, no. 8, pp. 2694–2706, 1998.
- [49] C. Fiorentini, M. C. Rizzetti, C. Busi et al., "Loss of synaptic D1 dopamine/N-methyl-D-aspartate glutamate receptor complexes in L-DOPA-induced dyskinesia in the rat," *Molecular Pharmacology*, vol. 69, no. 3, pp. 805–812, 2006.
- [50] C. Missale, S. R. Nash, S. W. Robinson, M. Jaber, and M. G. Caron, "Dopamine receptors: from structure to function," *Physiological Reviews*, vol. 78, no. 1, pp. 189–225, 1998.
- [51] C. Le Moine and B. Bloch, "D₁ and D₂ dopamine receptor gene expression in the rat striatum: sensitive cRNA probes demonstrate prominent segregation of D1 and D2 mRNAs in distinct neuronal populations of the dorsal and ventral striatum," *Journal of Comparative Neurology*, vol. 355, no. 3, pp. 418–426, 1995.
- [52] M. Morelli and G. DiChiara, "Agonist-induced homologous and heterologous sensitization to D-1- and D-2-dependent contraversive turning," *European Journal of Pharmacology*, vol. 141, no. 1, pp. 101–107, 1987.
- [53] D. Xiao, E. Bastia, Y. H. Xu et al., "Forebrain adenosine A_{2A} receptors contribute to L-3,4-dihydroxyphenylalanine-induced dyskinesia in hemiparkinsonian mice," *Journal of Neuroscience*, vol. 26, no. 52, pp. 13548–13555, 2006.
- [54] L. A. Smith, M. J. Jackson, L. Johnston et al., "Switching from levodopa to the long-acting dopamine D₂/D₃ agonist piribedil reduces the expression of dyskinesia while maintaining effective motor activity in MPTP-treated primates," *Clinical Neuropharmacology*, vol. 29, no. 3, pp. 112–125, 2006.
- [55] M. J. Jackson, L. A. Smith, G. Al-Barghouthy, S. Rose, and P. Jenner, "Decreased expression of l-dopa-induced dyskinesia by switching to ropinirole in MPTP-treated common marmosets," *Experimental Neurology*, vol. 204, no. 1, pp. 162–170, 2007.
- [56] G. C. Cotzias, P. S. Papavasiliou, and E. S. Tolosa, "Treatment of Parkinson's disease with aporphines. Possible role of growth hormone," *The New England Journal of Medicine*, vol. 294, no. 11, pp. 567–572, 1976.
- [57] M. A. Brodsky, B. S. Park, and J. G. Nutt, "Effects of a dopamine agonist on the pharmacodynamics of levodopa in parkinson disease," *Archives of Neurology*, vol. 67, no. 1, pp. 27–32, 2010.
- [58] K. A. Stockwell, D. K. A. Scheller, L. A. Smith et al., "Continuous rotigotine administration reduces dyskinesia resulting from pulsatile treatment with rotigotine or l-DOPA in

- MPTP-treated common marmosets," *Experimental Neurology*, vol. 221, no. 1, pp. 79–85, 2010.
- [59] G. Linazasoro, "Pathophysiology of motor complications in Parkinson disease: postsynaptic mechanisms are crucial," *Archives of Neurology*, vol. 64, no. 1, pp. 137–140, 2007.
 - [60] C. W. Olanow and J. A. Obeso, "Preventing levodopa-induced dyskinesias," *Annals of Neurology*, vol. 47, no. 4, pp. S167–S178, 2000.
 - [61] L. V. Metman, S. Konitsiotis, and T. N. Chase, "Pathophysiology of motor response complications in Parkinson's disease: hypotheses on the why, where, and what," *Movement Disorders*, vol. 15, no. 1, pp. 3–8, 2000.
 - [62] R. Constantinescu, M. Romer, M. P. McDermott, C. Kamp, and K. Kieburz, "Impact of pramipexole on the onset of levodopa-related dyskinesias," *Movement Disorders*, vol. 22, no. 9, pp. 1317–1319, 2007.
 - [63] R. K. B. Pearce, T. Banerji, P. Jenner, and C. D. Marsden, "De novo administration of ropinirole and bromocriptine induces less dyskinesia than levodopa in the MPTP-treated marmoset," *Movement Disorders*, vol. 13, no. 2, pp. 234–241, 1998.
 - [64] E. C. Maratos, M. J. Jackson, R. K. B. Pearce, and P. Jenner, "Antiparkinsonian activity and dyskinesia risk of ropinirole and levodopa combination therapy in drug naïve MPTP-lesioned common marmosets (*Callithrix jacchus*)," *Movement Disorders*, vol. 16, no. 4, pp. 631–641, 2001.
 - [65] P. J. Blanchet, F. Calon, M. Morissette et al., "Regulation of dopamine receptors and motor behavior following pulsatile and continuous dopaminergic replacement strategies in the MPTP primate model," *Advances in Neurology*, vol. 86, pp. 337–344, 2001.
 - [66] M. A. Hely, J. G. L. Morris, W. G. J. Reid et al., "The Sydney multicentre study of Parkinson's disease: a randomised, prospective five year study comparing low dose bromocriptine with low dose levodopa-carbidopa," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 57, no. 8, pp. 903–910, 1994.
 - [67] A. J. Lees, R. Katzenschlager, J. Head, and Y. Ben-Shlomo, "Ten-year follow-up of three different initial treatments in de-novo PD: a randomized trial," *Neurology*, vol. 57, no. 9, pp. 1687–1694, 2001.
 - [68] W. H. Oertel, E. Wolters, C. Sampaio et al., "Pergolide versus levodopa monotherapy in early Parkinson's disease patients: The PELMOPET study," *Movement Disorders*, vol. 21, no. 3, pp. 343–353, 2006.
 - [69] J. A. Obeso, M. C. Rodriguez-Oroz, and I. Zamarbide, "Clinical features, pathophysiology, and management of motor complications in Parkinson's disease," in *Principles of Treatment in Parkinson's Disease*, A. H. V. Schapira and C. W. Olanow, Eds., pp. 99–112, Butterworth Heinemann Elsevier, Philadelphia, Pa, USA, 2005.
 - [70] M. J. Nirenberg and S. Fahn, "The role of levodopa and catechol-o-methyltransferase inhibitors," in *Principles of Treatment in Parkinson's Disease*, A. H. V. Schapira and C. W. Olanow, Eds., pp. 3–16, Butterworth Heinemann Elsevier, Philadelphia, Pa, USA, 2005.
 - [71] C. W. Olanow, S. Fahn, M. Muentner et al., "A multicenter double-blind placebo-controlled trial of pergolide as an adjunct to sinemet in Parkinson's disease," *Movement Disorders*, vol. 9, no. 1, pp. 40–47, 1994.
 - [72] Parkinson Study Group, "A randomized controlled trial comparing pramipexole with levodopa in early Parkinson's disease: design and methods of the CALM-PD study," *Clinical Neuropharmacology*, vol. 23, no. 1, pp. 34–44, 2000.
 - [73] Parkinson Study Group, "Pramipexole versus levodopa as initial treatment for Parkinson disease: a randomized controlled trial," *The Journal of the American Medical Association*, vol. 284, no. 15, pp. 1931–1938, 2000.
 - [74] Parkinson Study Group CALM Cohort Investigators, "Long-term effect of initiating pramipexole versus levodopa in early Parkinson disease," *Archives of Neurology*, vol. 66, pp. 563–570, 2009.
 - [75] R. A. Hauser, O. Rascol, A. D. Korczyn et al., "Ten-year follow-up of Parkinson's disease patients randomized to initial therapy with ropinirole or levodopa," *Movement Disorders*, vol. 22, no. 16, pp. 2409–2417, 2007.
 - [76] R. L. Watts, K. E. Lyons, R. Pahwa et al., "Onset of dyskinesia with adjunct ropinirole prolonged-release or additional levodopa in early Parkinson's disease," *Movement Disorders*, vol. 25, no. 7, pp. 858–866, 2010.
 - [77] A. Colzi, K. Turner, and A. J. Lees, "Continuous subcutaneous waking day apomorphine in the long term treatment of levodopa induced interdose dyskinesias in Parkinson's disease," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 64, no. 5, pp. 573–576, 1998.
 - [78] A. J. Manson, K. Turner, and A. J. Lees, "Apomorphine monotherapy in the treatment of refractory motor complications of Parkinson's disease: long-term follow-up study of 64 patients," *Movement Disorders*, vol. 17, no. 6, pp. 1235–1241, 2002.
 - [79] W. H. Poewe, O. Rascol, N. Quinn et al., "Efficacy of pramipexole and transdermal rotigotine in advanced Parkinson's disease: a double-blind, double-dummy, randomised controlled trial," *The Lancet Neurology*, vol. 6, no. 6, pp. 513–520, 2007.
 - [80] R. Pahwa, M. A. Stacy, L. W. Elmer, and S. H. Isaacson, "Ropinirole 24-hour prolonged release provides efficacy as early as week 2 when used as adjunctive therapy to levodopa in patients with advanced Parkinson's disease," *Movement Disorders*, vol. 21, supplement 15, p. S595, 2006.
 - [81] R. Pahwa, M. A. Stacy, S. A. Factor et al., "Ropinirole 24-hour prolonged release: randomized, controlled study in advanced Parkinson disease," *Neurology*, vol. 68, no. 14, pp. 1108–1115, 2007.
 - [82] K. D. Sethi, R. A. Hauser, and N. L. Earl, "Ropinirole 24-hour prolonged release improves disease-specific and global symptoms when used as adjunctive therapy to levodopa in patients with advanced Parkinson's disease," *Movement Disorders*, vol. 21, supplement 15, p. S570, 2006.
 - [83] K. D. Sethi, F. Stocchi, and L. Giorgi, "Ropinirole 24-hour prolonged release in advanced Parkinson's disease: relationship between treatment response and disease severity," *Movement Disorders*, vol. 22, supplement 16, p. S92, 2007.
 - [84] M. A. Stacy, R. Pahwa, and N. L. Earl, "Ropinirole 24-hour prolonged release reduces 'off' time and the dose of levodopa needed when used as adjunctive therapy in patients with advanced Parkinson's disease," *Movement Disorders*, vol. 21, supplement 15, p. S596, 2006.
 - [85] F. Stocchi, B. P. Hersh, N. L. Earl, and B. L. Scott, "Safety and tolerability of ropinirole 24-hour prolonged release in patients with early and advanced Parkinson's disease," *Movement Disorders*, vol. 21, supplement 15, p. S572, 2006.
 - [86] F. Stocchi, N. P. Stover, and L. Giorgi, "Ropinirole 24-hour prolonged release as adjunct to levodopa in patients with advanced Parkinson's disease—efficacy according to baseline depression score," *Movement Disorders*, vol. 22, supplement 16, p. S90, 2007.

- [87] J. C. Möller, W. H. Oertel, J. Köster, G. Pezzoli, and L. Provinciali, "Long-term efficacy and safety of pramipexole in advanced Parkinson's disease: results from a European multicenter trial," *Movement Disorders*, vol. 20, no. 5, pp. 602–610, 2005.
- [88] G. König, O. Pogarell, J. C. Möller, M. Delf, and W. H. Oertel, "Pramipexole, a nonergot dopamine agonist, is effective against rest tremor in intermediate to advanced Parkinson's disease," *Clinical Neuropharmacology*, vol. 22, no. 5, pp. 301–305, 1999.
- [89] K. S. Wong, C. S. Lu, D. E. Shan, C. C. Yang, T. H. Tsoi, and V. Mok, "Efficacy, safety, and tolerability of pramipexole in untreated and levodopa-treated patients with Parkinson's disease," *Journal of the Neurological Sciences*, vol. 216, no. 1, pp. 81–87, 2003.
- [90] B. Musch and L. Bonura, "Cabergoline once a day as adjunctive therapy to levodopa in Parkinson's disease," *Movement Disorders*, vol. 15, supplement 3, 2000.
- [91] M. M. Pinter, O. Pogarell, and W. H. Oertel, "Efficacy, safety, and tolerance of the non-ergoline dopamine agonist pramipexole in the treatment of advanced Parkinson's disease: a double blind, placebo controlled, randomised, multicentre study," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 66, no. 4, pp. 436–441, 1999.
- [92] L. Wermuth, "A double-blind, placebo-controlled, randomized, multi-center study of pramipexole in advanced Parkinson's disease," *European Journal of Neurology*, vol. 5, no. 3, pp. 235–242, 1998.
- [93] A. Lieberman, A. Ranhosky, and D. Korts, "Clinical evaluation of pramipexole in advanced Parkinson's disease: results of a double-blind, placebo-controlled, parallel-group study," *Neurology*, vol. 49, no. 1, pp. 162–168, 1997.
- [94] W. J. Weiner, S. A. Factor, J. Jankovic et al., "The long-term safety and efficacy of pramipexole in advanced Parkinson's disease," *Parkinsonism and Related Disorders*, vol. 7, no. 2, pp. 115–120, 2001.
- [95] M. Guttman, "Double-blind comparison of pramipexole and bromocriptine treatment with placebo in advanced Parkinson's disease," *Neurology*, vol. 49, no. 4, pp. 1060–1065, 1997.
- [96] M. Kreider, S. Knox, D. Gardiner, and D. Wheadon, "A multicenter double-blind study of ropinirole as an adjunct to levodopa in Parkinson's disease," *Neurology*, vol. 46: A475, 1996.
- [97] A. Lieberman, C. W. Olanow, K. Sethi et al., "A multicenter trial of ropinirole as adjunct treatment for Parkinson's disease," *Neurology*, vol. 51, no. 4, pp. 1057–1062, 1998.
- [98] O. Rascol, A. J. Lees, J. M. Senard et al., "A placebo-controlled study of ropinirole, a new D2 agonist, in the treatment of motor fluctuations of L-DOPA-treated parkinsonian patients," *Advances in Neurology*, vol. 69, pp. 531–534, 1996.
- [99] O. Rascol, A. J. Lees, J. M. Senard, Z. Pirtosek, J. L. Montastruc, and D. Fuell, "Ropinirole in the treatment of levodopa-induced motor fluctuations in patients with Parkinson's disease," *Clinical Neuropharmacology*, vol. 19, no. 3, pp. 234–245, 1996.
- [100] M. J. Steiger, T. El-Debas, T. Anderson, L. J. Findley, and C. D. Marsden, "Double-blind study of the activity and tolerability of cabergoline versus placebo in parkinsonians with motor fluctuations," *Journal of Neurology*, vol. 243, no. 1, pp. 68–72, 1996.
- [101] J. T. Hutton, W. C. Koller, J. E. Ahlskog et al., "Multicenter, placebo-controlled trial of cabergoline taken once daily in the treatment of Parkinson's disease," *Neurology*, vol. 46, no. 4, pp. 1062–1065, 1996.
- [102] A. N. Lieberman and J. T. Hutton, "Levodopa sparing effect of cabergoline compared to placebo in patients affected by motor fluctuations, under levodopa therapy," *Movement Disorders*, vol. 11, supplement 1, p. 269, 1996.
- [103] J. Schoenfelder, J. Simons, D. E. Souder, and J. Bianchine, "A placebo-controlled study of the safety and efficacy of cabergoline in the treatment of Parkinson's disease," *Functional Capacity Evaluation Reports 21336/731i*, 1993, Clinical Reference 42.
- [104] C. W. Olanow, S. Fahn, M. Muenster et al., "A multicenter double-blind placebo-controlled trial of pergolide as an adjunct to sinemet® in Parkinson's disease," *Movement Disorders*, vol. 9, no. 1, pp. 40–47, 1994.
- [105] J. A. Temlett, A. Ming, M. Saling et al., "Adjunctive therapy with bromocriptine in Parkinson's disease," *South African Medical Journal*, vol. 78, no. 11, pp. 680–685, 1990.
- [106] Y. Toyokura, Y. Mizuno, and M. Kase, "Effects of bromocriptine on Parkinsonism. A nation-wide collaborative double-blind study," *Acta Neurologica Scandinavica*, vol. 72, no. 2, pp. 157–170, 1985.
- [107] E. Schneider and P. A. Fischer, "Bromocriptine in the treatment of progressive stages of Parkinson's disease," *Deutsche Medizinische Wochenschrift*, vol. 107, no. 5, pp. 175–179, 1982.
- [108] E. N. H. Jansen, "Bromocriptine in levodopa response-losing parkinsonism. A double blind study," *European Neurology*, vol. 17, no. 2, pp. 92–99, 1978.
- [109] F. Stocchi, S. Ruggieri, L. Vacca, and C. W. Olanow, "Prospective randomized trial of lisuride infusion versus oral levodopa in patients with Parkinson's disease," *Brain*, vol. 125, no. 9, pp. 2058–2066, 2002.
- [110] U. K. Rinne, F. Bracco, C. Chouza et al., "Early treatment of Parkinson's disease with cabergoline delays the onset of motor complications. Results of a double-blind levodopa controlled trial," *Drugs*, vol. 55, supplement 1, pp. 23–30, 1998.
- [111] C. E. Clarke and K. D. Deane, "Cabergoline versus bromocriptine for levodopa-induced complications in Parkinson's disease," *Cochrane Database of Systematic Reviews*, no. 1, Article ID CD001519, 2001.
- [112] R. Stowe, N. Ives, C. E. Clarke et al., "Evaluation of the efficacy and safety of adjuvant treatment to levodopa therapy in Parkinson's disease patients with motor complications," *Cochrane Database of Systematic Reviews*, vol. 7, p. CD007166, 2010.
- [113] J. Jankovic, "Motor fluctuations and dyskinesias in Parkinson's disease: clinical manifestations," *Movement Disorders*, vol. 20, supplement 11, pp. S11–S16, 2005.
- [114] L. V. Metman, P. Del Dotto, P. van den Munchkof, J. Fang, M. M. Mouradian, and T. N. Chase, "Amantadine as treatment for dyskinesias in Parkinson's disease," *Neurology*, vol. 50, no. 5, pp. 1323–1326, 1998.
- [115] E. Lugging, G. K. Wenning, S. Bosch et al., "Beneficial effects of amantadine on levodopa-induced dyskinesias in Parkinson's disease," *Movement Disorders*, vol. 15, pp. 873–878, 2000.
- [116] B. J. Snow, L. Macdonald, D. Mcauley, and W. Wallis, "The effect of amantadine on levodopa-induced dyskinesias in Parkinson's disease: a double-blind, placebo-controlled study," *Clinical Neuropharmacology*, vol. 23, no. 2, pp. 82–85, 2000.
- [117] N. J. Crosby, K. H. Deane, and C. E. Clarke, "Amantadine for dyskinesia in Parkinson's disease," *Cochrane Database of Systematic Reviews*, no. 2, p. CD003467, 2003.

- [118] F. Durif, B. Debilly, M. Galitzky et al., "Clozapine improves dyskinesias in Parkinson disease A double-blind, placebo-controlled study," *Neurology*, vol. 62, no. 3, pp. 381–388, 2004.
- [119] F. Pierelli, A. Adipietro, G. Soldati, F. Fattapposta, G. Pozzessere, and C. Scoppetta, "Low dosage clozapine effects on levodopa induced dyskinesias in parkinsonian patients," *Acta Neurologica Scandinavica*, vol. 97, no. 5, pp. 295–299, 1998.
- [120] V. C. Anderson, K. J. Burchiel, P. Hogarth, J. Favre, and J. P. Hammerstad, "Pallidal vs subthalamic nucleus deep brain stimulation in Parkinson disease," *Archives of Neurology*, vol. 62, no. 4, pp. 554–560, 2005.
- [121] B. Ford, L. Winfield, S. L. Pullman et al., "Subthalamic nucleus stimulation in advanced Parkinson's disease: blinded assessments at one year follow up," *Journal of Neurology, Neurosurgery and Psychiatry*, vol. 75, no. 9, pp. 1255–1259, 2004.
- [122] K. Østergaard, N. Sunde, and E. Dupont, "Effects of bilateral stimulation of the subthalamic nucleus in patients with severe Parkinson's disease and motor fluctuations," *Movement Disorders*, vol. 17, no. 4, pp. 693–700, 2002.
- [123] E. Moro, M. Scerrati, L. M. A. Romito, R. Roselli, P. Tonali, and A. Albanese, "Chronic subthalamic nucleus stimulation reduces medication requirements in Parkinson's disease," *Neurology*, vol. 53, no. 1, pp. 85–90, 1999.
- [124] F. J. G. Vingerhoets, J. G. Villemure, P. Temperli, C. Pollo, E. Pralong, and J. Ghika, "Subthalamic DBS replaces levodopa in Parkinson's disease: two-year follow-up," *Neurology*, vol. 58, no. 3, pp. 396–401, 2002.