Editorial

PPARs and Bone Metabolism

Beata Lecka-Czernik

Department of Geriatrics, Reynolds Institute on Aging, University of Arkansas for Medical Sciences, Little Rock, AR 72205, USA

Received 19 July 2006; Accepted 19 July 2006

Copyright © 2006 Beata Lecka-Czernik. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Welcome to the inaugural *special* issue of *PPAR Research: PPARs and Bone Metabolism*. In addition to the key roles PPARs play in numerous processes including glucose and fat metabolism, inflammation, cancer, and central nervous system maintenance, a new role for PPAR- γ has recently emerged: the maintenance of bone homeostasis during aging and disease. In this premier issue we have assembled what is close to a comprehensive overview of the role of PPAR- γ in the control of bone maintenance. This takes into account PPAR- γ 's role in mesenchymal stem cell lineage allocation, possible cross-talk with relevant nuclear receptors, examination of PPAR- γ gene polymorphisms and bone mineral density in humans, a role of PPAR- γ in bone loss due to skeletal disuse, evidence that human bone is vulnerable to antidiabetic therapies with PPAR- γ agonists, the thiazolidinediones, and evidence that the antiosteoblastic activity of PPAR- γ can be separated from its proadipocytic and antidiabetic activities by using selective modulators. We also present a novel hypothesis that PPAR- γ acts as a regulator of chondrocyte development and cartilage homeostasis. We realize that we have not covered all aspects of PPARs involvement in the control of bone maintenance; however, this introduction should serve as a competent first attempt to present these new aspects of bone biology to the broader audience of our readers.

Beata Lecka-Czernik