

Review Article

The Role of PPAR γ in the Cyclooxygenase Pathway in Lung Cancer

Saswati Hazra, Katherine A. Peebles, Sherven Sharma, Jenny T. Mao, and Steven M. Dubinett

UCLA Lung Cancer Research Program, Division of Pulmonary and Critical Care Medicine and Hospitalists, Department of Medicine, Jonsson Comprehensive Cancer Center, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095, USA

Correspondence should be addressed to Saswati Hazra, shazra@mednet.ucla.edu

Received 16 April 2008; Revised 18 June 2008; Accepted 8 July 2008

Recommended by Dipak Panigrahy

Decreased expression of peroxisome proliferator activated receptor- γ (PPAR γ) and high levels of the proinflammatory enzyme cyclooxygenase-2 (COX-2) have been observed in many tumor types. Both reduced (PPAR γ) expression and elevated COX-2 within the tumor are associated with poor prognosis in lung cancer patients, and recent work has indicated that these signaling pathways may be interrelated. Synthetic (PPAR γ) agonists such as the thiazolidinedione (TZD) class of anti-diabetic drugs can decrease COX-2 levels, inhibit growth of non-small-cell lung cancer (NSCLC) cells in vitro, and block tumor progression in xenograft models. TZDs alter the expression of COX-2 and consequent production of the protumorigenic inflammatory molecule prostaglandin E2 (PGE₂) through both (PPAR γ) dependent and independent mechanisms. Certain TZDs also reduce expression of PGE₂ receptors or upregulate the PGE₂ catabolic enzyme 15-prostaglandin dehydrogenase. As several COX-2 enzymatic products have antitumor properties and specific COX-2 inhibition has been associated with increased risk of adverse cardiac events, directly reducing the effects or concentration of PGE₂ may provide a more safe and effective strategy for lung cancer treatment. Understanding the mechanisms underlying these effects may be helpful for designing anticancer therapies. This article summarizes recent research on the relationship between (PPAR γ), TZDs, and the COX-2/PGE₂ pathways in lung cancer.

Copyright © 2008 Saswati Hazra et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Despite the many advances made in diagnostic and treatment strategies, lung cancer remains the leading cause of cancer-related mortality in the United States and is responsible for more deaths than prostate, colon, and breast cancers combined [1]. Investigating the molecular mechanisms underlying the pathogenesis of lung cancer provides opportunities to develop innovative therapies that may reduce suffering due to this devastating disease. Decreased expression of peroxisome proliferator activated receptor (PPAR γ) originally identified as a regulator of glucose metabolism and adipocyte differentiation [2] has been associated with poor prognosis in lung cancer patients [3]. PPAR γ affects inflammatory gene expression, cell division, apoptosis, invasion, release of proangiogenic cytokines, and differentiation in many cancer types including lung cancer [4–8]. These properties have prompted extensive research on PPAR γ in cancer treatment and prevention. Members of the thiazolidinedione (TZD) class of PPAR γ agonists are currently approved for treatment of diabetes, and elicit many of the antitumor properties of

PPAR γ activation through both PPAR γ dependent and independent pathways [9–14]. Several studies have demonstrated elevated constitutive expression of the inducible proinflammatory enzyme, cyclooxygenase-2 (COX-2) in human lung cancer [15–19]. Mounting evidence from investigations into the molecular effects of COX-2 over-expression in lung tumor cells indicates that this enzyme has a multifaceted role in conferring the malignant and metastatic phenotypes. The COX-2 enzymatic product prostaglandin E₂ (PGE₂) has been implicated in apoptosis resistance [20–22], angiogenesis [23, 24], decreased host immunity [25, 26], and enhanced invasion and metastasis [27–29]. This review summarizes investigations in the relationship between PPAR γ , its ligands, and COX-2 and PGE₂ in lung cancer.

The PPAR family consists of three isoforms: PPAR α , PPAR γ , and PPAR δ , each encoded by different genes. PPARs are members of the nuclear hormone class of receptors and are involved in energy metabolism through transcriptional regulation of specific gene sets. Observations

regarding high PPAR γ expression in adipose tissue in combination with its role in lipid and glucose homeostasis led to the development of the TZD class of PPAR γ agonists, including troglitazone, ciglitazone, rosiglitazone, and pioglitazone as antidiabetic and insulin-sensitizing drugs. Rosiglitazone and pioglitazone are currently approved for treatment of type 2 diabetes mellitus [30], and this class of drugs has been clinically available for approximately a decade. Some of the TZDs have been shown to exert anti-inflammatory [31], antiproliferative [32], and antiangiogenic effects [4]. The COX metabolite 15d-PGJ₂ is a natural PPAR γ ligand and is considered a negative regulator of inflammatory and immune responses [33]. More recent results indicating that PPAR γ activation may attenuate inflammatory responses and cancer progression have led to extensive investigation into the role of this protein in inflammation and carcinogenesis.

PPAR γ is expressed in human non-small-cell lung cancer (NSCLC) and small cell lung carcinoma [34], and the expression of PPAR γ has been correlated with tumor histological type and grade [35]. In NSCLC, decreased PPAR γ expression was correlated with poor prognosis [3]. TZDs inhibit tumor formation in a variety of animal models, including colon [36] and lung cancers [37], and PPAR γ over-expression protects against tumor development in a mouse model of lung tumorigenesis [38]. Further, increased PPAR γ activity promotes epithelial differentiation of NSCLC cells in 3D culture [5]. It has also been shown that PPAR γ inhibits the growth of NSCLC in vitro and in vivo [5, 39, 40].

Cyclooxygenase is the rate-limiting enzyme for production of prostaglandins and thromboxanes from free arachidonic acid [41, 42]. Two COX isoforms, COX-1 and COX-2, have been extensively studied. COX-1 is constitutively expressed in most cells and tissues. COX-2 is an inducible enzyme that acts to produce prostaglandins and/or thromboxanes during an acute inflammatory response. The direct enzymatic product of COX-2 and PGH₂ is converted to prostaglandins or thromboxanes by individual isomerases or prostaglandin synthases, and relative production of the various COX-2 products depends upon cellular concentrations of down-stream metabolic and catabolic enzymes within the COX-2 pathway. In NSCLC, the major eicosanoid produced is prostaglandin E₂ (PGE₂) through microsomal PGE₂ synthase (mPGES) activity. The nicotinamide adenine dinucleotide positive-dependent catabolic enzyme 15-hydroxyprostaglandin dehydrogenase (15-PGDH) metabolizes PGE₂ to biologically inactive 15-keto derivatives. The final PGE₂ concentration experienced by NSCLC cells depends upon expression of PGES and 15-PGDH. A large body of evidence indicates that increased PGE₂ production contributes to tumorigenesis. COX-2 over-expression is frequently observed in NSCLC, and the accompanying increased proliferation, invasion, angiogenesis, and resistance to apoptosis have been attributed in part to elevated PGE₂ production in the vicinity of the tumor. Thus, COX-2 and its downstream signaling pathways represent potential targets for lung cancer chemoprevention and therapy.

Studies indicate that COX-2 and PPAR γ signaling pathways are intertwined. PPAR γ ligands suppress COX-2 expres-


sion induced by LPS and PMA in macrophages, astrocytes, and epithelial cells [43–45]. The COX-2 metabolite 15d-PGJ₂ is an endogenous ligand for PPAR γ [46], and during resolution of inflammation elevated 15d-PGJ₂ production downregulates COX-2 through a negative feedback loop involving PPAR γ and NF- κ B [44, 47]. Synthetic and endogenous PPAR γ ligands decrease the high COX-2 expression associated with several malignancies including cervical [48] and liver cancers [49] and forced PPAR over-expression decreases COX-2 levels in lung cancer cells [38]. While PPAR γ agonists decrease COX-2 expression or prevent COX-2 induction in most settings, COX-2 expression is increased in some studies [50, 51]. For example, Ikawa et al. reported that rosiglitazone (also known as BRL49653) increases COX-2 expression in human colorectal carcinoma cells [52]. PPAR γ ligands also have been shown to induce COX-2 expression in mammary epithelial cells [53], monocytes [54], and human synovial fibroblasts [55]. The effect of PPAR γ agonists on COX-2 expression may vary based upon the cell type, the specific agonist molecule, and the presence of additional inflammatory mediators. Off-target effects of TZDs may also partially account for differences in the effects of these molecules on COX-2 expression.

Although TZDs are widely known as ligands for PPAR γ , they may mediate receptor-independent effects, as previously reported [56–58]. For example, by using embryonic stem cells from PPAR γ -null mice, Chawla et al. found that neither macrophage differentiation nor anti-inflammatory effects of synthetic PPAR γ ligands are PPAR γ receptor-dependent. To distinguish the effects of PPAR γ from off-target effects of PPAR γ ligands in lung cancer cells, Bren-Mattison et al. utilized a molecular approach to over-express PPAR γ in two NSCLC cell lines and assessed the direct effect of PPAR γ . Their goal was to determine whether the antitumorogenic effects of PPAR γ were mediated via COX-2 pathways in NSCLC. Their results clearly demonstrated that exogenously expressed PPAR γ suppresses COX-2 promoter activity and protein expression resulting in suppression of PGE₂ production [38]. The COX-2 promoter has binding sites for cAMP response element, NF-IL-6, and NF- κ B [59]. Although the COX-2 promoter contains multiple regulatory elements, their data indicate that the inhibitory effects of PPAR γ are mediated through NF- κ B. Several studies have demonstrated that constitutive activation of NF- κ B in tumor cells is mediated through activation of Akt [60, 61]. Bren-Mattison et al. demonstrated that the inhibitory effects of PPAR γ on COX-2 were mediated via increased activity of PTEN leading to decreased phospho-Akt and inhibition of NF- κ B [38]. These authors further demonstrated that transgenic mice over-expressing PPAR γ exhibited reduced COX-2 in type II alveolar epithelial cells of lung, and those mice were protected against lung cancer development in a chemical carcinogenesis mouse model [38]. In summary, these data indicate that COX-2 downregulation may mediate some of the antitumorogenic effects of PPAR γ over-expression.

The PPAR γ agonists may also affect COX-2 in a PPAR γ independent manner (see Table 1). For example, in A549 NSCLC cells troglitazone enhanced both COX-2 and mPGES expression in a concentration dependent manner, resulting

TABLE 1: Off-target effects of TZDs in NSCLC.

Thiazolidinediones	Molecular effects	Mechanisms	Reference
Troglitazone	↑ PGE ₂	↑ COX-2, ERK and PI3K phosphorylation	[62]
Pioglitazone, Rosiglitazone	↓ PGE ₂	↑ 15-PGDH	[14]
Ciglitazone	↓ PGE ₂	↓ COX-2	[13]

FIGURE 1: Effects of various TZDs on the PGE₂ pathway.

in a marked increase in PGE₂ production [62]. Cotreatment with the PPAR γ antagonists GW9662 and bisphenol A diglycidyl ether (BADGE) had no effect on COX-2 induction by troglitazone indicating that this event is PPAR γ independent. Troglitazone increased COX-2 expression at least in part by activating ERK and PI3K pathways, which have been previously demonstrated to influence COX-2 levels [63–65]. Combined troglitazone and TNF α stimulation resulted in additive enhancement of COX-2 expression. The COX-2 metabolite 15d-PGJ₂ had no detectable effects on COX-2 or mPGES expression or PGE₂ production in A549 cells. This is consistent with the hypothesis that PPAR γ -independent mechanisms can partially account for discrepancies in the effects of different TZD drugs on COX-2 expression. Thus, in lung cancer, PPAR γ agonists appear to regulate COX-2 expression and affiliated protumorigenic cellular phenotypes through both PPAR γ dependent and independent means.

We recently examined the effect of the pioglitazone and rosiglitazone on COX-2 and PGE₂ levels in A427 and A549 NSCLC cells. Both TZDs inhibited PGE₂ production in NSCLC cells via a COX-2 independent pathway. To define the mechanism underlying COX-2 independent suppression of PGE₂ production, we focused on the prostaglandin synthases that are responsible for the PGE₂ production and on 15PGDH the catabolic enzyme responsible for its degradation to biologically inactive 15-keto derivatives [66]. None of the three prostaglandin synthases (microsomal PGES1, PGES2, and cytosolic PGES) was downregulated by pioglitazone or rosiglitazone, however, 15-PGDH was induced by TZDs. TZD-mediated suppression of PGE₂

concentration was significantly inhibited by small interfering RNA to 15-PGDH. Studies using dominant-negative PPAR γ over-expression or GW9662 revealed that the induction of 15-PGDH by both pioglitazone and rosiglitazone is PPAR γ -independent. These findings indicate that it is possible to use a clinically available pharmacological intervention to suppress tumor-derived PGE₂ by enhancing catabolism rather than blocking synthesis. The potential benefits of inhibiting PGE₂ levels in a COX-2-independent manner include the following. First, promoting 15-PGDH activity could decrease PGE₂ without modifying other prostaglandins such as PGI₂. This is potentially important because the latter has been noted to have antitumor properties [67]. It has been suggested that a ratio of PGs may be important in regulating the malignant phenotype. Thus, inhibiting COX-2 activity would diminish both PGE₂ and PGI₂, whereas selective induction of 15-PGDH could lead to a more favorable PGI₂/PGE₂ ratio. Second, suppression of PGE₂ levels without alteration in COX-2 may limit some of the cardiovascular toxicities associated with COX-2 inhibition [68]. Finally, unlike COX-2 inhibition, which may lead to upregulation of certain leukotrienes that favor malignant progression [69], 15-PGDH induction may lead only to a decrement of PGE₂. This speculation will require further investigation.

Different TZDs have the capacity to modulate arachidonic acid metabolism by a variety of pathways (see Figure 1). Recent evidence indicates that ciglitazone induces differentiation and apoptosis in NSCLC [7]. The mechanisms of ciglitazone's capacity to modulate PGE₂ levels in lung adenocarcinoma cells were recently reported [13].

In contrast to pioglitazone and rosiglitazone, ciglitazone mediates COX-2 dependent suppression of PGE₂ in NSCLC. Ciglitazone treatment suppressed COX-2 mRNA expression and COX-2 promoter activity but did not modify the expression of enzymes downstream of COX-2 including PGES and 15-PGDH. Utilization of dominant-negative PPAR γ showed that suppression of COX-2 and PGE₂ by ciglitazone is mediated via non-PPAR pathways.

PPAR γ ligands may also interfere with protumorigenic signals derived from COX-2 by interrupting the function of PGE₂ G-protein coupled receptors (GPCRs) designated E-prostanoid (EP) receptors 1–4 [70]. Han and Roman found that in NSCLC cell lines, the PPAR γ ligands GW1929, 15 dPGJ₂, ciglitazone, troglitazone, and rosiglitazone significantly decreased EP2 mRNA and protein levels causing growth inhibition in NSCLC cells [71]. The inhibitory effects of rosiglitazone and ciglitazone but not 15d-PGJ₂ were suppressed by the PPAR γ antagonist GW9662 suggesting the involvement of PPAR γ -dependent and independent mechanisms.

Recently, a retrospective study by Govindarajan et al. demonstrated a significant reduction in lung cancer risk in diabetic patients using the TZD rosiglitazone [72]. Importantly, several clinical studies in diabetes patients have demonstrated an increased risk of cardiovascular events associated with rosiglitazone or pioglitazone treatment [73–75]. This is of particular significance in light of cardiovascular toxicity associated with COX-2 inhibition. Recently, several chemoprevention trials are being initiated using TZDs [76]. However, adverse cardiac events are associated with chronic TZD treatment [74]. Based on these findings, future clinical studies attempting to utilize TZDs in prevention of cancer will require selection of patient populations without cardiovascular risk. Prospective clinical studies specifically designed to address the effects of TZDs on cancer, and cardiac outcomes are required. If the anti-inflammatory and antitumor effects of TZDs are derived through pathways distinct from those leading to cardiovascular toxicity, more selective candidate drug molecules may be therapeutically effective, without leading to adverse cardiac events. Thus, more research is required to define opportunities to specifically interfere with PGE₂ production, metabolism, or downstream effects. This could ultimately lead to reduction in lung cancer growth or prevention while leaving the steady-state concentrations of desirable eicosanoids intact [77].

Both elevated COX-2 and reduced PPAR γ expression are associated with poor prognosis in lung cancer patients [3, 78–80] and recent work has revealed multiple interactions between PPAR γ signaling and the COX-2 pathway. The COX-2 product 15d-PGJ₂ is an endogenous ligand for PPAR γ , and PPAR γ activation as a result of elevated 15d-PGJ₂ results in COX-2 downregulation in an autoregulatory feedback loop that may contribute to natural resolution of the inflammatory response [46]. Forced expression of PPAR γ decreases COX-2 expression in cultured human NSCLC cells and mouse lungs and protects against lung tumor development in a murine model [5, 38]. Synthetic PPAR γ ligands, several of which are currently approved for treatment of diabetes, can interrupt several stages of the

COX-2/PGE₂ protumorigenic pathway, although in certain cases PPAR γ ligands may increase COX-2 expression. These effects are primarily mediated through PPAR γ -independent pathways (see Table 1). PPAR γ ligands may directly decrease COX-2 transcription in an NF- κ B-dependent manner [38], or they can interfere with downstream targets such as the PGE₂ receptor EP2 [71] or the enzyme responsible for PGE₂ catabolism, 15-PGDH [66]. The targets downstream of COX-2 may be useful in light of recent evidence that interfering with COX-2 enzymatic activity may increase risk of cardiovascular events [68]. The discovery that certain PPAR γ agonists can specifically reduce PGE₂ concentration or expression of EP receptors may aid in the design of strategies to reduce the effects of harmful prostaglandins without impacting production of critical cardioprotective eicosanoids.

REFERENCES

- [1] A. Jemal, R. Siegel, E. Ward, T. Murray, J. Xu, and M. J. Thun, "Cancer statistics, 2007," *CA: A Cancer Journal for Clinicians*, vol. 57, no. 1, pp. 43–66, 2007.
- [2] J. Berger and D. E. Moller, "The mechanisms of action of PPARs," *Annual Review of Medicine*, vol. 53, pp. 409–435, 2002.
- [3] H. Sasaki, M. Tanahashi, H. Yukiue, et al., "Decreased peroxisome proliferator-activated receptor gamma gene expression was correlated with poor prognosis in patients with lung cancer," *Lung Cancer*, vol. 36, no. 1, pp. 71–76, 2002.
- [4] V. G. Keshamouni, D. A. Arenberg, R. C. Reddy, M. J. Newstead, S. Anthwal, and T. J. Standiford, "PPAR- γ activation inhibits angiogenesis by blocking ELR+CXC chemokine production in non-small cell lung cancer," *Neoplasia*, vol. 7, no. 3, pp. 294–301, 2005.
- [5] Y. Bren-Mattison, V. Van Putten, D. Chan, R. Winn, M. W. Geraci, and R. A. Nemenoff, "Peroxisome proliferator-activated receptor- γ (PPAR γ) inhibits tumorigenesis by reversing the undifferentiated phenotype of metastatic non-small-cell lung cancer cells (NSCLC)," *Oncogene*, vol. 24, no. 8, pp. 1412–1422, 2005.
- [6] C. G. Su, X. Wen, S. T. Bailey, et al., "A novel therapy for colitis utilizing PPAR- γ ligands to inhibit the epithelial inflammatory response," *The Journal of Clinical Investigation*, vol. 104, no. 4, pp. 383–389, 1999.
- [7] T.-H. Chang and E. Szabo, "Induction of differentiation and apoptosis by ligands of peroxisome proliferator-activated receptor γ in non-small cell lung cancer," *Cancer Research*, vol. 60, no. 4, pp. 1129–1138, 2000.
- [8] S. Han and J. Roman, "Peroxisome proliferator-activated receptor γ : a novel target for cancer therapeutics?" *Anti-Cancer Drugs*, vol. 18, no. 3, pp. 237–244, 2007.
- [9] A. T. Coyle and B. T. Kinsella, "Synthetic peroxisome proliferator-activated receptor γ agonists rosiglitazone and troglitazone suppress transcription by promoter 3 of the human thromboxane A₂ receptor gene in human erythrocytes cells," *Biochemical Pharmacology*, vol. 71, no. 9, pp. 1308–1323, 2006.
- [10] M. J. Betz, I. Shapiro, M. Fassnacht, S. Hahner, M. Reincke, and F. Beuschlein, "Peroxisome proliferator-activated receptor- γ agonists suppress adrenocortical tumor cell proliferation and induce differentiation," *The Journal of Clinical*

- Endocrinology & Metabolism*, vol. 90, no. 7, pp. 3886–3896, 2005.
- [11] T. Satoh, M. Toyoda, H. Hoshino, et al., “Activation of peroxisome proliferator-activated receptor- γ stimulates the growth arrest and DNA-damage inducible 153 gene in non-small cell lung carcinoma cells,” *Oncogene*, vol. 21, no. 14, pp. 2171–2180, 2002.
- [12] S. J. Baek, L. C. Wilson, L. C. Hsi, and T. E. Eling, “Troglitazone, a peroxisome proliferator-activated receptor γ (PPAR γ) ligand, selectively induces the early growth response-1 gene independently of PPAR γ . A novel mechanism for its anti-tumorigenic activity,” *The Journal of Biological Chemistry*, vol. 278, no. 8, pp. 5845–5853, 2003.
- [13] S. Hazra and S. M. Dubinett, “Ciglitazone mediates COX-2 dependent suppression of PGE₂ in human non-small cell lung cancer cells,” *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 77, no. 1, pp. 51–58, 2007.
- [14] S. Hazra, R. K. Batra, H. H. Tai, S. Sharma, X. Cui, and S. M. Dubinett, “Pioglitazone and rosiglitazone decrease prostaglandin E₂ in non-small-cell lung cancer cells by up-regulating 15-hydroxyprostaglandin dehydrogenase,” *Molecular Pharmacology*, vol. 71, no. 6, pp. 1715–1720, 2007.
- [15] M. Huang, M. Stolina, S. Sharma, et al., “Non-small cell lung cancer cyclooxygenase-2-dependent regulation of cytokine balance in lymphocytes and macrophages: up-regulation of interleukin 10 and down-regulation of interleukin 12 production,” *Cancer Research*, vol. 58, no. 6, pp. 1208–1216, 1998.
- [16] T. Hida, Y. Yatabe, H. Achiwa, et al., “Increased expression of cyclooxygenase 2 occurs frequently in human lung cancers, specifically in adenocarcinomas,” *Cancer Research*, vol. 58, no. 17, pp. 3761–3764, 1998.
- [17] J. Brabender, J. Park, R. Metzger, et al., “Prognostic significance of cyclooxygenase 2 mRNA expression in non-small cell lung cancer,” *Annals of Surgery*, vol. 235, no. 3, pp. 440–443, 2002.
- [18] H. Achiwa, Y. Yatabe, T. Hida, et al., “Prognostic significance of elevated cyclooxygenase 2 expression in primary, resected lung adenocarcinomas,” *Clinical Cancer Research*, vol. 5, no. 5, pp. 1001–1005, 1999.
- [19] Y. Hosomi, T. Yokose, Y. Hirose, et al., “Increased cyclooxygenase 2 (COX-2) expression occurs frequently in precursor lesions of human adenocarcinoma of the lung,” *Lung Cancer*, vol. 30, no. 2, pp. 73–81, 2000.
- [20] M. Tsujii and R. N. DuBois, “Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase 2,” *Cell*, vol. 83, no. 3, pp. 493–501, 1995.
- [21] K. Krysan, F. H. Merchant, L. Zhu, et al., “COX-2-dependent stabilization of survivin in non-small cell lung cancer,” *The FASEB Journal*, vol. 18, no. 1, pp. 206–208, 2004.
- [22] K. Krysan, H. Dalwadi, S. Sharma, M. Pöld, and S. Dubinett, “Cyclooxygenase 2-dependent expression of survivin is critical for apoptosis resistance in non-small cell lung cancer,” *Cancer Research*, vol. 64, no. 18, pp. 6359–6362, 2004.
- [23] K. M. Leahy, A. T. Koki, and J. L. Masferrer, “Role of cyclooxygenases in angiogenesis,” *Current Medicinal Chemistry*, vol. 7, no. 11, pp. 1163–1170, 2000.
- [24] S. Gately and W. W. Li, “Multiple roles of COX-2 in tumor angiogenesis: a target for antiangiogenic therapy,” *Seminars in Oncology*, vol. 31, no. 2, supplement 7, pp. 2–11, 2004.
- [25] V. C. Liu, L. Y. Wong, T. Jang, et al., “Tumor evasion of the immune system by converting CD4⁺CD25⁻ T cells into CD4⁺CD25⁺ T regulatory cells: role of tumor-derived TGF- β ,” *The Journal of Immunology*, vol. 178, no. 5, pp. 2883–2892, 2007.
- [26] F. Baratelli, Y. Lin, L. Zhu, et al., “Prostaglandin E₂ induces FOXP3 gene expression and T regulatory cell function in human CD4⁺ T cells,” *The Journal of Immunology*, vol. 175, no. 3, pp. 1483–1490, 2005.
- [27] M. Dohadwala, J. Luo, L. Zhu, et al., “Non-small cell lung cancer cyclooxygenase-2-dependent invasion is mediated by CD44,” *The Journal of Biological Chemistry*, vol. 276, no. 24, pp. 20809–20812, 2001.
- [28] M. Dohadwala, R. K. Batra, J. Luo, et al., “Autocrine/paracrine prostaglandin E₂ production by non-small cell lung cancer cells regulates matrix metalloproteinase-2 and CD44 in cyclooxygenase-2-dependent invasion,” *The Journal of Biological Chemistry*, vol. 277, no. 52, pp. 50828–50833, 2002.
- [29] H. Sheng, J. Shao, M. K. Washington, and R. N. DuBois, “Prostaglandin E₂ increases growth and motility of colorectal carcinoma cells,” *The Journal of Biological Chemistry*, vol. 276, no. 21, pp. 18075–18081, 2001.
- [30] R. J. Durbin, “Thiazolidinedione therapy in the prevention/delay of type 2 diabetes in patients with impaired glucose tolerance and insulin resistance,” *Diabetes, Obesity & Metabolism*, vol. 6, no. 4, pp. 280–285, 2004.
- [31] A. Consoli and E. Devangelio, “Thiazolidinediones and inflammation,” *Lupus*, vol. 14, no. 9, pp. 794–797, 2005.
- [32] S. Schmidt, E. Moric, M. Schmidt, M. Sastre, D. L. Feinstein, and M. T. Heneka, “Anti-inflammatory and antiproliferative actions of PPAR- γ agonists on T lymphocytes derived from MS patients,” *Journal of Leukocyte Biology*, vol. 75, no. 3, pp. 478–485, 2004.
- [33] J. U. Scher and M. H. Pillinger, “15d-PGJ₂: the anti-inflammatory prostaglandin?” *Clinical Immunology*, vol. 114, no. 2, pp. 100–109, 2005.
- [34] K. Inoue, Y. Kawahito, Y. Tsubouchi, et al., “Expression of peroxisome proliferator-activated receptor (PPAR)- γ in human lung cancer,” *Anticancer Research*, vol. 21, no. 4A, pp. 2471–2476, 2001.
- [35] S. Theocharis, H. Kanelli, E. Politi, et al., “Expression of peroxisome proliferator activated receptor-gamma in non-small cell lung carcinoma: correlation with histological type and grade,” *Lung Cancer*, vol. 36, no. 3, pp. 249–255, 2002.
- [36] T. Yoshizumi, T. Ohta, I. Ninomiya, et al., “Thiazolidinedione, a peroxisome proliferator-activated receptor-gamma ligand, inhibits growth and metastasis of HT-29 human colon cancer cells through differentiation-promoting effects,” *International Journal of Oncology*, vol. 25, no. 3, pp. 631–639, 2004.
- [37] V. G. Keshamouni, R. C. Reddy, D. A. Arenberg, et al., “Peroxisome proliferator-activated receptor- γ activation inhibits tumor progression in non-small-cell lung cancer,” *Oncogene*, vol. 23, no. 1, pp. 100–108, 2004.
- [38] Y. Bren-Mattison, A. M. Meyer, V. Van Putten, et al., “Antitumorigenic effects of peroxisome proliferator-activated receptor- γ in non-small-cell lung cancer cells are mediated by suppression of cyclooxygenase-2 via inhibition of nuclear factor- κ B,” *Molecular Pharmacology*, vol. 73, no. 3, pp. 709–717, 2008.
- [39] M. Wick, G. Hurteau, C. Dessev, et al., “Peroxisome proliferator-activated receptor- γ is a target of nonsteroidal anti-inflammatory drugs mediating cyclooxygenase-independent inhibition of lung cancer cell growth,” *Molecular Pharmacology*, vol. 62, no. 5, pp. 1207–1214, 2002.

- [40] S. Han and J. Roman, "Rosiglitazone suppresses human lung carcinoma cell growth through PPAR γ -dependent and PPAR γ -independent signal pathways," *Molecular Cancer Therapeutics*, vol. 5, no. 2, pp. 430–437, 2006.
- [41] M. Katori and M. Majima, "Cyclooxygenase-2: its rich diversity of roles and possible application of its selective inhibitors," *Inflammation Research*, vol. 49, no. 8, pp. 367–392, 2000.
- [42] H. R. Herschman, "Prostaglandin synthase 2," *Biochimica et Biophysica Acta*, vol. 1299, no. 1, pp. 125–140, 1996.
- [43] K. Subbaramaiah, D. T. Lin, J. C. Hart, and A. J. Dannenberg, "Peroxisome proliferator-activated receptor γ ligands suppress the transcriptional activation of cyclooxygenase-2. Evidence for involvement of activator protein-1 and CREB-binding protein/p300," *The Journal of Biological Chemistry*, vol. 276, no. 15, pp. 12440–12448, 2001.
- [44] H. Inoue, T. Tanabe, and K. Umehara, "Feedback control of cyclooxygenase-2 expression through PPAR γ ," *The Journal of Biological Chemistry*, vol. 275, no. 36, pp. 28028–28032, 2000.
- [45] N. Janabi, "Selective inhibition of cyclooxygenase-2 expression by 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ in activated human astrocytes, but not in human brain macrophages," *The Journal of Immunology*, vol. 168, no. 9, pp. 4747–4755, 2002.
- [46] S. A. Kliewer, J. M. Lenhard, T. M. Willson, I. Patel, D. C. Morris, and J. M. Lehmann, "A prostaglandin J₂ metabolite binds peroxisome proliferator-activated receptor γ and promotes adipocyte differentiation," *Cell*, vol. 83, no. 5, pp. 813–819, 1995.
- [47] D. W. Gilroy, P. R. Colville-Nash, D. Willis, J. Chivers, M. J. Paul-Clark, and D. A. Willoughby, "Inducible cyclooxygenase may have anti-inflammatory properties," *Nature Medicine*, vol. 5, no. 6, pp. 698–701, 1999.
- [48] S. Han, H. Inoue, L. C. Flowers, and N. Sidell, "Control of COX-2 gene expression through peroxisome proliferator-activated receptor γ in human cervical cancer cells," *Clinical Cancer Research*, vol. 9, no. 12, pp. 4627–4635, 2003.
- [49] M.-Y. Li, H. Deng, J.-M. Zhao, D. Dai, and X.-Y. Tan, "PPAR γ pathway activation results in apoptosis and COX-2 inhibition in HepG2 cells," *World Journal of Gastroenterology*, vol. 9, no. 6, pp. 1220–1226, 2003.
- [50] G. He, Y. M. Sung, and S. M. Fischer, "Troglitazone induction of COX-2 expression is dependent on ERK activation in keratinocytes," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 74, no. 3, pp. 193–197, 2006.
- [51] E.-K. Yoon, W.-K. Lee, J.-H. Lee, S.-M. Yu, S.-G. Hwang, and S.-J. Kim, "ERK-1/-2 and p38 kinase oppositely regulate 15-deoxy- $\Delta^{12,14}$ -prostaglandinJ₂-induced PPAR- γ activation that mediates dedifferentiation but not cyclooxygenase-2 expression in articular chondrocytes," *Journal of Korean Medical Science*, vol. 22, no. 6, pp. 1015–1021, 2007.
- [52] H. Ikawa, H. Kameda, H. Kamitani, et al., "Effect of PPAR activators on cytokine-stimulated cyclooxygenase-2 expression in human colorectal carcinoma cells," *Experimental Cell Research*, vol. 267, no. 1, pp. 73–80, 2001.
- [53] E. A. Meade, T. M. McIntyre, G. A. Zimmerman, and S. M. Prescott, "Peroxisome proliferators enhance cyclooxygenase-2 expression in epithelial cells," *The Journal of Biological Chemistry*, vol. 274, no. 12, pp. 8328–8334, 1999.
- [54] A. V. Pontsler, A. St. Hilaire, G. K. Marathe, G. A. Zimmerman, and T. M. McIntyre, "Cyclooxygenase-2 is induced in monocytes by peroxisome proliferator activated receptor γ and oxidized alkyl phospholipids from oxidized low density lipoprotein," *The Journal of Biological Chemistry*, vol. 277, no. 15, pp. 13029–13036, 2002.
- [55] T. Kalajdzic, W. H. Faour, Q. W. He, et al., "Nimesulide, a preferential cyclooxygenase 2 inhibitor, suppresses peroxisome proliferator-activated receptor induction of cyclooxygenase 2 gene expression in human synovial fibroblasts: evidence for receptor antagonism," *Arthritis & Rheumatism*, vol. 46, no. 2, pp. 494–506, 2002.
- [56] A. Chawla, Y. Barak, L. Nagy, D. Liao, P. Tontonoz, and R. M. Evans, "PPAR- γ dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation," *Nature Medicine*, vol. 7, no. 1, pp. 48–52, 2001.
- [57] A. M. Lennon, M. Ramaugé, A. Dessouroux, and M. Pierre, "MAP kinase cascades are activated in astrocytes and preadipocytes by 15-deoxy- $\Delta^{12,14}$ -prostaglandinJ₂-prostaglandin J₂ and the thiazolidinedione ciglitazone through peroxisome proliferator activator receptor gamma-independent mechanisms involving reactive oxygenated species," *The Journal of Biological Chemistry*, vol. 277, no. 33, pp. 29681–29685, 2002.
- [58] R. A. Nemenoff, "Peroxisome proliferator-activated receptor- γ in lung cancer: defining specific versus "off-target" effectors," *Journal of Thoracic Oncology*, vol. 2, no. 11, pp. 989–992, 2007.
- [59] S. T. Reddy, D. J. Wadleigh, and H. R. Herschman, "Transcriptional regulation of the cyclooxygenase-2 gene in activated mast cells," *The Journal of Biological Chemistry*, vol. 275, no. 5, pp. 3107–3113, 2000.
- [60] J. A. Romashkova and S. S. Makarov, "NF- κ B is a target of AKT in anti-apoptotic PDGF signalling," *Nature*, vol. 401, no. 6748, pp. 86–90, 1999.
- [61] L. V. Madrid, C.-Y. Wang, D. C. Guttridge, A. J. G. Schottelius, A. S. Baldwin Jr., and M. W. Mayo, "Akt suppresses apoptosis by stimulating the transactivation potential of the RelA/p65 subunit of NF- κ B," *Molecular and Cellular Biology*, vol. 20, no. 5, pp. 1626–1638, 2000.
- [62] K. M. Patel, K. L. Wright, P. Whittaker, P. Chakravarty, M. L. Watson, and S. G. Ward, "Differential modulation of COX-2 expression in A549 airway epithelial cells by structurally distinct PPAR γ agonists: evidence for disparate functional effects which are independent of NF- κ B and PPAR γ ," *Cellular Signalling*, vol. 17, no. 9, pp. 1098–1110, 2005.
- [63] D. A. Bradbury, L. Corbett, and A. J. Knox, "PI 3-kinase and MAP kinase regulate bradykinin induced prostaglandin E₂ release in human pulmonary artery by modulating COX-2 activity," *FEBS Letters*, vol. 560, no. 1–3, pp. 30–34, 2004.
- [64] B.-C. Chen, Y.-S. Chang, J.-C. Kang, et al., "Peptidoglycan induces nuclear factor- κ B activation and cyclooxygenase-2 expression via Ras, Raf-1, and ERK in RAW 264.7 macrophages," *The Journal of Biological Chemistry*, vol. 279, no. 20, pp. 20889–20897, 2004.
- [65] S. A. Weaver, M. P. Russo, K. L. Wright, et al., "Regulatory role of phosphatidylinositol 3-kinase on TNF- α -induced cyclooxygenase 2 expression in colonic epithelial cells," *Gastroenterology*, vol. 120, no. 5, pp. 1117–1127, 2001.
- [66] H. Cho and H.-H. Tai, "Inhibition of NAD⁺-dependent 15-hydroxyprostaglandin dehydrogenase (15-PGDH) by cyclooxygenase inhibitors and chemopreventive agents," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 67, no. 6, pp. 461–465, 2002.
- [67] R. L. Keith, Y. E. Miller, T. M. Hudish, et al., "Pulmonary prostacyclin synthase overexpression chemoprevents tobacco smoke lung carcinogenesis in mice," *Cancer Research*, vol. 64, no. 16, pp. 5897–5904, 2004.
- [68] A. Finckh and M. D. Aronson, "Cardiovascular risk of cyclooxygenase-2 inhibitors: where we stand now," *Annals of Internal Medicine*, vol. 142, no. 3, pp. 212–214, 2005.

- [69] J. T. Mao, I.-H. Tsu, S. M. Dubinett, et al., "Modulation of pulmonary leukotriene B₄ production by cyclooxygenase-2 inhibitors and lipopolysaccharide," *Clinical Cancer Research*, vol. 10, no. 20, pp. 6872–6878, 2004.
- [70] R. N. DuBois, S. B. Abramson, L. Crofford, et al., "Cyclooxygenase in biology and disease," *The FASEB Journal*, vol. 12, no. 12, pp. 1063–1073, 1998.
- [71] S. Han and J. Roman, "Suppression of prostaglandin E₂ receptor subtype EP2 by PPAR γ ligands inhibits human lung carcinoma cell growth," *Biochemical and Biophysical Research Communications*, vol. 314, no. 4, pp. 1093–1099, 2004.
- [72] R. Govindarajan, L. Ratnasinghe, D. L. Simmons, et al., "Thiazolidinediones and the risk of lung, prostate, and colon cancer in patients with diabetes," *Journal of Clinical Oncology*, vol. 25, no. 12, pp. 1476–1481, 2007.
- [73] C. J. Rosen, "The rosiglitazone story—lessons from an FDA advisory committee meeting," *The New England Journal of Medicine*, vol. 357, no. 9, pp. 844–846, 2007.
- [74] "Thiazolidinediones and cardiovascular disease," *The Medical Letter on Drugs and Therapeutics*, vol. 49, no. 1265, pp. 57–58, 2007.
- [75] S. E. Nissen and K. Wolski, "Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes," *The New England Journal of Medicine*, vol. 356, no. 24, pp. 2457–2471, 2007.
- [76] R. A. Nemenoff, M. Weiser-Evans, and R. A. Winn, "Activation and molecular targets of peroxisome proliferator-activated receptor- γ ligands in lung cancer," *PPAR Research*, vol. 2008, Article ID 156875, 8 pages, 2008.
- [77] K. A. Peebles, J. M. Lee, J. T. Mao, et al., "Inflammation and lung carcinogenesis: applying findings in prevention and treatment," *Expert Review of Anticancer Therapy*, vol. 7, no. 10, pp. 1405–1421, 2007.
- [78] H.-S. Kim, H.-R. Youm, J.-S. Lee, K.-W. Min, J.-H. Chung, and C.-S. Park, "Correlation between cyclooxygenase-2 and tumor angiogenesis in non-small cell lung cancer," *Lung Cancer*, vol. 42, no. 2, pp. 163–170, 2003.
- [79] F. R. Khuri, H. Wu, J. J. Lee, et al., "Cyclooxygenase-2 overexpression is a marker of poor prognosis in stage I non-small cell lung cancer," *Clinical Cancer Research*, vol. 7, no. 4, pp. 861–867, 2001.
- [80] H. Tsubochi, N. Sato, M. Hiyama, et al., "Combined analysis of cyclooxygenase-2 expression with p53 and Ki-67 in nonsmall cell lung cancer," *The Annals of Thoracic Surgery*, vol. 82, no. 4, pp. 1198–1204, 2006.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

